

88847-2

2/3/2014

1/15



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Biopesticides and Pollution Prevention Division (7511P)
1200 Pennsylvania Avenue NW
Washington, DC 20460

NOTICE OF PESTICIDE REGISTRATION

Registration
 Reregistration

(under FIFRA, as amended)

EPA Reg. Number:
88847-2

Date of Issuance:
February 3, 2014

Unconditional

Term of Issuance:

Name of Pesticide Product:

VST-006330 EP

Name and Address of Registrant (include ZIP Code):

Vestaron Corporation
4717 Campus Drive, Ste. 1200
Kalamazoo, MI 49008

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his or her motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This registration does not eliminate the need for continual reassessment of the pesticide. If EPA determines at any time, that additional data are required to maintain in effect an existing registration, the Agency will require submission of such data under section 3(c)(2)(B) of FIFRA.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) and is subject to the following terms and conditions:

1. Revise the EPA Registration number on the label to read, "EPA Reg. No. 88847-2."
2. Submit two (2) copies of the revised final printed labeling before you release the product for shipment

A stamped copy of the label is enclosed for your records.

Signature of Approving Official:

Robert McNally, Director
Biopesticides and Pollution Prevention Division (7511P)

Date:

February 3, 2014

EPA Form 8570-6

CONCURRENCES

SYMBOL ▶		7511P	7511P				
SURNAME ▶		Cheri	McNally				
DATE ▶		2/3/14	2/3/14				

EPA Form 1320-1A (1/90)

OFFICIAL FILE COPY

2/15

VST-006330 EP

MASTER LABEL, containing:
Sublabel A: Greenhouse and Field Use
Sublabel B: Home & Garden Use

Active Ingredient: GS-omega/kappa-Hctx-Hv1a* 20.00%
 Other Ingredients: 80.00%
 Total: 100.00%

*Versitude™ peptide.

EPA Reg. No.: (pending as File Symbol 88847-E)
 Net Contents: XX
 (Batch)(Lot) No: XXXX

EPA Est. No.: XXXXX-XX-XXX

Manufactured by: Vestaron Corporation
 4717 Campus Drive, Ste. 1200
 Kalamazoo, MI 49008

ACCEPTED
 FEB 3 2014

Under the Federal Insecticide, Fungicide,
 and Rodenticide Act, as amended, for
 the pesticide registered under
 EPA Reg. No. 88847-2

3/15

Sublabel A: Greenhouse and Field Use

VST-006330 EP

Biological Insecticide

Active Ingredient: GS-omega/kappa-Htx-Hv1a* 20.00%
 Other Ingredients: 80.00%
 Total: 100.00%

*Versitude™ peptide.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 – 20 minutes. • Call a poison control center or doctor for treatment advice.
<p align="center">HOT LINE NUMBER</p> <p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.</p>	

EPA Reg. No.: (pending as File Symbol 88847-E)
 Net Contents: XX
 (Batch)(Lot) No: XXXX

EPA Est. No.: XXXXX-XX-XXX

Manufactured by: Vestaron Corporation
 4717 Campus Drive, Ste. 1200
 Kalamazoo, MI 49008

CAUTION

Under the Federal Insecticide, Fungicide, and Rodenticide Act, this pesticide is registered under EPA Reg. No. XXXXX-XX-XXX. The pesticide is labeled for use as a biological insecticide for the control of insects on greenhouse and field use.

PRECAUTIONARY STATEMENTS

Hazards to humans and domestic animals - CAUTION. Causes moderate eye and skin irritation. Avoid contact with eyes, skin or clothing. Wear protective eyewear and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear:

- long-sleeved shirt and long pants
- waterproof gloves
- shoes plus socks

Follow the manufacturer's instructions for cleaning / maintaining PPE. If no instructions are available, use detergent and hot water for washables. Keep and wash PPE separately from other laundry.

Engineering Controls: When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [10 CFR 170.240(d)(406)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards: For terrestrial uses: Exposure of adult bees to direct treatment or residues on blooming crops can lead to effects on honey bee larvae. See the "Directions for Use" section of this label for application instructions that minimize risk to honey bee larvae. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean highwater mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

To minimize potential exposure to bee larvae: **Do Not** apply this product at any time between 4 days prior to bloom and until after petal fall, or if bees are foraging in the treatment area.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water), is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

PRODUCT INFORMATION

VST-006330 EP is a biological insecticide containing the active ingredient GS-omega/kappa-Hctx-Hv1a (Versitude™ peptide) for use on ornamental plants, turf and edible crops against cabbage looper (*Tricoplusia ni*). VST-006330 EP functions primarily as a central nervous inhibitor to aid in the suppression of cabbage loopers infesting labeled crops. VST-006330 EP is mixed with water and applied as a foliar spray with ground or aerial equipment equipped for conventional insecticide spraying.

VST-006330 EP can be used in either the field or greenhouse.

USE INSTRUCTIONS

VST-006330 EP is a highly selective insecticide for use against the listed insect pest. Close scouting and early attention to infestations is highly recommended. Proper timing of application targeting newly hatched larvae is important for optimal results.

Thorough coverage of infested plant parts is necessary to aid suppression. VST-006330 EP does not have systemic activity. For some crops, directed drop nozzles by ground machine are required.

Under heavy pest populations shorten the spray interval, and/or increase the spray volume to improve coverage.

Repeat applications at an interval sufficient to aid suppression, usually 3-10 days depending upon plant growth rate, insect activity, and other factors.

Tank mix with contact insecticides/miticides, including *Bacillus thuringiensis* products, to enhance suppression. Refer to tank mix section.

6/15

To ensure compatibility of tank-mix combinations they must be evaluated prior to use. To determine the physical compatibility of this product with other products use a jar test. Using a restrictive of the labeling limitations and precautions of all products used in mixtures. physically compatible, effective, or non-injurious under your use conditions. Follow the most or fertilizers if there has been no previous experience or use of the combination to show it is Do not combine VST-006330 EP in the spray tank with other pesticides, surfactants, adjuvants, Tank mixing:

For conventional air and ground applications, use at least 10 gallons of total volume per acre in water based sprays. Spray volume:

not mix more VST-006330 EP than can be used in 24 hours. EP to the mix tank and continue circulation. Maintain circulation while loading and spraying. Do of water. Add water to the mix tank. Start the mechanical or hydraulic agitation to provide moderate circulation before adding VST-006330 EP. Add the desired volume of VST-006330 Important - Do not add VST-006330 EP to the mix tank before introducing the desired amount Mixing directions:

Avoiding spray drift is the responsibility of the applicator. Do not spray when wind speed favors drift beyond the area intended for use. depend upon crop development, weather, application equipment, and local experience. Apply VST-006330 EP in ground and aerial equipment with quantities of water sufficient to provide thorough coverage of infested plant parts. The amount of water needed per acre will

USE RESTRICTION: Do not apply more than 10 lbs VST-006330 EP per acre per year.

GROUND AND AERIAL APPLICATIONS

VST-006330 EP can be applied up to and including the day of harvest. Preharvest interval - agricultural use: developed for your crop(s) and location. the use of chemical insecticides. Consult local agricultural authorities for specific IPM strategies insecticides. This will result in reduced susceptibility to insect damage and overall reduction in necessary. Apply VST-006330 EP alone or in combination and / or rotation with chemical VST-006330 EP is an important tool in sound insect management whenever insecticide use is Integrated Pest Management (IPM):

VST-006330 EP has been evaluated for phytotoxicity on a variety of crops under various normal growing conditions. However, testing all crop varieties, in all mixtures and combinations is not feasible. Prior to treating entire crop, test a small portion of the crop for sensitivity. For hard-to-wet crops, consider using a spreader/sticker or an adjuvant that has been approved for targeted crop use to enhance the adhesion of VST-006330 EP to the crop. Examples of appropriate spreader/stickers or adjuvants are: 1) alkoxylated surfactants, 2) organic silicates, 3) vegetable oils, 4) methylated vegetable oils, 5) mineral oils, and/or 6) phospholipids.

7/5

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to backflow. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from

Maintain continuous agitation in the mix tank during mixing and application to insure a uniform suspension. Greater accuracy in calibration and distribution will be achieved by injecting a larger volume for a more dilute solution per unit time. Crop injury or lack of effectiveness can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation waters.

GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

Do not combine VST-006330 EP with pesticides, surfactants or fertilizers for application through chemigation equipment unless prior experience has shown the combination physically compatible, effective and non-injurious under conditions of use. VST-006330 EP has not been fully evaluated for compatibility with all adjuvants or surfactants. It is advisable to conduct a spray compatibility test if a mixture with adjuvants or surfactants is planned.

First prepare a suspension of VST-006330 EP in a mix tank. Fill tank 1/2 to 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of VST-006330 EP, and then the remaining volume of water. Then set the sprinkler to deliver a minimum of 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the suspension of VST-006330 EP into the irrigation water line so as to deliver the desired rate per acre. Inject the suspension of VST-006330 EP with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. Any questions on calibration should be directed to your State Extension Service Specialists, to equipment manufacturers or other experts.

Spray preparation

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation systems. Do not connect an irrigation system (including greenhouse systems) used for pesticide applications to a public water system.

CHEMIGATION USE DIRECTIONS

Add dry formulations first, then flowables, then emulsifiable concentrates last. After thoroughly mixing, let this mixture stand for 5 minutes. If the combination remains mixed or can be readily be remixed, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

5/8

prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Do not apply when wind speed favors drift, when system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.

AERIAL DRIFT REDUCTION ADVISORY INFORMATION

GENERAL: Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. Where states have more stringent regulations, they should be observed. Note: This section is advisory in nature and does not supersede the mandatory label requirements.

INFORMATION ON DROPLET SIZE: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that will provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE: Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets. Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When high flow rates are needed, use higher flow rate nozzles instead of increasing pressure. Number of nozzles - Use the minimum number of nozzles that provide uniform coverage. Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential. Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM WIDTH: For aerial applications, the boom width must not exceed 75% of the wingspan or 90% of the rotary blade. Use upwind swath displacement and apply only when wind speed is 3-10 mph as measured by an anemometer. Use medium or coarser spray according to ASAE

9/15

572 definition for standard nozzles or VMD for spinning atomizer nozzles. If application includes a no-spray zone, do not release spray at a height greater than 10 feet above the ground or crop canopy.

APPLICATION HEIGHT: Do not make application at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure to droplets to evaporation and wind.

SWATH ADJUSTMENT: When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

WIND: Drift potential is lowest between wind speeds of 2 - 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS: Do not apply during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SENSITIVE AREAS: The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas). Do not allow spray to drift from the application site and contact people, structures people occupy at any time and the associated property, parks and recreation areas, non-target crops, aquatic and wetland areas, woodlands, pastures, rangelands, or animals.

FOR USE ON THE FOLLOWING CROPS:

Pre-harvest Interval (PHI) = 0 days

Crop Group

I. Vegetable and Cole Crops:

Insect Pest	Crops
Cabbage looper	Artichokes
	Argula
	Asparagus
	Beans- including garden, dried, succulent,
	adzuki, fava, field, garbanzo, pinto, kidney,
	lentil, lima, moonbeam, navy, runner, snap,
	teary, wax, yard long
	Beets-including garden
	Bok Choy
	Broccoli
	Broccoli Raab
	Brussels Sprouts
	Cabbage
	Cardoni
	Carrots
	Cassava
	Cauliflower
	Cavalo
	Celertac
	Celery
	Chayote root
	Chervil
	Chick peas
	Chicory
	China greens
	Chinese artichoke
	Chinese broccoli
	Chinese cabbage (Napa)
	Chinese mustard cabbage
	Collards
	Corn salad
	Cress
	Cucumber
	Cucurbits including cantaloupe and
	watermelon
	Dandelion dock
	Dry bulb onions
	Edible burdock
	Edible chrysanthemum
	Edible gourds
	Eggplant

	<p>Escarole Endive Garlic Ginger Ginseng Green onions Greens: Beets, China, Dandelion, Mustard, Turnip, Rape Ground cherry Horseradish Jerusalem artichoke Kale Kohlrabi Leeks Lentils Lettuce: Head, Leaf, Romaine Lupine Manga Melons: Cantaloupe, Crenshaw, Honeydew, Muskmelon, Watermelon, etc. Mizuna Mustard spinach Napa Okra Olives Onions Parsley Parsnips Peas-including garden, dried, succulent, blackeyed, chickpea, cowpea, crowder, edible-pod, English, field, green, pigeon, snow, sugar snap, Pepino Peppers Potatoes Pumpkins Purslane Radicchio Radishes Rhubarb Rutabaga Salsify Shallots Skirret Soybean foliage Soybean Spinach Squash Sugar beets Sweet potatoes</p>
--	---

5/14

To the extent permitted by applicable law, the seller makes no warranty, expressed or implied, of merchantability, fitness or otherwise concerning use of this product. The user assumes all risks of use, storage or handling that are not in strict accordance with the accompanying directions.

WARRANTY

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment by shaking and tapping sides and bottom to loosen clinging particles. Then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

Pesticide Storage: Store in original container at less than 40° for up to 6 months after date of manufacture.

Pesticide Disposal: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

STORAGE AND DISPOSAL
 Do not contaminate water, food or feed by storage or disposal.

Rate/Acre: 4 pounds VST-006330 EP

Crops	Flowers: greenhouse and field Greenhouse vegetables
Insect Pest	Cabbage looper

III. Commercial Flowers and Ornamental Plants

Rate/Acre: 4 pounds VST-006330 EP

Crops	Cotton Soybeans Tobacco
Insect Pest	Cabbage looper Cabbage looper Cabbage looper

II. Field Crops

Rate/Acre: 4 pounds VST-006330 EP

	Swiss chard Tomatillo Tomatoes Turmeric Turnips Turnip-rooted chervil Turnip-rooted parsley Watercress Yams
--	---

13/15

Sublabel B: Home & Garden Uses

VST-006330 EP

Biological Insecticide

Active Ingredient: G-omega/kappa-Hxtx-Hv1a* 20.00%
Other Ingredients: 80.00%
Total: 100.00%

*Versitude™ peptide.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 – 20 minutes. • Call a poison control center or doctor for treatment advice.
<p>HOT LINE NUMBER</p> <p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.</p>	

EPA Reg. No.: (pending as File Symbol 88847-E)
EPA Est. No.: XXXXX-XX-XXX
Net Contents: XX
(Batch)(Lot) No: XXXX

Manufactured by: Vestaron Corporation
4717 Campus Drive, Ste. 1200
Kalamazoo, MI 49008

14/15

Garden plants (including):
Asparagus, beets, broccoli, Brussels sprouts, cabbage, cantaloupe, carrots, cauliflower, celery, collards, cucumbers, edible-podded legume vegetables including: snap bean, wax bean, yard long bean, jack bean, edible-pod pea, snow pea, sugar snap pea, dried shelled beans and peas

Apply to the following types home and garden plants:

WHERE TO APPLY

- Add 2 teaspoon – 4 tablespoons (0.3-2 ounces) of VST-006330 EP for each 1 gallon of water in a tank sprayer. Agitate prior to use.
- Use a coarse spray pattern to reduce drift to desirable plants.
- Spray to thoroughly wet foliage.

Do not apply more than 10 lbs of VST-006339 EP per acre per year.
Mix VST-006330 EP with water and apply as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for effective control. VST-006330 EP does not have systemic activity.

To minimize potential exposure to bee larvae: **Do Not** apply this product at any time between 4 days prior to bloom and until after petal fall, or if bees are foraging in the treatment area.

HOW TO APPLY

VST-006330 EP is a highly selective insecticide for use against cabbage looper (*Tricoplusia ni*). Close scouting and early attention to infestations is highly recommended. Proper timing of application targeting newly hatched larvae is important for optimal results.

HOW IT WORKS

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE

Environmental Hazards: Exposure of adult bees to direct treatment or residues on blooming crops can lead to effects on honey bee larvae. See the "Directions for Use" section of this label for application instructions that minimize risk to honey bee larvae. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems.

PRECAUTIONARY STATEMENTS

Hazards to humans and domestic animals - CAUTION. Causes moderate eye and skin irritation. Avoid contact with eyes, skin or clothing. Wear goggles or safety glasses, long sleeved shirt and long pants, gloves and shoes plus socks. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

15/15

To the extent permitted by applicable law, the seller makes no warranty, expressed or implied, of merchantability, fitness or otherwise concerning use of this product. The user assumes all risks of use, storage or handling that are not in strict accordance with the accompanying directions.

WARRANTY

STORAGE AND DISPOSAL
Do not contaminate water, food or feed by storage or disposal.
Pesticide Storage: Store in a cool dry place inaccessible to children.
Pesticide Disposal and Container Handling: If empty: Nonrefillable container. Do not reuse or refill this container. Place in trash and offer for recycling if available. **If partially filled:** Call your local solid waste agency or (800) 858-7378 (National Pesticide Information Center) for disposal instructions. Never place unused product down any indoor or outdoor drain.

Ornamentals-including annuals and perennials
To aid in the suppression of cabbage looper.
including: field bean, kidney bean, lima bean (dry), navy bean, pinto bean, adzuki bean, black-eyed pea, cowpea, mung bean, southern pea, lentil (dry); eggplant, herbs, kale, lettuce, melons, mustard greens, pepper, parsnip, potato, radish, rutabaga, safsiy, spinach, squash (winter and summer), sweet potato, tomatoes, turnip greens, turnips, watercress, and watermelon.