

SCRAMINATOR™ Crawling Insect Killer

KILLS: Ants, Asian Lady Beetle, Bed Bugs, Bees, Black Carpet Beetles, Box elder Bugs, Carpenter Ants, Centipedes, Cigarette Beetles, Cluster Flies, Cockroaches, Confused Flour Beetles, Crickets, Darkling Beetles, Drugstore Beetles, Earwigs, Firebrats, Fleas, Furniture Carpet Beetles, Ground Beetles, Hide Beetles, Hornets, Indian meal Moths, Lesser Grain Borers, Merchant Grain Beetles, Millipedes, Mites, Paper Wasps, Phorid Flies, Pillbugs, Red Flour Beetles, Rice Weevils, Rusty Grain Beetles, Sawtoothed Grain Beetles, Silverfish, Sow bugs, Spider Beetles, Spiders, Varied Carpet Beetles, Warehouse Beetles, Wasps, Webbing Clothes Moths and Yellow jackets.

ACTIVE INGREDIENTS:	BY WT.
Diatomaceous Earth consisting of:	
Silicon Dioxide	76.5%
Other Ingredients	23.5%
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

(See other [side] panel for additional precautionary statements)

Scraminator Crawling Insect Killer is effective and long lasting when kept dry. Scraminator Crawling Insect Killer is a mechanical action insect killer. Insects cannot become immune to its action. Insects that contact or ingest this powder die within 48 hours.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals:

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Use adequate ventilation and avoid breathing dust. For prolonged exposure to dust, use a suitable dust mask approved by NIOSH/MSHA.

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> -Hold eye open and rinse slowly and gently with water for 15-20 minutes. -Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. - Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the National Pesticide Information Center 1-800-858-7378 for emergency medical treatment advice.</p>	

ACCEPTED
FEB - 2 2016

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide Registered under EPA Reg. No. 89014-1

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Application Directions:

To dispense product, turn tab on top of container and shake gently. Close tab when finished. Scraminator Crawling insect killer can be applied directly from the box or a can be applied with a hand duster or powder duster to areas where crawling insects are found. Scraminator Crawling Insect Killer is intended for application with a hand duster or power duster to areas such as cracks, crevices, voids, hiding and runway areas, indoors and outdoors, where crawling insects are found. Apply lightly and uniformly.

Scraminator Crawling Insect Killer is a ready-for-use insecticide dust which has given effective knockdown and control of the pests listed. Scraminator Crawling Insect Killer is for use in residential and commercial pest control, in and around buildings and structures and their immediate surroundings, and on modes of transport (i.e. apartments, homes and other residential buildings; daycare centers and schools; hospitals and nursing homes; hotels and motels; industrial buildings; laboratories; office buildings; other commercial and industrial facilities; restaurants, food/feed handling establishments and food processing plants; warehouses; and on aircraft, buses, rail cars, trucks, trailers and vessels). In living areas, make applications in such a manner as to avoid deposits on exposed surfaces or introducing the material into the air. Treat a small area of the surface to be treated and look for staining or other adverse reaction before making regular application.

Apply insecticide directly into cracks and crevices, use a bulbous duster or other suitable equipment. Apply lightly and uniformly to infested areas. Pay particular attention to cracks, crevices and voids where insects may be harboring, traveling, breeding or entering the structure. Examples of these areas are service ducts; false floors and ceilings; wall voids; around electrical and telephone fittings and equipment; around water and sewer pipes; under and behind cabinets, refrigerators and sinks; around window and door frames; along baseboards; in attics and crawl spaces; landscaped (including lawns and turf) and mulched areas. The amount of product to be applied will vary with the site but stay in the range of 0.15 - 0.6 oz/10 sq. ft. or 1 - 5 oz/100 sq. ft. Apply once a week.

For control of Earwigs, Millipedes, Centipedes, Silverfish, Cockroaches, and Ants: Sprinkle around room perimeter and leave the product on the floor leaving a thin layer of Scraminator Crawling Insect Killer in areas where these pests are found or may hide, such as cracks and crevices, behind and beneath refrigerators, cabinets, stoves, garbage cans, and in and around sewer pipes and drains, window frames, and in attics and basements. Dust insects directly if possible. Apply once a week.

Carpet Beetles, Furniture and Varied) Webbing Clothes Moths and Spiders: Dust along baseboards, carpet edges, under carpeting, furniture, in closets and shelving where these insects are seen or suspected.

Fleas: Thoroughly treat floor, carpeting and bedding in and around pets' sleeping quarters. Treat surrounding cracks and crevices, baseboards, carpeting, and wherever fleas are suspected. Sprinkle material evenly and leave on the carpet for at least 15 minutes then vacuum. Apply once a week for four weeks as described above.

Mattresses (bed bugs and dust mites: Sprinkle material evenly and leave on the mattress for at least 15 minutes then vacuum. Sprinkle in mattress bag and leave material in the bag. Apply once a week for four weeks as described above.

Ants, Cockroaches, and Earwigs.: Lightly coat a thin layer of Scraminator Crawling Insect Killer in and around cracks, crevices or voids associated with windows, doors, porches, screens, eaves, patios, garages, under stairways, around foundations, sewers, animal burrows, mulched areas, manure or compost piles and in crawl spaces and other areas where insects may be harboring, traveling, breeding or entering the structure. Dust around foundations, along ant trails, and around sewer pipes and drains. Apply once a week.

Bees, Wasps, Hornets, Paper Wasps and Yellow jackets: It is generally advisable to treat wasp and bee nests in the evening when insects are less active and have returned to the nest. Wear protective clothing, if deemed necessary, to avoid stings. Using hand or power duster (with extension tubes if necessary), thoroughly dust nest, nest entrance and surrounding areas where insects alight. Nests in voids should be located and treated. If necessary, drill a hole into the void(s), blow dust in, and reseal. For ground wasps and yellow jackets, dust entrance and surrounding areas. For best results, carefully check nests 1 - 2 days after treatment to ensure complete kill. When treating overhead nests, apply in such a manner as to avoid drift onto the applicator.

Fruit and Phorid Flies: Apply where these insects harbor, travel, breed or enter the structure. For example: under cabinets or shelving; behind or under equipment or drains; protected exterior areas such as eaves, trash corrals and dumpsters; and other areas where organic debris may accumulate.

Asian Lady Beetles, Box elder Bugs and Cluster Flies: Apply where these insects may be harboring, traveling, breeding or entering the structure.

HOW TO TREAT EMPTY GRAIN, CEREAL OR FLOUR STORAGE AREAS

Apply lightly approximately one 1 pound per 1,000 square feet on all Interior wall surfaces of empty storage areas. Where possible, special attention should be given to applying to cracks, crevices, and other areas where insects hide or crawl. After application there is no need to remove the product or aerate the areas. Leave area undisturbed for two to three days to eliminate existing infestations before filling. It is desirable to reapply the product each time the storage area is emptied.

HOW TO APPLY TO GRAIN OR SEED

Treat grain and seed as soon after harvest as possible. Scraminator Crawling Insect Killer (Kills Insects Only) and may be applied to the grain at the time of storage, to protect it from further damage. Adjust the moisture content of grain to not over 12% by weight. Using standard milling equipment apply at the rate of 2 to 7 pounds directly onto each ton of dean, dry grain or seed to be treated.

Thoroughly mix with grain. Product may be dispensed into the grain either at the bottom of an elevator leg or as it is being moved into storage by auger or conveyor belt. Seal bin for storage. It is wise to check the status of stored grains regularly.

IMPORTANT: Treated grain will have a dry appearance, its movement speed will be reduced, and the grade and test weight may be lowered. Milling, baking, nutritive and germination qualities of grain or seed however are not affected by this treatment.

HOW TO PROTECT SMALL AMOUNTS OF GRAIN

Thoroughly mix 1 cupful (2 1/2 oz.) of Scraminator Crawling Insect Killer Into each 25 lbs. of grain so that substantially all kernels are lightly coated with the product. Place treated grain into an "open head" 5 or 6 gallon plastic pail for storage. Tightly replace lid for seal.

COMMERCIAL PEST CONTROL

Treatment of Commercial Transportation Facilities: Food Processing Plant Premise/Equipment; Eating Establishments Food Handling Areas (food contact); Eating Establishment Food Serving Areas (food Contact); and Food/Grocery Marketing/Storage/Distribution Facility Premise.

Apply to control the insect pests as listed below as a "Crack and Crevice" or a "Spot" treatment In areas where food or feed is - Received, Processed (including enclosed processing systems), Packaged (Bottled, Boxed, Canned, Wrapped), Transported, Stored (including waste), Prepared, or Served, as a crack and crevice treatment apply sparingly into cracks and crevices using equipment designed to place powder into specific locations, Limit spot treatments to wall or floor surfaces where pests have been seen or where pests are suspected of hiding. Individual spot treatments must not exceed 2 sq. ft. All food or feed must be removed or covered before treatment. Treat open spaces below floors, above ceilings, within walls, under stairs, underneath and behind equipment or appliances, around pipes, under sinks, in utility closets and around garbage and waste cans, for Non-Food Areas. Apply (as directed above) as a "Crack and Crevice" or a "Spot" treatment for control of Ants, Box elder Bugs, Crickets, Roaches and Silverfish to areas where pests hide, run or land, such as attics and spaces above suspended rooms, basements, and crawl spaces, cracks in baseboards, moldings, and door frames, corners, in closets, cabinets, and shelves, window sills and frames, under and behind furniture and appliances, etc. Work into and apply under rugs, door mats, couches, etc...

STORED GRAIN and SEED PEST CONTROL

Scraminator Crawling Insect Killer controls listed insect attack against Feed Grains, Seeds, Stored Grains; Barley, Beans, Buckwheat, Corn, Flax, Oats, Pea, Rice, Rye, Soybean and Sorghum (Milo) and Wheat. Scraminator Crawling Insect Killer kills listed exposed grain infesting insects including: Rice Weevils, Lesser Grain Borers, Flat Grain Beetles, Saw Toothed Grain Beetles, Granary Weevils, Angoumois Grain Moths, Mediterranean Flour Moths, Indian Meal Moth (larvae), Red Flour Beetles, and Confused Flour Beetle.

Food Areas of Food/Feed Handling Establishments: Food/feed handling establishments are places other than private residences in which food/feed is held, processed, prepared or served. In food handling areas, or areas where food or feed products are prepared or stored, apply as a crack & crevice or spot treatment.

575

As a C&C treatment, apply sparingly into cracks and crevices using equipment designed to place powder into specific locations. Limit spot treatments to walls or floors where pests are suspected of hiding. No individual spot treatments should exceed 2 sq. ft. Remove or cover all food or feed items in the treatment area before applying product. Treat open spaces below floors, above ceilings, within walls, under stairs, underneath and behind equipment or appliances, around pipes and sinks, in utility closets and around garbage and waste cans. For best results, apply to dry areas, once a week, to maintain control of problem pests.

Buyer assumes all responsibility for safety and use not in accordance with label directions.

Note: Seller warrants that this product complies with the specifications expressed in this label. To the extent consistent with applicable law, Seller makes no other warranties, and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for the intended purpose. To the extent consistent with applicable law, Seller's liability or default, breach or failure under this label shall be limited to the amount of the purchase price. To the extent consistent with applicable law, Seller shall have no liability for consequential damages.

STORAGE & DISPOSAL

PESTICIDE STORAGE: Store in a cool dry area away from heat or open flame and out of reach of children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. If Empty: Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PO Box 498303
Cincinnati, OH 45249

For Information Call 1-513-225-5615
EPA Reg. No. 89014 - 1
EPA Est. No. 089014-OH-001
™ Trade Mark 2012 SWAN Properties LLC