



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Antimicrobials Division (7510P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

89110-19

Date of Issuance:

7/30/18

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

BIONIX ® TP8

Name and Address of Registrant (include ZIP Code):

ISOMERIC Industries
 361 17th Street NW #1224
 Atlanta, GA 30363

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Antimicrobials Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the following label changes before you release the product for shipment:
3. Revise the EPA Registration Number to read, "EPA Reg. No. 89110-19."
4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Signature of Approving Official:

Eric Miederhoff, Product Manager 31
 Regulatory Management Branch I
 Antimicrobials Division (7510P)
 Office of Pesticide Programs

Date:

7/30/18

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated April 20, 2018
- Alternate CSF #1 April 24, 2018
- Alternate CSF #2 April 24, 2018

If you have any questions, please contact Zebora Johnson by phone at (703)308-7080 or via email at johnson.zebora@epa.gov.

Enclosure: Accepted Label

BIONIX TP8

INDUSTRIAL ANTIMICROBIAL

ACCEPTED

07/30/2018

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 89110-19

Active Ingredient:

Tetrakis(hydroxymethyl) phosphonium sulfate.....7.5%

Other Ingredients: 92.5%

Total: 100.0%

EPA Reg. No. 89110-RO EPA Est. _____ **NET CONTENTS:** As Marked on Container

KEEP OUT OF REACH OF CHILDREN DANGER

FIRST AID	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice• Have person sip a glass of water if able to swallow Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center for further treatment advice. If breathing is difficult, oxygen should be administered by qualified personnel.
Note to physician: Probable mucosal damage may contraindicate the use of gastric lavage.	
General Information: Have the product container or label with you when calling a poison control center or doctor or going for treatment. For non-emergency and general information on product use, etc., information pertaining to this product, call the National Pesticides Information Center at 1-800-858-7378 (NPIC web site: www.npic.orst.edu). For emergencies, call the Poison Control Center 1-800-222-1222.	

[See side panels for additional Precautionary Statements]

In case of emergency, call INFOTRAC: (800)-535-5053

Manufactured for:
ISOMERIC INDUSTRIES
361 17TH Street NW #1224
Atlanta, GA 30363
678-665-4275

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

DANGER: Corrosive: Causes irreversible eye damage. Fatal if swallowed or absorbed through skin or harmful if inhaled. Do not get in eyes, on skin, or on clothing. Wear goggles, face shield, or safety glasses. Wear coveralls over long-sleeved shirts and long pants, socks, chemical-resistant foot-wear and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothes before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Not for use in paint.

INDUSTRIAL AND/OR COMMERCIAL RECIRCULATING COOLING WATER SYSTEMS (for control of bacteria, fungi and algae) [Not for use in California]

Note: For cooling water systems of equal to or greater than 4000 gallons, do not apply by open pouring of liquid to cooling water systems; a metering pump delivery system is required for this use and application method.

Initial Slug Dose: Add 533.3-3496.6 ppm of BIONIX TP8 (40-262.5 ppm THPS) for 2-6 hours based on total water volume. Repeat until control is obtained. Thereafter, add either **Intermittently** 350-1400 ppm of BIONIX TP8 (26-105 ppm THPS) or **Continuously** 186.6-653.3 ppm of BIONIX TP8 (14-49 ppm THPS) per day. Dirty systems must be cleaned prior to treatment.

[or:]

[The following set of directions may be used in lieu of corresponding directions above:]

Note: For cooling water systems of equal to or greater than 4000 gallons, do not apply by open pouring of liquid to cooling water systems; a metering pump delivery system is required for this use and application method.

INDUSTRIAL AND/OR COMMERCIAL RECIRCULATING COOLING WATER SYSTEMS (for control of bacteria, fungi and algae)

Initial Slug Dose: Add 933.3-3496.6 ppm of BIONIX TP8 (70-262.5 ppm THPS) based on total water volume. Repeat until control is obtained. Thereafter, add either **Intermittently** 350-1400 ppm of BIONIX TP8 (26-105 ppm THPS) or **Continuously** 186.6-653.3 ppm of BIONIX TP8 (14-49 ppm THPS) per day. Dirty systems must be cleaned prior to treatment.

HEAT TRANSFER SYSTEMS (Evaporative Condensers, Dairy Sweetwater Systems, Hydrostatic Sterilizers and Retorts, Brewery and Other Pasteurizers, and Warmers)

Add BIONIX TP8 at the same application rates, and in the same manner as described above. It should be added to the system at a point of uniform mixing such as a basin area, sump area, or other reservoir or collecting area from which the treated water will be circulated uniformly throughout the system.

SERVICE WATER AND AUXILIARY SYSTEMS

BIONIX TP8 should be added to service water and auxiliary systems at the same application rates, and in the same manner as described above. It should be added to the system at a point of uniform mixing such as a basin area, sump area, or other reservoir or collecting area from which the treated water will be circulated uniformly throughout the system.

AIR WASHER SYSTEMS (For control of bacteria and fungi) [Not for use in California]

This product may be used only in air washer systems which have mist eliminating components. Pre-clean the system with detergent and allow air washer to run with fan on for two hours. Flush and check nozzles, manually cleaning as necessary. Add 350-1400 ppm of BIONIX TP8 (26-70 ppm THPS) at a point where uniform mixing and even distribution will occur. Repeat as needed to maintain control.

PAPER AND PAPERBOARD MANUFACTURING (for control of bacteria, fungi and algae)

a) For use as a slimicide in the manufacture of paper and paperboard products and adhesives that do not contact food.

Dosing: Additions should be made at a point in the system where mixing action is good, e.g. raw stock chest beater or mixing unit. Add intermittently or continuously depending on mill conditions.

Intermittent Dosing: Add 326.6-4666.6 ppm of BIONIX TP8 (24.5-350 ppm THPS) based on total water volume or an equivalent based on dry weight of paper produced.

Continuous Dosing: Add 186.6-653.3 ppm of BIONIX TP8 (14-49 ppm THPS) based on total water volume or an equivalent based on dry weight of paper produced.

b) For use as a preservative to retard microbial growth in water-based coatings, starches, pigments and filler slurries. Do not use in paper and paperboard and adhesives that will contact food. [Not for use in California]

The treatment rate necessary to retard spoilage of the additive will vary with the extent of contamination of make-up water and the length of storage.

Dosing: Apply from 2333.3-6666.6 ppm of BIONIX TP8 (175-500 ppm THPS) to the additive to be preserved based on the total weight of the additive and water.

INDUSTRIAL FRESH WATER SYSTEMS [Not for use in California]

Do not use in freshwater used in the manufacture of paper and paperboard products that may contact food. BIONIX TP8 is effective in controlling algae in holding ponds and in controlling bacteria and fungi in holding and processing tanks of industrial fresh water systems supplying water to pulp and paper mills, textile mills, and other manufacturing plants. In pulp and paper mills, treatment of the fresh water with BIONIX TP8 can make an important contribution to slime control. The use of BIONIX TP8 as described will reduce the development of slime in fresh water pipes and other equipment, and on the pulp and paper mill machine parts contacted by fresh water.

For the control of algae in industrial fresh water systems, BIONIX TP8 should be added to provide a concentration of 13.3-133.3 ppm of product (1-10 ppm of THPS). Treatment should be based on the amount of water entering a pond or reservoir or leaving the pond or reservoir and entering the immediate processing operations. While treatment can be made continuously, regular slug-dosing treatment will provide adequate control.

INDUSTRIAL WASTEWATER SYSTEMS (Wastewater Systems, Wastewater Sludge and Wastewater Holding Tanks) [Not for use in California]

BIONIX TP8 should be added to a wastewater system or sludge at a convenient point of uniform mixing such as digester.

Slug Dosing: Add 533.3-3500 ppm of BIONIX TP8 (40-262.5 ppm THPS) per 1,000 gallons of wastewater or sludge.

Continuous Dosing: Add 13.3-266.6 ppm of BIONIX TP8 (1-20 ppm THPS) per 1,000 gallons of wastewater or sludge.

MACROFOULING CONTROL [Not for use in California]

BIONIX TP8 should be added continuously to maintain a level of 20 ppm active ingredient (THPS) in the system for a period of at least 96 hours.

Initial Dose: When macrofouling is present in the system, apply 266.6 ppm of BIONIX TP8 (20 ppm THPS) based on total water volume. Continue to add BIONIX TP8 as needed to maintain the 20 ppm active ingredient (THPS) level for a period of at least 96 hours.

FIRE PROTECTION SYSTEMS

BIONIX TP8 is effective at controlling microbial growth in waters and on pipe surfaces in fire protection systems. Such microbial growth when combined with other forms of corrosion can lead to accelerated corrosion rates and pitting corrosion, commonly referred to as microbiologically influenced corrosion. BIONIX TP8 also helps to remove free oxygen from the water, thus eliminating an important nutrient for bacteria and an important reactant in many corrosion reactions.

BIONIX TP8 should be added to a fire protection system using a chemical metering pump capable of variable pump rates. The BIONIX TP8 should be injected at a point, such as a riser, manifold or makeup feed water line, where uniform mixing and distribution will occur. Add 1000-4000 ppm BIONIX TP8 (75-300 ppm THPS) depending on severity of microbial contamination in the system. Repeat as needed.

SOLUTIONS / EMULSIONS

Not for use in manufacture of paper and paperboard products and adhesives that may come in contact with food. For the preservation of solutions, emulsions, adhesives and other aqueous liquid products, the addition of 0.233%-2.33% of BIONIX TP8 (0.0175%-0.175% THPS) is effective. Add at a point in the processing system where there will be sufficient time and agitation for good mixing and dispersion. The exact amount of BIONIX TP8 to be added for the preservation of given formulations will depend on the components as well as local storage time and requirements.

OIL FIELD AND PETROCHEMICAL OPERATIONS

BIONIX TP8 is effective in controlling sulfate reducing bacteria, general aerobic bacteria, including microorganisms that contribute to biofilm formation in oil field recovery, processing and distribution applications and supporting systems; such as injection water, water holding tanks, disposal well water, recirculating water handling systems, and pipelines. BIONIX TP8 has been shown to dissolve iron sulfide and sequester iron when used under these conditions, leading to improved filter life and well injectivity, and reduction of hydrocarbon sheen. BIONIX TP8 is also effective for use in controlling microbial growth in fluids used for drilling and stimulation of oil wells.

Water Floods [Not for use in California]

BIONIX TP8 should be added to a water flood system at a point where uniform mixing will occur.

Initial Treatment: For a noticeably fouled system, add 933.3-3496.6 ppm BIONIX TP8 (70-262.5 ppm THPS). When added to a flowing system, slug dose for 2-6 hours based on flow rates. Repeat as necessary until control is achieved.

Subsequent Treatment: Once control has been achieved, add 66.6-980 ppm BIONIX TP8 (5-73.5 ppm THPS) weekly or as needed to maintain control. When added to a flowing system, slug dose for 2-6 hours based on flow rates.

Continuous Treatment: BIONIX TP8 can be dosed continuously at a level of 66.66-666.6 ppm (5-50 ppm THPS).

[or:]

[The following set of directions may be used in lieu of corresponding directions above:]

Water Floods

BIONIX TP8 should be added to a water flood system at a point where uniform mixing will occur.

Initial Treatment: For a noticeably fouled system, add 933.3-3496.6 ppm BIONIX TP8 (70-262.5 ppm THPS). When added to a flowing system, slug dose for 2-6 hours based on flow rates. Repeat as necessary until control is achieved.

Subsequent Treatment: Once control has been achieved, add 140-980 ppm BIONIX TP8 (10.5-73.5 ppm THPS) weekly or as needed to maintain control. When added to a flowing system, slug dose for 2-6 hours based on flow rates.

Continuous Treatment: BIONIX TP8 can be dosed continuously at a level of 140-666.6 ppm (10.5-50 ppm THPS).

Hydraulic Fracturing [Not for use in California]

BIONIX TP8 should be added to the frac water storage tanks or directly into the well head injection pipeline as the water is being pumped down-hole. Add 933.3-3496.6 ppm of BIONIX TP8 (50-262.5 ppm THPS), depending on the degree of bacterial fouling in the source water.

Oil and Gas Production and Transmission Pipelines and Systems

BIONIX TP8 should be added at a point in the pipeline where uniform mixing will occur. The application should be conducted to ensure maximum distribution of BIONIX TP8 through the entire internal surface of the pipeline by adding an amount of biocide which eventually comes out the other end of the pipeline. Criteria for success of the treatment will be reduction in bacterial count and/or corrosion rates.

Slug Dosing: Follow instructions for water flood treatment.

Continuous Dosing: BIONIX TP8 can be dosed continuously at a level of 140-1002.6 ppm (10.5-75 ppm THPS).

Drilling Muds, Packer Fluids, Completion and Workover Fluids

BIONIX TP8 should be added to these fluids at a point where uniform mixing will occur. Add 326.6-14000 ppm of BIONIX TP8 (24.5-1050 ppm THPS) to a freshly prepared fluid depending on severity of contamination.

Gas Storage Well and Systems [Not for use in California]

Individual injection wells should be treated with BIONIX TP8 at the same application rates, and in the same manner as described under Water Floods. Injections should be repeated as needed to maintain control.

Individual drips should be treated with a sufficient quantity of BIONIX TP8 to produce a concentration of 333.3-1333.3 ppm BIONIX TP8 (25-100 ppm THPS) when diluted by the water present in the drip. Injections should be repeated as needed to maintain control.

Well Remediation Operations [Not for use in California]

Individual production or injection wells may be bullheaded with BIONIX TP8 to control bacteria and simultaneously dissolve iron sulfide deposits. The BIONIX TP8 will be pumped into the well as a solution in water containing from 50% to 100% BIONIX TP8 (10%-20% THPS). The well is shut-in for a period of time (at least 6 hours) then put back into operation.

Hydrotesting

Water used to hydrotest pipelines or vessels should contain 666.6-6666.6 ppm BIONIX TP8 (50-500 ppm THPS), depending on water quality and length of time the equipment will remain idle.

Pipeline Pigging and Scraping Operation

Add BIONIX TP8 to a slug of water immediately following the scraper (ideally this water volume can be kept to a minimum and contained between the scraper and a trailing pig). Sufficient BIONIX TP8 should be added to produce a concentration of 0.025% to 0.25% (50-500 ppm THPS) in the water at the discharge point or pig trap, depending on the length of the pipeline and the severity of the biofouling.

[[Information in brackets] is optional or instructional; italics in brackets indicate instructions not part of labeling.]

STORAGE AND DISPOSAL

[For rigid non refillable containers 5 gallons or less:]

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store this product in a cool, dry area away from direct sunlight and heat to avoid deterioration. In case of a spill, flood the area with large quantities of water.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING AND DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling or reconditioning if appropriate. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[For rigid nonrefillable containers greater than 5 gallons:]

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store this product in a cool, dry area away from direct sunlight and heat to avoid deterioration. In case of a spill, flood the area with large quantities of water.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING AND DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling or reconditioning if appropriate. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[For refillable containers:]

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store this product in a cool, dry area away from direct sunlight and heat to avoid deterioration. In case of a spill, flood the area with large quantities of water.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING AND DISPOSAL: Refillable container. Refill this container with THPS only. Do not reuse this container for any other purpose. Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. Pressure rinse the container for final disposal as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[For Bulk Shipment Transport Vehicles: Note: In accordance with 40CFR 156 (e) bulk shipment transport vehicles are exempt from the requirements for container handling and disposal, therefore, only pesticide storage and disposal is required on bulk transport vehicle labeling]

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store this product in a cool, dry area away from direct sunlight and heat to avoid deterioration. In case of a spill, flood the area with large quantities of water.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

[In the text "this product" or alternate brand name may be substituted for Tolcide®PS20A product name]

[[Information in brackets] is optional or instructional; italics in brackets indicate instructions not part of labeling.]

[All products must bear a batch code. This is a lot number or other code used by the registrant or producer to identify the batch of the pesticide product distributed and sold. Location may be on the container or the label.]

Note: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, buyer assumes all risk of use and/or handling of this material, when such use and/or handling is contrary to label directions.