

89167-17

9/26/2012

1/20



U S ENVIRONMENTAL PROTECTION AGENCY

**Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave , NW
Washington, D C 20460**

EPA Registration
Number

89167-17

Date of Issuance

SEP 26 2012

NOTICE OF PESTICIDE

Registration
 Reregistration

(under FIFRA as amended)

Term of Issuance **Unconditional**

Name of Pesticide Product

AX SU 50-25 Herbicide

Name and Address of Registrant (include ZIP Code)

**Axion Ag Products, LLC c/o Lighthouse Product Services
410 North Boyland Ave 3937 Cedarwood Lane
Raleigh, NC 27603 Johnstown, CO 80534**

Note Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant the above named pesticide is hereby registered/reregistered under the Federal Insecticide Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment the Administrator on his motion may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA provided that you submit and/or cite all data required for registration review/reregistration of your product when the Agency requires all registrants of similar products to submit data.

The basic confidential statement of formula (CSF) dated April 26, 2012 is acceptable.

A stamped copy of the label is enclosed for your records. Submit one (1) copy of the final printed label before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions regarding this Notice, please contact Mindy Ondish at (703)605-0723 or at ondish.mindy@epa.gov

Signature of Approving Official

**Kable Bo Davis
Product Manager 25
Herbicide Branch
Registration Division (7505P)**

Date

SEP 26 2012

2/20

GROUP 2 HERBICIDE

AX SU 50-25 Herbicide

DRY FLOWABLE

FOR USE ON WHEAT (INCLUDING DURUM), BARLEY, OAT, TRITICALE, AND FALLOW

ACTIVE INGREDIENTS

Thifensulfuron-methyl - Methyl 3-[[[(4-methoxy-6 methyl-1 3 5 triazin-2 yl) amino]carbonyl]amino]sulfonyl]-2 thiophenecarboxylate	50 0%
Tribenuron-methyl Methyl 2-[[[N-(4-methoxy-6-methyl-1 3 5 triazin 2 yl) methylamino]carbonyl]amino]sulfonyl]benzoate	25 0%

OTHER INGREDIENTS

25 0%

TOTAL

100 0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION - PRECAUCION**

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle
(If you do not understand the label find someone to explain it to you in detail)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

**For Chemical Emergency, Spill, Leak, Fire, Exposure, or Accident
Call CHEMTREC Day or Night
Within USA and Canada 1 800-424-9300
Outside USA and Canada +1 703-527-3887 (collect calls accepted)**

EPA Reg No 89167-17 EPA EST NO
NET CONTENTS _____ OZS _____ LBS

Manufactured for
Axion Ag Products LLC
3937 Cedarwood Lane
Johnstown CO 80534

ACCEPTED

SEP 26 2012

Under the Federal Insecticide
Fungicide and Rodenticide Act
as amended for the pesticide
registered under
EPA Reg No 89167-17

092012

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION - PRECAUCION**

Causes moderate eye irritation Harmful if absorbed through skin Avoid contact with eyes skin or clothing

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below If you want more options follow the instructions for Category A on an EPA chemical resistance category selection chart

Applicators and other handlers must wear

- Long sleeved shirt and long pants
- Chemical-resistant gloves Category A (such as butyl rubber natural rubber neoprene rubber or nitrile rubber) all ≥14 mils
- Shoes plus socks

Follow manufacturer s instructions for cleaning/maintaining PPE If no such instructions for washables exist use detergent and hot water Keep and wash PPE separately from other laundry

USER SAFETY RECOMMENDATIONS	
Users Should	<ul style="list-style-type: none"> • Wash hands before eating drinking chewing gum using tobacco or using the toilet • Remove clothing/PPE immediately if pesticide gets inside Then wash thoroughly and put on clean clothing • Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing

FIRST AID	
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15 to 20 minutes • Call a poison control center or doctor for treatment advice
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes • Remove contact lenses if present after the first 5 minutes then continue rinsing eye • Call a poison control center or doctor for treatment advice

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center doctor or going for treatment For emergency information concerning this product call the National Pesticides Information Center (NPIC) at 1-800 858-7378 (NPIC Web site www.npic.orst.edu) Or call your poison control center at 1 800 222-1222

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate

PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site
- Make scheduled checks of spray equipment
- Ensure that all operation employees accurately measure pesticides
- Mix only enough product for the job at hand - Avoid over filling of spray tank
- Do not discharge excess material on the soil at a single spot in the field/grove or mixing/loading station
- Dilute and agitate excess solution and apply at labeled rates/uses
- Avoid storage of pesticides near well sites
- When triple rinsing the pesticide container be sure to add the rinsate to the spray mix

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Do not apply this product in a way that will contact workers or other persons either directly or through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe consult the agency responsible for pesticide regulation

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170 This standard contains requirements for the protection of agricultural workers on farms forests nurseries and greenhouses and handlers of agricultural pesticides It also contains requirements for training decontamination notification and emergency assistance It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants soil or water is

Coveralls

Chemical resistant gloves category A (such as butyl rubber natural rubber neoprene rubber or nitrile rubber) all ≥14 mils

Shoes plus socks

PRODUCT INFORMATION

This product is a dry flowable granule that is used for selective post-emergence weed control in wheat (including durum) barley oat triticale and fallow The best control is obtained when this product is applied to young actively growing weeds The use rate will depend on weed spectrum and size of weed at time of application The degree and duration of control may depend on the following

- weed spectrum and infestation intensity
- weed size at application
- environmental conditions at and following treatment

This product is noncorrosive nonflammable nonvolatile and does not freeze This product should be mixed in water and applied as a uniform broadcast spray

BIOLOGICAL ACTIVITY AND ENVIRONMENTAL CONDITIONS

This product is absorbed primarily through the foliage of plants rapidly inhibiting the growth of susceptible weeds One to 3 weeks after application to weeds (2 to 5 weeks for wild garlic) leaves of susceptible plants appear chlorotic and the growing point subsequently dies

This product provides the best control in vigorously growing crops that shade competitive weeds Weed control in areas of thin crop stand or seeding skips may not be as satisfactory However a crop canopy that is too dense at application can intercept spray and reduce weed control

The herbicidal action of this product may be affected in crops stressed from adverse environmental conditions (such as extreme temperatures or moisture) abnormal soil conditions cultural practices or variations in crop variety In warm moist conditions the expression of herbicide symptoms is accelerated in cold dry conditions expression of herbicide symptoms is delayed In addition weeds hardened off by drought stress are less susceptible to this product

WEEDS CONTROLLED - ALL USES

Annual knawel
Annual sowthistle
Black mustard

Blue/Purple mustard
Broadleaf dock
Bur buttercup

Bushy wallflower/Treacle
mustard
Clasping pepperweed

Coast fiddleneck	Filaree (redstem Texas)	Redroot pigweed
Common buckwheat	Flixweed	Russian thistle*
Common chickweed	Green smartweed	Scentless chamomile/mayweed
Common cocklebur*	Kochia	Shepherd s-purse
Common groundsel	Ladysthumb	Slimleaf lambsquarters
Common lambsquarters	Lanceleaf sage*	Smallflower buttercup
Common radish	London rocket	Smallseed false fax
Common ragweed*	Marshelder	Stinking chickweed
Common sunflower	Mayweed chamomile	Stinking mayweed /Dogfennel
Corn chamomile	Miners lettuce	Swinecress
Corn gromwell*	Narrowleaf lambsquarters	Tansymustard
Corn spurry	Night flowering catchfly	Tarweed fiddleneck
Cowcockle	Pennsylvania smartweed	Tumble/Jim Hill mustard
Cress (mouse-ear)	Pineappleweed	Volunteer lentils
Curly dock	Prickly lettuce*	Volunteer peas
False chamomile	Prostrate knotweed	Volunteer sunflower
Field chickweed	Prostrate pigweed	Wild buckwheat*
Field pennycress	Redmaids	Wild chamomile
		Wild garlic*
		Wild mustard
		Wild radish*

WEEDS PARTIALLY CONTROLLED**

Canada thistle*	Mallow (common little)
Carolina geranium	Nightshade (cutleaf hairy)
Catchweed bedstraw	Vetch*
Cutleaf evening primrose	

* See SPECIFIC WEED PROBLEMS in the Cereals section below for more information

**Partial control A visual reduction of weed population as well as a significant loss of vigor for individual weed plants For better results use 0.5 or 0.6 ounce of this product per acre and include a tank mix partner such as 2,4-D MCPA Buctril Bison Bronate or Bronate Advanced) or dicamba (such as Diablo/Clarity) refer to the TANK MIXTURES section of this label

†Naturally occurring resistant biotypes of kochia prickly lettuce and Russian thistle are known to occur See the TANK MIXTURES and SPECIFIC WEED PROBLEMS sections of this label for additional details

Pre Plant or At Planting Burndown in Cotton, Field Corn, Grain Sorghum, Rice and Soybeans

Application of this product may be applied for burndown of emerged weeds before planting or at planting of cotton field corn grain sorghum rice and soybeans

This product may be used as part of a pre-plant or at-planting burndown treatment in combination with other suitable registered herbicides Read and follow all manufacturers label recommendations for the companion herbicide If those recommendations conflict with this label do not tank mix the herbicide with this product

In fields to be planted to cotton apply this product at 0.3 to 0.5 ounce per acre In fields to be planted to field corn grain sorghum rice or soybeans apply this product at 0.3 to 0.6 ounce per acre for control or partial control of the weeds listed on the EPA registered label Allow at least 14 days between application and planting of cotton corn soybeans or grain sorghum Include a nonionic surfactant petroleum based crop oil concentrate or vegetable-seed oil-based product (methylated seed oils are considered a vegetable seed-based oil)

If another herbicide is tank mixed with this product to increase the broadleaf weed spectrum select adjuvants based on the adjuvant limitations of the other companion herbicide

SPRAY ADJUVANTS

Nonionic Surfactant (NIS)

Apply at a rate (concentration) of 0.25 to 0.5% v/v (1-2 qt per 100 gal spray solution) Use the higher rate in hot and dry conditions to enhance control

Crop Oil Concentrate

Under dry conditions or during cool weather a petroleum based crop oil concentrate or vegetable-seed oil based product may be used in place of a nonionic surfactant at 1-2 gallon/100 gal of spray solution (1-2% v/v) to enhance weed control Use a petroleum-based crop oil concentrate with at least 14% emulsifiers/surfactant and 80% oil

Ammonium Nitrogen Fertilizer

An ammonium nitrogen fertilizer can be added to a surfactant or a crop oil concentrate to enhance control Alternatively a high quality sprayable grade of ammonium sulfate (21-0-0) may be used

IMPORTANT PRECAUTIONS

Seedling disease nematodes cold weather deep planting (more than 2") excessive moisture high salt concentration and/or drought may weaken cotton seedlings and increase the possibility of crop injury Cotton resumes normal growth once favorable growing conditions return

RESTRICTIONS

- DO NOT apply after planting field corn grain sorghum rice or soybeans
- DO NOT apply later than 14 days before planting cotton corn soybeans or grain sorghum
- DO NOT allow livestock to graze on or feed forage hay or straw from treated soybean fields
- DO NOT make more than one pre-plant or at-planting application of this product to field corn rice sorghum or soybeans per growing season
- DO NOT apply more than 0.6 oz of this product to rice field corn sorghum or soybeans pre-plant or at planting

FALLOW

APPLICATION TIMING

This product may be used as a fallow treatment in the spring or fall when the majority of weeds have emerged and are actively growing

USE RATES

Apply this product at 0.3 to 0.6 oz per acre to fallow Two applications of this product may be made provided the total amount applied does not exceed 1.0 oz per acre per crop season

This product should be applied in combination with other suitable registered fallow herbicides such as Landmaster II Fallow Master Credit plus 2.4-D (ester formulations work best) Credit plus Diablo/Banvel SGF/Clarity 2.4-D Diablo/Diablo SGF/Clarity

TANK MIXTURES IN FALLOW

This product when used as a fallow treatment should be tank mixed with other herbicides that are registered for use in fallow Read and follow all manufacturer's label instructions for the companion herbicide If those instructions conflict with this label do not tank mix the herbicide with this product

PREPLANT BURNDOWN

USE RATES

Wheat (including durum), Barley, Triticale and Oat

Apply this product at 0.3 to 0.6 oz per acre as a burndown treatment to wheat (including durum) barley triticale and oat to control emerged weeds prior to or shortly after planting (prior to emergence) Make applications when the majority of weeds have emerged and are actively growing

Cotton

This product may be applied for burndown of emerged weeds prior to the emergence of cotton

This product may be used as part of a pre-plant burn-down treatment in combination with other suitable registered pre-plant herbicides

Read and follow all manufacturer's label instructions for the companion herbicide If those instructions conflict with this label do not tank mix the herbicide with this product Apply this product at 0.3 to 0.5 oz

per acre for control or partial control of the weeds listed on this label. Allow at least 14 days between application of this product and planting of cotton. Include a non-ionic surfactant, petroleum based crop oil concentrate, or vegetable-seed oil based product (methylated seed oils are considered a vegetable seed-based oil).

If another herbicide is tank mixed with this product to increase the broadleaf weed spectrum, select adjuvants based on the adjuvant limitations of the companion herbicide.

SPRAY ADJUVANTS

Nonionic Surfactant (NIS)

Apply at a rate (concentration) of 0.25-0.5% v/v (1-2 qt per 100 gal spray solution). Use the higher rate in hot and dry conditions to enhance control.

Crop Oil Concentrate

Under dry conditions or during cool weather, a petroleum based crop oil concentrate, or vegetable seed oil based product may be used in place of a nonionic surfactant at 1-2 gallon/100 gal of spray solution (1-2% v/v) to enhance weed control. Use a petroleum-based crop oil concentrate with at least 14% emulsifiers/surfactant and 80% oil.

Ammonium Nitrogen Fertilizer

An ammonium nitrogen fertilizer can be added to a surfactant or a crop oil concentrate to enhance control. Alternatively, a high-quality sprayable grade of ammonium sulfate (21-0-0) may be used.

IMPORTANT PRECAUTIONS

Seedling disease, nematodes, cold weather, deep planting (more than 2"), excessive moisture, high salt concentration, and/or drought may weaken cotton seedlings and increase the possibility of crop injury. Cotton resumes normal growth once favorable growing conditions return.

RESTRICTION DO NOT apply later than 14 days before planting cotton.

Cotton

Apply this product at 0.3 to 0.5 oz per acre as a burndown treatment to cotton. Allow at least 14 days between application of this product and planting of cotton. Include a nonionic surfactant, petroleum based crop oil concentrate, or vegetable seed oil-based product (methylated seed oils are considered a vegetable seed based oil). If another herbicide is tank mixed with this product to increase the broadleaf weed spectrum, select adjuvants based on the adjuvant limitations of the companion herbicide.

Sugarbeets, Winter Rape and Canola

Apply this product at 0.3 to 0.6 oz per acre as a burndown treatment to sugarbeets, winter rape, and canola. Allow at least 60 days between application of this product and planting of sugarbeets, winter rape, and canola.

Any other crop (such as corn, rice, grain sorghum or soybeans)

Apply this product at 0.3 to 0.6 oz per acre as a burndown treatment to any other crop (such as corn, rice, grain sorghum, or soybeans). Allow at least 45 days between application of this product and planting of any other crop (such as corn, rice, grain sorghum, or soybeans).

Sequential treatments of this product may also be made provided the total amount of product applied during one fallow/pre plant cropland season does not exceed 1.0 oz per acre, for example, 0.5 oz in the fall followed by 0.5 oz in the spring. Use the 0.6 oz per acre rate when weed infestation is heavy and predominantly consists of those weeds listed under PARTIAL CONTROL, or when application timing and environmental conditions are marginal. (See APPLICATION TIMING Section for restriction on planting intervals.)

PREPLANT APPLICATION FOR BROADLEAF WEED AND WILD GARLIC CONTROL PRIOR TO PLANTING CORN, COTTON, RICE, GRAIN SORGHUM OR SOYBEANS

In the states of AL, AR, DE, GA, IA, IN, IL, KY, LA, MD, MO, MS, NC, OH, PA, SC, TN, TX, and VA, this product may be applied prior to planting corn, cotton, rice, grain sorghum, or soybeans for the control of certain broadleaf weeds and wild garlic. This product may be applied from late fall through early spring.

Do not apply within 45 days prior to planting the above listed crops. Do not use this product through any type of irrigation equipment.

APPLICATION INFORMATION

This product provides postemergence activity for short term control or partial control of labeled weeds. Degree of control and duration of effect are dependent on rate used, sensitivity and size of target weed, environmental conditions at the time of and following application, and spray coverage.

Best results are obtained when this product is applied to young, actively growing weeds. The use rate will depend on weed spectrum and size of weed at time of application. However, typical symptoms of dying weeds (discoloration) may not be noticeable for 1-3 weeks after application (2-5 weeks for wild garlic) depending on the environmental conditions and weed susceptibility. Warm, moist conditions following treatment promote the activity of this product while cold, dry conditions delay the activity. Weeds hardened off by cold weather, drought stress, or too wet conditions will be less susceptible.

Some naturally occurring weed biotypes* resistant to this product are known to exist. If weeds listed on this label are not satisfactorily controlled, respray problem areas in a timely and effective manner using a broadleaf herbicide having a different mode of action.

*Biotypes are naturally occurring individuals of the species which have a slightly different make up. Resistant biotypes may look exactly the same as susceptible biotypes. Herbicide resistant biotypes are able to survive a use rate several times higher than needed to control susceptible biotypes.

APPLICATION TIMING

Apply this product at least 45 days prior to planting corn, cotton, rice, grain sorghum, or soybeans. Annual broadleaf weeds should be past the cotyledon stage, actively growing, but less than 4 inches tall or across. See SPECIFIC WEED PROBLEMS for information on vetch (hairy common), wild garlic, and wild radish.

WEED CONTROL Application Rates

Apply this product at 0.5 - 0.6 oz per acre to control emerged weeds. Use 0.6 oz per acre rate when weed infestation is heavy and predominantly consists of those weeds listed under PARTIAL CONTROL or when application timing and environmental conditions are marginal. Reduce the application rate to 0.5 ounces for fields to be planted to cotton.

Sequential treatments of this product to control emerged weeds may also be made provided the total amount of this product applied during one fallow cropland season does not exceed 1.0 oz per acre. For example, 0.5 oz in the fall followed by 0.5 oz in the spring. At least one of the sequential treatments should include a tank mix partner herbicide having a different mode of action than this product.

Foliar absorption is the primary means of this product uptake by plants; therefore, thorough coverage of all target weeds is essential.

Conditions which are conducive to healthy, actively growing plants optimize this product's weed control performance. Ideal conditions include warm temperatures and adequate soil moisture before, during, and immediately after application.

TANK MIXTURES IN PRE-PLANT BURNDOWN

This product may be used as a pre-plant burndown treatment alone or tank mixed with other herbicides that are registered for use as a pre-plant burndown product, such as Landmaster II, Fallow Master, Credit plus, Diablo/Diablo SGF/Clarity, or Diablo/Diablo SGF/Clarity alone.

Read and follow all manufacturers' label instructions for the companion herbicide. If those instructions conflict with this label, follow the most restrictive labeling (such as planting interval after application) or do not tank mix the herbicide with this product.

CEREALS

USE RATES

Do not use less than 0.3 oz of this product per acre.

Wheat (including Durum), Barley and Triticale

Apply this product at 0.3 to 0.6 oz per acre to wheat (including durum), barley, or triticale. Two applications of this product may be made provided the total amount applied does not exceed 1.0 oz per acre per crop season.

Use 0.3 to 0.4 oz of this product per acre for light infestation of the weeds listed under Weeds Controlled. Conditions at application should be optimum for effective treatment of these weeds.

Use 0.5 oz of this product per acre for heavy infestation of the weeds listed under Weeds Partially Controlled

Use 0.6 oz of this product per acre for heavy infestation of the weeds listed under Weeds Partially Controlled when application timing and environmental conditions are marginal (refer to Environmental Conditions and Biological Activity for best performance)

This product may be used as postemergence applications to triticale anytime after crop is in the 2-3 leaf stage but before the flag leaf is visible. Follow the postemergence use rate instructions listed for wheat.

Other suitable herbicides, fungicides, and insecticides registered for use on triticale may be tank mixed or used sequentially with these products providing the recommended application timing is the same. Read and follow all manufacturers' label instructions for the tank mix partner prior to use. The most restrictive provisions on either label apply.

Oat (Spring and Winter)

Apply this product at 0.3 to 0.4 oz per acre for control of the weeds listed in Weeds Controlled table. Do not make more than one application of this product per crop season on oat.

APPLICATION TIMING

Wheat (Including Durum), Barley, Winter Oat and Triticale

Make applications after the crop is in the 2 leaf stage but before the flag leaf is visible.

Spring Oat

Make applications after the crop is in the 3 leaf stage but before jointing. Do not use on Ogle, Porter or Premier varieties as crop injury can occur.

Since this product has very little or no soil activity, it controls only those weeds that have germinated; therefore, apply this product when all or most of the weeds have germinated. Annual broadleaf weeds should be past the cotyledon stage, actively growing, and less than 4" tall or wide. Wild garlic plants should be less than 12" tall with 2 to 4" of new growth. See Specific Weed Problems for more information.

Rainfall immediately after treatment can wash this product off of weed foliage, resulting in reduced weed control. Several hours of dry weather are needed to allow this product to be sufficiently absorbed by weed foliage.

SPECIFIC WEED PROBLEMS - CEREALS

Canada thistle For control in wheat, barley and triticale, use 0.6 oz per acre plus surfactant when all thistles are 4 to 8" with 2 to 6" of new growth. Make the application in the spring. Control will be improved by using this product in combination with 2,4-D (refer to TANK MIXTURES).

For control in oat, use 0.4 oz of this product per acre plus 2,4-D (refer to TANK MIXTURES).

Common cocklebur, Common ragweed, Lanceleaf sage In wheat, barley and triticale, apply this product at 0.4 to 0.5 oz per acre in combination with 2,4-D at rates from 1/4 to 3/8 lb active ingredient (ester formulations work best) when weeds are small and actively growing. When using 1/4 lb active ingredient of 2,4-D, be sure to add surfactant at the rate of 1/4 to 1/2 quart per 100 gallons of spray solution (0.06 to 0.125% v/v - use the higher rate under stress conditions).

For control in oat, use 0.4 oz of this product per acre plus 2,4-D. Refer to the Tank Mixtures sections of this label for additional details.

Corn gromwell, Wild buckwheat For control in wheat, barley and triticale, use 0.5 to 0.6 oz of this product per acre plus surfactant. For control in oat, use 0.4 oz of this product per acre plus 2,4-D, MCPA or Buctril (refer to TANK MIXTURES).

Kochia, Russian thistle, Prickly lettuce Naturally occurring resistant biotypes of these weeds are known to occur. For best results, use this product in a tank mix with dicamba (such as Diablo/Diablo SGF/Clarity) and 2,4-D or Bromoxynil (such as Buctril) and 2,4-D (3/4 - 1 pt Buctril + 1/4 - 3/8 lb active ingredient 2,4-D ester). This product should be applied in the spring when weeds are less than 2" tall or 2" across and are actively growing. Refer to the Tank Mixtures section of this label for additional details.

Vetch (common and hairy) For control in wheat, barley and triticale, use 0.5 to 0.6 oz of this product per acre plus surfactant when vetch is less than 6" in length. For severe infestations of vetch or when vetch is greater than 6" in length, use this product in combination with 2,4-D or MCPA (refer to the Tank Mixtures section of this label).

For control in oat use 0.4 oz of this product per acre plus 2.4-D or MCPA (refer to TANK MIXTURES)

Wild garlic For control in wheat, barley and triticale use 0.5 to 0.6 oz of this product per acre plus surfactant when wild garlic plants are less than 12" tall with 2 to 4" of new growth. For severe infestations use the 0.6 oz per acre rate of this product. Plants hardened-off by cold weather and/or drought stress may be more difficult to control. Thorough spray coverage of all garlic plants is essential. Typical symptoms of dying garlic plants may not be noticeable for 2 to 5 weeks.

For control in oat use 0.4 oz of this product per acre plus 2.4-D or MCPA (refer to TANK MIXTURES)

Wild radish For best results in wheat, barley and triticale apply 0.4 to 0.6 oz of this product per acre plus surfactant either in the fall or spring to wild radish rosettes less than 6 inches in diameter. Applications made later than 30 days after weed emergence will result in partial control. For increased control of severe wild radish infestations or wild radish emerged greater than 30 days apply this product at 0.3 oz per acre in combination with MCPA at 1/4 lb active ingredient per acre. Surfactant is required when tank mixing with MCPA add 1 quart per 100 gallons of spray solution (0.25% vol/vol). Fall applications should be made prior to hardening off of plants.

For control in oat use 0.4 oz of this product per acre plus 2.4 D or MCPA (refer to TANK MIXTURES)

TANK MIXTURES - CEREALS

This product may be tank mixed with other suitable registered herbicides to control weeds listed as suppressed weeds resistant to this product or weeds not listed under Weeds Controlled. Read and follow all manufacturers label instructions for the companion herbicide. If those instructions conflict with this label do not tank mix the herbicide with this product.

This product can also be mixed with registered fungicides, insecticides or liquid fertilizer for use on wheat, barley, triticale, oat or fallow.

With 2,4 D (amine or ester) or MCPA (amine or ester)

This product may be tank mixed with the amine and ester formulations 2.4 D and MCPA herbicides for use on wheat, barley, triticale and oat.

For best results in the Red River Valley and adjacent areas of North Dakota and Minnesota add the ester formulations of 2.4 D or MCPA herbicides to the tank at 3/8 lb active ingredient (such as 3/4 pt of a 4 lb/gal product, 1/2 pt of a 6 lb/gal product). No additional surfactant is needed with this mixture.

For best results in other areas add the ester formulations of 2.4 D or MCPA herbicides to the tank at 1/4 to 3/8 lb active ingredient (such as 1/2 - 3/4 pt of a 4 lb/gal product, 1/3 - 1/2 pt of a 6 lb/gal product). Surfactant may be added to the mixture at 1/2 to 1 qt per 100 gal of spray solution (0.125 to 0.25% v/v) however adding surfactant may increase the potential for crop injury especially at the higher phenoxy rates.

Higher rates of 2.4-D or MCPA may be used but do not exceed the highest rate allowed by those respective labels. Read and follow all label instructions on timing, precautions and warnings for these herbicides before using these tank mixtures.

With dicamba (such as Diablo/Diablo SGF/Clarity)

This product may be tank mixed with 1/16 to 1/8 lb active ingredient dicamba (such as 2.4 fluid oz Diablo, 4.8 fluid oz Diablo SGF, 2-4 fluid oz Clarity). Use higher rates when weed infestation is heavy. Surfactant may be added to the mixture at 1/2 to 1 qt per 100 gal of spray solution (0.125 to 0.25% v/v) however adding surfactant may increase the potential for crop injury. Refer to the specific dicamba label for application timing and restrictions.

Tank mixes of this product plus dicamba may result in reduced control of some broadleaf weeds.

With 2,4 D (amine or ester) and Diablo/Clarity

This product may be applied in a 3-way tank mix with formulations of dicamba and 2.4 D. Make application of this product + 1/16 to 1/8 lb active ingredient dicamba (such as 2-4 fluid oz Diablo, 4.8 fluid oz Diablo SGF, 2-4 fluid oz Clarity) + 1/4 - 3/8 lb active ingredient 2.4-D Ester or Amine per acre. Use higher rates when weed infestation is heavy. Surfactant may be added to the mixture at 1/2 to 1 qt per 100 gal of spray solution (0.125 to 0.25% v/v) however adding surfactant may increase the potential for crop injury. Consult the specific 2.4-D label, dicamba label or local recommendations for more information and restrictions.

Apply this 3 way combination to winter wheat and winter oat after the crop is tillering and prior to jointing (first node) In Spring wheat (including durum) and Spring oat apply after the crop is tillering and before it exceeds the 5 leaf stage

In Spring barley apply after the crop is tillering and before it exceeds the 4 leaf stage

With bromoxynil (such as Buctril, Bronate, Bronate Advanced, or Rhino)

This product may be tank mixed with bromoxynil containing herbicides registered for use on wheat barley triticale or fallow For best results add bromoxynil containing herbicides to the tank at 3/16 to 3/8 lb active ingredient per acre (such as Bronate or Buctril at 3/4 - 1-1/2 pt per acre)

Read and follow all label instructions on timing precautions and warnings for these herbicides before using these tank mixtures Follow the most restrictive labeling Tank mixes of this product plus Buctril may result in reduced control of Canada thistle

With Express or Express XP Herbicide

This product may be tank mixed with Express or Express XP based on local recommendations Read and follow all label instructions on timing precautions and warnings for these herbicides before using this tank mixture

With Ally or Ally XP Herbicide

This product may be tank mixed with Ally or Ally XP based on local recommendations Read and follow all label instructions on timing precautions and warnings for these herbicides before using this tank mixture

With Starane

For improved control of Kochia (2 - 4 tall) Russian thistle mustard species and wild buckwheat This product may be tank mixed with 1/3 to 1 1/3 pints per acre of Starane Refer to the The most restrictive provisions on either label will apply Do not use the tank mix if any restrictions on the Starane label conflict with recommendations on this herbicide label

Other suitable registered herbicides fungicides and insecticides registered for use on small grains or fallow may be tank mixed or used sequentially with this mixture Read and follow all manufacturers label instructions for the companion herbicide The most restrictive provisions on either label will apply

With Starane + Salvo

This product can be tank mixed with Starane + Salvo herbicides for improved control of broadleaf weeds in wheat barley fallow and oats

For improved control of Kochia (2 - 4 tall) Russian thistle mustard species and wild buckwheat this product may be tank mixed with 2/3 to 2 2/3 pints per acre of Starane + Salvo Refer to this product's label and the Starane and Salvo labels for information regarding use restrictions labeled crops rotational cropping recommendations sprayer cleanup use precautions and other information The most restrictive provisions on either label will apply Do not use the tank mix if any restrictions on the Starane or Salvo label conflict with instructions on this product's herbicide label

With Starane + Sword

This product can be tank mixed with Starane + Sword herbicides for improved control of broadleaf weeds in wheat barley fallow and oats

For improved control of Kochia (2 - 4 tall) Russian thistle mustard species and wild buckwheat this product may be tank mixed with 3/4 to 2-3/4 pints per acre of Starane + Sword Refer to this product's label and the Starane and Sword labels for information regarding use restrictions labeled crops rotational cropping recommendations sprayer cleanup use precautions and other information The most restrictive provisions on either label will apply Do not use the tank mix if any restrictions on the Starane or Sword label conflict with instructions on this product's label

Other suitable registered herbicides fungicides and insecticides registered for use on cereal grains or fallow may be tank mixed or used sequentially with this mixture Read and follow all manufacturers label instructions for the companion herbicide The most restrictive provisions on either label will apply

With Aim

This product can be tank mixed with Aim herbicide for improved control of weeds in wheat barley and triticale

With Stinger or Cutback or Cutback M or Widematch

This product can be tank mixed with Stinger Cutback Cutback M or Widematch herbicides for improved control of weeds in wheat barley and triticale

With Other Broadleaf Herbicides

Tank mixes of this product plus metribuzin may result in reduced control of wild garlic

With Hoelon Herbicide

This product may be used in combination with Hoelon 3EC and Buctril herbicides in accordance with the Hoelon 3EC label For best results use the three way tank mix of this product at 0.4 oz per acre plus Hoelon 3EC at 2 2/3 pt per acre plus Buctril at 1 1/2 pt per acre Apply only to winter wheat This tank mix should only be used under good soil conditions when wild oat is in the 1.4 leaf stage If conditions are not ideal for the performance of Hoelon 3EC wild oat control may be reduced Be sure to follow all warnings and cautions on the Hoelon 3EC and Buctril labels

With Assert Herbicide or Avenge Herbicide

This product can be tank mixed with Avenge or Assert When tank mixing with Assert always include another broadleaf weed herbicide with a different mode of action (for example 2.4 D ester MCPA ester Buctril or Bronate) Tank-mixed applications of this product plus Assert may cause temporary crop discoloration stunting or injury when heavy rainfall occurs shortly after application

With Discover NG

This product can be tank mixed with Discover NG herbicide for improved control of weeds in spring wheat

With Everest

This product can be tank mixed with Everest herbicide for improved control of weeds in spring wheat

With Maverick

This product can be tank mixed with Maverick herbicide for improved control of weeds in wheat

With Puma

This product can be tank mixed with Puma 1 EC for control of some annual grass weeds This tank mix may also include MCPA ester bromoxynil or bromoxynil/MCPA for greater spectrum of broadleaf control see Puma IEC label for specific use directions and restrictions on tank mixes

With other grass control products

Tank mixtures of this product and grass control products may result in poor grass control AXION AG PRODUCTS LLC recommends that you first consult your state experiment station university or extension agent Agricultural dealer or AXION AG PRODUCTS LLC representative as to the potential for antagonism before using the mixture If no information is available limit the initial use of this product and the grass product to a small area

With Insecticides

This product may be tank mixed or used sequentially with insecticides (or fungicides) registered for use on cereal grains However under certain conditions (drought stress or if the crop is in the 2.4 leaf stage) tank mixes or sequential applications of this product with organophosphate insecticides (such as parathion) may produce temporary crop yellowing or in severe cases crop injury Test these mixtures in a small area before treating large areas

Do not use this product plus Malathion as crop injury will result

With Liquid Nitrogen Solution Fertilizer

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water Run a tank mix compatibility test before mixing this product in fertilizer solution This product must first be slurried with water and then added to liquid nitrogen solutions (e.g. 28-0-0 32 0-0) Ensure that the agitator is running while the product is added Use of this mixture may result in temporary crop yellowing and stunting

If using low rates of liquid nitrogen fertilizer in the spray solution (less than 50% of the spray solution volume) the addition of surfactant is necessary Add surfactant at 1/4 qt - 1 qt per 100 gal of spray solution (0.06 - 0.25% v/v) based on local recommendations

When using high rates of liquid nitrogen fertilizer solution in the spray solution adding surfactant increases the risk of crop injury Consult your agricultural dealer consultant fieldman or AXION AG PRODUCTS LLC representative for a specific recommendation before adding an adjuvant to these tank mixtures

If 2 4 D or MCPA is included with this product and fertilizer mixture ester formulations tend to be more compatible (See manufacturer's label) Additional surfactant is not needed when using this product in tank mix with 2 4-D ester or MCPA ester and liquid nitrogen fertilizer solutions

Note In certain areas east of the Mississippi river unacceptable crop response may occur with use of straight or dilute nitrogen fertilizer carrier solutions where cold temperatures or widely fluctuating day/night temperatures exist In these areas consult your agricultural dealer consultant field advisor or AXION AG PRODUCTS LLC representative for a specific recommendation before using nitrogen fertilizer carrier solutions

Liquid nitrogen fertilizer solutions that contain sulfur can increase crop response

Do not use low rates of liquid nitrogen fertilizer solution as a substitute for a surfactant

Do not use with liquid fertilizer solutions with a pH less than 3 0

SPRINKLER CHEMIGATION WITH THIS PRODUCT AND BRONATE FOR POSTEMERGENCE WEED CONTROL IN WINTER & SPRING WHEAT & SPRING BARLEY IN IDAHO

HOW TO USE

Use 0 4 to 0 5 oz of this product per acre in combination with 3/4 to 1 1/2 pint Bronate per acre Apply to wheat barley and triticale after the 3-leaf stage but before the flag leaf is visible Make only one chemigation application of this tank mixture per crop year

For best results apply to broadleaf weeds up to the 4 leaf stage or 2 inches in height or 1 inch in diameter whichever comes first Consult this product and Bronate package labels for list of weeds controlled/suppressed

SPRINKLER IRRIGATION APPLICATION

Apply this tank mix through sprinkler irrigation systems including center pivot lateral move side (wheel) roll solid set or hand move irrigation systems only Do not apply these herbicides through any other type of irrigation system

Crop injury lack of effectiveness or illegal pesticide residues in the crop can result from nonuniform distribution of treated water If you have questions about calibration you should contact State Extension Service specialists equipment manufacturers or other experts Do not connect an irrigation system (including greenhouse systems) used for this product's application to any public water system A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise

The sprinkler chemigation system must contain a functional check valve vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump The pesticide injection pipeline must also contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected Systems must use a metering pump such as a positive displacement injection pump (e.g diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock Do not apply when wind speed favors drift beyond the area intended for treatment

SPECIFIC REQUIREMENTS FOR APPLICATION THROUGH SPRINKLER IRRIGATION SYSTEMS

- 1 In center pivot and continuous lateral move systems this product + Bronate should be applied continuously for the duration of the water application In solid set systems application of the tank mix should be made during the last 30 to 45 minutes of the irrigation set
- 2 Set the sprinkler system to deliver approximately 0 5 inch or less of water per acre for best product performance
- 3 Fill the supply tank with half of the water amount desired add this product and agitate it well Add the Bronate and then add the remaining water amount with agitation Bronate requires a dilution with at least 4 parts water to 1 part Bronate

- 4 Agitation is recommended in the pesticide supply tank when applying this tank mix
- 5 The use of a surfactant is not recommended with this tank mix application
- 6 Inject this product + Bronate solution at least 8 feet ahead of a right angle turn of irrigation pipe to insure adequate mixing Allow sufficient time for the herbicide mixture to be flushed through the lines before turning off irrigation water
- 7 Follow both this product and Bronate label instructions for spray tank cleanout both before and after application Flush lines with clean water following application
- 8 Do not apply when wind speed favors drift beyond the area intended for treatment Avoiding spray drift is the responsibility of the applicator

MIXING INSTRUCTIONS

- 1 Fill the tank 1/4 to 1/3 full of water
- 2 While agitating add the required amount of this product
- 3 Continue agitation until the product is fully dispersed at least 5 minutes
- 4 Once the product is fully dispersed maintain agitation and continue filling tank with water This product should be thoroughly mixed with water before adding any other material
- 5 As the tank is filling add tank mix partners (if desired) then add the required volume of nonionic surfactant Always add surfactant last Do not use with spray additives that alter the pH of the spray solution below pH 5.0 or above pH 9.0 as rapid product degradation can occur Spray solutions of pH 6.0-8.0 allow for optimum stability of this product
- 6 If the mixture is not continuously agitated settling will occur If settling occurs thoroughly re-agitate before using
- 7 Apply product spray mixture within 24 hours of mixing to avoid product degradation
- 8 If this product and a tank mix partner are to be applied in multiple loads pre slurry this product in clean water prior to adding to the tank This will prevent the tank mix partner from interfering with the dissolution of this product

PRODUCT MEASUREMENT

This product is measured using the product volumetric measuring cylinder The degree of accuracy of this cylinder varies by ± 7.5% For more precise measurement use scales calibrated in ounces

CROP ROTATION - ALL USES

Wheat (including durum) Barley Triticale and Oat may be replanted anytime after the application of this product

Cotton can be planted 14 days after the application of this product Sugarbeets Winter Rape and Canola can be planted 60 days after the application of this product Any other crop may be planted 45 days after the application of this product

SURFACTANTS - ALL USES

Unless otherwise specified add a AXION AG PRODUCTS LLC recommended nonionic surfactant having at least 80% active ingredient at 1 to 2 qt per 100 gal of spray solution (0.25 to 0.5% v/v -refer to TANK MIXTURES for specific adjuvant instructions when this product is used in a tank mix)

For pre plant burndown in Cotton include a nonionic surfactant petroleum based crop oil concentrate or a vegetable-seed oil based product (methylated seed oils are considered a vegetable seed based oil) If another herbicide is tank mixed with this product to increase the broadleaf weed spectrum select adjuvants based on the adjuvant limitations of the companion herbicide

Consult your agricultural dealer applicator or AXION AG PRODUCTS LLC representative for a listing of recommended surfactants Antifoaming agents may be used if needed

Do not use low rates of liquid nitrogen fertilizer solution as a substitute for surfactant

GROUND APPLICATION ALL USES

For optimum spray distribution and thorough coverage use flat-fan or low-volume flood nozzles

For flat fan nozzles use a spray volume of at least 5 gal per acre (GPA)

For flood nozzles on 30 spacings use at least 10 GPA flood nozzles no larger than TK10 (or the equivalent) and a pressure of at least 30 psi For 40 nozzle spacings use at least 13 GPA for 60 spacings use at least 20 GPA It is essential to overlap the nozzles 100% for all spacings

Raindrop RA nozzles are not recommended for product applications as weed control performance may be reduced. Use screens that are 50-mesh or larger.

AERIAL APPLICATION - ALL USES

Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage at 2 to 5 GPA. Use at least 3 GPA in Idaho, Oregon, or Utah.

Do not apply this product by air in the state of New York. See the Spray Drift Management section of this label.

GRAZING

Do not graze livestock in treated areas. In addition, do not feed forage or hay from treated areas to livestock (harvested straw may be used for bedding and/or feed).

SPRAY EQUIPMENT

For specific application equipment, refer to the manufacturer's recommendations for additional information on GPA, pressure, speed, nozzle types and arrangements, nozzle heights above the target canopy, etc.

Be sure to calibrate air or ground equipment properly before application. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern with minimum drift. Use higher spray volumes to obtain better coverage when crop canopy is dense. Avoid swath overlapping and shut off spray booms while starting, turning, slowing, or stopping to avoid injury to the crop.

Do not make applications using equipment and/or spray volumes or during weather conditions that might cause spray to drift onto nontarget sites. For additional information on spray drift, refer to Spray Drift Management section of label.

Continuous agitation is required to keep this product in suspension.

SPRAYER CLEANUP

The spray equipment must be cleaned before this product is sprayed. Follow the cleanup procedures specified on the labels of the previously applied products. If no directions are provided, follow the six steps outlined in After Spraying This Product.

AT THE END OF THE DAY

It is recommended that during periods when multiple loads of this product are applied, at the end of each day of spraying, the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits which can accumulate in the application equipment.

AFTER SPRAYING THIS PRODUCT AND BEFORE SPRAYING CROPS OTHER THAN WHEAT, BARLEY, TRITICALE AND OAT

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of this product as follows:

- 1 Drain tank, thoroughly rinse spray tanks, boom, and hoses with clean water. Loosen and physically remove any visible deposits.
- 2 Fill the tank with clean water and 1 gal of household ammonia* (contains 3% active) for every 100 gal of water. Flush the hoses, boom, and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 min. Flush the hoses, boom, and nozzles again with the cleaning solution, and then drain the tank.
- 3 Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
- 4 Repeat step 2.
- 5 Rinse the tank, boom, and hoses with clean water.
- 6 If only Ammonia is used as a cleaner, the rinsate solution may be applied back to the crop(s) listed on this label. Do not exceed the maximum labeled use rate. If other cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

*Equivalent amounts of an alternate-strength ammonia solution or a AXION AG PRODUCTS LLC approved cleaner can be used in the cleanout procedure. Carefully read and follow the individual cleaner instructions. Consult your Ag dealer, applicator, or AXION AG PRODUCTS LLC representative for a listing of approved cleaners.

Notes

- 1 CAUTION Do not use chlorine bleach with ammonia as dangerous gases will form Do not clean equipment in an enclosed area
- 2 Steam cleaning aerial spray tanks is recommended prior to performing the above cleanout procedure to facilitate the removal of any caked deposits
- 3 When this product is tank mixed with other pesticides all cleanout procedures should be examined and the most rigorous procedure should be followed
- 4 In addition to this cleanout procedure all pre cleanout guidelines on subsequently applied products should be followed as per the individual labels
- 5 Where routine spraying practices include shared equipment frequently being switched between applications of this product and applications of other pesticides to product sensitive crops during the same spray season it is recommended that a sprayer be dedicated to this product to further reduce the chance of crop injury

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift The applicator is responsible for considering all these factors when making application decisions
AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control The presence of sensitive species nearby the environmental conditions and pest pressure may affect how an applicator balances drift control and coverage APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS! See Wind Temperature and Humidity and Temperature Inversions sections of this label

Controlling Droplet Size General Techniques

- Volume Use high flow rate nozzles to apply the highest practical spray volume Nozzles with higher rated flows produce larger droplets
 - Pressure Use the lower spray pressures recommended for the nozzle Higher pressure reduces droplet size and does not improve canopy penetration WHEN HIGHER FLOW RATES ARE NEEDED USE A HIGHER CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE
- Nozzle Type Use a nozzle type that is designed for the intended application With most nozzle types narrower spray angles produce larger droplets Consider using low drift nozzles

Controlling Droplet Size - Aircraft

- Number of Nozzles Use the minimum number of nozzles with the highest flow rate that provide uniform coverage
- Nozzle Orientation Orienting nozzles so that the spray is emitted backwards parallel to the airstream will produce larger droplets than other orientations
- Nozzle Type - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types
 - Boom Length - The boom length should not exceed 3/4 of the wing or rotor length longer booms increase drift potential
- Application Height Application more than 10 ft above the canopy increases the potential for spray drift

BOOM HEIGHT

Setting the boom at the lowest referenced height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind For ground equipment the boom should remain level with the crop and have minimal bounce

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph However many factors including droplet size and equipment type determine drift potential at any given wind speed AVOID GUSTY AND WINDLESS CONDITIONS

Note Local terrain can influence wind patterns Every applicator should be familiar with local wind patterns and how they affect spray drift

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions set up equipment to produce larger droplets to reduce effects of evaporation

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion Temperature inversions restrict vertical air mixing which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind They begin to form as the sun sets and often continue into the morning Their presence can be indicated by ground fog however if fog is not present inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion while smoke that moves upward and rapidly dissipates indicates good vertical air mixing

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind However it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product

AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream Some may reduce the potential for drift but if a sprayer is unsuitable for the application and/or set up improperly high drift potential can result It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application is configured properly and that drift is not occurring

Note Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration Consult the spray equipment section of this label to determine if use of an air assist sprayer is recommended

RESISTANCE

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field naturally occurring resistant biotypes may survive a correctly applied herbicide treatment propagate and become dominant in that field Adequate control of these resistant weed biotypes cannot be expected If weed control is unsatisfactory it may be necessary to retreat the problem area using a product affecting a different site of action

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage retreatment tank mix partners and/or sequential herbicide applications that have a different site of action Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes Consult your agricultural dealer consultant applicator and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological cultural and genetic practices aimed at preventing economic pest damage IPM principles and practices include field scouting or other detection methods correct target pest identification population monitoring and treating when target pest populations reach locally determined action thresholds Consult your state cooperative extension service professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area

RESTRICTIONS

- Do not apply drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots
- Do not use on lawns walks driveways tennis courts or similar areas Prevent drift of spray to desirable plants
- Do not graze treated fields or feed treated forage or hay Harvested straw may be used for bedding and/or feed
Do not harvest sooner than 45 days after the last application of this product
- Do not apply by air in the State of New York
- Do not apply to wheat barley triticale or oat crops underseeded with another crop
- This product is only registered on wheat barley oat triticale and fallow Do not use on any other crop
- Do not exceed the total rate of this product for wheat (including durum) barley and triticale of 1.0 ounce product per acre applied to any one crop during one growing season
- Do not exceed the total rate of this product for oat (spring and winter) of 0.4 ounces product per acre applied to any one crop during one growing season

PRECAUTIONS

- Take all necessary precautions to avoid all direct or indirect contact (such as spray drift) with non-target plants or areas
- Carefully observe all sprayer cleanup instructions both prior to and after using this product as spray tank residue may damage crops other than wheat barley triticale or oat
- Dry dusty field conditions may result in reduced control in wheel track areas
- Varieties of wheat (including durum) barley and triticale may differ in their response to various herbicides AXION AG PRODUCTS LLC recommends that you first consult your state experiment station university or extension agent as to sensitivity to any herbicide If no information is available limit the initial use to a small area
- Under certain conditions such as heavy rainfall prolonged cold weather or wide fluctuations in day/night temperatures prior to or soon after product application temporary discoloration and/or crop injury may occur To reduce the potential of crop injury tank mix this product with 2,4-D (ester formulations perform best—see Tank Mixtures) and apply after the crop is in the tillering stage of growth
- This product should not be applied to wheat barley triticale or oat that is stressed by severe weather conditions drought low fertility water saturated soil disease or insect damage as crop injury may result Risk of injury is greatest when crop is in the 2 to 5- leaf stage Severe winter stress drought disease or insect damage following application also may result in crop injury

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage or disposal

PESTICIDE STORAGE Store product in original container only Do not contaminate water other pesticides fertilizer food or feed in storage Store in a cool dry place

PESTICIDE DISPOSAL Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility

CONTAINER DISPOSAL

For Plastic Containers Nonrefillable container Do not reuse or refill this container Offer for recycling if available Triple rinse container (or equivalent) promptly after emptying

Triple rinse as follows Empty the remaining contents into application equipment or a mix tank Fill the container half full with water and recap Shake for 10 seconds Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure 2 more times

For Fiber Sacks Nonrefillable container Do not reuse or refill this container Completely empty sack by shaking and tapping sides and bottom to loosen clinging particles Empty residue into manufacturing or application equipment Then offer for recycling if available or dispose of sack in a sanitary landfill or by or inCineration or if allowed by State and local authorities by burning If burned stay out of smoke

For Fiber Drums with Liners Nonrefillable container Do not reuse or refill this container Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles Empty residue into manufacturing or application equipment Then offer for recycling if available or dispose of liner in a sanitary landfill or by or inCineration or if allowed by State and local authorities by burning If burned stay out of smoke

For Paper and Plastic Bags Nonrefillable container Do not reuse or refill this container Completely empty bag into application equipment Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or if allowed by State and local authorities by burning If burned stay out of smoke For minor spills leaks etc follow all precautions indicated on this label and clean up immediately

Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes In the event of a major spill fire or other emergency contact CHEMTREC 1-800-424 9300

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product If the terms are not acceptable return the product at once unopened and the purchase price will be refunded

The Directions for Use of this product must be followed carefully It is impossible to eliminate all risks inherently associated with the use of this product Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application weather presence of other materials or other influencing factors in the use of the product which are beyond the control of AXION AG PRODUCTS LLC or Seller To the extent consistent with applicable law all such risks shall be assumed by Buyer and User and Buyer and User agree to hold AXION AG PRODUCTS LLC and Seller harmless for any claims relating to such factors

AXION AG PRODUCTS LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use subject to the inherent risks referred to above when used in accordance with directions under normal use conditions This warranty does not extend to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or AXION AG PRODUCTS LLC and Buyer and User assume the risk of any such use **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW AXION AG PRODUCTS LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE**

To the extent consistent with applicable law neither AXION AG PRODUCTS LLC nor Seller shall be liable for any incidental consequential or special damages resulting from the use or handling of this product **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF AXION AG PRODUCTS LLC AND SELLER FOR ANY AND ALL CLAIMS LOSSES INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY CONTRACT NEGLIGENCE TORT STRICT LIABILITY OR OTHERWISE) RESULTING**

FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR AT THE ELECTION OF AXION AG PRODUCTS LLC OR SELLER THE REPLACEMENT OF THE PRODUCT

AXION AG PRODUCTS LLC and Seller offer this product and Buyer and User accept it subject to the foregoing conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of AXION AG PRODUCTS LLC

Credit Cutback Cutback M and Diablo are registered trademarks of Nufarm Americas Inc
Express Ally Assure STS and Glean are trademarks or registered trademarks of E I duPont de Nemours & Company
Fallow Master Landmaster and Maverick are registered trademarks of Monsanto

Assert and Diablo are registered trademarks of Micro Flo Company LLC

Clarity is a registered trademark of BASF Ag Products

Buctril Bronate Bronate Advanced Hoelon and Puma are registered trademarks of Bayer Crop Science

Starane Stinger and Widematch are registered trademarks of Dow AgroSciences LLC

Salvo and Sword are registered trademarks of Loveland Products Inc

Aim is a registered trademark of FMC Corp

Discover is a registered trademark of Syngenta Crop Protection Inc

Everest is a registered trademark of Arvesta Corp

Raindrop RA is a registered trademark of Delavan

Avenge is a registered trademark of Amvac Chemical Corporation

Rhino is a registered trademark of The McGregor Company