



2/15

GROUP 2 HERBICIDE

# AX SU 2525

## HERBICIDE

WATER DISPERSIBLE GRANULE  
FOR USE ON WHEAT, BARLEY AND FALLOW

**ACTIVE INGREDIENTS**

Thifensulfuron-methyl - Methyl 3 [[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]amino]sulfonyl]-2-thiophenecarboxylate	25 0%
Tribenuron-methyl - Methyl 2-[[[N-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)methylamino]carbonyl]amino]sulfonyl]benzoate	25 0%

**OTHER INGREDIENTS**

50 0%

**TOTAL**

100 0%

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
**SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS**

For Chemical Emergency, Spill, Leak, Fire, Exposure, or Accident  
 Call CHEMTREC Day or Night  
 Within USA and Canada 1-800-424 9300  
 Outside USA and Canada +1 703-527 3887 (collect calls accepted)

EPA Reg No 89167-18

EPA EST NO

NET CONTENTS \_\_\_\_\_ LBS

Manufactured For  
 AXION AG PRODUCTS LLC  
 3937 Cedarwood Lane  
 Johnstown CO 80534

092012

**ACCEPTED**

SEP 26 2012

Under the Federal Insecticide  
 Fungicide, and Rodenticide Act  
 as amended for the pesticide  
 registered under  
 EPA Reg No 89167-18

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Causes moderate eye irritation Avoid contact with eyes skin or clothing

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical resistant to this product are listed below If you want more options follow the instructions for Category A on an EPA chemical resistance category selection chart

Applicators and other handlers must wear

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as natural rubber
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE If no such instructions for washables exists use detergent and hot water Keep and wash PPE separately from other laundry

**USER SAFETY RECOMMENDATIONS**

**Users Should**

- Wash hands before eating drinking chewing gum using tobacco or using the toilet
- Remove clothing/PPE immediately if pesticide gets inside Then wash thoroughly and put on clean clothing
- Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing

**FIRST AID**

**IF IN EYE** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eye Call a poison control center or doctor for treatment advice

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center doctor or going for treatment For emergency information concerning this product call the National Pesticides Information Center (NPIC) at 1 800-858-7378 (NPIC Web site [www.npic.orst.edu](http://www.npic.orst.edu)) Or call your poison control center at 1-800-222-1222

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate

**PESTICIDE HANDLING**

- Calibrate sprayers only with clean water away from the well site
- Make scheduled checks of spray equipment
- Ensure that all operation employees accurately measure pesticides
- Mix only enough product for the job at hand
- Avoid overfilling of spray tank
- Do not discharge excess material on the soil at a single spot in the field grove or mixing/loading station
- Dilute and agitate excess solution and apply at labeled rates or uses
- Avoid storage of pesticides near well sites
- When triple rinsing the pesticide container be sure to add the rinsate to the spray mix

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling Do not apply this product in a way that will contact workers or other persons either directly or through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe consult the agency responsible for pesticide regulation

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170 This Standard contains requirements for the protection of agricultural workers on farms forests nurseries and greenhouses and handlers of agricultural pesticides It contains requirements for training decontamination notification and emergency assistance It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants soil or water is

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

This product is for use on wheat barley and fallow in many states Check with your state extension or Dept of Agriculture before use to be certain this product is registered in your state AXION AG PRODUCTSLLC will not be responsible for losses or damages resulting from the use of this product in any manner not in accordance with instructions on this label

**PRODUCT INFORMATION**

This product is a water dispersible granule that is used for selective postemergence weed control in wheat (including durum) barley and fallow The best control is obtained when this product is applied to young actively growing weeds The use rate will depend on weed spectrum and size of weed at time of application The degree and duration of control may depend on the following

- weed spectrum and infestation intensity
- weed size at application
- environmental conditions at and following treatment

This product is noncorrosive nonflammable nonvolatile and does not freeze This product should be mixed and completely dissolved in water and applied as a uniform broadcast spray

**ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY**

This product is absorbed primarily through the foliage of plants rapidly inhibiting the growth of susceptible weeds One to three weeks after application to weeds leaves of susceptible plants appear chlorotic and the growing point subsequently dies

This product provides the best control in vigorously growing crops that shade competitive weeds Weed control in areas of thin crop stand or seeding skips may not be as satisfactory However a crop canopy that is too dense at application can intercept spray and reduce weed control

The herbicidal action of this product may be affected in crops stressed from adverse environmental conditions (such as extreme temperatures or moisture) abnormal soil conditions cultural practices or variations in crop variety In warm moist conditions the expression of herbicide symptoms is accelerated in cold dry conditions expression of herbicide symptoms is delayed In addition weeds hardened off by drought stress are less susceptible to this product

**APPLICATION INFORMATION**

**USE RATE**

Apply this product at a rate of 0.4 to 1 ounce per acre When applying 0.4 to 0.6 ounce per acre this product must be used in a tank- mix combination with other registered herbicides

**WHEAT (INCLUDING DURUM) AND BARLEY**

Apply 0.4 to 1 ounce of this product per acre to wheat (including durum) or barley. The total amount of this product cannot exceed 1 ounce per acre per crop season.

**FALLOW**

Apply 0.4 to 1 ounce of this product per acre to fallow. The total amount of this product cannot exceed 1 ounce per acre per crop season. This product should be applied in combination with other suitable registered fallow herbicides such as glyphosate plus 2,4-D (ester formulations work best) or glyphosate plus dicamba.

When this product is applied at a rate of 0.4 to 0.6 ounce per acre, this product must be used in a tank mix combination with other registered fallow herbicides.

**PRE-PLANT BURNDOWN**

Apply 0.4 to 1 ounce of this product per acre as a burndown treatment prior to or shortly after planting (prior to emergence). The total amount of this product cannot exceed 1 ounce per acre per crop season. Apply up to 0.6 ounces per acre of this product as a pre-plant burndown to Cotton. Allow at least 14 days from time of application to time of planting cotton.

**POST HARVEST**

Apply this product at 0.4 to 1.0 ounce per acre to crop stubble after harvest. Use the 1.0 ounce per acre rate when weed infestation is heavy and predominantly consists of those weeds listed under the WEEDS PARTIALLY CONTROLLED section of this label or when application timing and environmental conditions are marginal. (See the APPLICATION TIMING section of this label for restriction on planting intervals). This product should be applied in combination with other suitable registered burndown herbicides. (See the TANK MIXTURES section of this label for additional information).

Sequential treatments of this product may also be made provided the total amount of this product applied during one fallow/pre plant cropland season does not exceed 1.0 ounce per acre.

**APPLICATION TIMING**

**WHEAT (INCLUDING DURUM) AND BARLEY** Make applications after the crop is in the 2-leaf stage but before the flag leaf is visible.

**FALLOW**

This product may be used as a fallow treatment in the spring, summer or fall when the majority of weeds have emerged and are actively growing.

**PRE-PLANT BURNDOWN**

Apply this product as a burndown treatment to wheat (including durum) and barley to control emerged weeds prior to or shortly after planting (prior to emergence). Make applications when the majority of weeds have emerged and are actively growing. Wheat and Barley may be replanted anytime after the application of this product.

Allow at least 14 days between application of this product and planting of cotton. Allow at least 60 days between application of this product and planting of sugar beets, winter rape and canola. Allow at least 45 days between application of this product and planting of any other crop (such as corn, rice, grain sorghum or soybeans).

**POST HARVEST**

This product may be used as a burndown treatment to crop stubble when the majority of weeds have emerged and are actively growing. (See the CROP ROTATION section of this label for additional information).

Since this product has very little or no soil activity, it controls only those weeds that have germinated. Therefore, apply this product when all or most of the weeds have germinated. Annual broadleaf weeds should be past the cotyledon stage, actively growing, and less than 4" tall or wide. Rainfall immediately

after treatment can wash this product off of weed foliage resulting in reduced weed control Several hours of dry weather are needed to allow this product to be sufficiently absorbed by weed foliage

**CROP ROTATION**

Wheat (including durum) and barley may be replanted anytime after the application of this product

Cotton can be planted 14 days after the application of this product

Sugar beets Winter Rape and Canola can be planted at 60 days after the application of this product Any other crop may be planted 45 days after the application of this product

**WEEDS CONTROLLED**

This product effectively controls the following weeds when used according to label directions

Annual knawel	Common sunflower	London rocket	Stimleaf lambsquarters
Annual sowthistle	Corn chamomile	Marshelder	Smallf lower buttercup
Black mustard	Corn gromwell *	Mayweed chamomile	Smallseed falsef lax
Blue/Purple mustard	Corn spurry	Miners lettuce	Stinking chickweed
Broadleaf dock	Cowcockle	Narrowleaf	Stinking mayweed/
Bur buttercup	Cress (mouse-ear)	Nightflowering catchfly	Doofennel
Bushy wallflower/	Curly dock	Pennsylvania	Sunflower
Treacle mustard	False chamomile	Pineappleweed	Swinecress
Canada thistle *	Field chickweed	Prickly lettuce*	Tansymustard
Clasping pepperweed	Field pennycress	Prostrate knotweed	Tarweed fiddleneck
Coast fiddleneck	Filaree (redstem)	Prostrate pigweed	Tumble/Jim Hill mustard
Common buckwheat	Flixweed	Redmaids	Volunteer canola
Common chickweed	Green smartweed	Redroot pigweed	Volunteer lentils
Common cocklebur *	Henbit	Russian thistle*	Volunteer peas
Common groundsel	Kochia *	Scentless chamomile/	Wild buckwheat*
Common	Ladvstthumb	mayweed	Wild chamomile
Common ragweed *	Lanceleaf sage *	Shepherd s-purse	Wild mustard

**WEEDS PARTIALLY CONTROLLED\*\***

This product partially controls the following weeds when used according to label directions

Catchweed bedstraw Mallow (common little) Marestail Nightshade (cutleaf hairy)

\*See SPECIFIC WEED PROBLEMS for more information

\*\* Partial control A visual reduction of weed population as well as a significant loss of vigor For better results use the highest rate of this product per acre and include a tank mix partner such as 2 4-D MCPA Buctril or Banvel/Clarity (refer to TANK MIXTURES)

**SPECIFIC WEED PROBLEMS**

**Canada thistle** For control in wheat and barley use 0.8 ounce per acre plus surfactant when all thistles are 4 to 8 with 2 to 6 of new growth Make the application in the spring Control will be improved by using this product in combination with 2 4 D or dicamba (refer to TANK MIXTURES)

**Common cocklebur, Common ragweed, Lanceleaf sage** In wheat and barley apply this product at 0.4 to 0.8 ounce per acre in combination with 2 4-D at rates from 1/4 to 3/8 pound active ingredient (ester formulations work best) when weeds are small and actively growing When using 1/4 pound active ingredient of 2 4 D be sure to add surfactant at the rate of 1/4 to 1/2 quart per 100 gallons of spray solution (0.06 to 0.125% v/v--use the higher rate under stress conditions)

**Corn gromwell, Wild buckwheat** For control in wheat and barley use 0.8 ounce this product per acre plus surfactant

**Kochia, Russian thistle, Prickly lettuce** Naturally occurring resistant biotypes of these weeds are known to occur For best results use this product in a tank mix with Starane Starane + Sword Starane + Salvo dicamba (such as Banvel/Clarity) and 2 4-D or Bromoxynil (such as Buctril) and 2 4 D (3/4 - 1 pt Buctril + 1/4 3/8 pound active ingredient 2 4-D ester) This product should be applied in the spring when weeds are 2 to 4 tall or 2 to 4 across and are actively growing Refer to the Tank Mixtures section of this label for additional details

**SPRAY ADJUVANTS**

Always include a spray adjuvant with applications of this product. In addition to a spray adjuvant, an ammonium nitrogen fertilizer may be used.

Consult your Ag dealer or applicator, local AXION AG PRODUCTS LLC fact sheets, technical bulletins, and service policies prior to using an adjuvant system. If another herbicide is tank mixed with this product, select adjuvants authorized for use with both products. Products must contain only EPA-exempt ingredients.

**NONIONIC SURFACTANT (NIS)**

- Apply 0.06 to 0.50% volume/volume (1/2 to 4 pints per 100 gallon of spray solution). Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12. See the Tank Mixtures section of this label for additional information.

**PETROLEUM CROP OIL CONCENTRATE (COC) OR MODIFIED SEED OIL (MSO)**

Apply at 1% volume/volume (1 gallon per 100 gallon spray solution) or 2% volume/volume under arid conditions.

- Oil adjuvants must contain at least 80% high quality petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

**SPECIAL ADJUVANT TYPES**

- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO, and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions.
- In addition to the adjuvants specified above, other adjuvant types may be used if they provide the same functionality and have been evaluated and approved by AXION AG PRODUCTS LLC product management. Consult separate AXION AG PRODUCTS LLC technical bulletins for detailed information before using adjuvant types not specified on this label.

**AMMONIUM NITROGEN FERTILIZER**

- Use 2 qt /acre of a high-quality urea ammonium nitrate (UAN) such as 28%N or 32%N or 2 lb /acre of a spray grade ammonium sulfate (AMS). Use 4 qt /acre UAN or 4 lb /acre AMS under arid conditions.

**GROUND APPLICATION**

For optimum spray distribution and thorough coverage, use flat-fan or low volume flood nozzles. For flat-fan nozzles, use a spray volume of at least 5 gallon per acre (GPA). For flood nozzles on 30" spacings, use at least 10 GPA. Flood nozzles no larger than TK10 (or the equivalent) and a pressure of at least 30 psi. For 40" nozzle spacings, use at least 13 GPA. For 60" spacings, use at least 20 GPA. It is essential to overlap the nozzles 100% for all spacings. Raindrop RA nozzles are not recommended for this product applications, as weed control performance may be reduced. Use screens that are 50 mesh or larger.

**AERIAL APPLICATION**

Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage.

- Use 2 to 5 GPA.
- Use at least 3 GPA in Idaho, Oregon, or Utah.

Do not apply this product by air in the state of New York. See the **Spray Drift Management** section of this label.

**CHEMIGATION**

Do not apply this product through any irrigation system.

**PRODUCT MEASUREMENT**

This product is measured using the this product volumetric measuring cylinder The degree of accuracy of this cylinder varies by +/- 7 5% For more precise measurement use scales calibrated in ounces

**TANK MIXTURES**

This product may be tank mixed with other suitable registered herbicides to control weeds listed as suppressed weeds resistant to this product or weeds not listed under **Weeds Controlled** Read and follow all manufacturer s label instructions for the companion herbicide If those instructions conflict with this label do not tank mix the herbicide with this product

This product can also be mixed with registered fungicides insecticides or liquid fertilizer for use on wheat barley or fallow

**WITH 2,4 D (AMINE OR ESTER) OR MCPA (AMINE OR ESTER)**

This product may be tank mixed with the amine or ester formulations of 2 4-D or MCPA herbicides for use on wheat and barley For best results in the Red River Valley and adjacent areas of North Dakota and Minnesota add the ester formulations of 2 4-D or MCPA herbicides to the tank at 3/8 pound active ingredient (such as 3/4 pt of a 4 lb/gal product or 1/2 pt of a 6 lb/gal product) No additional surfactant is needed with this mixture

For best results in other areas add the ester formulations of 2 4-D or MCPA herbicides to the tank at 1/4 to 3/8 pound active ingredient (such as 1/2 3/4 pt of a 4 lb/gal product or 1/3-1/2 pt of a 6 lb/gal product) Surfactant may be added to the mixture at 1/2 to 1 quart per 100 gallon of spray solution (0 125 to 0 25% v/v) however adding surfactant may increase the potential for crop injury especially at the higher phenoxy rates

Higher rates of 2 4 D or MCPA may be used but do not exceed the highest rate allowed by those respective labels Read and follow all label instructions on timing precautions and warnings for these herbicides before using these tank mixtures

**WITH DICAMBA**

This product may be tank mixed with 1/16 to 1/8 pound active ingredient dicamba (such as 2-4 fluid oz Banvel or 2-4 fluid oz Clarity)

Use higher rates when weed infestation is heavy Surfactant may be added to the mixture at 1/2 to 1 quart per 100 gallon of spray solution (0 125 to 0 25% v/v) however adding surfactant may increase the potential for crop injury Refer to the specific dicamba label for application timing and restrictions Tank mixes of this product plus dicamba may result in reduced control of some broadleaf weeds

**WITH 2,4 D (AMINE OR ESTER) AND DICAMBA**

This product may be applied in a 3-way tank mix with formulations of dicamba and 2 4-D Make application of this product + 1/16 to 1/8 pound active ingredient dicamba (such as 2 4 fluid oz Banvel or 2-4 fluid oz Clarity) + 1/4 3/8 pound active ingredient 2 4 D ester or amine per acre Use higher rates when weed infestation is heavy Surfactant may be added to the mixture at 1/2 to 1 quart per 100 gallon of spray solution (0 125 to 0 25% v/v) however adding surfactant may increase the potential for crop injury Consult the specific 2 4-D label dicamba label or local recommendations for more information and restrictions

Apply this 3-way combination to winter wheat after the crop is tillering and prior to jointing (first node) In Spring Wheat (including Durum) apply after the crop is tillering and before it exceeds the 5-leaf stage In Spring Barley apply after the crop is tillering and before it exceeds the 4-leaf stage

**WITH BROMOXYNIL**

This product may be tank mixed with bromoxynil containing herbicides registered for use on wheat barley or fallow For best results add bromoxynil containing herbicides to the tank at 3/16 to 3/8 pound active ingredient per acre (such as Bronate or Buctril at 3/4 to 1 1/2 pint per acre)

Read and follow all label instructions on timing precautions and warnings for these herbicides before using these tank mixtures Follow the most restrictive labeling Tank mixes of this product plus Buctril may result in reduced control of Canada thistle

**WITH STARANE, STARANE + SWORD, STARANE + SALVO**

For improved control of Kochia (2-4 tall) Russian thistle mustard species and wild buckwheat this product may be tank mixed with 1/3 to 1-1/3 pints per acre of Starane 2/3 to 2-2/3 pints per acre of

Starane + Salvo or 3/4 to 2- 3/4 pints per acre of Starane + Sword Additional 2 4-D or MCPA can be added based on local recommendations (refer to 2 4 D and MCPA labels for maximum amount that can be applied to the crop) Refer to the Starane Starane + Salvo or Starane + Sword label for information regarding use restrictions labeled crops rotational cropping recommendations sprayer cleanup use precautions and other information The most restrictive provisions on any label will apply Do not use the tank mix if any restrictions on the labels conflict with instructions on this product label

**WITH HOELON HERBICIDE**

This product may be used in combination with Hoelon 3EC and Buctril herbicides in accordance with the Hoelon 3EC label For best results use the three-way tank mix of this product at 0 4 ounce per acre plus Hoelon 3EC at 2 2/3 pint per acre plus Buctril at 1-1/2 pints per acre Apply only to winter wheat This tank mix should only be used under good soil conditions when wild oat is in the 1 4 leaf stage If conditions are not ideal for the performance of Hoelon 3EC wild oat control may be reduced Be sure to follow all warnings and cautions on the Hoelon 3EC and Buctril labels

**WITH ASSERT HERBICIDE**

This product can be tank mixed with Assert When tank mixing this product with Assert always include another broadleaf weed herbicide with a different mode of action (for example 2 4-D ester MCPA ester Buctril or Bronate) Tank-mixed applications of this product plus Assert may cause temporary crop discoloration stunting or injury when heavy rainfall occurs shortly after application

**WITH OTHER GRASS CONTROL PRODUCTS**

Tank mixtures of this product and grass control products may result in poor grass control AXION AG PRODUCTSLLC recommends that you first consult your state experiment station university or extension agent Agricultural dealer or AXION AG PRODUCTSLLC representative as to the potential for antagonism before using the mixture If no information is available limit the initial use of this product and the grass product to a small area Do not tank mix with Achieve herbicide

**WITH INSECTICIDES OR FUNGICIDES**

This product may be tank mixed or used sequentially with insecticides (or fungicides) registered for use on cereal grains However under certain conditions (drought stress or if the crop is in the 2 4 leaf stage) tank mixes or sequential applications of this product with organophosphate insecticides (such as parathion) may produce temporary crop yellowing or in severe cases crop injury Test these mixtures in a small area before treating large areas However review all insecticide and fungicide labels for restrictions Do not use this product plus Malathion as crop injury will result

**WITH LIQUID NITROGEN SOLUTION FERTILIZER**

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water Run a tank mix compatibility test before mixing this product in fertilizer solution Do not add this product directly to liquid nitrogen fertilizer the granules will not dissolve This product must be thoroughly mixed with clean water before it is added to liquid nitrogen fertilizer If granules remain when the mixture is poured out add more clean water and mix until all granules have disappeared Ensure that the agitator is running when the this product premix is added Use of this mixture may result in temporary crop yellowing and stunting If using low rates of liquid nitrogen fertilizer in the spray solution (less than 50% of the spray solution volume) the addition of surfactant is necessary Add surfactant at 1/4 - 1 quart per 100 gallon of spray solution (0 06 - 0 25% v/v) based on local recommendations When using high rates of liquid nitrogen fertilizer solution in the spray solution adding surfactant increases the risk of crop injury Consult your agricultural dealer consultant fieldman or AXION AG PRODUCTSLLC representative for a specific recommendation before adding an adjuvant to these tank mixtures If 2 4-D or MCPA is included with this product and fertilizer mixture ester formulations tend to be more compatible (see manufacturer s label) Additional surfactant is not needed when using this product in tank mix with 2 4 D ester or MCPA ester and liquid nitrogen fertilizer solutions Do not use low rates of liquid nitrogen fertilizer solution as a substitute for a surfactant Do not use with liquid fertilizer solutions with a pH less than 3 0

**TANK MIXTURES IN FALLOW**

This product may be used as a fallow treatment and should be tank mixed with other herbicides that are

registered for use in fallow Read and follow all manufacturer s label instructions for the companion herbicide If those instructions conflict with this label do not tank mix the herbicide with this product

**TANK MIXTURES IN PRE-PLANT BURNDOWN**

This product may be used as a pre plant burndown treatment alone or tank mixed with other herbicides that are registered for use as a pre-plant burndown product such as glyphosate plus 2 4-D (ester formulations work best) or glyphosate plus dicamba

Read and follow all manufacturer s label instructions for the companion herbicide If those instructions conflict with this label follow the most restrictive labeling (such as planting interval after application) or do not tank mix the herbicide with this product

**TANK MIXTURES IN POST HARVEST APPLICATIONS**

This product may be used as a post harvest treatment to crop stubble and should be tank mixed with other herbicides that are registered for use in fallow

**MIXING INSTRUCTIONS**

Do not use with spray additives that alter the pH of the spray solution below pH 5 0 or above pH 9 0 as rapid product degradation can occur Spray solutions of pH 6 0-8 0 allow for optimum stability of this product

- 1 Fill the tank 1/4 to 1/3 full of water
- 2 While agitating add the required amount of this product
- 3 Continue agitation until the this product is fully dissolved at least 5 minutes
- 4 Once the this product is fully dissolved maintain agitation and continue filling tank with water
- 5 As the tank is filling add tank mix partners and then add the required volume of spray adjuvant Always add spray adjuvant last Antifoaming agents may be used Do not use with spray additives that alter the pH of the spray solution below pH 6 0 as rapid product degradation can occur Spray solutions of pH 7 0 and higher allow for optimum stability of this product
- 6 Dispersed tank mix partners can settle if the tank mixture is not continually agitated If settling occurs thoroughly re-agitate before using
- 7 Apply this product spray mixture within 24 hours of mixing to avoid product degradation
- 8 If this product and a tank mix partner are to be applied in multiple loads fully dissolve this product in clean water prior to adding to the tank

**GRAZING**

Do not graze livestock in treated areas In addition do not feed forage or hay from treated areas to livestock (straw harvested after grain harvest may be used for bedding and/or feed)

**SPRAY EQUIPMENT**

For specific application equipment refer to the manufacturer's recommendations for additional information on GPA pressure speed nozzle types and arrangements nozzle heights above the target canopy etc Be sure to calibrate air or ground equipment properly before application Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern with minimum drift Use higher spray volumes to obtain better coverage when crop canopy is dense Avoid swath overlapping and shut off spray booms while starting turning slowing or stopping to avoid injury to the crop

Do not make applications using equipment and/or spray volumes or during weather conditions that might cause spray to drift onto nontarget sites For additional information on spray drift refer to Spray Drift Management section of label

Continuous agitation may be required to keep this product and tank-mix partners in solution or suspension Refer to tank-mix partner labels for additional information

**SPRAYER CLEANUP**

The spray equipment must be cleaned before this product is sprayed Follow the cleanup procedures specified on the labels of the previously applied products If no directions are provided follow the six steps outlined in After Spraying in this label

**AT THE END OF THE DAY**

It is recommended that during periods when multiple loads of this product are applied at the end of each day of spraying the interior of the tank be rinsed with fresh water and then partially filled and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits which can accumulate in the application equipment.

**AFTER SPRAYING AND BEFORE SPRAYING CROPS OTHER THAN WHEAT AND BARLEY**

To avoid subsequent injury to desirable crops thoroughly clean all mixing and spray equipment immediately following applications of this product as follows:

- 1 Drain tank thoroughly rinse spray tanks boom and hoses with clean water. Loosen and physically remove any visible deposits.
- 2 Fill the tank with clean water and 1 gallon of household ammonia\*(contains 3% active) for every 100 gal of water. Flush the hoses boom and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 minutes. Flush the hoses boom and nozzles again with the cleaning solution and then drain the tank.
- 3 Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
- 4 Repeat Step 2.
- 5 Rinse the tank boom and hoses with clean water.
- 6 If only Ammonia is used as a cleaner the rinsate solution may be applied back to the crop(s) listed on this label. Do not exceed the maximum labeled use rate. If other cleaners are used consult the cleaner label for rinsate disposal instructions. If no instructions are given dispose of the rinsate on site or at an approved waste disposal facility.

\* Equivalent amounts of an alternate-strength ammonia solution can be used in the cleanout procedure. Carefully read and follow the individual cleaner instructions. Consult your Ag dealer for a listing of approved cleaners.

**Notes**

- 1 CAUTION Do not use chlorine bleach with ammonia as dangerous gases will form. Do not clean equipment in an enclosed area.
- 2 Steam-cleaning aerial spray tanks is recommended prior to performing the above cleanout procedure to facilitate the removal of any caked deposits.
- 3 When this product is tank mixed with other pesticides all cleanout procedures should be examined and the most rigorous procedure should be followed.
- 4 In addition to this cleanout procedure all pre-cleanout guidelines on subsequently applied products should be followed as per the individual labels.
- 5 Where routine spraying practices include shared equipment frequently being switched between applications of this product and applications of other pesticides to sensitive crops during the same spray season it is recommended that a sprayer be dedicated to this product to further reduce the chance of crop injury.

**SPRAY DRIFT MANAGEMENT**

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

**IMPORTANCE OF DROPLET SIZE**

**AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR**

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **Wind, Temperature and Humidity** and

Temperature Inversions sections of this label

**CONTROLLING DROPLET SIZE - GENERAL TECHNIQUES**

- **Volume** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types narrower spray angles produce larger droplets. Consider using low-drift nozzles

**CONTROLLING DROPLET SIZE AIRCRAFT**

- **Number of Nozzles** Use the minimum number of nozzles with the highest flow rate that provide uniform coverage
- **Nozzle Orientation** Orienting nozzles so that the spray is emitted backwards parallel to the airstream will produce larger droplets than other orientations
- **Nozzle Type** Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types
- **Boom Length** The boom length should not exceed 3/4 of the wing or rotor length - longer booms increase drift potential
- **Application Height** - Application more than 10 feet above the canopy increases the potential for spray drift

**BOOM HEIGHT**

Setting the boom at the lowest referenced height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment the boom should remain level with the crop and have minimal bounce

**WIND**

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However many factors including droplet size and equipment type determine drift potential at any given wind speed. **AVOID GUSTY AND WINDLESS CONDITIONS**

**Note** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift

**TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions set up equipment to produce larger droplets to reduce effects of evaporation

**TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog. However if fog is not present inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion while smoke that moves upward and rapidly dissipates indicates good vertical air mixing

**SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce the effects of wind. However it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product

**AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS**

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift but if a sprayer is unsuitable for the application and/or set up improperly high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application is configured properly and that drift is not occurring

**Note** Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the spray equipment section of this label to determine if use of an air assist sprayer is recommended.

**RESISTANCE**

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons, such as using a combination of tillage, retreatment, tank mix partners, and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

**INTEGRATED PEST MANAGEMENT**

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants, or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

**RESTRICTIONS**

- Do not apply, drain, or flush equipment on or near desirable trees or other plants or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants.
- Do not harvest sooner than 45 days after the last application of this product. Do not graze treated fields or feed treated forage or hay. Harvested straw may be used for bedding and/or feed.
- Do not apply by air in the State of New York.
- Do not apply to wheat or barley crops underseeded with another crop.
- Do not exceed the maximum application rate of 1 ounce per acre per crop season.

**PRECAUTIONS**

- Take all necessary precautions to avoid all direct or indirect contact (such as spray drift) with non-target plants or areas.
- Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than wheat or barley.
- Dry, dusty field conditions may result in reduced control in wheel track areas.
- Wheat and barley may differ in their response to various herbicides. AXION AG PRODUCTS LLC recommends that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use of this product to a small area.

Under certain conditions, such as heavy rainfall, prolonged cold weather (daily high temperature less

than 50°F) or wide fluctuations in day/night temperatures prior to or soon after this product's application temporary discoloration and/or crop injury may occur To reduce the potential of crop injury tank mix this product with 2 4-D (ester formulations perform best—see Tank Mixtures section of this label) and apply after the crop is in the tillering stage of growth

- This product should not be applied to wheat and barley that is stressed by severe weather conditions drought (including low levels of subsoil moisture) low fertility water saturated soil disease or insect damage as crop injury may result Risk of injury is greatest when crop is in the 2 to 5 leaf stage Severe winter stress drought disease or insect damage following application also may result in crop injury

**STORAGE AND DISPOSAL**

Do not contaminate water food or feed by storage or disposal

**PESTICIDE STORAGE** Store product in original container only Store in a cool dry place

**PESTICIDE DISPOSAL** Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility

**CONTAINER DISPOSAL For Plastic Containers** Nonrefillable container Do not reuse or refill this container Triple rinse container (or equivalent) promptly after emptying Triple rinse as follows Empty the remaining contents into application equipment or a mix tank Fill the container 1/4 full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after flow begins to drip Repeat this procedure two more times Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration or if allowed by State and local authorities by burning If burned stay out of smoke

**For Fiber Sacks** Nonrefillable container Do not reuse or refill this container Completely empty fiber sack by shaking and tapping sides and bottom to loosen clinging particles Empty residue into manufacturing or application equipment Then dispose of sack in a sanitary landfill or by incineration or if allowed by State and local authorities by burning If burned stay out of smoke

**For Paper and Plastic Bags** Nonrefillable container Do not reuse or refill this container Completely empty bag into application equipment Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or if allowed by State and local authorities by burning If burned stay out of smoke

For minor spills leaks etc follow all precautions indicated on this label and clean up immediately Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes In the event of a major spill fire or other emergency contact CHEMTREC 1-800-424-9300

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

NOTICE Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product If the terms are not acceptable return the product at once unopened and the purchase price will be refunded The Directions for Use of this product must be followed carefully It is impossible to eliminate all risks inherently associated with the use of this product Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application weather presence of other materials or other influencing factors in the use of the product which are beyond the control of AXION AG PRODUCTSLLC or Seller To the extent consistent with applicable law all such risks shall be assumed by Buyer and User and Buyer and User agree to hold AXION AG PRODUCTSLLC and Seller harmless for any claims relating to such factors AXION AG PRODUCTSLLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use subject to the inherent risks referred to above when used in accordance with directions under normal use conditions This warranty does not extend to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or AXION AG PRODUCTSLLC and Buyer and User assume the risk of any such use TO THE EXTENT CONSISTENT WITH APPLICABLE LAW AXION AG PRODUCTSLLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE

To the extent consistent with applicable law neither AXION AG PRODUCTSLLC nor Seller shall be liable for any incidental consequential or special damages resulting from the use or handling of this product TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF AXION AG PRODUCTSLLC AND SELLER FOR ANY AND ALL CLAIMS LOSSES INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY CONTRACT NEGLIGENCE TORT STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR AT THE ELECTION OF ARCANA LLC OR SELLER THE REPLACEMENT OF THE PRODUCT

AXION AG PRODUCTSLLC and Seller offer this product and Buyer and User accept it subject to the foregoing conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of AXION AG PRODUCTSLLC

Ally® Express® Finesse® and Glean® are registered trademarks of E I DuPont de Nemours and Company  
Fallow™ Master™ Landmaster™ Roundup™ are trademarks of Monsanto Co  
Assert™ Banvel™ are registered trademarks of MicroFlo Company  
Clarity™ is a registered trademark of BASF  
Bronate™ Buctrij™ and Hoelon™ are trademarks of Bayer CropScience  
Raindrop RA® is a registered trademark of Delavan  
Salvo™ and Sword™ are trademarks of Loveland Products Inc  
Starane™ is a trademark of Dow AgroSciences LLC  
Achieve® is a registered trademark of Syngenta Participations AG