

89167-40

6/26/2014

1/61



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (7505P)  
1200 Pennsylvania Ave., NW  
Washington, D.C. 20460

EPA Registration  
Number:

89167-40

Date of Issuance:

JUN 26 2014

NOTICE OF PESTICIDE:

Registration  
 Reregistration

(under FIFRA, as amended)

Term of Issuance: Un-Conditional

Name of Pesticide Product:

AX S-MET Herbicide

Name and Address of Registrant (include ZIP Code):

Axion Ag Products, LLC  
1966 W. 15th Street, Suite 6  
Loveland, CO 80539

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA §3(c)(5)

The basic confidential statement of formula (CSF) dated October 09, 2013 is acceptable.

A stamped copy of the label is enclosed for your records. Submit one (1) copy of the final printed label before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions regarding this Notice, please contact Shanta Adeeb at (703) 347-0502 or at Adeeb.shanta@epa.gov.

Signature of Approving Official:

Kathryn Montague  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)

Date:

JUN 26 2014

2/61

**Note:** It is illegal to sell, use or distribute this product within, or into, Nassau County or Suffolk County, New York.

# AX S-MET Herbicide

For weed control in corn; cotton; grasses grown for seed; horseradish; peanuts; beans, peas and lentils; potatoes; pumpkin; rhubarb; safflowers; sugar beets; sunflowers; sweet, grain or forage sorghum; soybean; soybean, immature seed; and tomatoes

Active Ingredient:

|  |        |
|--|--------|
| S-metolachlor (CAS No. 87392-12-9) ..... | 83.7%  |
| Other Ingredients:                       | 16.3%  |
| Total:                                   | 100.0% |

AX S-MET contains 7.62 lbs. of active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 89167-


EPA Est.:

Product of Switzerland - Formulated in the USA

Net Contents: \_\_\_ Gallons (\_\_\_ L)

MANUFACTURED FOR:  
Axion Ag Products, LLC  
1966 W 15<sup>th</sup> Street, Suite 6  
Loveland, CO 80538

061714



**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**CAUTION**

Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin, or clothing. May cause skin sensitization reactions in certain individuals.

| <b>FIRST AID</b>   |   |
|--|---|
| <b>If in eyes</b>  | <ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>  |
| <b>If on skin or clothing</b>  | <ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>   |
| <b>If swallowed</b>  | <ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Do not give any liquid to the person.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul> |
| <b>If inhaled</b>  | <ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>  |
| <p><b>Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.</b></p>   |   |
| <p><b>HOT LINE NUMBER</b></p> <p>For 24 Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident), Call <b>1-800-888-8372</b></p> |   |

**Personal Protective Equipment (PPE)**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or Viton®
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Control Statements**

Mixers and loaders supporting aerial applications are required to use closed systems. The closed system must be used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)]. When using the closed system, the mixers' and loaders' PPE requirements may be reduced or modified as specified in the WPS.

4/  
61

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### **User Safety Recommendations**

#### **Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **Environmental Hazards**

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

### **Ground Water Advisory**

The active ingredient in AX S-MET has the potential to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

### **Surface Water Advisory**

The active ingredient in AX S-MET has the potential to contaminate surface water through ground spray drift. Under some conditions, the active ingredient may also have a high potential for runoff into surface water (primarily via dissolution in runoff water) for several months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

### **Mixing/Loading Instructions**

Care must be taken when using this product to prevent back-siphoning into wells, spills, or improper disposal of excess pesticide, spray mixtures, or rinsates.

Check-valves or antisiphoning devices must be used on all mixing and/or irrigation equipment.

This product may not be mixed or loaded within 50 ft. of perennial or intermittent streams and rivers, natural or impounded lakes and reservoirs. This product may not be mixed/loaded or used within 50 ft. of all wells, including abandoned wells, drainage wells, and sink holes. Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 ft. of any well are prohibited, unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rain water that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all

times. The above-specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

AX S-MET must be used only as described on this label or in separately published EPA accepted supplemental labeling for this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.** Exception: If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or Viton
- Shoes plus socks

#### FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result.

**Restriction: Not for sale, use or distribution in Nassau County or Suffolk County, New York**

#### PRODUCT INFORMATION

Observe all precautions and limitations on the labels of each product used in tank mixtures. Tank mixtures are permitted only in those states where the tank mix partner is registered. Refer to and follow the label for each tank mix product used for precautionary statements, directions for use, geographic and other restrictions.

AX S-MET is a selective herbicide that can be applied as a Preplant surface-applied, preplant incorporated, or preemergence treatment in water or fluid fertilizer for control of most annual grasses and certain broadleaf weeds in corn (all types); cotton; grasses grown for seed; peanuts;

6/61

beans, peas and lentils; potatoes; safflowers; sugar beets; sunflowers; grain or forage sorghum; soybeans; soybean, immature seed; and tomatoes.

- Do not use in nurseries, turf, or landscape plantings.
- Do not apply under conditions which favor runoff or wind erosion of soil containing this product to nontarget areas.

To prevent off-site movement due to runoff or wind erosion:

- Avoid treating powdery dry or light sand soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.
- Do not apply to impervious substrates, such as paved or highly compacted surfaces.
- Do not use tailwater from the first flood or furrow irrigation of treated fields to treat nontarget crops, unless at least 1/2 inch of rainfall has occurred between application and the first irrigation.

Where directions specify a AX S-MET tank mixture with AAtrex® formulations, other brands of atrazine may be used. Follow the rates, recommendations, and limitations on the AAtrex or respective atrazine product label if other brands of atrazine are used.

Certain states may have established rate limitations for atrazine within specific geographical areas. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.

If AX S-MET is incorporated, any supplemental tillage before planting must not exceed the depth of incorporation.

Dry weather following preemergence application of AX S-MET or a tank mixture may reduce effectiveness. Cultivate if weeds develop.

Where reference is made to weeds partially controlled, partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial weed control.

**Precaution:** Crop injury may occur following the use of AX S-MET under abnormally high soil moisture conditions during early development of the crop.

**SOIL TEXTURES AND HERBICIDE RATES**

Where rates are based on coarse-, medium-, or fine-textured soils, it is understood that soil textural classes are generally categorized as follows:

| <b>Coarse</b> | <b>Medium</b> | <b>Fine</b>     |
|---------------|---------------|-----------------|
| Sand          | Loam          | Sandy clay loam |
| Loamy sand    | Silt loam     | Silty clay loam |
| Sandy loam    | Silt          | Clay loam       |
| Sandy clay    |               |                 |
| Silty clay    |               |                 |
|               |               | Clay            |

Within rate ranges in the rate tables and elsewhere on this label, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter.

AX S-MET may be applied preemergence alone, or in combination with tank mix partners specified on this label, following preplant incorporated herbicides when used according to their label requirements, provided that such use is not prohibited on the respective labels.

7/61

Thoroughly clean sprayer or other application device before using. Dispose of cleaning solution in a responsible manner. Do not use a sprayer or applicator contaminated with any other materials, or crop damage or clogging of the application device may result.

### **AX S-MET APPLIED ALONE WEEDS CONTROLLED**

AX S-MET is taken up by the shoots and/or roots of emerging weeds. This uptake results in the inhibition of shoot and root tissue growth soon after weed germination. Because of this, AX S-MET will not control emerged weeds and should be applied prior to weed emergence.

If AX S-MET is incorporated, do not exceed a 2-3 inch depth. Any tillage after the AX S-MET incorporation and before planting should not exceed 2-3 inches.

Dry weather following application of AX S-MET may reduce weed control. Cultivate if weeds develop.

Where reference is made to weeds partially controlled, partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial weed control. Control of these weeds can be erratic, due partially to variable weather conditions. The following procedures may improve the control of weeds listed as partially controlled in Table 1:

- Thoroughly till soil to destroy germinating and emerged weeds.
- Plant crop into moist soil immediately after tillage. If AX S-MET is to be used preemergence, apply at planting or immediately after planting.
- If available, sprinkler irrigate within 2 days after application. Apply ½ -1 inch of water. Use lower water volume (1/2 inch) on coarse-textured soils and higher volume (1 inch) on fine-textured soils. Also, refer to the section on **Center Pivot Irrigation Application** for this method of applying AX S-MET.
- If irrigation is not possible and rain does not occur within 2 days after planting and application, weed control may be decreased. Under these conditions, make a uniform, shallow cultivation as soon as weeds emerge.

8/61

**Table 1: Weeds Controlled or Partially Controlled by AX S-MET Applied Prior to Weed Emergence**  
**PART 1**

| Common Name             | Scientific Name                 | Weed Type | Control (C) or Partial Control (PC) |
|-------------------------|---------------------------------|-----------|-------------------------------------|
| Barnyardgrass           | <i>Echinochloa crus-galli</i>   | Grass     | C                                   |
| Crabgrass, large        | <i>Digitaria ischaemum</i>      | Grass     | C                                   |
| Crabgrass, smooth       | <i>Digitaria sanguinalis</i>    | Grass     | C                                   |
| Crowfootgrass           | <i>Dactyloctenium aegyptium</i> | Grass     | C                                   |
| Cupgrass, Southwestern  | <i>Eriochloa acuminata</i>      | Grass     | C                                   |
| Cupgrass, woolly        | <i>Eriochloa villosa</i>        | Grass     | PC <sup>1</sup>                     |
| Foxtail, bristly        | <i>Setaria verticillata</i>     | Grass     | C                                   |
| Foxtail, giant          | <i>Setaria faberi</i>           | Grass     | C                                   |
| Foxtail, green          | <i>Setaria viridis</i>          | Grass     | C                                   |
| Foxtail, millet         | <i>Setaria italica</i>          | Grass     | C                                   |
| Foxtail, yellow         | <i>Setaria pumila</i>           | Grass     | C                                   |
| Goosegrass              | <i>Eleusine indica</i>          | Grass     | C                                   |
| Johnsongrass (seedling) | <i>Sorghum halepense</i>        | Grass     | PC                                  |
| Millet, wild-proso      | <i>Panicum miliaceum</i>        | Grass     | PC <sup>1</sup>                     |
| Panicum, fall           | <i>Panicum dichotomiflorum</i>  | Grass     | C                                   |
| Panicum, Texas          | <i>Panicum texanum</i>          | Grass     | PC                                  |
| Rice, red               | <i>Oryza punctata</i>           | Grass     | C                                   |
| Sandbur, field          | <i>Cenchrus spinifex</i>        | Grass     | PC                                  |
| Ryegrass, Italian       | <i>Lolium multiflorum</i>       | Grass     | C                                   |
| Sandbur, Southern       | <i>Cenchrus echinatus</i>       | Grass     | PC                                  |
| Shattercane             | <i>Sorghum bicolor</i>          | Grass     | PC                                  |
| Signalgrass, broadleaf  | <i>Urochloa platyphylla</i>     | Grass     | C                                   |
| Sorghum (volunteer)     | <i>Sorghum bicolor</i>          | Grass     | PC                                  |
| Witchgrass              | <i>Panicum capillare</i>        | Grass     | C                                   |



9/61

**PART 2**

| Common Name               | Scientific Name                | Weed Type | Control (C) or Partial Control (PC) |
|---------------------------|--------------------------------|-----------|-------------------------------------|
| Amaranth, Palmer          | <i>Amaranthus palmeri</i>      | Broadleaf | C                                   |
| Amaranth, Powell          | <i>Amaranthus powellii</i>     | Broadleaf | C                                   |
| Beggarweed, Florida       | <i>Desmodium tortuosum</i>     | Broadleaf | PC                                  |
| Carpetweed                | <i>Mollugo verticillata</i>    | Broadleaf | C                                   |
| Eclipta                   | <i>Eclipta prostrata</i>       | Broadleaf | PC                                  |
| Galinsoga, hairy          | <i>Galinsoga quadriradiata</i> | Broadleaf | C                                   |
| Galinsoga, smallflower    | <i>Galinsoga parviflora</i>    | Broadleaf | C                                   |
| Nightshade, Eastern black | <i>Solanum ptychanthum</i>     | Broadleaf | C                                   |
| Nightshade, hairy         | <i>Solanum physalifolium</i>   | Broadleaf | PC                                  |
| Pigweed, prostrate        | <i>Amaranthus blitoides</i>    | Broadleaf | C                                   |
| Pigweed, redroot          | <i>Amaranthus retroflexus</i>  | Broadleaf | C                                   |
| Pigweed, smooth           | <i>Amaranthus hybridus</i>     | Broadleaf | C                                   |
| Pigweed, tumble           | <i>Amaranthus albus</i>        | Broadleaf | C                                   |
| Purslane, common          | <i>Portulaca oleracea</i>      | Broadleaf | PC                                  |
| Pusley, Florida           | <i>Richardia scabra</i>        | Broadleaf | C                                   |
| Spiderwort, tropical      | <i>Commelina benghalensis</i>  | Broadleaf | C                                   |
| Waterhemp, common         | <i>Amaranthus rudis</i>        | Broadleaf | C                                   |
| Waterhemp, tall           | <i>Amaranthus tuberculatus</i> | Broadleaf | C                                   |
|                           |                                |           |                                     |
| Nutsedge, yellow          | <i>Cyperus esculentus</i>      | Sedge     | C                                   |

<sup>1</sup> Refer to the corn section of this label for additional use directions.

**PREPLANT AND ROTATIONAL CROPS SECTION**

**Replanted Crop Directions:**

This section covers replant crops that may be planted following a lost crop that has had an application of AX S-MET.

If a crop treated with AX S-MET is lost, any crop on this label, or on a supplemental AX S-MET label, may be replanted immediately provided that the rate of AX S-MET applied to the previous crop was not greater than the labeled rate for the crop to be replanted. If the first application was banded and the replant crop is planted in the center of the untreated bands, a second banded treatment may be applied at the rate for the use-pattern for the replant crop, provided the application does not overlap the first application band.

**Rotational Crop Restrictions:**

Do not rotate to food or feed crops other than those listed below. For all crops not listed, wait at least 12 months following the last application of AX S-MET before planting.

10/  
61

Barley, oats, rye, or wheat may be planted 4 ½ months following treatment. Alfalfa may be planted 4 months following application. Clover may be seeded 9 months following application.

**Restrictions:** 1) Do not apply more than 1.9 lb. active ingredient per acre (2.0 pt. of AX S-MET) in the previous crop, and 2) do not make lay-by or other postemergence applications of AX S-MET in the previous crop.

Tobacco, buckwheat, and rice, may be planted in the next spring following treatment.

Below in the rotational crop subsections A through C is a listing of rotational crop options that are made possible through S-metolachlor tolerances which were established by the EPA as crop groupings.

**A. If not more than 1.33 pt./A of AX S-MET was applied to the field, the following crops (as well as any listed under subsections B or C below) may be planted 60 days after the last application. A second application of an S-metolachlor containing product to the following crops is prohibited within 60 days of the original application.**

**Crop Subgroup 1B Root Vegetables** - garden beet, edible burdock, carrot, celeriac, turnip-rooted chervil, chicory, ginseng, horseradish, turnip-rooted parsley, parsnip, radish, oriental radish, rutabaga, salsify, black salsify, Spanish salsify, skirret, and turnip.

**Crop Group 3 Bulb Vegetables (if to be harvested green)** - garlic, great-headed garlic, leek, green onion, Welsh onion, shallot, Winter squash (including pumpkins)

**B. If not more than 1.67 pt./A of AX S-MET was applied to the field, the following crops (as well as any listed under Subsection C below) may be planted 60 days after the last application. A second application of an S-metolachlor containing product to the following crops is prohibited within 60 days of the original application.**

**Crop Group 8 Fruiting Vegetables, except Cucurbits** - eggplant, groundcherry (*Physalis* spp.), pepino, peppers (bell, chili, cooking, pimento and sweet), tomatillo and tomato.

**C. If not more than 2.0 pt./A of AX S-MET was applied to the field, the following crops may be planted 60 days after the last application. A second application of an S-metolachlor containing product to the following crops is prohibited within 60 days of the original application.**

**Crop Subgroup 1C Tuberos and Corm Vegetables** - arracacha, arrowroot, Chinese artichoke, jerusalem artichoke, edible canna, bitter and sweet cassava, chayote (root), chufa, dasheen (taro), ginger, leren, potato, sweet potato, tanager, tumeric, yam bean, yam, true.

**Crop Group 3 Bulb Vegetables (if to be harvested dry)** - garlic, great-headed garlic, leek, dry bulb and green onion, Welsh onion, shallot.

**Crop Subgroup 4A Leaf Petiole Vegetables** - cardoon, celery, Chinese celery, celtuce, Florence fennel, rhubarb, and Swiss chard.

**Crop Subgroup 5A Head and Stem Brassica Vegetables** - broccoli, Chinese broccoli, brussel sprouts, cabbage, Chinese (napa) cabbage, Chinese mustard, cauliflower, cavalo broccolo and kohlrabi.

**Precautions:** 1) Rotating to crops within these crop groupings at less than 60 days may result in crop injury. 2) If the rate of AX S-MET applied in the previous crops was greater than the rate listed here (Sections A-C below), these crops cannot be planted until the following spring.

11/61

**APPLICATION PROCEDURES**

**Application Timing**

AX S-MET alone or in tank mixtures with other labeled herbicides may be applied for weed control in certain crops at various times. Refer to the given crop section of the label to determine which of the following application timings are allowed.

- **Preplant Surface-Applied:** For minimum-tillage or no-tillage systems only, AX S-MET alone and some AX S-MET tank mixtures may be applied up to 45 days before planting certain crops. Use only split applications for treatments made 30-45 days before planting, with 2/3 the recommended broadcast rate for the crop and soil texture applied initially and the remaining 1/3 at planting. Treatments less than 30 days before planting may be made either as a split or a single application. Refer to individual crop section on this label to determine if early preplant surface application is recommended. If weeds are present at the time of treatment, apply in a tank mixture combination with a contact herbicide (for example, Gramoxone Inteon®, Touchdown® brands, or Roundup® brands). Observe directions for use, precautions, and restrictions on the label of the contact herbicide. To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.
- **Preplant Incorporated:** Apply AX S-MET to the soil and incorporate into the top 2 inches of soil within 14 days before planting, using a finishing disk, harrow, rolling cultivator, or similar implement capable of providing uniform 2-inch incorporation. Use a preplant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. If crop will be planted on beds, apply and incorporate AX S-MET after bed formation, unless specified otherwise.
- **Preemergence:** Apply AX S-MET during planting (behind the planter) or after planting, but before weeds or crops emerge.
- **Postemergence:** AX S-MET will not control emerged weeds so it must be applied to a weed-free soil surface or in tank mixture with products that provide postemergence control of weeds present at the time of application. Refer to the individual crop section of this label to determine if a postemergence application may be made for that crop.

**Special Application Procedures**

- **CA Only (Corn, Safflowers, Beans, Peas and lentils):**  
**Preplant Incorporated:** Broadcast AX S-MET alone or with tank mix partners listed on this label to the soil and thoroughly incorporate with a disk or similar implement set to till 4-6 inches deep. For more thorough incorporation, till the soil in 2 different directions (cross-till). Crops may be planted on flat surface or on beds. Caution must be used when forming the beds that only soil from the AX S-MET treated zone is used (i.e., untreated soil must not be brought to soil surface). If the application is made to preformed beds, incorporate AX S-MET with a tillage implement set to till 2-4 inches deep. Care should be taken during tilling to keep the tilled (AX S-MET treated) soil on the beds. **Preemergence:** Apply AX S-MET after planting. Water with sprinkler or flood irrigation within 7-10 days.
- **Fall Application for Spring Weed Control (Only in IA, MN, ND, SD, WI, and portions of NE and IL - See specific instructions in the Corn, Soybeans, and Beans, Peas, Lentils sections of this label for timing of application and other information):** Do not apply to frozen ground. Use on medium and fine soils with greater than 2.5% organic matter that will be planted to corn or soybeans the next spring. Ground may be tilled before or after application. Do not exceed a 2 to 3-inch incorporation depth if tilled after treatment. **Restriction:** If a spring application is

12/61

made, the total rate of the fall plus spring applications must not exceed the maximum total rate for the specific crop, or illegal residues may result.

• **Fall Application for Italian Ryegrass Control (Corn, Cotton, Grain and Forage Sorghum, and Soybean Only - See specific instructions in the Corn, Cotton, Grain and Forage Sorghum, and Soybean sections of this label for timing of application and other information):** AX S-MET may be applied in the fall (September 1 – December 1) for residual control of glyphosate-resistant Italian ryegrass (*Lolium multiflorum*). A tillage operation may precede the application. Do not incorporate to a depth greater than 2-3 inches if tillage follows the application of AX S-MET.

**Restrictions:** 1) Do not apply AX S-MET to frozen ground. All crops on the AX S-MET label may be planted the following spring after application. 2) If a spring application is made, the combined total amount of AX S-MET applied in the fall plus the spring must not exceed the maximum seasonal S-metolachlor rate for the specific crop planted or illegal residues may result. Refer to the crop sections on this label for specific directions.

**Ground Application:** Apply AX S-MET alone or in tank mixtures by ground equipment in a minimum of 10 gals. of spray mixture per acre, unless otherwise specified. Use sprayers that provide accurate and uniform application. For AX S-MET tank mixtures with wettable powder or dry flowable formulations, screens and strainers must be no finer than 50-mesh. Rinse sprayer thoroughly with clean water immediately after use.

Calculate the amount of herbicide needed for band treatment by the formula:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

For information on applying in lower volumes of carrier, see **Low Carrier Application** section.

For application by air or through center pivot systems, see **Aerial Drift Management** and **Aerial Drift Reduction Advisory Information** sections.

For information on impregnating dry fertilizer, see **Dry Bulk Granular Fertilizers** section.

For information on application using variable-rate technologies, see **Variable-Rate Application** section.

**SPRAY EQUIPMENT**

**LOW CARRIER APPLICATION  
For Broadcast Ground Application Only**

Use sprayers, such as Ag-Chem RoGator®, Hagie, John Deere Hi-Cycle™, Melroe Spra-Coupe, Tyler Patriot™, or Willmar Air Ride®, that provide accurate and uniform application. **Only water may be used as a carrier.** Screens in suction and in-line strainers should be 50-mesh. Manufacturers may require that tip screens as fine as 100-mesh be used with some nozzles. Use a pump with capacity to: (1) maintain up to 35-40 psi at the nozzles, and (2) provide sufficient agitation in tank to keep mixture in suspension. Use a minimum of 5.0 gals. of spray mixture per acre. Maximum recommended sprayer speed is 15 mph. Rinse sprayer thoroughly with clean water immediately after each use.

**Note:** Use low pressure nozzles to reduce drift and increase application accuracy. Care should be taken when using automatic rate controlling devices to spray the material within the rated working pressure and flow ranges of the nozzles selected. Nozzle screens must be used when directed by

the manufacturer. All nozzles should be placed on 20-inch centers, except flooding types which should be placed on 40-inch centers. When Flat Fan-type nozzles are used, angles of 80° or 110° must be used. Always read and follow the manufacturer's directions for optimum setup and performance of their nozzles or tips.

**AERIAL APPLICATION**

Apply AX S-MET in water alone or in tank mixtures with AAtrex, Lorox®, or Sencor® in a minimum total volume of 2.0 gals./A by aircraft. AX S-MET may also be applied by air in combination with Balan®, Prowl®, or Treflan®. Avoid application under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft., using low-drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply AX S-MET alone or AX S-MET + AAtrex by aircraft at a minimum upwind distance of 400 ft. from sensitive plants, or apply AX S-MET Lorox, or Sencor at a minimum upwind distance of 300 ft. from sensitive plants.

**Aerial Drift Management**

The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses, or to applications using dry formulations.

- The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
- Nozzles must always point backward parallel with the air stream and never be pointed downward more than 45 degrees.

Where states have more stringent regulations, they must be observed.

The applicator must be familiar with and take into account the information covered in the **Aerial Drift Reduction Advisory Information** section below.

**Aerial Drift Reduction Advisory Information**

**Information on Droplet Size**

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see **Wind, Temperature and Humidity, and Temperature Inversions**).

**Controlling Droplet Size**

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's maximum pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

### **Application Height**

Applications must not be made at a height greater than 10ft. above the top of the largest plants, unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

### **Swath Adjustment**

When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the upwind and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

### **Wind**

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Do not make applications when wind speed is below 2 mph due to variable wind direction and high inversion potential. **Note:** Local terrain can influence wind patterns.

### **Temperature and Humidity**

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is greatest when conditions are both hot and dry.

### **Temperature Inversions**

Do not make applications during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

### **Sensitive Areas**

The pesticide must only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Avoid application to humans or animals. Flagmen and loaders must avoid inhalation of spray mist and prolonged contact with skin.

### **CENTER PIVOT IRRIGATION APPLICATION**

AX S-MET alone or in tank mixture with other herbicides on this label, which are registered for center pivot application, may be applied in irrigation water preemergence (after planting, but before weeds or crop emerge) at rates listed on this label. AX S-MET also may be applied postemergence to the crop and preemergence to weeds in crops where postemergence applications are allowed on this label. Follow all restrictions (height, timing, rate, etc.) to avoid illegal residues. Apply this product only through a center pivot irrigation system. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension specialists, equipment manufacturers, or other experts. Do not

connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

### Operating Instructions

- The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water- source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump or piston pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.
- Prepare a mixture with a minimum of 1 part water to 1 part, herbicide(s) and inject this mixture into the center pivot system. Injecting a larger volume of a more dilute mixture per hour will usually provide more accurate calibration of metering equipment. Maintain sufficient agitation to keep the herbicide in suspension.
- Meter into irrigation water during entire period of water application.
- Apply in 1/2-1 inch of water. Use the lower water volume (1/2 inch) on coarse- textured soils and the higher volume (1 inch) on fine-textured soils. More than 1 inch of water at application may reduce weed control by moving the herbicide below the effective zone in the soil.

**Precaution** for center pivot applications: Where sprinkler distribution patterns do not overlap sufficiently, unacceptable weed control may result. Where sprinkler distribution patterns overlap excessively, crop injury may result.

### DRY BULK GRANULAR FERTILIZERS

Many dry bulk granular fertilizers may be impregnated or coated with AX S-MET alone or selected AX S-MET tank mixtures which are registered for preplant incorporated or preplant surface application which are used to control weeds in crops on the AX S-MET label and are not prohibited from use on dry bulk granular fertilizers.

When applying AX S-MET or AX S-MET mixtures with dry bulk granular fertilizers, follow all directions for use and precautions on the respective product labels, regarding target crops, rates per acre, soil texture, application methods (including timing of application), and rotational crops.

All individual state regulations relating to dry bulk granular fertilizer blending, registration, labeling, and application are the responsibility of the individual and/or company selling the herbicide/fertilizer mixture.

16/61

Prepare the herbicide/fertilizer mixtures by using any closed drum, belt, ribbon, or other commonly used dry bulk fertilizer blender. Nozzles used to spray AX S-MET and AX S-MET mixtures onto the fertilizer must be placed to provide uniform spray coverage. Care must be taken to aim the spray directly onto the fertilizer only and to avoid spraying the walls of the blender.

If the herbicide/fertilizer mixture is too wet, add a highly absorptive material, such as Agsorb® or Celatom MP-79®, or similar granular clay or diatomaceous earth materials, to obtain a dry, free-flowing mixture. Absorptive materials should be added only after the herbicide has been thoroughly blended into the fertilizer mixture. Best application results will be obtained by using a granule of 6/30 particle size or of a size similar to that of the fertilizer material being used. Generally, less than 2% by weight of absorptive material will be needed. Avoid using more than 5% absorptive material by weight.

Calculate amounts of AX S-MET, AAtrex, AAtrex + Princep®, Balance® Pro, Princep, Sencor, or Sonalan® by the following formula:

|   |   |                                      |   |  |
|---|---|--------------------------------------|---|--|
| $\frac{2000}{\text{lbs. of fertilizer per acre}}$ | X | pts./A of liquid or flowable product | = | pts. of liquid or flowable product per ton of fertilizer |
| $\frac{2000}{\text{lbs. of fertilizer per acre}}$ | X | lbs./A of dry product                | = | lbs. of dry product per ton of fertilizer                |

**Pneumatic (Compressed Air) Application (AX S-MET Alone):** High humidity, high urea concentrations, low fertilizer use rates, and dusty fertilizer may cause fertilizer mixture to build up or plug the distributor head, air tubes, or nozzle deflector plates. To minimize buildup, premix AX S-MET with Exxon Aromatic 200 at a rate of 1.0-4.0 pts./gal. of AX S-MET. Aromatic 200 is a noncombustible/nonflammable petroleum product. Aromatic 200 may be used in either a fertilizer blender or through direct injection systems. Drying agents must not be used when using Aromatic 200.

**Precautions:** (1) Mixtures of AX S-MET and Aromatic 200 must be used on dry fertilizer only. Poor results or crop injury may result if these mixtures are used in water or liquid fertilizer solutions for spraying applications. (2) When impregnating AX S-MET in a blender before application, a drier mixture can be attained by substituting a drying agent for Aromatic 200. Agsorb FG or drying agents of 6/30 particle size may be used. (3) Drying agents must not be used with On-The-Go impregnation equipment.

**Restrictions:** (1) Do not impregnate AX S-MET or AX S-MET mixtures on ammonium nitrate, potassium nitrate, or sodium nitrate, either alone or in blends with other fertilizers. (2) Do not use AX S-MET or AX S-MET mixtures on straight limestone, since absorption will not be achieved. Fertilizer blends containing limestone can be impregnated.

**Application**

Apply 200-700 lbs. of the herbicide/fertilizer mixture per acre. For best results, apply the mixture uniformly to the soil with properly calibrated equipment immediately after blending. Uniform application of the herbicide/fertilizer mixture is essential to prevent possible crop injury. Nonuniform application may also result in unsatisfactory weed control. In areas where conventional tillage is practiced, a shallow incorporation of the mixture into the soil may improve weed control. On fine- or medium-textured soils in areas where soil incorporation is not planned, i.e., reduced tillage situations or in some conventional till situations, make applications approximately 30 days before



planting to allow moisture to move the herbicide/fertilizer mixture into the soil. On coarse-textured soils, make applications approximately 14 days prior to planting.

17/  
61

**Precaution:** To avoid crop injury, do not use the herbicide/fertilizer mixture on crops where bedding occurs.

### MIXING INSTRUCTIONS

**AX S-MET Alone:** Mix AX S-MET with water or fluid fertilizer and apply as a spray. Fill the spray tank  $\frac{1}{2}$ - $\frac{3}{4}$  full with water or fluid fertilizer, add the proper amount of AX S-MET, and then add the rest of the water or fluid fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform emulsion.

**Tank Mixtures:** Fill the spray tank  $\frac{1}{4}$  full with water, and start agitation; add 2,4-D, AAtrex, Balance Pro, Balan, Banvel®, Basagran®, Butoxone®, Butyrac®, Canopy®, Caparol® 4L, Command®, Cotoran®, Eptam®, Liberty® Herbicide, Liberty ATZ Herbicide, Lorox, Marksman®, MSMA, Princep, Prowl, Pursuit®, AAtrex + Princep, Scepter®, Sencor, Sonalan, or Treflan, and allow it to become dispersed; then add AX S-MET; then add Gramoxone Inteon, Landmaster® BW, Touchdown, or Roundup (glyphosate products) if these products are being used; and finally the rest of the water. For tank mixtures with AAtrex, Balance, Banvel, Canopy, Caparol 4L, Command, Cotoran\*, Eptam, Lorox, Marksman, Princep, Prowl\*, Pursuit, AAtrex + Princep, Scepter, Sencor, Sonalan, or Treflan, fluid fertilizers may replace all or part of the water as carrier, except in the AAtrex postemergence and the Banvel postemergence tank mixes. For tank mixtures with AAtrex, see additional mixing instructions on the AAtrex label. For each mixture, check compatibility with fluid fertilizer, as described below, before mixing in spray tank. For all tank mixtures, agitate during mixing and application to maintain a uniform suspension.

\*See **Special Mixing Instructions** for tank mixtures with Cotoran, and with AAtrex or Princep + Prowl under the appropriate tank mixture section.

For directions on how to conduct a compatibility test, see the **Compatibility Test** section.

### COMPATIBILITY TEST

A jar test is recommended before tank mixing to ensure compatibility of AX S-MET with other pesticides. The following test assumes a spray volume of 25 gals./A. For other spray volumes, make appropriate changes in the ingredients.

**Note:** Nitrogen solutions or complete fluid fertilizers may replace all or part of the water in the spray. Because liquid fertilizers vary, even within the same analysis, **always check compatibility with pesticide(s) before use.** Incompatibility of tank mixtures is more common with suspensions of fertilizer and pesticides.

### Test Procedure

1. Add 1.0 pt. of carrier (fertilizer or water) to each of 2 one qt. jars with tight lids. Use the same source of water that will be used for the tank mix and conduct the test at the temperature the tank mix will be applied.
2. To one of the jars, add  $\frac{1}{4}$  tsp. or 1.2 milliliters of a compatibility agent approved for this use, such as Envelop (1/4 tsp. is equivalent to 2.0 pts./100 gals. spray). Shake or stir gently to mix.
3. To both jars, add the appropriate amount of pesticide(s) in their relative proportions based on listed label rates. If more than one pesticide is used, add them separately with dry pesticides first, flowables next, and emulsifiable concentrates last. After each addition, shake or stir gently to thoroughly mix.

18/  
61

4. After adding all ingredients, put lids on and tighten, and invert each jar ten times to mix. Let the mixtures stand 15-30 minutes and then look for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the two jars. If either mixture separates, but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility: (a) Slurry the dry pesticide(s) in water before addition, or (b) add 1/2 the compatibility agent to the fertilizer or water and the other 1/2 to the emulsifiable concentrate or flowable pesticide before addition to the mixture. If incompatibility is still observed, do not use the mixture.

5. After compatibility testing is complete, dispose of any pesticide wastes in accordance with the **Storage and Disposal** section in this label.

**CROP USE DIRECTIONS**

**CORN (ALL TYPES) - AX S-MET ALONE**

Apply AX S-MET, either preplant surface, preplant incorporated, preemergence, or lay-by, using the appropriate rate specified below.

**PREPLANT SURFACE-APPLIED**

Refer to instructions for use of AX S-MET alone under **Application Procedures**.

**Use Directions and Restrictions for Fall Application for Spring Weed Control:**

- Apply after September 30 in ND, SD, MN, WI, and north of Route 30 in IA.
- Apply after October 15 north of Route 91 in NE and south of Route 30 in IA.
- Apply after October 31 north of Route 136 in IL.

In all locations, apply to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55°F and falling. In minimum-till or no-tillage systems on soils having greater than 2.5% organic matter, use 1.67-2.0 pts./A on medium-textured and 2.0 pts./A on fine-textured soils. Do not apply to frozen ground. A tillage operation may precede the application. A fall and/or a spring tillage may follow application, but do not exceed an incorporation depth greater than 2-3 inches. Minimize furrow and ridge formation in the tillage operations.

**Restriction:** If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for corn, or illegal residues may result.

**Fall Application for Italian Ryegrass Control:** AX S-MET may be applied for residual control of glyphosate-resistant Italian ryegrass (*Lolium multiflorum*). Apply AX S-MET at 1.33-1.67 pints/Acre in the fall (September 1-December 1) after harvest of the previous crop and prior to Italian ryegrass emergence. Use the lower AX S-MET rate for coarse textured soils and the higher rate for fine textured soils. A tillage operation may precede the application. Do not incorporate to a depth greater than 2-3 inches if tillage follows the application of AX S-MET. For fall applications after emergence of glyphosate-resistant Italian ryegrass, Gramoxone Inteon can be tank mixed with AX S-MET to control emerged ryegrass. Refer to the Gramoxone Inteon label for specific rates, application instructions and restrictions. Other registered herbicides may be tank mixed with AX S-MET for control or improved control of other weeds present at the time of application.

**Restrictions:** (1) Do not apply AX S-MET to frozen ground. (2) If a spring application is made, the combined total amount of AX S-MET applied in the fall plus the spring must not exceed the maximum seasonal S-metolachlor rate for corn (3.9 pt/A depending on soil texture), or illegal residues may result.

19/  
61

**Fall Application for Control or Suppression of Yellow Nutsedge (ID, OR and WA only):** For pre-emergent control or suppression of yellow nutsedge the following spring apply 1.33 pt/A of AX S-MET in the fall after the harvest of the previous crop but before freeze-up. Fall applications of AX S-MET can be surface-applied or incorporated.

**Restrictions:** (1) Make no more than one fall application per crop. (2) Apply not more than 1.33 pts./A in a single fall preplant application. (3) Do not apply to frozen ground. (4) If a spring application is made, the combined total amount of AX S-MET applied in the fall plus the spring must not exceed the maximum seasonal S-metolachlor rate for corn (3.9 pt/A depending on soil texture).

**Early Pre-plant Applications**

A. Use on medium- and fine-textured soils with minimum-tillage or no-tillage systems in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, TN, WI, and WY. Apply 2/3 of the listed rate of AX S-MET (1.67 pts./A on medium soils and 2.0 pts./A on fine soils) as a split treatment 30-45 days before planting and the remainder at planting. Applications made less than 30 days prior to planting may be as either a split or single treatment. Apply 1.33 pts./A on coarse soils not more than 2 weeks prior to planting. **Restriction:** If a spring application is made, the total rate of the fall plus spring application must not exceed the maximum total rate for corn, or illegal residues may result.

B. On medium- and fine-textured soils with minimum- or no-tillage systems in CT, DE, MA, MD, ME, MI, NH, NY, OH, PA, RI, VA, VT, and WV, preplant surface applications may be applied following the directions for use above. If the amount of rainfall results in unsatisfactory length of weed control following the earlier treatment, a postemergence application of an appropriately labeled broadleaf and/or grass weed herbicide may be used, i.e., AAtrex, Beacon®, Bicep Magnum, Bicep II Magnum®, Exceed®, Accent®, Banvel, Basagran, bromoxynil (Buctril®), or 2,4-D. If the postemergence treatment includes the herbicide used preplant surface-applied, do not exceed the total labeled rate for corn on a given soil texture. Observe all directions for use, precautions, and limitations on the label of the postemergent herbicide.

**PREPLANT INCORPORATED OR PREEMERGENCE**

Follow instructions for use of AX S-MET alone under **Application Procedures**. On coarse soils, apply 1.0-1.33 pts./A of AX S-MET if organic matter content is less than 3%, or 1.33 pts./A if organic matter content is 3% or greater. On medium soils, apply 1.33-1.67 pts./A of AX S-MET. On fine soils, apply 1.33-1.67 pts./A of AX S-MET if organic matter content is less than 3%, or 1.67-2.0 pts./A if organic matter content is 3% or greater.

**POSTEMERGENCE OR LAY-BY**

To extend the duration of weed control in corn, a maximum rate of 2.0 pts./A of AX S-MET may be applied after corn emergence until the corn plants reach 40 inches in height, following any preplant surface-applied, preplant incorporated, or preemergence herbicide application, including AX S-MET. For best results, applications must be made to soil free of emerged weeds and directed toward the base of corn plants in excess of 5 inches tall. The total AX S-MET rate applied on corn during any one crop year should not exceed 3.9 pts./A, depending on soil texture.

**Restrictions for all applications to corn:** (1) do not graze or feed forage from treated areas for 30 days following application and (2) do not harvest sweet corn ears from treated areas for 30 days following application.

**PROBLEM WEED CONTROL DIRECTIONS**

**Shattercane, Wild Proso Millet, Woolly Cupgrass, and Eclipta - Partial Control:** For more consistent partial control of shattercane, wild proso millet, woolly cupgrass, or eclipta, apply 1.0-1.33 pts./A of AX S-MET preplant incorporated followed by 1.0-1.33 pts./A of AX S-MET preemergence. Make the preemergence application during or after planting, but before weeds and

20/61

corn emerge. Apply the 1.33 pts./A rate of AX S-MET when a heavy infestation of shattercane, wild proso millet, woolly cupgrass, or eclipta is expected. A shallow cultivation may be needed to control any late emerging weeds.

**Woolly Cupgrass and Wild Proso Millet Control Program:** For control of these species, use the following 3-step program: (1) Apply AX S-MET early preplant, preplant incorporated, or preemergence at 1.67 pts./A on medium soils and 2.0 pts./A on fine-textured soils, up to the maximum label rate. Lightly incorporate with a rotary hoe if rainfall does not occur within 5-7 days; (2) Apply a postemergence tank mix of Beacon at 0.38 oz./A or Exceed at 1 packet per 4 acres plus Accent SP at 0.33 oz./A plus 1.0 qt. of crop oil concentrate plus 1.0 gal./A of 28% nitrogen, or the equivalent amount of ammonium sulfate, when grasses are 2-3 inches tall and the corn is at least 4 inches tall; and (3) Cultivate 14-21 days after the postemergence application.

**Restrictions:** (1) Do not apply more than the labeled application rate for a given soil texture per year, either as a single or split treatment, or illegal residues may result. (2) In corn, AX S-MET may be used up to 2.6 pts./A as either a preplant surface, preplant incorporated, or preemergence treatment on soils having an organic matter content between 6% and 20%. (3) In the event of escape of annual weeds following a preplant surface, preplant incorporated, or preemergence treatment of AX S-MET, follow with a postemergence application of an appropriately labeled broadleaf and/or grass weed herbicide, i.e., AAtrex, Beacon, Bicep II Magnum, Exceed, Accent, Banvel, Basagran, Buctril, or 2,4-D. If the postemergence treatment includes the herbicide used in the earlier treatment, do not exceed the total labeled rate for corn on a given soil texture. (4) Buctril may be applied postemergence alone or in tank mix combination with AAtrex. Do not exceed 1.2 lbs. a.i./A of AAtrex in tank mix combination with Buctril postemergence. Refer to the AAtrex, and Buctril labels for specific rates and precautions. (5) Do not use AX S-MET on peat or muck soils.

**CORN - AX S-MET COMBINATIONS**

AX S-MET in any tank mixture for corn may be applied in water or fluid fertilizer before corn emerges. Use only water as a carrier when AX S-MET is applied after corn emergence.

**Restrictions:** For all applications to corn, (1) do not graze or feed forage from treated areas for 30 days following application or possible illegal residues may result and (2) do not harvest sweet corn ears from treated areas for 30 days following application.

**IMPORTANT: FOR TANK MIXTURES WITH AATREX (OR OTHER BRANDS OF ATRAZINE) -** If applying AX S-MET in tank mixture with AAtrex, all the restrictions and rate limitations on the AAtrex label must be followed if more restrictive/protective than those on this label. In addition, if AAtrex is/must be applied at rates lower than those listed on this label, broadleaf weed control may be affected. Refer to the AAtrex label for weeds controlled at the reduced rates.

**Table 2: AX S-MET Tank Mixtures for Corn - Additional Weeds Controlled and Special Instructions**

|                             | AX S-MET +AAtrex and/or Princep (Preplant Surface,PPI, PRE) | AX S-MET + AAtrex (Post) | AX S-MET + Banvel (Field Corn) | AX S-MET + AAtrex + Lorox | AX S-MET + AAtrex or Princep+ Prowl | AX S-MET + Marksman | AX S-MET + Balance Pro |
|-----------------------------|---|--------------------------|--------------------------------|---------------------------|-------------------------------------|---------------------|------------------------|
| Special Mixing Instructions |   |                          |                                |                           | 1                                   |                     |                        |
| Comments                    | 2,3,4,5,7,8   | 2,3,4,5                  |                                | 2,3,4,5,6                 | 2,3,4,5                             | 7                   | 2,3,7                  |
| Browntop panicum            | X   |                          |                                | X                         | X                                   |                     |                        |

21/  
61

|                  |   |   |   |   |   |   |     |
|------------------|---|---|---|---|---|---|-----|
| Cocklebur        | X | 0 | 0 | X | X | X |     |
| Common purslane  | X |   |   | X | X | X | X   |
| Hairy nightshade | X |   |   | X | X | X |     |
| Jimsonweed       |   | X | 0 |   |   | X | X   |
| Kochia           |   | X |   |   |   | X | X   |
| Lambsquarters    | X | X | X | X | X | X | X   |
| Morningglory     | X | 0 | 0 | X | X | X |     |
| Mustard          |   | X |   |   |   | X | X   |
| Pigweed          |   |   |   | X | X | X | X   |
| Prickly sida     |   | X |   |   |   | X |     |
| Ragweed          | X | X | X | X | X | X | X   |
| Smartweed        | X | X | X | X | X | X | X   |
| Velvetleaf       | X | X | 0 | X | X | X | 0-X |

x = control; 0 = partial control; 0-X = partial to full control depending on ratio of products used or on weed population  
Field corn only

### Comments

#### 1) Special Mixing Instructions for AX S-MET + AAtrex or Princep and Prowl

- a) Fill the spray tank  $\frac{1}{2}$  full with water or fluid fertilizer and start agitation.
- b) To aid compatibility, add a compatibility agent, such as Envelop at 4.0 pts./100 gals. of spray mixture.
- c) Then add the AAtrex or Princep and allow it to become dispersed.
- d) Then add AX S-MET and Prowl 4E.
- e) Add the rest of the water.

2) Although a single formulation for AAtrex or Princep is listed in the rate tables, other formulations may be substituted, using the following formula:

- a) 1.0 lb. of AAtrex Nine-O® or Princep Caliber 90® = 1.8 pts. of AAtrex 4L or Princep 4L.

3) Although directions specify AAtrex formulations in tank mixture with AX S-MET, other brands of atrazine may be used. Follow the rates, and other use directions and restrictions on the atrazine label.

4) See additional mixing instructions on the AAtrex label.

5) Do not exceed a total of 2.5 lbs. a.i. of atrazine per acre per year. However, certain states may have established rate limitations for atrazine within specific geographical areas. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.

6) Other formulations of Lorox can be used: 1.0 lb. of Lorox DF = 1.0 pt. of Lorox L.

7) In Minimum-Tillage and No-Tillage systems, mix with Gramoxone Inteon for control of most emerged annual weeds and suppression of perennial weeds; or with Landmaster BW for

22/61

suppression of emerged field bindweed and control or suppression of annual weeds; or with Touchdown brands or Roundup brands for control of most emerged annual and perennial weeds.

8) Refer to the **Corn- AX S-MET Combinations-Tank Mixture with AAtrex; or AAtrex + 2,4-D; or AAtrex + 2,4-D + Banvel for Minimum Tillage or No- Tillage Systems** section for specific directions for 2,4-D or Banvel burndown combinations with Minimum-Tillage and No-Tillage systems.

AX S-MET in any tank mixture for corn may be applied in water or fluid fertilizer, except as restricted.

**Restrictions:** (1) For all applications to corn, do not graze or feed forage from treated areas for 30 days following application and do not harvest sweet corn ears from treated areas for 30 days following application, or possible illegal residues may result. (2) When applying AX S-MET in tank mixture with AAtrex, do not exceed a total of 2.5 lbs. a.i. of atrazine per acre per year. (3) Refer to **Corn (All Types) - AX S-MET Alone** directions for sequential postemergence treatments if escape weeds develop.

**TANK MIXTURE WITH AATREX OR PRINCEP, OR AATREX + PRINCEP- PREPLANT SURFACE, PREPLANT INCORPORATED, OR PREEMERGENCE**

In addition to the weeds controlled by AX S-MET alone, AX S-MET + AAtrex or Princep, or AX S-MET +AAtrex+Princep, applied preplant surface, preplant incorporated or preemergence, and also controls the following weeds: browntop panicum, cocklebur, common purslane, hairy nightshade, lambsquarters, morningglory, ragweed, smartweed, and velvetleaf.

Apply AX S-MET + AAtrex or Princep, or AX S-MET + AAtrex + Princep either preplant surface, preplant incorporated, or preemergence.

**Preplant Surface-Applied:** Follow instructions for use of AX S-MET alone under Application Procedures and under application instructions for AX S-MET alone on corn. Apply AX S-MET + AAtrex or Princep, or AX S-MET + AAtrex + Princep on medium soils (1.67 pts./A of AX S-MET + 3.2-4.0 pts./A of AAtrex 4L or Princep 4L, or AAtrex 4L + Princep 4L combined) and on fine soils (1.67-2.0 pts./A of AX S-MET + 4.0 pts./A of AAtrex 4L or 4.0-5.0 pts./A of Princep 4L, or AAtrex 4L + Princep 4L combined) in minimum-tillage and no-tillage systems in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, TN, WI, and WY. Apply the tank mixtures as a split or single treatment in those states and as indicated in the **AX S-MET Alone – Preplant Surface-Applied** section of the label for corn. On coarse soils, apply 1.33 pts./A of AX S-MET and 3.2 pts./A of AAtrex 4L or Princep 4L, or AAtrex 4L + Princep 4L combined.

**Preplant Incorporated or Preemergence:** Follow instructions for use of AX S-MET alone under **Application Procedures**. Apply AX S-MET + AAtrex or Princep, or AX S-MET + AAtrex + Princep, using the appropriate rates from Table 3.

**Restriction:** Do not apply more than the labeled rate for a given soil texture per year, either as a split or single treatment.

**Shattercane and Wild Proso Millet - Partial Control**

For more consistent partial control of shattercane or wild proso millet, where AX S-MET is applied in tank mixture or sequentially with other registered corn herbicides, the following applications may be made:

- Apply 1.0-1.33 pts./A of AX S-MET + 2.0 lbs. a.i./A of AAtrex or Princep preplant incorporated, followed by 1.0-1.33 pts./A of AX S-MET preemergence. Make the preemergence application during or after planting, but before weeds and corn emerge.

23/61

- Apply AX S-MET at 1.33 pts./A alone or in tank mix combination with up to 2.0 lbs. a.i./A of AAtrex, or Princep, preplant incorporated. Do not exceed the total rate of triazine herbicide listed in combination with AX S-MET for corn grown on a given soil texture. Follow with a post-directed application of Evik® BOW at 2.5 lbs./A. Refer to the Evik BOW label for specific directions for the post-directed application.
- Apply Eradicane® or (or equivalent EPTC or butylate formulations) at labeled rates preplant incorporated, followed by a preemergence application of AX S-MET at 1.0-1.33 pts./A. Do not use Eradicane on soils where rapid degradation has been shown to occur. Make the preemergence application during or after planting, but before weeds and corn emerge.

**Precaution:** When following the application regimes in numbers 1-3 above, a shallow cultivation may be needed after the pre emergence or postemergence application to help control any late emerging shattercane or wild proso millet plants.

**Restriction:** Do not exceed a total of 1.9 lbs. a.i./A (2.0 pts. of AX S-MET) in the preplant incorporated plus preemergence application on soils with less than 6% organic matter.

**Table 3: AX S-MET + AAtrex or Princep, or AX S-MET + AAtrex + Princep, Preplant Incorporated or Preemergence - Corn (All Types)**

| Soil Texture                                  | < 3% Organic Matter  |    |   | 3% Organic Matter or Greater   |    |   |
|---|--|----|---|--|----|---|
|   | AX S-MET + AAtrex Nine-O <sup>®</sup> or Princep Caliber 90 <sup>®</sup> | Or | AX S-MET + AAtrex Nine-O <sup>®</sup> + Princep Caliber 90 <sup>®</sup> | AX S-MET + AAtrex Nine-O <sup>®</sup> or Princep Caliber 90 <sup>®</sup> | Or | AX S-MET + AAtrex Nine-O <sup>®</sup> + Princep Caliber 90 <sup>®</sup> |
| Coarse  | 0.8-1.0 pt.<br>+<br>1.1-2.2 lbs.   |    | 0.8-1.0 pt.<br>+<br>0.6-1.1 lbs.<br>+<br>0.6-1.1 lbs.                   | 1.0 pt.<br>+<br>1.3-2.2 lbs.   |    | 1.0 pt.<br>+<br>0.7-1.1 lbs.<br>+<br>0.7-1.1 lbs.                       |
| Medium  | 1.0-1.33 pts.<br>+<br>1.3-2.2 lbs.                                       |    | 1.0-1.33 pts.<br>+<br>0.7-1.1 lbs.<br>+<br>0.7-1.1 lbs.                 | 1.33 pts.<br>+<br>1.8-2.2 lbs.   |    | 1.33 pts.<br>+<br>0.9-1.1 lbs.<br>+<br>0.9-1.1 lbs.                     |
| Fine  | 1.33 pts.<br>+<br>1.8-2.2 lbs.   |    | 1.33 pts.<br>+<br>0.9-1.1 lbs.<br>+<br>0.9-1.1 lbs.                     | 1.33-1.67 pts.<br>+<br>1.8-2.2 lbs.***                                   |    | 1.33-1.67 pts.<br>+<br>0.9-1.1 lbs.***<br>+<br>0.9-1.1 lbs.***          |
| Muck or Peat (soils with >20% organic matter) | <b>DO NOT USE</b>  |    |   |  |    |   |

\* Use Princep in preference to AAtrex when heavy infestations of crabgrass or fall panicum are expected. On soils having between 6% and 20% organic matter, AX S-MET may be used up to 2.33 pts./A in tank mix combination with 2.2 lbs./A of AAtrex Nine-O, or equivalent rates of AAtrex 4L. Refer to the AAtrex label for weeds controlled at this reduced rate.

\*\* When using the tank mixture of AX S-MET + AAtrex Nine-O + Princep Caliber 90, use equal rates of each as shown when heavy broadleaf weed infestations are expected. When heavy infestations of crabgrass or fall panicum are expected, use a 1:2 ratio of AAtrex + Princep instead of the 1:1 ratio given in Table 3.

(Example: Total AAtrex Nine-O + Princep Caliber 90 = 1.2 lbs./A, use 0.4 lb. of AAtrex + 0.8 lb. of Princep, respectively.) Refer to Comment No.2 following Table 2 for AAtrex 4L and Princep 4L conversions.

\*\*\* For cocklebur, yellow nutsedge, and velvetleaf control on fine-textured soils above 3% organic matter, apply 2.25 lbs./A of AAtrex Nine-O, or equivalent rates of AAtrex 4L, or the same total amount of AAtrex + Princep with 1.33-1.67 pts./A of AX S-MET.

**TANK MIXTURE WITH AATREX - POSTEMERGENCE**

| <u>Weeds Controlled</u>    |               | <u>Weeds Partially Controlled</u> |
|----------------------------|---------------|-----------------------------------|
| Barnyardgrass (watergrass) | lambsquarters | cocklebur                         |
| crabgrass                  | mustard       | morningglory                      |
| crowfootgrass              | pigweed       | yellow nutsedge                   |
| fall panicum               | prickly sida  |                                   |
| giant foxtail              | purslane      |                                   |
| green foxtail              | ragweed       |                                   |
| yellow foxtail             | smartweed     |                                   |
| jimsonweed                 | velvetleaf    |                                   |
| kochia                     |               |                                   |

Apply 1.0 pt./A of AX S-MET + 1.3 lbs./A of AAtrex Nine-O\* on coarse soils, 1.33 pts./A of AX S-MET + 1.8 lbs./A of AAtrex Nine-O on medium soils, or 1.33-1.67 pts./A of AX S-MET + 1.8-2.2 lbs./A\*\* of AAtrex Nine-O on fine soils. Apply this tank mixture before grass and broadleaf weeds pass the 2-leaf stage and before corn exceeds 5 inches in height. Application to weeds larger than the 2-leaf stage will generally result in unsatisfactory control.

**Lay-by:** Apply to corn plants not more than 12 inches tall. Applications to corn in excess of 5 inches should be directed to the base of the corn plants; whereas, applications to corn plants less than 5 inches tall may be made over the top. Occasionally, some corn leaf burn may result, but this should not affect later growth or yield.

**Precaution:** TO AVOID CROP INJURY, Do not apply this postemergence tank mixture in fluid fertilizer.

\*When using AAtrex 4L, use equivalent rates. One lb. of AAtrex Nine-O = 1.8 pts. Of AAtrex 4L.

\*\*For better control of cocklebur, morningglory, velvetleaf, and yellow nutsedge on fine-textured soils above 3% organic matter, apply 2.2 lbs./A of AAtrex Nine-O, or equivalent rate of AAtrex 4L, with 1.33-1.67 pts./A of AX S-MET.

Tank mixtures of AX S-MET + AAtrex may be applied following use of any registered preplant surface-applied, preplant incorporated, or preemergence corn herbicide, including AX S-MET + AAtrex.

**Restriction:** The total AX S-MET rate must not exceed 3.9 pts., or the AAtrex rate more than 2.5 lbs. a.i./A during any one crop year. Refer to the AAtrex label for geographic, soil-texture, and rotational restrictions.

**TANK MIXTURE WITH BANVEL**

**Preemergence:** Use this tank mixture only on field corn which is flat-planted (no furrows) in CO, IA, IL, IN, KS, MN, NE, OH, SD, and WI.

In addition to the weeds controlled by AX S-MET alone, AX S-MET + Banvel, applied preemergence, also controls lambsquarters, ragweed, smartweed, cocklebur\*, jimsonweed\*, morningglory\*, and velvetleaf\*.

\*Partially controlled.



25/61

Apply AX S-MET + Banvel preemergence. Broadcast 1.0 pt./A of Banvel with 1.33 pts./A of AX S-MET on medium soils, or with 1.33-1.67 pts./A of AX S-MET on fine soils. Do not apply on coarse soils or on soils with less than 2.5% organic matter. Apply this tank mixture to the soil surface at planting or after planting, but before corn emerges. Plant corn at least 1.5 inches deep and apply behind planting equipment, avoiding incorporation by the planter wheel or other seed covering device. Do not incorporate before corn emergence. If it is necessary to rotary hoe to break the soil crust, do not disturb the soil more than 1/2 inch deep.

**Postemergence for Control of Pigweed (Mid-Atlantic states, including DE, MD, PA, VA, and WV):** Apply 1.0-1.5 pts. of AX S-MET + 0.5-1.0 pt./A of Banvel or Clarity® by ground equipment when pigweed plants are less than 3 inches tall and before corn exceeds 5 inches in height in a minimum of 20 gals. of spray per acre. Use the lower rate on coarse-textured and low organic matter soils. Use the higher rate on fine-textured and high organic matter soils.

**Precaution:** TO AVOID CROP INJURY, Avoid drift to sensitive nontarget plants, such as soybeans, during application.

**Restriction:** Do not apply with aircraft.

**TANK MIXTURE WITH AATREX OR PRINCEP + PROWL FOR PROLONGED CONTROL OF LAMBSQUARTERS AND PIGWEED IN FIELD CORN ONLY (NORTHEAST U.S., INCLUDING MI, IN, KY, AND STATES EAST OF THESE)**

For prolonged control of lambsquarters and pigweed, in addition to a broad spectrum of annual broadleaf and grass weeds, AX S-MET in tank mix combination with AAtrex\* or Princep + Prowl 4E may be applied after planting, but before corn or weeds emerge. Apply by ground equipment in a minimum of 10 gals. of water or 20 gals. of liquid fertilizer. Apply by air in a minimum of 5.0 gals. of water. Refer to Table 3 of this label for rates of AX S-MET, AAtrex, or Princep to be applied. Apply Prowl 4E according to the following rates in Table 4.

**Restriction:** \*Do not apply AX S-MET in tank mix combination with AAtrex 80W + Prowl, as this combination is not compatible. Other AAtrex formulations may be used.

**Mixing Instructions:** See Comment No.1 following Table 2.

**Table 4: Prowl 4E - Broadcast Rates Per Acre**  
**Percent Organic Matter in Soil**

| Soil Texture | Less Than 1.5% | 1.5-3%   | Over 3%  |
|--------------|----------------|----------|----------|
| Coarse       | 1.5-2.0 pts.   | 2.0 pts. | 3.0 pts. |
| Medium       | 2.0 pts.       | 3.0 pts. | 3.0 pts. |
| Fine         | 2.0 pts.       | 3.0 pts. | 3.0 pts. |

Observe all directions for use, precautions, and limitations on the respective product labels when applying these products in tank mix combination. Refer to the Prowl 4E label for replanting instructions in the event of crop loss.

**TANK MIXTURE OF AX S-MET WITH AATREX OR PRINCEP, OR AATREX + PRINCEP WITH GRAMOXONE INTEON, LANDMASTER BW, TOUCHDOWN OR ROUNDUP FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS**

In minimum-tillage or no-tillage systems where corn is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides Gramoxone Inteon,

26/61

Landmaster BW, Touchdown brands or Roundup brands should be tank mixed with AX S-MET + AAtrex, AX S-MET + Princep, or AX S-MET + AAtrex + Princep. See Comment No. 7 following Table 2. The AX S-MET, AX S-MET + AAtrex or Princep, or AX S-MET + AAtrex + Princep portion of the tank mixture provides preemergence control of the weeds listed on this label in the tank mixture section for **AX S-MET, AX S-MET + AAtrex or Princep, or AX S-MET + AAtrex + Princep - Preplant Surface, Preplant Incorporated, or Preemergence.**

**Application:** Apply before, during, or after planting, but before the corn emerges. Add Gramoxone Inteon, Landmaster BW, Touchdown brands or Roundup brands and apply as directed on the product label.

**Gramoxone Inteon:** Apply as directed on the product label. This treatment will not control weeds taller than 6 inches.

Do not apply combinations containing Gramoxone Inteon in suspension-type liquid fertilizers, because the activity of paraquat will be reduced.

**Landmaster BW:** 27-54 oz./A depending on weed species and size. See the Landmaster BW label for weeds controlled, listed rates for specific weeds, and other information concerning use.

**Touchdown Brands or Roundup Brands:** See the Touchdown brand or Roundup brand labels for weeds controlled, listed rates, and other use directions.

Apply in 20-60 gals. of water or fluid fertilizer per acre with ground equipment.

On coarse soils, apply 1.0 pt./A of AX S-MET with 1.3 lbs. of AAtrex Nine-O\* or Princep Caliber 90\*, or with 0.7 lb. of AAtrex Nine-O\*\* + 0.7 lb. of Princep Caliber 90\*\*. On medium soils, apply 1.33 pts./A of AX S-MET with 1.8 lbs. of AAtrex Nine-O or Princep Caliber 90, or with 0.9 lb. of AAtrex Nine-O + 0.9 lb. of Princep Caliber 90. On fine soils\*\*\*, apply 1.33-1.67 pts./A of AX S-MET with 1.8-2.2 lbs. of AAtrex Nine-O or Princep Caliber 90, or with 0.9-1.1 lbs. of AAtrex Nine-O + 0.9-1.1 lbs. of Princep Caliber 90.

\* Use Princep in preference to AAtrex when heavy infestations of crabgrass or fall panicum are expected.

\*\* When using the tank mixture of AX S-MET + AAtrex Nine-O + Princep Caliber 90, use equal rates of AAtrex and Princep as shown when heavy broadleaf weed infestations are expected. When heavy infestations of crabgrass or fall panicum are expected, use a 1:2 ratio of AAtrex + Princep instead of the 1:1 ratio given.

(Example: Total AAtrex Nine-O + Princep Caliber 90 = 1.8 lbs./A, use 0.6 lb. of AAtrex + 1.2 lbs. of Princep, respectively.) Refer to Comment No: 2 following Table 2 for AAtrex 4L and Princep 4L conversions.

\*\*\*For cocklebur, yellow nutsedge, and velvetleaf control on fine-textured soils above 3% organic matter, apply 2.25 lbs./A of AAtrex Nine-O, or equivalent rate of AAtrex 4L, or the same total amount of AAtrex + Princep, with 1.33-1.67 pts./A of AX S-MET.

**TANK MIXTURE WITH AATREX; OR AATREX + 2,4-D; OR AATREX + 2,4-D + BANVEL FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS**

In minimum-tillage or no-tillage systems where corn is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, AX S-MET applied in combination with AAtrex will kill most emerged small annual weeds. Apply AX S-MET + AAtrex before, during, or after planting, but before corn emerges, according to the rates in Table 3.

Where heavy crop residues exist, add 0.8-1.6 pts./A of an appropriately labeled 3.8 lbs. a.i./gal. of 2,4-D amine (such as Weedar 64, Weedar 64A, DMA-4 Herbicide, Weedone® 638, or Formula 40) to the spray tank last and apply in a minimum of 25 gals. of carrier per acre.

As carriers, nitrogen solutions and complete liquid fertilizers, applied before corn emergence, enhance burn down of existing weeds, and therefore are recommended instead of water. Add a surfactant such as Voyager 90/10 or Voyager 80/20 surfactant or another appropriate surfactant at its specified rate, or add crop oil concentrate plus 28% liquid nitrogen (or equivalent). Apply before weeds exceed 3 inches in height.

If alfalfa is present, add Banvel to the spray mixture at 0.33-0.5 pt./A and apply before alfalfa exceeds 6 inches in height.

For fields with existing sod grasses (e.g., bromegrass, orchardgrass, rye, or timothy), when existing weeds exceed 3 inches in height or when very dry conditions exist, add Gramoxone Inteon at the rate indicated on the product label in place of or in addition to 2,4-D as indicated above. Do not apply Gramoxone Inteon in suspension-type liquid fertilizer. Observe all directions for use, precautions, and limitations on the respective product labels when applying these products in tank mix combination. Use Balance combinations only on field corn.

**TANK MIXTURE WITH MARKSMAN IN CONSERVATION TILLAGE - FIELD AND SILAGE CORN**

In conservation tillage systems where corn is planted directly into a cover crop or previous crop residue, AX S-MET + Marksman will kill most emerged small annual weeds. Apply AX S-MET + Marksman before, during, or after planting, but before corn emergence on medium and fine soils with greater than 2.5% organic matter. For fields with existing vegetation exceeding 3 inches in height or when very dry conditions exist, add Gramoxone Inteon at its standard rate. AX S-MET + Marksman may be applied postemergence to corn less than 3 inches tall and before weedy grasses exceed the 2-leaf stage.

As carriers, nitrogen solutions and complete liquid fertilizers, applied before corn emergence, enhance burn down of existing weeds. Do not apply Gramoxone Inteon in suspension-type liquid fertilizer or use on emerged corn.

Refer to the Marksman label and follow all directions, limitations, precautions, and information regarding application and use in corn.

**TANK MIXTURE WITH BALANCE PRO - FIELD CORN ONLY**

AX S-MET and Balance PRO have a complementary crop response and weed control profile which allows various tank mix rate combinations to be considered. The addition of Balance PRO will improve the control of certain problem weeds including Texas panicum, woolly cupgrass, and wild proso millet. AX S-MET improves both the duration and spectrum of annual grass and small seeded broadleaf weed control, in particular foxtails (yellow foxtail), witchgrass, and yellow nutsedge.

**Restriction:** The Balance PRO label does not allow applications to coarse textured soils with less than 1.5% organic matter and warns about applications to all soils with less than 1.5% organic matter or with pH greater than 7.5, as well as applications made to areas in fields with clay knolls, eroded hillsides, and exposed subsoil.

Listed below are compensating rate options for combinations of AX S-MET and Balance PRO, i.e. higher rates of AX S-MET are combined with lower rates of Balance PRO, and vice versa. Select a rate option for AX S-MET plus Balance PRO by weighing the intensity of problem weed pressure (population presence and density) and your acceptance for risk of an adverse crop response. For

example, where Texas panicum, woolly cupgrass, or wild proso millet are a primary target weed, use a tank mix combination with a higher Balance PRO rate for the given soil type.

Where your acceptance of an adverse crop response risk is low and/or a more general weed spectrum is targeted (especially yellow foxtail, witchgrass or yellow nutsedge), use a tank mix combination with a higher AX S-MET rate for the given soil type. Where a target weed is listed as controlled on both product labels, a tank mix combination option including intermediate rates of both products may be used. Where a target weed is listed as controlled on only one product label, do not apply a rate of that product below what is listed for that weed on the individual product label, or unacceptable control may result. Follow all other directions for use, rate limitations, precautions and restrictions on both the AX S-MET and Balance PRO product labels.

AX S-MET plus Balance PRO tank mix rate options when applied pre-plant (incorporated or surface applied) up to 7 days before planting or preemergence in field corn:

**For coarse textured soils**, where 1.5 or 1.88 oz./A of Balance PRO is used, 1.0-1.33 pts./A of AX S-MET may be applied. Do not use Balance PRO on coarse textured soils with less than 1.5% organic matter.

**For medium textured soils**, where 1.5 oz./A of Balance PRO is used, rates as low as 1.33 pts./A of AX S-MET may be applied. Where 1.88 or 2.25 oz./A of Balance PRO is used, rates as low as 1.0 pts./A of AX S-MET may be applied. AX S-MET can be used in combinations with Balance PRO at rates up to 1.67 pts./A on medium textured soils.

**For fine textured soils**, where 1.5 oz./A of Balance PRO is used, rates as low as 1.33 pts./A of AX S-MET may be applied if the soil organic matter is less than 3% - if the soil organic matter content is 3% or greater, 1.67 pts./A of AX S-MET should be applied. Where 1.88 or 2.25 oz./A of Balance PRO is used, rates as low as 1.33 pts./A of AX S-MET may be applied. Where 3.0 oz./A or more of Balance PRO are used, rates as low as 1.0 pts./A of AX S-MET may be applied. AX S-MET can be used in combinations with Balance PRO at rates up to 2.0 pts./A on fine textured soils if the soil organic matter content is 3% or greater.

**TANK MIXTURES FOR POSTEMERGENCE SALVAGE WEED CONTROL IN FIELD CORN ONLY**

For postemergence control of weeds in specific types of field corn, the AX S-MET combinations listed below may be used. Full season weed control from early preplant, preplant incorporated or preemergence treatments can lead to maximum yield potential under competition-free conditions. However, if control of emerged weeds is needed, a postemergence program listed below can be applied to provide residual control for the remainder of the season.

- (1) Follow all label directions, instructions, precautions, and limitations for each product used. (2) Do not use fluid fertilizer with these mixtures or corn injury may occur. (3) For each tank mixture with AX S-MET, apply only to the specific field corn type specified on the tank mix product label. (4) In-row weed control may be reduced because of lack of coverage when applied to corn over 4 inches tall.

**AX S-MET + Liberty Herbicide or Ignite® 280 SL Herbicide: Postemergence Use in LibertyLink® Corn or Corn Warranted by Bayer CropScience as Being Tolerant to Liberty Herbicide or Ignite 280 SL Herbicide**

These tank mixtures can be applied postemergence to weeds and corn from seed designated as LibertyLink or corn warranted by Bayer CropScience as being tolerant to Liberty Herbicide or Ignite 280 SL Herbicide. Liberty provides postemergence control of a broad spectrum of grass and broadleaf weeds and the AX S-MET provides residual control of grasses and certain broadleaf weeds listed in the label section **AX S-MET Applied Alone - Weeds Controlled**. Refer to the **AX S-MET Alone - Preplant Incorporated or Preemergence** section and use the minimum rate per

soil texture and organic matter classification for season-long residual control from this tank mix combination with Liberty. Refer to the Liberty Herbicide or Ignite 280 SL Herbicide labels for the postemergence application rates according to weed species and their maximum height at the time of postemergence application. Where multiple weed species are present, use the highest Liberty rate listed to control the species and growth stages present.

Follow all applicable use directions, limitations, precautions, and information regarding application to corn on the AX S-MET, Liberty Herbicide, or Ignite 280 SL Herbicide labels. Where difficult species and/or severe weed populations are expected, use the maximum rate where rate ranges are listed.

**AX S-MET + Touchdown Brands or Roundup Brands for Postemergence Application to Glyphosate Tolerant Corn (e.g., Roundup Ready® or Agrisure™ GT)**

The tank mixture of AX S-MET + Touchdown or Roundup brands can be applied postemergence to weeds and to corn designated as glyphosate tolerant. Application may be applied postemergence to glyphosate tolerant corn from emergence until corn reaches 30 inches tall or the V8 stage (8 leaves with collars), whichever comes first. This mixture will provide postemergence control of weed species on the Touchdown brand or Roundup brand label and residual control of weed species on the AX S-MET label. Use the minimum AX S-MET rate postemergence with Touchdown or Roundup in glyphosate tolerant corn as specified in the **Corn - AX S-MET Alone - Preplant Incorporated or Preemergence** section of this label according to soil texture and organic matter. Refer to the Touchdown brand or Roundup brand label and follow appropriate use directions, application procedures, precautions, and limitations. Refer to the Touchdown brand or Roundup brand label for directions for control of problem species.

Follow all applicable use directions, limitations, precautions, and information regarding application to corn on the AX S-MET and Touchdown brand or Roundup Ultra brand labels, and on the **Supplemental Labeling of Roundup Ultra for Postemergence Application to Corn with the Roundup Ready Gene**. Where difficult species and/or severe weed populations are expected, use the maximum rate where rate ranges are listed.

**AX S-MET + Touchdown Brands or Roundup Brands + AAtrex for Postemergence Application to Glyphosate Tolerant Corn (i.e., Roundup Ready or Agrisure GT)**

The tank mixture of AX S-MET + AAtrex + Touchdown brands or Roundup brands can be applied postemergence to weeds and to corn designated as glyphosate tolerant. Application may be applied postemergence to glyphosate tolerant corn from emergence up to 12 inches in height. This mixture will provide postemergence control of weed species on the Touchdown brand or Roundup brand label and residual control of weed species on the AX S-MET + AAtrex label. Use the minimum AX S-MET + AAtrex rate postemergence with Touchdown or Roundup in glyphosate tolerant corn as specified in the **Corn - AX S-MET Combinations - Tank Mixture With AAtrex or Princep, or AAtrex + Princep - Preplant Incorporated or Preemergence** section and Table 3 of this label according to soil texture and organic matter.

Follow all applicable use directions, limitations, precautions, and information regarding application to corn on the AX S-MET, AAtrex, and Touchdown brand or Roundup brand labels for application to glyphosate tolerant corn. Where difficult species and/or severe weed populations are expected, use the maximum rate where rate ranges are listed.

**COTTON - AX S-MET ALONE**

**Application:** Apply AX S-MET preemergence only in Area 1\* at the rate of 0.5-1.0 pt./A on sandy loams, 0.66-1.33 pts./A on medium soils, or 1.0-1.33 pts./A on fine soils. Apply AX S-MET preplant incorporated or preemergence in Area 2\*\* at 1.0 pt./A on sandy loams, 1.0-1.33 pts./A on medium soils, or 1.33 pts./A on fine soils. Apply AX S-MET postemergence to cotton and

preemergence to weeds at 0.5-1.33 pts./A, according to the state rate limitations in the following **Postemergence** section. **Do not use on sands and loamy sand.**

\*Area 1 = AR, KS, LA, MS, TN, and Bootheel of MO

\*\*Area 2 = NM, OK, and TX

**Fall Application for Italian Ryegrass Control:** AX S-MET may be applied for residual control of glyphosate-resistant Italian ryegrass (*Lolium multiflorum*). Apply AX S-MET at 1.33-1.67 pints/Acre in the fall (September 1 - December 1) after harvest of the previous crop and prior to Italian ryegrass emergence. Use the lower AX S-MET rate for coarse textured soils and the higher rate for fine textured soils. A tillage operation may precede the application. Do not incorporate to a depth greater than 2-3 inches if tillage follows the application of AX S-MET. For fall applications after emergence of glyphosate-resistant Italian ryegrass, Gramoxone Inteon can be tank mixed with AX S-MET to control emerged ryegrass. Refer to the Gramoxone Inteon label for specific rates, application instructions and restrictions. Other registered herbicides may be tank mixed with AX S-MET for control or improved control of other weeds present at the time of application.

**Restrictions:** (1) Do not apply AX S-MET to frozen ground. (2) If a spring application is made, the combined total amount of AX S-MET applied in the fall plus the spring must not exceed the maximum seasonal S-metolachlor rate for cotton (2.6 pt/A, depending on soil texture).

**Preplant Incorporated (NM, OK, and TX Only):** Apply to the soil and incorporate into the top inch of soil immediately before planting, at planting, or after planting, but before crop or weeds emerge. Use a rolling cultivator or similar implement to uniformly incorporate not more than 1 inch deep. Use a preplant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. Where furrow irrigation is used, wet the top of the bed for best results. If the crop is to be planted on beds, apply and incorporate after bed formation. Cotton should be planted below the zone of incorporation; i.e., at least 1 inch on fine soils and 1.5 inches on coarse and medium soils. If incorporated prior to planting, use a planter that will result in a minimum of soil disturbance.

**Note:** For best control of yellow nutsedge and suppression of seedling johnsongrass, apply AX S-MET preplant incorporated at the maximum rate for the soil texture, whether applied alone or mixed with Caparol 4L.

**Preemergence:** Apply to the soil surface at planting or after planting, but before weeds or crop emerge.

**Postemergence:** Apply AX S-MET broadcast over-the-top or directed to the soil surface according to the rate limitations listed below by state. Over-the-top postemergence application may be made not later than 100 days before harvest, and directed-postemergence application may be made not later than 80 days before harvest. Application before weeds emerge or after clean cultivation to remove existing weeds is necessary since AX S-MET will not control emerged weeds. AX S-MET postemergence may be applied over any previous registered herbicide treatment. In sprinkler-irrigated areas, sprinkler irrigate after application with ½-1 inch of water (½ inch on coarse-textured soils to 1 inch on fine-textured soils) to incorporate AX S-MET. In furrow-irrigated areas, apply AX S-MET, incorporate with a rolling cultivator or similar implement that provides uniform shallow incorporation (2 inches or less), and then irrigate. In nonirrigated areas, if at least ½ inch of rainfall does not occur within 10 days after application, cultivate with a rolling cultivator or similar implement that provides uniform shallow incorporation of AX S-MET.

**VA, NC, SC, GA, FL, and AL:** Apply AX S-MET postemergence at 1.0-1.33 pts./A.

**TN, AR, KS, MS, MO, and LA:** Apply AX S-MET postemergence at 0.5-1.33 pts./A.

**TX, OK, NM, AZ, CA, and Clay Soils in AR:** Apply AX S-MET postemergence at 1.0-1.33 pts./A before August 1.

**Multiple Applications:** Where weed pressure is heavy, difficult to control species are expected, or reinfestation may occur, and a weed control program is used, multiple applications of AX S-MET are effective when used as part of the weed control program. Apply as a preplant incorporated or preemergence treatment and follow with an application postemergence to cotton before weeds emerge or after clean cultivation to remove existing weeds, since AX S-MET will not control emerged weeds. Apply AX S-MET postemergence over a previous preplant or preemergence AX S-MET application as shown in the following table.

**Multiple AX S-MET Applications to Cotton**

| State                  | Preplant Incorporated or Preemergence Pts./A | + | Postemergence Pts./A     |
|------------------------|--|---|--------------------------|
| MS, LA, TN, AR, KS, MO | 0.5-1.33 (Preemergence Only)                 | + | 0.5-1.33                 |
| TX, OK, NM             | 1.0-1.33                                     | + | 1.0-1.33 before August 1 |
| NC, VA                 | 1.0-1.33 (Preemergence Only)                 | + | 1.0-1.33                 |

In sprinkler-irrigated areas, sprinkler irrigate after application with 1/2-1 inch of water (1/2 inch on coarse-textured soils to 1 inch on fine-textured soils) to incorporate AX S-MET. In furrow-irrigated areas, apply AX S-MET, incorporate with a rolling cultivator or similar implement that provides uniform shallow incorporation (2 inches or less), and then irrigate. In non-irrigated areas, if at least 1/2 inch of rainfall does not occur within 10 days after application, cultivate with a rolling cultivator or similar implement that provides uniform shallow incorporation of AX S-MET.

**Notes:** For best control of yellow nutsedge and suppression of seedling johnsongrass, apply AX S-MET preplant incorporated, preemergence, or postemergence to cotton and preemergence to weeds at the maximum rate for the soil texture, whether applied alone or in combinations. Do not apply more than a total of 2.0 pts./A on coarse soils or 2.6 pts./A of AX S-MET on medium and fine soils during a growing season. These treatments may be applied over previous registered herbicide treatments.

**Restrictions:** (1) Do not apply on Taloka silt loam. (2) Do not use in Gaines County, TX. (3) Do not graze or feed forage or fodder from cotton to livestock.

**Precautions:** TO AVOID CROP INJURY (1) Do not apply AX S-MET on sand or loamy sand soils, or in areas where water is likely to "pond" over the bed; (2) To avoid concentration in the seed furrow, do not make broadcast applications of AX S-MET to cotton planted in furrows more than 2 inches deep. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width must not exceed the width of the bottom of the furrow; (3) In furrow-planted cotton, to avoid concentration in the furrow and potential injury, do not apply AX S-MET postemergence until after first "knifing" or cultivation to level soil surface. (4) Do not apply over-the-top in fluid fertilizer or any other adjuvant, surfactant, oil, or other pesticide not listed in the cotton section of this label.

**COTTON - DUAL MAGNUM COMBINATIONS**

**TANK MIXTURE WITH CAPAROL 4L**

AX S-MET tank mixtures with Caparol 4L may be applied preplant incorporated or preemergence in water or fluid fertilizer. When fluid fertilizer is used as a carrier for AX S-MET, either alone or in combination with Caparol 4L, mix only the amount that will be sprayed in one operation. These mixtures should not be allowed to stand without agitation. Only water may be used as a carrier for postemergence-directed application.

In addition to those weeds controlled by AX S-MET alone, AX S-MET + Caparol 4L, applied preplant incorporated or preemergence also controls the following weeds: junglerice, wild oats, annual morningglory, groundcherry, hairy nightshade, lambsquarters, malva, mustard, prickly sida (teaweed), purslane, ragweed, and shallow-germinating seedlings of cocklebur and coffeeweed. As a postemergence-directed application, Caparol provides postemergence control and residual control of weeds on its label, while AX S-MET provides residual control of weed species on its label. AX S-MET will not control emerged weeds.

**Preplant Incorporated or Preemergence:** Apply AX S-MET + Caparol 4L, either preplant incorporated or preemergence, using the appropriate rate from Table 5. Cotton should be planted below the zone of incorporation; i.e., at least 1.0 inch on fine soils and 1.5 inches on coarse and medium soils. If incorporated before planting, use a planter that will result in a minimum of soil disturbance.

**Table 5: AX S-MET + Caparol 4L - Cotton (NM, OK, TX)**

**Broadcast Rates Per Acre**

| Use Areas   | Soil Texture     | AX S-MET      | Caparol 4L |
|---|------------------|---------------|------------|
| ALL   | Sand, loamy sand | DO NOT USE    | DO NOT USE |
| OK, and Blacklands and Gulf Coast of TX                                   | Loams            | 0.8-1.33 pts. | 2.4 pts.   |
|   | Clays            | 1.33 pts.     | 4.8 pts.   |
| Rio Grande Valley of TX   | Loams            | 0.8-1.33 pts. | 3.2 pts.   |
|   | Clays            | 1.33 pts.     | 4.8 pts.   |
| NM; High Plains, Rollings Plains, Edwards Plateau of TX; and Southwest TX | Sandy loam       | 0.8-1.0 pt.   | 1.6 pts.   |
|   | Loams            | 0.8-1.33 pts. | 2.4 pts.   |
|   | Sandy clay loams | 1.33 pts.     | 2.4 pts.   |
|   | Other Clay soils | 1.33 pts.     | 3.2 pts.   |

**Postemergence-Directed (AR, AZ, CA, LA, MS, NM, OK, TN, TX, and MO):** AX S-MET may be tank mixed with Caparol 4L in water and applied postemergence-directed in cotton for control of emerged weeds listed on the Caparol 4L label and residual preemergence control of weeds controlled by AX S-MET and Caparol 4L. Or application may be made after cultivation for residual preemergence control. These treatments may be applied over previous registered treatments, including AX S-MET, provided the maximum label rate of any product is not exceeded. Do not apply over-the-top of cotton, or injury may occur.

Apply AX S-MET + Caparol 4L in a minimum of 20 gals. of spray volume per acre. Follow the directions, limitations, and precautions on the Caparol 4L label when Caparol is applied as a postemergence-directed application. Refer to the directions, limitations, and precautions for use of AX S-MET under the **Cotton - AX S-MET Alone – Postemergence** section.

**Precautions:** TO AVOID CROP INJURY; (1) Do not apply on sand or loamy sand soils, or in areas where water is likely to "pond" over the bed; (2) Do not apply in cut areas of newly leveled fields, or in areas of excess salt; (3) Do not apply to glandless cotton varieties; (4) To avoid concentration in the seed furrow, do not make broadcast applications of AX S-MET + Caparol 4L to cotton planted in furrows more than 2 inches deep. Band applications may be made to cotton



planted in furrows deeper than 2 inches, but band width must not exceed the width of the bottom of the furrow.

**Restrictions:** (1) Do not apply on Taloka silt loam. (2) Do not use in Gaines County, TX. (3) Do not graze or feed forage or fodder from cotton to livestock.

Refer to the Caparol 4L label for further instructions and limitations.

**TANK MIXTURE WITH COTORAN DF**

AX S-MET may be applied in tank mixture with Cotoran DF preemergence for control of those weeds controlled by AX S-MET alone and those as listed on the Cotoran DF label. This combination will also control spotted spurge, hyssop spurge, nodding spurge, and prostrate spurge. Apply to the soil surface at planting or after planting, but before weeds or crops emerge, using the appropriate rates from Table 6. The tank mixture may be applied postemergence to cotton, but preemergence to weeds, or it may be applied postemergence to both cotton and broadleaf weeds for control of weeds on the Cotoran label. Apply as a directed, semi-directed, or over-the-top spray AX S-MET will not control emerged weeds, but will provide preemergence control of species on its label.

**Mixing Instructions:** Incompatibility may occur when tank mixing AX S-MET and Cotoran DF. To help overcome this condition, fill the spray tank 1/4 full with water or fluid fertilizer and start agitation, add the Cotoran DF and allow it to become dispersed. Add a surfactant such as VOYAGER 90/10 OR VOYAGER 80/20 at 0.5% volume/volume final spray (4.0 pts./100 gals.), then add the AX S-MET and finally the rest of the water or fluid fertilizer. Agitate during mixing and application to maintain a uniform suspension. Do not use fluid fertilizer as a carrier for postemergence applications.

**Table 6: AX S-MET + Cotoran DF - Cotton**  
**Broadcast Rates Per Acre**

|                       | AX S-MET (pts.) | AX S-MET (pts.) | Cotoran DF*** (lbs.) |
|-----------------------|-----------------|-----------------|----------------------|
| Soil Texture          | Area 1*         | Area 2**        |                      |
| Sand, loamy sand      | DO NOT USE      | DO NOT USE      | DO NOT USE           |
| Sandy loam            | 0.5-1.0         | 0.8-1.0         | 1.2                  |
| Loam, silt loam, silt | 0.66-1.33       | 1.0-1.33        | 1.2-1.9              |
| Fine soil             | 1.0-1.33        | 1.33            | 1.9-2.4              |

\*Area 1=AR, LA, MS, Bootheel of MO and TN

\*\*Area 2=Eastern OK, Gulf Coast, Rio Grande Valley, and Eastern TX

\*\*\*When using Cotoran 4L, use equivalent rates. Multiply lbs. of Cotoran DF by 1.7 to get pts. of Cotoran 4L.

**Postemergence:** This tank mixture may be applied postemergence to cotton, but preemergence to weeds or postemergence to both cotton and weeds for control of weeds on the Cotoran label. Apply as a directed, semi-directed, or over-the-top spray. AX S-MET will not control emerged weeds, but will provide preemergence control of species on its label. Where rate ranges are given for Cotoran DF, use the higher rate when applying postemergence to weeds that are 2 inches or less. These treatments may be applied over previous registered treatments, including AX S-MET, provided the maximum label rate of any product is not exceeded.

34/61

**Precautions:** (1) Do not apply AX S-MET + Cotoran on sand or loamy sand soils, or in areas where water is likely to "pond" over the bed, or crop injury may occur. (2) To avoid concentration in the seed furrow, do not make broadcast applications of AX S-MET + Cotoran to cotton planted in furrows more than 2 inches deep. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width must not exceed the width of the bottom of the furrow. (3) The use of Cotoran following the use of a systemic insecticide at planting may result in crop injury.

**Restrictions:** (1) Do not feed treated forage or gin trash to livestock, or graze treated areas. (2) Do not use on Taloka silt loam, or crop injury may occur. (3) Do not use in Gaines County, TX;

Refer to the Cotoran labels for further instructions, precautions, and limitations.

**TANK MIXTURE OF AX S-MET OR AX S-MET + COTORAN WITH GRAMOXONE INTEON, TOUCHDOWN BRANDS OR ROUNDUP BRANDS FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS**

In minimum-tillage or no-tillage systems where cotton is planted directly into a cover crop, stale seedbed, or previous crop residues, the contact herbicides Gramoxone Inteon, Touchdown brands or Roundup brands may be added to a tank mix of either AX S-MET or AX S-MET + Cotoran. When used as directed, the Gramoxone Inteon portion of the tank mixture controls most emerged weeds and suppresses many perennial weeds. Touchdown or Roundup combinations will control emerged annual and perennial weeds when applied as directed on the Touchdown or Roundup label. The AX S-MET and AX S-MET + Cotoran portion of the tank mixture provides preemergence control of the weeds listed on this label in the AX S-MET and AX S-MET + Cotoran sections, respectively.

Refer to the label of each product used in combination and observe the planting details, information regarding application, geographical restrictions, and all other precautions and limitations. Refer to **Mixing Instructions under Tank Mixture with Cotoran DF** section.

**Application:** Apply before, during, or after planting, but before the cotton emerges. Apply AX S-MET at 0.8-1.0 pt./A on sandy loams, medium-, and fine-textured soils. Refer to Table 6 for the Cotoran DF rates.

**Gramoxone Inteon:** Apply as directed on the product label. This treatment will not control weeds taller than 6 inches.

Do not apply combinations containing Gramoxone Inteon in suspension-type liquid fertilizers, as the activity of paraquat will be reduced.

**Touchdown Brands or Roundup Brands:** See the Touchdown or Roundup label for weeds controlled, use rates, and other use directions.

Do not apply AX S-MET + Cotoran 4L + Roundup in tank mixture because of compatibility problems.

Apply in 20-60 gals. of water or fluid fertilizer per acre with ground equipment.

**Precautions:** (1) If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stands for several days, or where the seeding slit has not been properly closed. (2) Refer to the Cotoran labels and the **Tank Mixture with Cotoran DF** section of this label for further instructions, precautions, and limitations.

**Restriction:** Do not use in Gaines County, TX.

**TANK MIXTURE WITH MSMA, MSMA + CAPAROL, OR MSMA + COTORAN**

AX S-MET may be tank mixed with MSMA in water and applied postemergence directed for control of emerged weeds listed on the MSMA product label and residual preemergence control of weeds controlled by AX S-MET. The addition of Caparol or Cotoran will add control of weed species on their respective labels.

**Postemergence-Directed (AL, AR, AZ, CA, FL, GA, LA, MS, NC, NM, OK, SC, TN, TX, VA, and Bootheel of MO):** Apply AX S-MET + MSMA postemergence-directed to cotton at least 3 inches tall according to the directions, limitations, and precautions on the MSMA product label, as well as the directions, limitations, and precautions for use of AX S-MET in the section for **Cotton - AX S-MET Alone - Postemergence**. Do not apply after first cotton bloom. These treatments may be applied over previous registered treatments, including AX S-MET, provided the maximum label rate of any product is not exceeded. Cotoran or Caparol may be added to the AX S-MET + MSMA tank mixture according to the respective label directions for application to cotton at least 3 inches tall. When these mixtures are used, follow the mixing instructions for AX S-MET + Caparol or Cotoran and then add the MSMA product.

**Restriction:** Do not use AX S-MET in tank mix with premixes of MSMA plus herbicides other than those registered for use in tank mixture with AX S-MET on cotton.

**TANK MIXTURE WITH TREFLAN FOR POST-DIRECTED FOLLOWED BY SOIL INCORPORATION APPLICATIONS**

AX S-MET may be applied as a tank mixture with Treflan in cotton for improved late season weed control when used as an incorporated layby type application. This combination may be applied after the cotton is at least 3 inches tall and has reached the 4 true-leaf stage. Make the application directed to the soil surface and away from the crop foliage. Incorporate using a sweep or rolling type cultivator to provide uniform and shallow mixing into the top 2 inches of soil. Refer to each product label for the appropriate application rates by soil type and for this application timing -- and follow all product use limitations and restrictions.

**TANK MIXTURE WITH TOUCHDOWN BRANDS OR ROUNDUP BRANDS FOR USE ON ROUNDUP READY COTTON ONLY**

Apply AX S-MET as a tank mixture with Touchdown or Roundup in water postemergence over-the-top or postemergence-directed for control of emerged weeds listed on the Touchdown or Roundup labels and for residual preemergence control of weeds listed on the AX S-MET label. See the **Cotton - AX S-MET Alone - Postemergence** section of this label for rates and timings of AX S-MET and follow the Touchdown or Roundup label for their respective rates, application method, and application timing restrictions. Do not add additional spray adjuvants, surfactants, fertilizer additives, or pesticides to this tank mixture if applied postemergence over-the-top or unacceptable injury may occur. Refer to the Touchdown brand or Roundup brand label and follow appropriate use directions, application procedures, precautions, and limitation.

**Precaution:** Postemergence over-the-top applications of this tank mixture may cause temporary injury in the form of necrotic spotting to exposed cotton leaves, which will not affect normal plant development.

**Restrictions:** (1) Do not apply this tank mixture postemergence to any cotton variety unless it is designated Roundup Ready and unless the Touchdown or Roundup formulation is used is registered for postemergence use in Roundup Ready Cotton. (2) Do not apply Touchdown or Roundup postemergence over the-top to cotton past the growth stage limit specified on their respective labels. (3) Do not use on sand or loamy sand soils in Gaines County, TX.

36/61

**SOYBEAN, IMMATURE SEED**

AX S-MET may be applied preplant or pre-emergence for the control or suppression of grass and small seeded broadleaf weeds in soybean, immature seed or other food grade soybeans. For specific rates, see the rate table listed below.

**Preplant Surface-Applied:** For minimum-tillage or no-tillage systems only, AX S-MET alone may be applied up to 45 days before planting. Use only split applications for treatments made 30-45 days before planting, with 2/3 the recommended broadcast rate for the crop and soil texture applied initially and the remaining 1/3 applied at planting. Treatments less than 30 days before planting may be made either as a split or a single application. If weeds are present at the time of treatment, apply in a tank mixture combination with a contact herbicide (for example, Gramoxone Inteon, Touchdown, or Roundup). Observe directions for use, precautions, and restrictions on the label of the contact herbicide. To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

**Preplant Incorporated:** Apply AX S-MET to the soil and incorporate into the top 2 inches of soil within 14 days before planting; using a finishing disk, harrow, rolling cultivator, or similar implement capable of providing uniform 2-inch incorporation. Use a preplant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. If crop will be planted on beds, apply and incorporate AX S-MET after bed formation, unless specified otherwise.

**Preemergence:** Apply AX S-MET during planting (behind the planter) or after planting, but before weeds emerge.

**AX S-MET Broadcast Rates Per Acre**

**Percent Organic Matter in Soil**

| Soil Texture | < 3%         | ≥ 3%         |
|--------------|--------------|--------------|
| Coarse       | 1-1.33 pt    | 1.33 pt      |
| Medium       | 1.33-1.67 pt | 1.33-1.67 pt |
| Fine         | 1.33-1.67 pt | 1.67-2.0 pt  |

**Precaution:** AX S-MET will not control emerged weeds.

**Restrictions:** (1) Do not cut for hay within 120 days following a AX S-MET application. (2) Do not use for forage within 60 days following a AX S-MET application. (3) Do not apply more than 2.0 pt/A of AX S-MET during any one crop year.

**GRASSES GROWN FOR SEED (ID, OR, WA) - AX S-MET APPLIED ALONE**

To control weeds and volunteer grasses in established grasses grown for seed, apply AX S-MET to established stands of tall fescue, orchardgrass, perennial ryegrass, fine fescue, bentgrass, and Kentucky bluegrass just before, during, or immediately following the first fall rains or just before or during a late summer or early fall irrigation, but before target grasses emerge. The seed crop must have had one seed harvest or been established at least one year. The postharvest residue (straw) must be evenly spread, removed, or burned before applying AX S-MET. Rainfall or irrigation is required after application and before weed emergence for best control. AX S-MET will provide preemergence control/suppression of volunteer seedlings of perennial ryegrass, fine fescue spp., tall fescue, orchardgrass, bentgrass and Kentucky bluegrass. AX S-MET will control those weed species listed in the **AX S-MET Alone** section of the AX S-MET label and will suppress or control raitail fescue, annual bluegrass, Italian ryegrass, California brome, downy brome, and roughstalk bluegrass.

37/61

Apply AX S-MET by ground equipment in a minimum of 10 gallons of water per acre using the rate listed below according to grass species.

| Established Grass Crop Grown for Seed | Pts./A   |
|---------------------------------------|----------|
| Fine fescue spp.                      | 1.0      |
| Perennial ryegrass                    | 1.0      |
| Bentgrass                             | 1.0-1.33 |
| Kentucky bluegrass                    | 1.0-1.33 |
| Orchardgrass                          | 1.0-1.33 |
| Tall fescue                           | 1.0-1.33 |

**Precautions:** TO AVOID CROP INJURY or POOR CONTROL (1) Do not apply after the 15th of November. (2) Tank mixtures with other pesticides, or the addition of an adjuvant, can increase the risk of crop injury. (3) Application to perennial ryegrass and fine fescue stands under stress may cause crop injury. (4) If weed escapes occur following an AX S-MET application, an application of a postemergence herbicide may be necessary to control escapes. When making such an application, follow all directions, precautions, and limitations on the label of the postemergence herbicide. (5) Control may be decreased if excessive straw from the previous harvest is present at application and/or insufficient rainfall/irrigation occurs.

**Restrictions:** (1) Do not graze forage regrowth for 60 days following application west of the Cascades. (2) In areas east of the Cascades, do not graze forage regrowth for 150 days following application. (3) Hay may be harvested anytime between seed harvest and the next application of S-metolachlor. (4) Apply AX S-MET only once per crop year.

**HORSERADISH**

Apply a single application of AX S-MET at a broadcast rate of 1.0-1.33 pt./A to the soil surface after planting, but before weeds or crop emergence (i.e., preemergence). Use lower rates on soils relatively coarse-textured and higher rates on fine-textured soils. A band application may also be used, applying proportionally less spray mixture on the area actually treated. AX S-MET will not control emerged weeds. Control emerged weeds with an appropriate registered foliar herbicide or by mechanical means. AX S-MET alone may be applied as directed after any of the following preplant incorporated herbicides when used according to their label directions: Balan at 3.0-4.0 qts./A; Treflan E.C. at 1.0 pt./A; Sonalan at 1.25-3.0 pts./A; Pursuit at 0.25 pt./A; or Prowl at 1.0-2.0 pts./A

**Restrictions:** (1) Make only one application of AX S-MET per crop. (2) Do not apply more than 1.33 pt./A of AX S-MET per crop. (3) Harvest horseradish at normal timing.

**PEANUTS - AX S-MET ALONE**

Apply AX S-MET, either preplant incorporated, postplant incorporated, preemergence, or lay-by, using the appropriate rate specified below. **Preplant Incorporated or Preemergence:** Follow instructions for use of AX S-MET alone under **Application Procedures**. AX S-MET alone may be applied as directed after any of the following preplant incorporated herbicides when used according to their label recommendations: Balan at 3.0-4.0 qts./A; Treflan E.C. at 1.0 pt./A; Sonalan at 1.25-3.0 pts./A; Pursuit at 0.25 pt./A; or Prowl at 1.0-2.0 pts./A.

38/61

**Postplant Incorporated:** Apply and shallowly incorporate AX S-MET into the soil after planting, but before peanut germination. Incorporation depth and incorporating implements must be kept above the seed, or seed will be damaged.

**Lay-by:** Apply AX S-MET to the soil immediately after the last normal cultivation.

Apply AX S-MET alone, preplant incorporated, postplant incorporated, preemergence, or lay-by, at a broadcast rate of 1.0-1.33 pts./A in the Southeast\* and 0.8-1.33 pts./A in NM, OK, and TX.

\*In the Southeast, use 1.33-2.0 pts./A and apply preemergence for partial control of Florida beggarweed.

**Restrictions:** (1) Do not graze or feed peanut forage or fodder to livestock for 30 days following application, and (2) Do not apply within 90 days of harvest.

**PEANUTS - AX S-MET COMBINATIONS**

**TANK MIXTURE WITH BALAN L.C.**

AX S-MET + Balan tank mixture applied preplant incorporated controls those weeds listed under **AX S-MET Applied Alone** and those weeds as listed on the Balan label.

Apply 1.0-1.33 pts./A of AX S-MET + 3.0-4.0 qts./A of Balan in a minimum of 10 gals. of spray volume per acre for ground application or in a minimum of 5.0 gals. of spray volume per acre for aerial application. Follow the use directions on the Balan label for soil preparation and incorporation of this tank mix. Apply and incorporate AX S-MET + Balan up to 14 days prior to planting.

Follow all restrictions and precautions on the Balan label.

**Multiple Applications:** Where weed pressure is heavy or where species difficult to control are expected, AX S-MET is most effective when used as follows:

**Southeast Only (AL, FL, GA, NC, SC, VA)**

**Preplant Incorporated:** Apply AX S-MET preplant incorporated as directed under **Peanuts - AX S-MET Alone** or apply AX S-MET + Balan preplant incorporated as directed previously in this section. Refer to the respective section for weeds controlled.

**OR**

**Preemergence before "ground cracking":** Apply AX S-MET any time from preemergence up to "ground cracking" at 1.0-2.0 pts./A for extended control of weeds not yet emerged. Refer to the **AX S-MET Applied Alone** section for a list of weeds controlled.

Follow the PPI or PRE application by:

**Lay-by:** Apply AX S-MET at lay-by as directed under **Peanuts - AX S-MET Alone**. Use only when late germinating weeds are expected to be a problem. Refer to the **AX S-MET Applied Alone** section for a list of weeds controlled.

**Restrictions:** (1) Do not apply more than the equivalent of 2.67 lbs. of active ingredient of AX S-MET per acre during any one year, or illegal residues may result. If Dual II Magnum is used as a sequential treatment, the lbs. of active ingredient (1.0 pt. = 0.95 lb.) plus the lbs. of active ingredient of AX S-MET must not exceed 2.67 lbs. **Do not use Dual II Magnum or Dual IIG**

**Magnum after peanuts have emerged.** (2) Do not graze or feed peanut forage or fodder to livestock for 30 days following application, and (3) Do not apply within 90 days of harvest.

**Southwest Only (NM, OK, TX)**

**1st Application:** Apply AX S-MET preplant incorporated or preemergence or at cracking as directed previously in this section. Refer to the respective section for weeds controlled.

**2nd Application:** Apply AX S-MET at lay-by as directed under **Peanuts – AX S-MET Alone** on that label. Use only when late germinating weeds are expected to be a problem. Refer to the **AX S-MET Applied Alone** section for a list of weeds controlled.

**Restrictions:** (1) Do not apply more than the equivalent of 2.67 lbs. of active ingredient of AX S-MET per acre during any one year. If Dual II Magnum® is used as a sequential treatment, the lbs. of active ingredient (1.0 pt. = 0.95 lb.) plus the lbs. of active ingredient of AX S-MET must not exceed 2.67 lbs. **Do not use Dual II Magnum or Dual IIG Magnum after peanuts have emerged.** (2) Do not graze or feed peanut forage or fodder to livestock for 30 days following application, and (3) Do not apply within 90 days of harvest.

**TANK MIXTURE OR SEQUENTIALLY WITH PURSUIT**

The tank mixture or sequential treatment of AX S-MET and Pursuit controls all weeds controlled by AX S-MET alone and by Pursuit alone. Refer to the **AX S-MET Applied Alone** section for weeds controlled by AX S-MET and to the Pursuit label for weeds controlled by Pursuit.

Refer to the respective labels for application methods, timing, rates, restrictions, and precautions; and use in accordance with the more restrictive label. Do not exceed the label rate of either product. AX S-MET will not control emerged weeds.

**TANK MIXTURE WITH SONALAN**

The tank mixture controls all weeds controlled by AX S-MET alone and by Sonalan alone. Refer to the **AX S-MET Applied Alone** section for weeds controlled by AX S-MET and to the Sonalan label for weeds controlled by Sonalan.

Apply AX S-MET + Sonalan preplant incorporated using the appropriate rate from Table 7. Follow directions for soil preparation procedures for Sonalan.

**Table 7: AX S-MET + Sonalan - Peanuts**

**Broadcast Rates Per Acre**

|               | Southeast     | Southeast     | NM, OK, TX    | NM, OK, TX    |
|---------------|---------------|---------------|---------------|---------------|
| Soil Texture  | AX S-MET      | Sonalan       | AX S-MET      | Sonalan       |
| <b>COARSE</b> | 1.0-1.33 pts. | 1.25-2.0 pts. | 0.8-1.33 pts. | 1.25-2.0 pts. |
| <b>MEDIUM</b> | 1.0-1.33 pts. | 1.75-2.5 pts. | 0.8-1.33 pts. | 1.75-2.5 pts. |
| <b>FINE</b>   | 1.0-1.33 pts. | 2.25-3.0 pts. | 0.8-1.33 pts. | 2.25-3.0 pts. |

Follow all use directions, limitations, precautions, and information regarding application to peanuts on the AX S-MET and Sonalan labels.

**TANK MIXTURE WITH PROWL**

AX S-MET + Prowl applied preplant incorporated controls all weeds controlled by AX S-MET alone plus Texas panicum, field sandbur, johnsongrass from seed, lambsquarters, kochia, annual

40/  
61

spurge, and other species on the Prowl label. Apply AX S-MET + Prowl by ground or by aerial equipment within 14 days before planting. Incorporate into the top 1-2 inches of soil before planting and within 7 days of application, using a finishing disk or similar implement capable of providing uniform incorporation. If, peanuts will be planted on beds, apply and incorporate after bed formation. Refer to the **Incorporation** instructions of the respective labels for additional directions.

Apply AX S-MET + Prowl preplant incorporated, using the appropriate rates from Table 8.

**Table 8: AX S-MET + Prowl - Peanuts**

**Broadcast Rates Per Acre**

|                     | <b>NM, OK, TX</b>       | <b>Other Peanut Growing States</b> |
|---------------------|-------------------------|------------------------------------|
| <b>Soil Texture</b> | <b>AX S-MET + Prowl</b> | <b>AX S-MET + Prowl</b>            |
| Sand, loamy sand    | 0.8 + 1.0-1.5 pts.      | 1.0-1.33 + +1.5-2.0 pts.           |
| Sandy loam          | 0.8-1.0 + 1.0-1.5 pts.  | 1.0-1.33 + 1.5-2.0 pts.            |
| Fine soil           | 1.33 + 1.0-1.5 pts.     | 1.33 + 1.5-2.0 pts.                |

Follow all use directions, limitations, precautions, and information regarding application to peanuts on the AX S-MET and Prowl labels.

**TANK MIXTURE OR SEQUENTIALLY WITH GRAMOXONE INTEON**

AX S-MET + Gramoxone Inteon applied at ground cracking or sequentially will control or suppress small (1-6 inch)-emerged annual grass and broad leaf weeds and provide residual control of weed species listed in the **AX S-MET Applied Alone** section of this label. Apply Gramoxone Inteon plus the appropriate AX S-MET rate from the **Peanuts - AX S-MET Alone** section in a minimum spray volume of 20 gals./A with ground equipment. A second application of AX S-MET + Gramoxone Inteon may be made 28 days after ground cracking. (Refer to the **Peanuts - AX S-MET Combinations - Multiple Applications** section of this label for geographical areas where multiple applications are allowed.) Refer to the Gramoxone Inteon label and follow all directions, limitations, and restrictions.

**TANK MIXTURE OR SEQUENTIALLY WITH GRAMOXONE INTEON + BASAGRAN**

The addition of Basagran to the AX S-MET + Gramoxone Inteon mixture will result in improved control of such problem broadleaf weeds as prickly sida, cocklebur, smartweed, and bristly starbur. AX S-MET + Gramoxone Inteon + Basagran applied at ground cracking or sequentially will control or suppress small (1-6 inch) emerged annual grass and broadleaf weeds and provide residual control of weed species listed in the **AX S-MET Applied Alone** section of this label. Apply Basagran + Gramoxone Inteon with the appropriate AX S-MET rate from the **Peanuts - AX S-MET Alone** section in a minimum spray volume of 20 gals./A with ground equipment. A second application of AX S-MET + Gramoxone Inteon + Basagran may be made 28 days after ground cracking. (Refer to the **Peanuts - AX S-MET Combinations - Multiple Applications** section of this label for geographical areas where multiple applications are allowed.) Refer to the Gramoxone Inteon and Basagran labels and follow all directions, limitations, and restrictions.

**TANK MIXTURE OR SEQUENTIALLY WITH GRAMOXONE INTEON + BUTYRAC 200 OR BUTOXONE 200**

The addition of Butyrac 200 or Butoxone 200 to the AX S-MET + Gramoxone Inteon mixture will result in improved control of such problem broadleaf weeds as sicklepod, morningglory, and cocklebur. AX S-MET + Gramoxone Inteon + Butyrac 200 or Butoxone 200 applied at ground cracking or sequentially will control or suppress small (1-6 inch) emerged annual grass and broadleaf weeds and provide residual control of weed species listed in the **AX S-MET Applied**



41/  
61

**Alone** section of this label. Apply Gramoxone Inteon + Butyrac 200 or Butoxone 200 with the appropriate AX S-MET rate from the **Peanuts - AX S-MET Alone** section in a minimum spray volume of 20 gals./A with ground equipment. A second application of AX S-MET + Gramoxone Inteon + Butyrac 200 or Butoxone 200 may be made 28 days after ground cracking. (Refer to the Peanuts - **AX S-MET Combinations - Multiple Applications** section of this label for geographical areas where multiple applications are allowed.) Refer to the Gramoxone Inteon, Butyrac 200 or Butoxone 200 labels and follow all directions, limitations, and restrictions.

**TANK MIXTURE OR SEQUENTIALLY WITH BASAGRAN**

AX S-MET + Basagran applied at ground cracking or sequentially will control species on the Basagran label and provide residual control of species listed in the **AX S-MET Applied Alone** section of this label. Apply 1.0-2.0 pts./A of Basagran in 20 gals./A, depending on weed species and stage of growth as specified on the Basagran label, with the appropriate AX S-MET rate from the **Peanuts - AX S-MET Alone** section. A second application of the combination may be made before peanut pegging. (Refer to the **Peanuts - AX S-MET Combinations - Multiple Applications** section of this label for geographical areas where multiple applications are allowed.) A second Basagran application may be made in all peanut growing areas, if needed. Refer to the respective labels and follow all directions, limitations, and restrictions for each product.

**TANK MIXTURE OR SEQUENTIALLY WITH BASAGRAN + BUTYRAC 200 OR BUTOXONE 200**

AX S-MET + Basagran + Butyrac 200 or Butoxone 200 applied at ground cracking or sequentially will control species on the Basagran label and on the Butyrac or Butoxone labels, especially morningglories. Apply 1.5-2.0 pts./A of Basagran + 8.0 fl. oz./A of Butyrac 200 or Butoxone 200 in 20 gals./A, depending on weed species and stage of growth as specified on the Basagran label, with the appropriate AX S-MET rate from the **Peanuts - AX S-MET Alone** section. A second application of the combination may be made before peanut pegging. (Refer to the **Peanuts - AX S-MET Combinations - Multiple Applications** section of this label for geographical areas where multiple applications are allowed.) A second Basagran + Butyrac 200 or Butoxone 200 application may be made in all peanut growing areas, if needed. Refer to the respective labels and follow all directions, limitations, and restrictions for each product.

**TANK MIXTURE OR SEQUENTIALLY WITH STORM®**

AX S-MET + Storm applied at ground cracking through 2 expanded tetrafoliate leaves or AX S-MET applied according to the directions for **AX S-MET Alone** and followed with an at-cracking through postemergence treatment of Storm as specified on its label will control species on the Storm label and provide residual control of species listed in the **AX S-MET Applied Alone** section of this label. AX S-MET will not control emerged weeds. Refer to the **Peanuts - AX S-MET Alone** section and to the Storm label and follow all directions, limitations, and restrictions for each product.

**BEANS, PEAS AND LENTILS - AX S-MET ALONE**

Beans, peas, and lentils (including garbanzo, great northern beans, kidney beans, lima beans, mung beans, navy beans, peas (English\*; southern peas, such as blackeye, pinkeye, crowder, etc.), pinto beans, snap beans (green, wax, string), lentils, and lupines (sweet, white, white sweet, and grain).

**Use Directions and Restrictions for Fall Application:**

- Apply after September 30 in ND, SD, MN, WI, and north of Route 30 in IA.
- Apply after October 15 north of Route 91 in NE and south of Route 30 in IA.
- Apply after October 31 north of Route 136 in IL.

In all locations, apply to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55°F and falling. In minimum-till or no-tillage systems on soils having greater

42/  
61

than 2.5% organic matter, use 1.67-2.0 pts./A on medium-textured and 2.0 pts./A on fine-textured soils. Do not apply to frozen ground. A tillage operation may precede the application. A fall and/or a spring tillage may follow application, but do not exceed an incorporation depth greater than 2-3 inches. Minimize furrow and ridge formation in the tillage operations.

**Restriction:** If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for beans, peas, and lentils.

**Use Directions and Restrictions for Spring Application:**

Apply AX S-MET, either preplant incorporated or preemergence, using the appropriate rate specified below. **Preplant Incorporated or Preemergence:** Follow instructions for use of AX S-MET alone under **Application Procedures**. On coarse soils with less than 3% organic matter, apply 1.0-1.33 pts./A of AX S-MET or 1.33 pts./A if organic matter is 3% or greater. On medium soils, apply 1.33-1.67 pts./A of AX S-MET. On fine soils, apply 1.33-1.67 pts./A of AX S-MET if organic matter content is less than 3%, or 1.67-2.0 pts./A if organic matter content is 3% or greater.

\*On English peas, use only preemergence applications. If soils are cold and wet during pea germination and emergence, the use of AX S-MET may delay maturity and/or reduce yields.

**Restrictions:** To avoid possible illegal residues, (1) Do not cut for hay within 120 days following a AX S-MET application, (2) Do not use for forage within 60 days following a Dual II Magnum application, and (3) Do not apply more than 2.0 pts./A of AX S-MET during any one crop year.

**BEANS, PEAS AND LENTILS - AX S-MET COMBINATIONS**

**Restriction:** When applying AX S-MET in combination on beans, peas, and lentils do not cut for hay within 120 days following application, or illegal residues may result.

**TANK MIXTURE AND SEQUENTIAL APPLICATIONS WITH EPTAM - BEANS (GREEN OR DRY)**

This mixture controls all weeds controlled by AX S-MET alone and by Eptam alone. Refer to the **AX S-MET Applied Alone** section of this label for weeds controlled by AX S-MET alone and to the Eptam label for weeds controlled by Eptam.

**Preplant Incorporated:** Follow instructions for use of AX S-MET alone under **Application Procedures**. **Sequential:** Apply Eptam alone preplant incorporated, as specified on that label. Follow with a preemergence application of AX S-MET, at rates specified for AX S-MET alone, during planting (behind the planter) or after planting, but before the weeds or crop emerge.

Refer to the **Product Information** section of this label and to the Eptam label for weather, cultural practices, and all other precautions and limitations that affect performance of these products.

Apply 2.5-4.5 pts./A of Eptam 7E\* with AX S-MET as specified. On coarse soils, apply 0.8 pt./A of AX S-MET if organic matter content is less than 3%, or 1.0 pt./A if organic matter content is 3% or greater. On medium soils, apply 1.0 pt./A of AX S-MET if organic matter content is less than 3%, or 1.33 pts./A if organic matter content is 3% or greater. On fine soils, apply 1.33 pts./A of AX S-MET if organic matter is less than 3%, or 1.33-1.67 pts./A if organic matter is 3% or greater.

\*Refer to the Eptam label for rate limitations depending on geographical area, and for species and varietal restrictions.

**Restriction:** Do not exceed 3.5 pts./A of Eptam 7E on small white beans or green beans grown on coarse-textured soils.

**TANK MIXTURE WITH TREFLAN - BEANS (DRY - KIDNEY, NAVY, PINTO, ETC.; LIMA; AND SNAP)**

AX S-MET + Treflan tank mix applied preplant incorporated controls those weeds listed under **AX S-MET Applied Alone** and those weeds listed for Treflan alone on the Treflan label. AX S-MET + Treflan may be applied by ground or by aerial equipment and incorporated up to 14 days prior to planting. Follow the most restrictive procedures on this label and on the respective Treflan label, using equipment that provides uniform 2-inch incorporation.

Apply AX S-MET + Treflan tank mix using the appropriate AX S-MET rate specified for AX S-MET alone, and the Treflan rate from the Dry Beans, and the Lima and Snap Beans sections of the respective Treflan label. Choose the product rate for the specific soil texture/organic matter classification and weed species expected.

Follow all restrictions and precautions on the respective Treflan label and in the **Beans, Peas, and Lentils - AX S-MET Alone** section of this label.

**POTATOES - AX S-MET ALONE**

Apply AX S-MET, either incorporated, preemergence, or postemergence to potatoes after hilling/lay-by, according to directions specified below for control of weeds listed under the **Product Information** section. Within a rate range, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter. Effectiveness will be reduced if later cultural practices expose untreated soil. For applications by center pivot irrigation, see the Center Pivot Irrigation Application section of this label.

**Incorporated:** Apply AX S-MET at 1.0-2.0 pts./A to the soil and incorporate into the top 3 inches before planting, using a finishing disk, harrow, rolling cultivator, or similar implement. Planting and later cultural practices should not bring untreated soil to the surface. Postplant incorporated application may be made any time after planting to drag-off, but before potato emergence. Use an implement that evenly distributes AX S-MET in the top 2 inches of soil. Do not damage potato seed pieces or sprouts with incorporation equipment.

**Preemergence:** Apply AX S-MET at 1.0-2.0 pts./A, either after planting as a preemergence, delayed preemergence, after drag-off or hilling treatment, but before weeds emerge. Up to 2.6 pts./A of AX S-MET alone may be used where soil organic matter is between 6% and 20%.

**Postemergence After Hilling/Lay-by:** Apply 1.67 pts./A of AX S-MET postemergence to potatoes through after hilling/at lay-by to control AX S-MET - sensitive species for remainder of the growing season. This application will not control emerged weeds. It may be applied over a previous AX S-MET application, but do not apply more than 3.6 pts./A of AX S-MET in a single crop season.

**Restrictions:** (1) Do not use on muck or peat soils. If cool, wet soil conditions occur after application, AX S-MET may delay maturity and/or reduce yield of Superior and other early maturing potato varieties. (2) Do not apply both as a preemergence and an incorporated treatment.

**Pre Harvest Interval** - Do not harvest Potatoes treated with AX S-MET within 60 days after the at-planting to drag-off application, or within 40 days after a lay-by application, or illegal residues may result. (2) These directions for use do not apply to sweet potatoes or yams.

**POTATOES - AX S-MET COMBINATIONS**

**TANK MIXTURE WITH SENCOR**

In addition to those weeds controlled by AX S-MET alone, AX S-MET applied in tank mix combination with, or sequentially with, any of the registered Sencor formulations, also controls the

following broadleaf weeds: cocklebur\*, hairy nightshade\*, hemp sesbania, jimsonweed\*, lambsquarters, prickly sida, ragweed, smartweed, velvetleaf, Venice mallow, and wild mustard.

\*Partially controlled.

AX S-MET at 1.0-2.0 pts./A plus the labeled Sencor use rate may be used preemergence or postemergence to potatoes through after last hilling. Apply 1.0-1.33 pts./A of AX S-MET on coarse soils and 1.33-2.0 pts./A on other soil textures. Within this rate range, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter. Effectiveness will be reduced if later cultural practices expose untreated soil. AX S-MET will not control emerged weeds.

Refer to the Sencor label for precautionary statements, restrictions, application information, center pivot irrigation application, weeds controlled, and varietal limitations.

**Restrictions:** (1) Do not harvest potatoes treated with AX S-MET in tank mixture with Sencor within 60 days after application. (2) Potatoes may not be harvested within 40 days after a lay-by application of AX S-MET. (3) Do not use this tank mixture on muck or peat soils. (4) Postemergence applications to potatoes, with the exception of center pivot application, must be made only as a directed or semi-directed spray to avoid chlorosis, minor necrosis, or leaf distortion. (5) These directions for use do not apply to sweet potatoes or yams.

**AX S-MET + LOROX TANK MIXTURE (EAST OF ROCKY MOUNTAINS)**

AX S-MET may be applied in a tank mix combination with any of the registered Lorox formulations as a preemergence broadcast application to potatoes. Apply to the soil surface after planting and before emergence of the crop or after final drag-off according to the rates specified in Table 9.

**Table 9: AX S-MET + Lorox - Potatoes (East of Rocky Mountains)  
Broadcast Rates Per Acre**

|  | 1% to Less Than 3% Organic Matter | 1% to Less Than 3% Organic Matter | 3-5% Organic Matter | 3-5% Organic Matter |
|--|-----------------------------------|-----------------------------------|---------------------|---------------------|
| Soil Texture                           | AX S-MET                          | Lorox*                            | AX S-MET            | Lorox*              |
| <b>COARSE</b><br>Sandy loam            | 1.0 pt.                           | 1.0-1.5 lbs.                      | 1.33 pts.           | 1.5-2.0 lbs.        |
| <b>MEDIUM</b><br>Loam, silt loam, silt | 1.33 pts.                         | 1.5-2.0 lbs.                      | 1.67-2.0 pts.       | 2.0-2.5 lbs.        |

\*When using Lorox L or Lorox OF, use equivalent rates. One pt. of Lorox L equals 1.0 lb. of Lorox DF.

**Restrictions:** (1) Do not use on sands or loamy sands, and (2) Do not incorporate or spray over the top of emerged potatoes.

Refer to the **Product Information** section of this label and to the Lorox label for precautionary statements, restrictions, application information, and weeds controlled.

**TANK MIXTURE WITH PROWL 4E**

In addition to the weeds controlled by AX S-MET alone, this tank mixture with Prowl 4E controls such problem species as kochia, lambsquarters, purslane, annual spurge, stinging nettle, and others specified on the Prowl 4E Alone label. Apply AX S-MET + Prowl 4E preemergence,

preemergence incorporated, or early postemergence according to the specific directions on the Prowl4E label, using the rates in Table 10.

**Table 10: AX S-MET + Prowl 4E - Potatoes**

**Broadcast Rates Per Acre**

|                     | <b>Less than 3% Organic Matter</b> | <b>More Than 3% Organic Matter</b> |
|---------------------|------------------------------------|------------------------------------|
| <b>Soil Texture</b> | <b>AX S-MET + Prowl 4E*</b>        | <b>AX S-MET + Prowl 4E*</b>        |
| <b>COARSE</b>       | 1.0-1.33 pts. + 1.0-1.5 pts.       | 1.0-1.33 pts. + 1.0-1.5 pts.       |
| <b>MEDIUM</b>       | 1.33 pts. + 1.5-2.0 pts.           | 1.33-1.67 pts. + 2.0-3.0 pts.      |
| <b>FINE</b>         | 1.33-1.67 pts. + 2.0-3.0 pts.      | 1.67-2.0 pts. + 3.0 pts.           |

\*When using other formulations of Prowl, use equivalent rates of active ingredient.

Refer to the AX S-MET and Prowl 4E labels and observe all directions, timings, limitations, precautions, and restrictions concerning the use of these products on potatoes and follow the most restrictive.

**TANK MIXTURE WITH PROWL 4E + EPTAM**

In addition to the weeds controlled by AX S-MET alone, this tank mixture will control those species on the Prowl 4E and Eptam labels. Refer to the AX S-MET + Prowl 4E labels for rates of those products and add Eptam 7E at 3.5-7.0 pts./A, depending on geographical area. Refer to the respective AX S-MET, Prowl 4E, and Eptam labels and observe all directions, limitations, precautions, and restrictions concerning the use of these products on potatoes and follow the most restrictive.

**PUMPKIN - AX S-MET ALONE**

**Preemergence**

Apply AX S-MET preemergence (before the weeds have emerged) at 1.0 to 1.33 pint/A as an inter-row or inter-hill application in pumpkin. Leave 1 foot of untreated area over the row, or 6 inches to each side of the planted hill and/or any emerged pumpkin foliage (inter-row or inter-hill means not directly over the planted seed or young pumpkin plants). Use the lower AX S-MET rate on soils light in texture (loamy sand or lighter) and low in soil organic matter (less than 3%). AX S-MET applied as a broadcast spray over the planted row or hill, or applications made directly to crop foliage will increase the risk of injury to the pumpkin crop such as stand loss, delayed maturity, and loss of yield.

**Restriction:** Do not apply AX S-MET closer than 30 days before pumpkin harvest.

AX S-MET will not control emerged weeds, and thus should be applied before the weeds emerge. Weeds that are present should be controlled by another means, i.e. by mechanical means or by another herbicide.

**RHUBARB**

Apply AX S-MET at a broadcast rate of 0.67-1.33 pt./A to the soil surface in early spring, prior to crop emergence. Use lower rates on soils relatively coarse-textured and higher rates on fine-textured soils. A band application may also be used, applying proportionally less spray mixture on the area actually treated. AX S-MET will not control emerged weeds. Control emerged weeds with an appropriate registered foliar herbicide or by mechanical or physical means.

**Restrictions:** (1) Make only one application of AX S-MET per crop. (2) Do not apply more than 1.33 pt./A of AX S-MET per crop. (3) Do not harvest rhubarb within 62 days of the AX S-MET application.

46/61

### SAFFLOWERS - AX S-MET ALONE

**Preplant Incorporated or Preemergence:** Follow instructions for use of AX S-MET alone under **Application Procedures**.

On coarse soils, apply 1.0-1.33 pts./A of AX S-MET if organic matter content is less than 3%, or 1.33 pts./A if organic matter is 3% or greater. On medium soils, apply 1.33-1.67 pts./A of AX S-MET. On fine soils, apply 1.33-1.67 pts./A of AX S-MET if organic matter content is less than 3%, or 1.67-2.0 pts./A if organic matter content is 3% or greater.

### GRAIN OR FORAGE SORGHUM (SEED TREATED WITH CONCEP®) - AX S-MET ALONE

Apply AX S-MET preplant surface, preplant incorporated, preemergence or postemergence using the appropriate rate specified below. Apply AX S-MET alone only when the sorghum seed has been properly treated with Concep seed treatment. Preplant or preemergence applications of AX S-MET to sorghum not treated with Concep seed treatment will result in crop death.

**Fall Application for Italian Ryegrass Control:** AX S-MET may be applied for residual control of glyphosate-resistant Italian ryegrass (*Lolium multiflorum*). Apply AX S-MET at 1.33-1.67 pints/Acre in the fall (September 1-December 1) after harvest of the previous crop and prior to Italian ryegrass emergence. Use the lower AX S-MET rate for coarse textured soils and the higher rate for fine textured soils. A tillage operation may precede the application. Do not incorporate to a depth greater than 2-3 inches if tillage follows the application of AX S-MET. For fall applications after emergence of glyphosate-resistant Italian ryegrass, Gramoxone Inteon can be tank mixed with AX S-MET to control emerged ryegrass. Refer to the Gramoxone Inteon label for specific rates, application instructions and restrictions. Other registered herbicides may be tank mixed with AX S-MET for control or improved control of other weeds present at the time of application.

**Restrictions:** (1) Do not apply AX S-MET to frozen ground. (2) If a spring application is made, do not apply AX S-MET or any other product containing S-metolachlor the following spring to grain or forage sorghum.

**Preplant Surface-Applied:** Refer to instructions for use of AX S-MET under **Application Procedures section** on this label. For minimum-tillage or no-tillage systems only, AX S-MET may be applied up to 45 days before planting in CO, IA, IL, KS, MO, NE, and SD. Use only split applications for treatments made 30-45 days prior to planting, with 2/3 of the broadcast rate applied initially and the remaining 1/3 at planting. Apply 1.5 pts./A of AX S-MET on medium soils or 1.67 pts./A on fine soils. Treatments less than 30 days prior to planting may be made either as a split or single application. Apply 1.33 pts./A of AX S-MET on coarse soils not more than 2 weeks prior to planting. Under dry conditions, irrigation may be made after application to move AX S-MET into the soil.

**Preplant Incorporated or Preemergence:** Refer to instructions for use of AX S-MET under **Application Procedures section** on this label. Broadcast 1.0-1.33 pts./A of AX S-MET on coarse soils, 1.33-1.5 pts./A on medium soils, or 1.33-1.67 pts./A on fine soils.

**Postemergence:** Refer to instructions for use of AX S-MET under **Application Procedures section** on this label. AX S-MET may be applied broadcast postemergence at 1.0-1.33 pt./A on coarse soils, 1.33-1.5 pt./A on medium soils, or 1.33-1.67 pt./A on fine soils. AX S-MET will not control emerged weeds. Therefore, emerged weeds must be controlled by cultural or chemical

47/  
61

means. When applied alone, AX S-MET will be safe to emerged sorghum. The risk of sorghum injury increases when adjuvants (e.g. non-ionic, crop oil), Nitrogen sources (e.g. AMS, UAN) or fertilizers are applied with AX S-MET.

**Restrictions:** (1) Except for the split preplant surface treatment, do not make more than one application per year, or illegal residues may result. (2) Do not apply AX S-MET postemergence within 75 days of harvest.

**Precautions:** TO AVOID CROP INJURY (1) If sorghum seed is not properly treated with Concep seed treatment, preplant and preemergence applications of AX S-MET will severely injure the crop. (2) Under high soil moisture conditions prior to sorghum emergence, injury may occur following preplant and preemergence application AX S-MET. The crop will normally outgrow this effect. (3) Do not use AX S-MET on sorghum grown under dry mulch tillage, or injury may occur.

**GRAIN OR FORAGE SORGHUM (SEED TREATED WITH CONCEP) – AX S-MET TANK MIXTURES**

AX S-MET preplant or preemergence (prior to sorghum emergence) tank mixtures with AAtrex may be applied in water or fluid fertilizer. Apply AX S-MET preplant or preemergence tank mixtures only when the sorghum seed has been properly treated with Concep seed treatment. Preplant or preemergence applications of AX S-MET to sorghum not treated with Concep seed treatment will result in crop death.

**IMPORTANT: FOR TANK MIXTURES WITH AATREX (OR OTHER BRANDS OF ATRAZINE) - If applying AX S-MET in tank mixture with AAtrex, all the restrictions and rate limitations on the AAtrex label must be followed if more restrictive/protective than those on this label. In addition, if AAtrex is/must be applied at rates lower than those listed on this label, broadleaf weed control may be affected. Refer to the AAtrex label for weeds controlled at the reduced rates.**

**Precautions:** TO AVOID CROP INJURY (1) Applications of AX S-MET + AAtrex on highly alkaline soils or on eroded areas where calcareous subsoils are exposed may cause sorghum injury. (2) If sorghum seed is not properly treated with Concep, preplant and preemergence applications of AX S-MET + AAtrex may severely injure the crop: (3) Under high soil moisture conditions prior to sorghum emergence, injury may occur following the use of preplant and preemergence applications of AX S-MET + AAtrex. The crop will normally outgrow this effect. (4) Do not use AX S-MET + AAtrex on sorghum grown under dry mulch tillage, or injury may occur.

**Restriction:** Except for the split preplant surface treatment, do not make more than one application per year.

**TANK MIXTURE WITH AATREX**

In addition to the weeds controlled by AX S-MET alone, AX S-MET + AAtrex also controls the following broadleaf weeds when applied either preplant surface, preplant incorporated, or preemergence: cocklebur, common purslane, hairy nightshade, lambsquarters, morningglory, ragweed, smartweed, and velvetleaf.

**Preplant Surface-Applied:** Refer to instructions for use of AX S-MET under **Application Procedures** on this label. For minimum-tillage or no-tillage systems only, AX S-MET + AAtrex may be applied up to 45 days prior to planting in IA, IL, eastern KS, MO, NE, and SD. Use only split applications for treatments made 30-45 days prior to planting, with 2/3 of the broadcast rate applied initially and the remaining 1/3 at planting. Apply 1.5 pts./A of AX S-MET + 1.7-2.0 lbs./A of AAtrex Nine-O\* on medium soils with 1.5% organic matter or greater. Apply 1.5 pts./A of AX S-MET + 1.7-2.0 lbs./A of AAtrex Nine-O on fine soils with less than 1.5% organic matter, or apply 1.67 pts./A of AX S-MET + 2.0-2.2 lbs./A of AAtrex Nine-O on fine soils with 1.5% organic matter

48/  
61

or greater. Treatments less than 30 days prior to planting may be made either as a split or single application. Under dry conditions, irrigation may be made after application to move AX S-MET + AAtrex into the soil.

**Precautions:** To avoid crop injury, (1) Do not use on coarse soils, and (2) Do not use on medium soils with less than 1.5% organic matter.

**Preplant Incorporated or Preemergence:** Refer to instructions for use of AX S-MET under Application **Procedures** on this label. On medium soils with 1.5% organic matter or greater, apply 1.0 pt./A of AX S-MET + 1.3 lbs./A of AAtrex Nine-O\*. On line soils with less than 1.5% organic matter, apply 1.0 pt./A of AX S-MET + 1.3 lbs./A of AAtrex Nine-O; on fine soils with 1.5% organic matter or greater, apply 1.2-1.33 pts./A of AX S-MET + 1.6-1.8 lbs./A of AAtrex Nine-O.

\*When using AAtrex 4L, use equivalent rates. One lb. of AAtrex Nine-O = 1.8 pts. of AAtrex 4L.

Precautions: To avoid crop injury, (1) Do not use on coarse soils; (2) Do not use on medium soils with less than 1.5% organic matter.

**Regional Restrictions** (1) Do not use in NM, OK, or TX, except in northeast OK and the TX Gulf Coast and Blacklands areas; and (2) Do not apply preplant incorporated in AZ or the Imperial Valley of CA.

**TANK MIXTURE OF AX S-MET OR AX S-MET + AATREX, WITH GRAMOXONE INTEON, LANDMASTER BW, TOUCHDOWN BRANDS OR ROUNDUP BRANDS FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS**

In minimum-tillage or no-tillage systems where sorghum (seed treated with Concep) is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides Gramoxone Inteon, Landmaster BW, Touchdown or Roundup may be tank mixed with AX S-MET or AX S-MET + AAtrex. See Comment No. 7 following Table 2. The AX S-MET or AX S-MET + AAtrex portion of the tank mixture provides preemergence control of the weeds listed on this label under the respective sections.

Refer to the label of each product used in combination and observe the planting details, restrictions, and all other precautions and limitations.

**Application:** Apply before, during, or after planting, but before sorghum emerges. Add Gramoxone Inteon, Landmaster BW, Touchdown brands or Roundup brands and apply as directed on the product labels.

**Gramoxone Inteon:** Apply as directed on the product label. This treatment will not control weeds taller than 6 inches.

**Landmaster BW:** 27-54 oz./A depending on weed species and size. See the Landmaster BW label for weeds controlled, listed rates for specific weeds, and other information concerning use.

**Touchdown Brands or Roundup Brands:** See the Touchdown brand or Roundup brand label for weeds controlled, listed rates, and other use directions.

**SWEET SORGHUM (SEED TREATED WITH CONCEP)**

Apply AX S-MET, preplant surface, preplant incorporated, preemergence or postemergence using the appropriate rate specified below. Apply AX S-MET only when the sweet sorghum seed has been properly treated with Concep seed treatment. Preplant or preemergence applications of AX S-MET to sweet sorghum not treated with Concep seed treatment will result in crop death.



49/61

**Soil-Applied:** AX S-MET may be applied up to 45 days before planting. Use only split applications for treatments made 30-45 days prior to planting, with 2/3 of the broadcast rate applied initially and the remaining 1/3 at planting. Treatments less than 30 days prior to planting may be made either as a split or single application. Under dry conditions, irrigation after application may be made to move AX S-MET into the soil.

**AX S-MET rates for soil applications to sweet sorghum**

| Soil Type | 30-45 days prior to planting <sup>1</sup> | < 30 days prior to planting | At Planting <sup>2</sup> |
|-----------|---|-----------------------------|--------------------------|
| Coarse    | DO NOT APPLY                              | 1.33 pts                    | 1.0-1.33 pts             |
| Medium    | 1.5 pts                                   | 1.5 pts                     | 1.33-1.5 pts             |
| Fine      | 1.67 pts                                  | 1.67 pts                    | 1.33-1.67 pts            |

<sup>1</sup> Use only as a split application with 2/3 of the broadcast rate applied initially and the remaining 1/3 applied at planting.

<sup>2</sup> Preplant Incorporated or pre-emergence

**Post-Applied:** AX S-MET may be applied post-emergence to sweet sorghum for residual control of grasses and small seeded broadleaf weeds. Postemergence application to sweet sorghum may be made to crop up to 5 inches in height. AX S-MET will not control emerged weeds. Therefore, emerged weeds must be controlled by cultural or chemical methods. When applied alone, AX S-MET will be safe to emerged sweet sorghum. Use of adjuvants is prohibited on sweet sorghum.

**AX S-MET rates for post-emergence applications to sweet sorghum.**

| Soil Type | Post-emergence Rate |
|-----------|---------------------|
| Coarse    | 1.0-1.33 pts        |
| Medium    | 1.33 pts            |
| Fine      | 1.33 pts            |

**Precautions:** TO AVOID CROP INJURY (1) If sweet sorghum seed is not properly treated with Concep seed treatment, soil applications of AX S-MET prior to sorghum emergence will severely injure the crop. (2) Under high soil moisture conditions prior to sweet sorghum emergence, injury may occur following soil applications of AX S-MET. The crop will normally outgrow this effect. (3) Do not use AX S-MET on sorghum grown under dry mulch tillage, or injury may occur.

**Restriction:**(1) Only one application per season is allowed. AX S-MET may be applied either as a soil applied treatment or a post-emergence treatment but not both. (2) Do not apply AX S-MET post-emergence within 90 days of harvest.

**SOYBEANS - AX S-MET ALONE**

Apply AX S-MET, preplant surface-applied, preplant incorporated, preemergence, or postemergence using the appropriate rate specified below. Follow instructions for use of AX S-MET alone under **Application Procedures** section of this label.

**Fall Application for Spring Weed Control**

- Apply after September 30 in ND, SD, MN, WI, and north of Route 30 in IA.
- Apply after October 15 north of Route 91 in NE and south of Route 30 in IA.
- Apply after October 31 north of Route 136 in IL.

In all locations, apply to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55° F and falling. In minimum-till or no-tillage systems on soils having greater than 2.5% organic matter, use 1.67-2.0 pts./A of AX S-MET on medium-textured and 2.0 pts./A of AX S-MET on fine-textured soils. Do not apply to frozen ground. A tillage operation may precede the application. A fall and/or a spring tillage may follow application, but do not exceed an incorporation depth greater than 2-3 inches. Minimize furrow and ridge formation in the tillage operations. If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for soybeans of 2.6 pts./A depending on soil texture or illegal residues may result.

**Fall Application for Italian Ryegrass Control:** AX S-MET may be applied for residual control of glyphosate-resistant Italian ryegrass (*Lolium multinode*). Apply AX S-MET at 1.33-1.67 pints/Acre in the fall (September 1-December 1) after harvest of the previous crop and prior to Italian ryegrass emergence. Use the lower AX S-MET rate for coarse textured soils and the higher rate for fine textured soils. A tillage operation may precede the application. Do not incorporate to a depth greater than 2-3 inches if tillage follows the application of AX S-MET. For fall applications after emergence of glyphosate-resistant Italian ryegrass, Gramoxone Inteon can be tank mixed with AX S-MET to control emerged ryegrass. Refer to the Gramoxone Inteon label for specific rates, application instructions and restrictions. Other registered herbicides may be tank mixed with AX S-MET for control or improved control of other weeds present at the time of application.

**Restrictions:** (1) Do not apply AX S-MET to frozen ground. (2) If a spring application is made, the combined total amount of AX S-MET applied in the fall plus the spring must not exceed the maximum seasonal S-metolachlor rate for soybeans (2.6 pt./A, depending on soil texture), or illegal residues may result.

**Preplant Surface - Spring Applied:** Use on medium and fine soils with minimum-tillage or no-tillage systems in CO, CT, DE, IA, IL, IN, KS, KY, MA, MD, ME, MI, MN, MO, MT, ND, NE, NH, NY, OH, PA, RI, SD, TN, VA, VT, WI, WV, and WY. Apply 2/3 the labeled rate of AX S-MET (1.67 pts./A on medium soils and 2.0 pts./A on fine soils) as a split treatment 30-45 days prior to planting and the remainder at planting. Applications made less than 30 days before planting may be as either a split or single treatment. Apply 1.33 pts./A of AX S-MET on coarse soils not more than 2 weeks prior to planting.

**Restrictions:** (1) On soybeans, AX S-MET may be used up to 2.6 pts./A as a preplant surface treatment on soils having organic matter content between 6% and 20%. (2) The total AX S-MET rate applied to soybeans during anyone crop must not exceed 2.6 pts./A (3) Do not graze or feed treated soybean forage, hay, or straw to livestock 30 days following treatment.

**Preplant Incorporated or Preemergence:** On coarse soils, apply 1.0-1.33 pts./A of AX S-MET if organic matter content is less than 3%, or 1.33 pts./A if organic matter content is 3% or greater. On medium soils, apply 1.33-1.67 pts./A of AX S-MET. On fine soils, apply 1.33-1.67 pts./A of AX S-MET if organic matter content is less than 3%, or 1.67-2.0 pts./A if organic matter content is 3% or greater.

**Restrictions:** (1) On soybeans, AX S-MET may be used up to 2.6 pts./A as a preplant incorporated or preemergence treatment on soils having an organic matter content between 6% and 20%. (2) The total AX S-MET rate applied to soybeans during any one crop must not exceed 2.6 pts./A. (3) Do not graze or feed treated soybean forage, hay, or straw to livestock 30 days following treatment.

**Postemergence:** Apply 1.0-1.33 pts./A of AX S-MET as a postemergence treatment to soybeans from emergence up through the third trifoliate leaf stage. AX S-MET will not control emerged weeds so it must be applied to a weed-free soil surface or in a tank mixture with products that provide postemergence control of weeds present at the time of application.

AX S-MET can also be applied as part of a sequential soybean weed control program. If AX S-MET was applied as a preplant surface, preplant incorporated, or a preemergence treatment, a second treatment of AX S-MET can be applied postemergence provided that the total AX S-MET rate during any one crop does not exceed 2.6 pts./A.

**Restrictions:** (1) Do not apply more than 1.33 pts./A of AX S-MET postemergence to soybeans. (2) The total AX S-MET rate applied preplant, preemergence or postemergence to soybeans during any one crop must not exceed 2.6 pts./A (2.5 lb. a.i./A of S-metolachlor). (3) Make postemergence applications at least 90 days before harvest. (4) Do not graze or feed treated forage or hay from soybeans to livestock following a postemergence application of AX S-MET.

**SOYBEANS - AX S-MET COMBINATIONS**

Water or fluid fertilizer may be used as carrier for AX S-MET in combination with Sencor, Lorox, Canopy, Pursuit, Scepter, Sonalan, or Command.

**Restrictions:** The total AX S-MET rate applied to soybeans during any one crop year must not exceed 2.6 pts./A. For all of the following combinations, AX S-MET may be used up to 2.33 pts. /A on soils having an organic matter content between 6% and 20%.

**TANK MIXTURE WITH SENCOR**

In addition to those weeds controlled by AX S-MET alone, AX S-MET + Sencor, when applied as directed, also controls the following broadleaf weeds: cocklebur\*, hairy nightshade, hemp sesbania, jimsonweed\*, lambsquarters, prickly sida, ragweed, smartweed, velvetleaf, Venice mallow, and wild mustard.

\*Partially controlled.

Apply AX S-MET and Sencor preplant incorporated or preemergence, using the appropriate rates from Table 11. **Preplant Incorporated or Preemergence:** Follow instructions for use of AX S-MET alone under **Application Procedures**.

**Sequential:** Apply AX S-MET alone Preplant Incorporated, as specified in Table 11 for this tank mixture. Follow with a preemergence application of Sencor during planting (behind the planter) or after planting, but before weeds or soybeans emerge.

Refer to the Sencor labels for planting details and soybean variety restrictions.

**Table 11: AX S-MET + Sencor - Soybeans**

**Broadcast Rates Per Acre**

| Soil Texture**   | 0.5% to Less Than 3% Organic Matter |            | 3% Organic Matter or Greater |            |
|--|-------------------------------------|------------|------------------------------|------------|
|  | AX S-MET                            | Sencor DF* | AX S-MET                     | Sencor DF* |
| <b>COARSE</b><br>Loamy sand (over 2% organic matter), sandy loam | 0.8-1.0 pt.                         | + 0.33 lb. | 1.0 pt.                      | + 0.5 lb.  |
| <b>MEDIUM</b>  | 1.0-1.33 pts.                       | + 0.5 lb.  | 1.33 pts.<br>lb.***          | + 0.67     |

52/61

|  |                      |                           |
|--|----------------------|---------------------------|
| <b>FINE</b>  | 1.33 pts. + 0.67 lb. | 1.33-1.67 pts. + 0.67 lb. |
| <b>Mississippi Delta Only</b><br>Silty clay, clay      | 1.33 pts. + 1.0 lb.  | 1.33-1.67 pts. + 1.0 lb.  |
| Muck or Peat (soils with more than 20% organic matter) | DO NOT USE           | DO NOT USE                |

\*When using Sencor 4 multiply lbs. of DF by 1.5 to get pts./A.

\*\*On all sand and on loamy sand with less than 2% organic matter, do not use this tank mixture preemergence, or the sequential treatment. Do not use the tank mixture preplant incorporated on any sand, loamy sand, or sandy loam, or crop injury may occur.

\*\*\*Use 0.5 lb./A if applied preplant incorporated.

**Precautions:** TO AVOID CROP INJURY (1) Do not use the tank mix or sequential application on soil with less than 0.5% organic matter or on alkaline soil with a pH over 7.4, or crop injury may occur. (2) If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stands for several days.

Follow most restrictive limitations and precautions on the **AX S-MET – Soybeans Alone** section of the AX S-MET label and the Soybean directions on the Sencor label.

**TANK MIXTURE WITH LOROX**

In addition to those weeds controlled by AX S-MET alone, AX S-MET + Lorox, applied preemergence, also controls the following broadleaf weeds: cocklebur\*, jimsonweed\*, lambsquarters, morningglory\*, prickly sida, ragweed, smartweed, velvetleaf\*, Venice mallow, and wild mustard.

\*Partially controlled.

**Preemergence:** Apply during planting (behind planter) or after planting, but before weeds or soybeans emerge. Refer to the Lorox label for planting details. Apply the appropriate rates from Table 12.

**Precaution:** TO AVOID CROP INJURY, Do not use on soil with less than 0.5% organic matter.

**Table 12: AX S-MET + Lorox - Soybeans**  
**Broadcast Rates Per Acre**

|  | 0.5% to Less Than 3% Organic Matter |                | 3% Organic Matter or Greater |                |
|--|-------------------------------------|----------------|------------------------------|----------------|
| Soil Texture*  | AX S-MET DF***                      | + Lorox        | AX S-MET                     | + Lorox DF***  |
| <b>COARSE**</b>  | 0.8 pt.                             | + 1.0 lb.      | 1.0 pt.                      | + 1.0-1.5 lbs. |
| <b>MEDIUM</b>  | 1.0 pt.                             | + 1.0-1.5 lbs. | 1.33 pts.                    | + 1.5-2.0 lbs. |
| <b>FINE</b>  | 1.33 pts.                           | + 2.0 lbs.     | 1.33-1.67 pts.               | + 2.5-3.0 lbs. |
| Muck or Peat (Soils with more than 20% organic matter) | DO NOT USE                          |                | DO NOT USE                   |                |

\*Do not use on sand, gravelly soils, or exposed subsoils.

\*\*Do not use on loamy sand, except in the northeastern U.S. on loamy sand with over 1% organic matter.

\*\*\*When using Lorox L or Lorox DF, use equivalent rates. One pt. of Lorox L equals 1.0 lb. of Lorox DF.

Follow the most restrictive limitations and precautions on the **AX S-MET - Soybeans Alone** section of the AX S-MET label and the Soybean directions on the Lorox labels.

**TANK MIXTURE WITH TREFLAN**

AX S-MET + Treflan tank mix applied preplant incorporated controls those weeds listed under the **AX S-MET Applied Alone** section and those weeds listed for Treflan Alone on the Treflan label. AX S-MET + Treflan may be applied by ground or by aerial equipment and incorporated up to 14 days before planting. Follow the use directions on the Treflan and AX S-MET labels, using equipment that provides uniform 2-inch incorporation.

Apply AX S-MET + Treflan tank mix using the appropriate rate from the **Soybeans - AX S-MET Alone** section of this label and the Treflan Alone section of the Treflan label for the specific soil texture/organic matter classification and weed species expected.

To control DNA-resistant goosegrass\* and other species on the respective labels where the soil organic matter is 3% or less, apply the rate in Table 13.

**Table 13: AX S-MET + Treflan - Organic Matter Content Less than 3%  
Broadcast Rates Per Acre**

|                     | <b>AX S-MET</b>                    | <b>Treflan E.C.**</b>              | <b>Treflan E.C.**</b>      |
|---------------------|------------------------------------|------------------------------------|----------------------------|
| <b>Soil Texture</b> | <b>Organic Matter Less Than 3%</b> | <b>Organic Matter Less than 2%</b> | <b>Organic Matter 2-3%</b> |
| <b>COARSE*</b>      | 0.8-1.0 pt.                        | 1.0 pt.                            | 1.5 pts.                   |
| <b>MEDIUM</b>       | 1.0 pt.                            | 1.5 pts.                           | 1.5 pts.                   |
| <b>FINE</b>         | 1.33 pts.                          | 2.0 pts.                           | 2.0 pts.                   |

\*Where a range of rates is given for AX S-MET, use the minimum rate where DNA-resistant goosegrass is the predominant species.

\*\*When Treflan MTF or Treflan 5 is used, use comparable rates. Multiply pts. of Treflan E.C. by 1 for Treflan MTF and by 0.8 for Treflan 5.

Follow the most restrictive limitations and precautions on the **AX S-MET - Soybeans Alone** section of the AX S-MET label and the Soybean directions on the Treflan labels.

**TANK MIXTURE WITH SCEPTER**

This tank mixture controls all weeds controlled by AX S-MET alone and by Scepter alone. Refer to the **AX S-MET Applied Alone** section for weeds controlled by AX S-MET and to the Scepter label for weeds controlled by Scepter. Refer to the Scepter label for geographical locations where this tank mixture may be applied.

Apply AX S-MET + Scepter preplant incorporated or preemergence, using rates in Table 14. Follow use directions under **Application Instructions** on the Scepter label. For preplant incorporated applications, apply and incorporate within 30 days before planting. Observe all other precautions and limitations on the Scepter labels.

**Table 14: AX S-MET + Scepter - Soybeans**  
**Broadcast Rates Per Acre**

|  | Less Than 3% Organic Matter | Less Than 3% Organic Matter | 3% or More Organic Matter | 3% or More Organic Matter |
|--|-----------------------------|-----------------------------|---------------------------|---------------------------|
| Soil Texture   | AX S-MET                    | Scepter                     | AX S-MET                  | Scepter                   |
| <b>COARSE</b>  | 0.8 pt.                     | 0.67 pt.                    | 1.0 pt.                   | 0.67 pt.                  |
| <b>MEDIUM</b>  | 1.0 pt.                     | 0.67 pt.                    | 1.33 pts.                 | 0.67 pt.                  |
| <b>FINE</b>  | 1.33 pts.                   | 0.67 pt.                    | 1.33-1.67* pts.           | 0.67 pt.                  |
| Muck or Peat (soils with more than 20% organic matter) | DO NOT USE                  | DO NOT USE                  | DO NOT USE                | DO NOT USE                |

\*Use the higher rate of AX S-MET if heavy weed infestations are expected.

Follow the most restrictive limitations and precautions on the **AX S-MET - Soybeans Alone** section of the AX S-MET label and the Soybean directions on the Scepter labels.

**TANK MIXTURE WITH CANOPY**

This tank mixture controls all weeds controlled by AX S-MET alone and by Canopy alone. Refer to the **AX S-MET Applied Alone** section for weeds controlled by AX S-MET and to the Canopy label for weeds controlled by Canopy.

Apply preplant incorporated or preemergence, using the appropriate rates from Table 15. **Preplant Incorporated:** Apply within 2 weeks of planting. Uniformly incorporate into the top 1-2 inches of soil before planting soybeans. **Preemergence:** Apply after planting, but before soybeans emerge.

Follow the most restrictive limitations and precautions on the **AX S-MET - Soybeans Alone** section of the AX S-MET label and the Soybean directions on the Canopy labels including varietal restrictions.

**Table 15: AX S-MET + Canopy - Soybeans**  
**Broadcast Rates Per Acre**

|               | Less Than 3% Organic Matter | 3% or More Organic Matter | 3% or More Organic Matter |
|---------------|-----------------------------|---------------------------|---------------------------|
| Soil Texture  | AX S-MET                    | AX S-MET                  | Canopy                    |
| <b>COARSE</b> | 0.8 pt.                     | 1.0 pt.                   | *                         |
| <b>MEDIUM</b> | 1.0 pt.                     | 1.33 pts.                 | *                         |
| <b>FINE</b>   | 1.33 pts.                   | 1.33-1.67 pts.            | *                         |

\*Refer to the Canopy label for appropriate rate, according to geographical location, soil and organic matter classification, and pH limitations.

**Restriction:** Do not apply to sand, or to any soil with less than 0.5% organic matter, or to any soil with pH greater than 7.0, except as noted on the Canopy label.

**TANK MIXTURE WITH COMMAND**

This tank mixture controls all weeds controlled by AX S-MET alone and by Command alone. Refer to the **AX S-MET Applied Alone** section for weeds controlled by Dual Magnum and to the Command label for weeds controlled by Command.

Apply AX S-MET + Command preplant incorporated, using rates in Table 16, Follow all Command application instructions as to incorporation interval, geographical location, equipment operation, soil moisture conditions, etc.

Follow the most restrictive limitations and precautions on the Dual Magnum - Soybeans Alone section of the AX S-MET label and the Soybean directions on the Command labels including rotational restrictions.

**Table 16: AX S-MET + Command - Soybeans**  
**Broadcast Rates Per Acre**

|               | AX S-MET              | AX S-MET                       | Command 4E    | Command 4E    |
|---------------|-----------------------|--------------------------------|---------------|---------------|
| Soil Texture  | 0.5-3% Organic Matter | Greater Than 3% Organic Matter | Northern Area | Southern Area |
| <b>COARSE</b> | 0.8 pt.               | 1.0 pt.                        | 1.5-2.0 pts.  | 2.0-2.5 pts.  |
| <b>MEDIUM</b> | 1.0 pt.               | 1.33 pts.                      | 1.5-2.0 pts.  | 2.0-2.5 pts.  |
| <b>FINE</b>   | 1.33 pts.             | 1.33-1.67 pts.                 | 1.5-2.0 pts.  | 2.0-2.5 pts.  |

**TANK MIXTURE WITH SONALAN**

This tank mixture controls all weeds controlled by AX S-MET alone and by Sonalan alone. Refer to the **AX S-MET Applied Alone** section for weeds controlled by AX S-MET and to the Sonalan label for weeds controlled by Sonalan.

Apply AX S-MET and Sonalan preplant incorporated, using the appropriate rates from Table 17.

**Preplant Incorporated:** Follow use directions for soil preparation procedures for Sonalan as described on the Sonalan label.

**Sequential:** Apply Sonalan alone preplant incorporated as specified on the Sonalan label. Follow with a preemergence application of AX S-MET during planting (behind the planter) or after planting, but before weeds or soybeans emerge.

**Table 17: AX S-MET + Sonalan - Soybeans**  
**Broadcast Rates Per Acre**

|               | Less Than 3% Organic Matter | Less Than 3% Organic Matter | 3% or More Organic Matter | 3% or More Organic Matter |
|---------------|-----------------------------|-----------------------------|---------------------------|---------------------------|
| Soil Texture  | AX S-MET                    | Sonalan                     | AX S-MET                  | Sonalan                   |
| <b>COARSE</b> | 1.0-1.33 pts.               | 1.25-2.0 pts.               | 1.33 pts.                 | 1.25-2.0 pts.             |
| <b>MEDIUM</b> | 1.33-1.67 pts.              | 1.75-2.5 pts.               | 1.33-1.67 pts.            | 1.75-2.5 pts.             |

56/61

|  |                |               |               |               |
|--|----------------|---------------|---------------|---------------|
| <b>FINE</b>  | 1.33-1.67 pts. | 2.25-3.0 pts. | 1.67-2.0 pts. | 2.25-3.0 pts. |
| Muck or Peat (soils with more than 20% organic matter) | DO NOT USE     | DO NOT USE    | DO NOT USE    | DO NOT USE    |

\*For eastern black nightshade on these soils, apply Sonalan at 3.0 pts./A on medium- and 3.5 pts./A on fine-textured soils, and follow with 2 incorporation passes.

Follow the most restrictive limitations and precautions on the **AX S-MET - Soybeans Alone** section of the AX S-MET label and the Soybean directions on the Sonalan labels.

**TANK MIXTURE WITH PURSUIT**

This tank mixture controls all weeds controlled by AX S-MET alone and- by Pursuit alone. Refer to the **AX S-MET Applied Alone** section for weeds controlled by AX S-MET and to the Pursuit label for weeds controlled by Pursuit. Refer to the Pursuit label for geographical locations where this tank mixture may be applied.

Apply AX S-MET + Pursuit early preplant, preplant incorporated, or preemergence after planting, using rates in Table 18. Application can be made in water or liquid fertilizer. Follow all use directions under Soil Applications on the Pursuit label. For early preplant and preplant incorporated applications, apply within 30 days before planting.

Follow the most restrictive limitations and precautions on the **AX S-MET -Soybeans Alone** section of the AX S-MET label and the Soybean directions on the Pursuit labels including rotational restrictions.

**Table 18: AX S-MET + Pursuit - Soybeans**

**Broadcast Rates Per Acre**

|                     | <b>Less Than 3% Organic Matter</b> | <b>3% or More Organic Matter</b> | <b>Less Than 3% - 3% or More Organic Matter</b> |
|---------------------|------------------------------------|----------------------------------|---|
| <b>Soil Texture</b> | <b>AX S-MET</b>                    | <b>AX S-MET</b>                  | <b>Pursuit</b>                                  |
| <b>COARSE</b>       | 0.8 pt.                            | 1.0 pt.                          | 0.25 pt.  |
| <b>MEDIUM</b>       | 1.0 pt.                            | 1.33 pts.                        | 0.25 pt.  |
| <b>FINE</b>         | 1.33 pts.                          | 1.33-1.67 pts.                   | 0.25 pt.  |

**Sequential:** Apply AX S-MET early preplant, preplant incorporated or preemergence after planting at 0.8 pt./A on coarse soils and 1.0 pt./A on medium- and fine-textured soils. Follow with a sequential postemergence application of Pursuit to control emerged weeds according to the Pursuit label. AX S-MET will improve the consistency and level of control from Pursuit on most grass species. Refer to the Pursuit postemergence label for a listing of weeds controlled, application rate, and growth stage limitations.

**TANK MIXTURE WITH SENCOR, SCEPTER, LOROX, CANOPY, OR PURSUIT, PLUS GRAMOXONE INTEON, TOUCHDOWN BRANDS OR ROUNDUP BRANDS FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS**

In minimum-tillage or no-tillage systems where soybeans are planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides Gramoxone



Inteon, Touchdown brands or Roundup brands may be added to a tank mix of either AX S-MET + Sencor, AX S-MET + Scepter, AX S-MET + Lorox, AX S-MET + Canopy, or AX S-MET + Pursuit. When used as directed, the Gramoxone Inteon portion of the tank mixture controls most emerged weeds and suppresses many perennial weeds. Touchdown or Roundup combinations will control emerged annual and perennial weeds when applied as directed on the Touchdown or Roundup label. The AX S-MET + Sencor, Scepter, Lorex, Canopy, or Pursuit portion of the tank mixture provides preemergence control of the weeds listed on this label in the tank mixture section for AX S-MET + Sencor/Lexone, AX S-MET + Scepter, AX S-MET + Lorox, AX S-MET + Canopy, and AX S-MET + Pursuit, respectively.

Refer to the label of each product used in combination and observe the planting details, soybean variety restrictions, information regarding application to soybeans, geographical restrictions, and all other precautions and limitations.

**Application:** Apply before, during, or after planting, but before the soybeans emerge. Add Gramoxone Inteon, Touchdown brands or Roundup brands and apply as directed on the product labels.

**Gramoxone Inteon:** Apply as directed on the product label. This treatment will not control weeds taller than 6 inches.

**Restriction:** Do not apply combinations containing Gramoxone Inteon in suspension-type liquid fertilizers, as the activity of paraquat will be reduced.

**Touchdown or Roundup:** See the Touchdown brand or Roundup brand label for weeds controlled, labeled rates, and other use directions.

Apply in 20-60 gals. of water or fluid fertilizer per acre with ground equipment.

**AX S-MET + Sencor + Gramoxone Inteon, Touchdown Brands or Roundup Brands**

On loamy sand with over 2% organic matter, apply 1.0 pt./A of AX S-MET + 0.33-0.5 lb./A of Sencor DF. On medium soils, apply 1.33 pts./A of AX S-MET + 0.5-0.67 lb./A of Sencor DF. On fine soils, apply 1.33-1.67 pts./A of AX S-MET + 0.67 lb./A of Sencor DF.

\*When using Sencor 4L, multiply lbs. of DF by 1.5 to get pts./A.

**Precautions:** TO AVOID CROP INJURY, (1) Do not use this tank mixture on soil with less than 0.5% organic matter, on alkaline soil with a pH over 7.4, or on all sand and on loamy sand with less than 2% organic matter. (2) If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stands for several days, or where the seeding slit has not been properly closed.

**AX S-MET + Scepter + Gramoxone Inteon, Touchdown Brands or Roundup Brands**

On coarse soils, apply 1.0 pt./A of AX S-MET + 0.67 pt./A of Scepter. On medium soils, apply 1.33 pts./A of AX S-MET + 0.67 pt./A of Scepter. On fine soils, apply 1.67 pts./A of AX S-MET + 0.67 pt./A of Scepter.

**Restriction:** (1) Do not apply within 90 days of harvest, and (2) Do not graze or feed treated soybean forage, hay, or straw to livestock.

**AX S-MET + Lorox + Gramoxone Inteon, Touchdown Brands or Roundup Brands**

On coarse soils\*, apply 1.0 pt./A of AX S-MET + 1.0-1.5 lbs./A of Lorox DF\*\*. On medium soils, apply 1.33 pts./A of AX S-MET + 1.0-2.0 lbs./A of Lorox DF. On fine soils, apply 1.33-1.67 pts./A of AX S-MET + 2.0-3.0 lbs./A of Lorox DF.

\*Do not use on loamy sand, except in the northeastern U.S. on loamy sand with over

58/61

1% organic matter. Do not use on sand, gravelly soils, or exposed subsoils.

\*\*When using Lorox DF, use equivalent rates. One pt. of Lorox L equals 1.0 lb. of Lorox DF.

**Precaution:** TO AVOID CROP INJURY, Do not use on soil with less than 0.5% organic matter.

**AX S-MET + Canopy + Gramoxone Inteon, Touchdown Brands or Roundup Brands**

Use only where soils have 0.5-5% organic matter. On coarse soils (except sand), apply 1.0 pt./A of AX S-MET, on medium soils, apply 1.33 pts./A of AX S-MET, and on fine soils, apply 1.33-1.67 pts./A of Dual Magnum. Refer to the Canopy label for appropriate rate, according to geographical location, soil and organic matter classification, pH limitations, and all other use directions.

**Restriction:** Do not apply to sand, or to any soil with less than 0.5% organic matter, or to any soil with pH greater than 7.0, except as noted on the Canopy label.

**AX S-MET + Pursuit + Gramoxone Inteon, Touchdown Brands or Roundup Brands**

On coarse soils, apply 1.0 pt./A of AX S-MET + 0.25 pt./A of Pursuit. On medium soils, apply 1.33 pts./A of AX S-MET + 0.25 pt./A of Pursuit. On fine soils, apply 1.67 pts./A of AX S-MET + 0.25 pt./A of Pursuit.

**POSTEMERGENCE USE ON SOYBEANS - AX S-MET TANK MIXTURES**

**Tank Mixture with Glyphosate Products (e.g., Touchdown Brands or Roundup Brands)**

AX S-MET at 1.0-1.33 pts./A may be tank mixed with glyphosate products at labeled rates and applied from emergence up through the third trifoliolate leaf stage of Roundup Ready or glyphosate-tolerant soybeans. AX S-MET alone will not control emerged weeds. Use this treatment only on soybeans designated for use with glyphosate (e.g., Roundup Ready or glyphosate-tolerant soybeans). The glyphosate product must be registered for postemergence use in Roundup Ready or glyphosate-tolerant soybeans.

**Tank Mixture with Pursuit**

AX S-MET at 1.0-1.33 pts./A may be tank mixed with Pursuit at labeled rates and applied from emergence up through the third trifoliolate leaf stage of soybeans. AX S-MET alone will not control emerged weeds.

**Tank Mixture with Liberty Herbicide or Ignite 280 SL Herbicide**

AX S-MET at 1.0-1.33 pts./A may be tank mixed with Liberty Herbicide or Ignite 280SL Herbicide at labeled rates and applied from emergence up through the third trifoliolate leaf stage of soybeans. AX S-MET alone will not control emerged weeds. Use this treatment only on soybeans designated for use with glufosinate (e.g., Liberty Link).

**Restriction:** Follow the tank mix product label for adjuvant use directions. The use of COC or UAN with AX S-MET may result in temporary crop injury. To avoid possible illegal residues when AX S-MET is applied postemergence to soybeans: (1) Do not apply more than 1.33 pts./A postemergence. (2) Make postemergence applications at least 90 days before harvest. (3) Do not graze or feed treated forage or hay from soybeans to livestock following a postemergence application of AX S-MET.

**SUGAR BEETS - AX S-MET ALONE**

**Postemergence Applications**

AX S-MET may be applied postemergence to sugar beets after the sugar beets have reached the first true leaf stage. However, because AX S-MET is primarily a soil-active herbicide, it must be applied prior to weed emergence in order to provide consistent control of listed weeds. As such, weeds that are emerged with or before the crop, or that are present at the time AX S-MET is

59/61

applied, must be controlled with another appropriately labeled herbicide. Apply AX S-MET at 1 pt./A on coarse soils, 1.33 pts./A on medium soils, and 1.67 pts./A on fine soils. More than one postemergence application may be applied, but the total should not exceed 2.6 pts./A. Weeds present at the time of application will not be controlled.

**Restriction:** (1) Do not apply more than 2.67 pts./A postemergence. (2) Do not harvest within 60 days after the last application.

**Precaution:** TO AVOID CROP INJURY, In coarse soils, AX S-MET applied before emergence of sugar beets (i.e., other than postemergence) may cause injury.

**SUGAR BEETS - AX S-MET TANK MIX COMBINATIONS**

AX S-MET may tank mixed with Assure® II, Betanix®, Betanex®, Poast®, Progress®, Select®, Stinger™, or Upbeet® and applied to sugar beets. Tank mixtures of these products with AX S-MET will increase the risk of crop injury over that of either product applied alone, as the AX S-MET formulation has some adjuvant properties. The addition of a spray adjuvant such as crop oil concentrates (COC's) or methylated seed oils (MSO's) can further increase the risk of crop injury. Injury risk can be reduced by using the lowest effective rate of the tank mix partner(s) and/or adjuvant and by avoiding applications under adverse growing conditions or high soil or air humidity. Refer to the individual product labels and follow all use restrictions and limitations.

**SUNFLOWERS - AX S-MET ALONE**

**Preplant Incorporated or Preemergence**

Within the rate ranges given below, use the higher rate of AX S-MET if heavy weed infestations are expected. On coarse soils with organic matter of less than 3%, apply 1.0-1.33 pts./A of AX S-MET; 1.33 pts./A if organic matter is 3% or greater. On medium soils, apply 1.33-1.67 pts./A of AX S-MET. On fine soils with organic matter of less than 3%, apply 1.33-1.67 pts./A of AX S-MET; 1.67-2.0 pts./A if organic matter content is 3% or greater.

**Restrictions:** (1) Do not allow livestock to graze or feed in treated area. (2) Do not exceed the maximum label rates given above for sunflowers for the soil type.

**TOMATOES - AX S-MET ALONE**

**Transplanted**

AX S-MET may be applied preplant incorporated or preplant before transplanting. If the latter method is used, keep soil disturbance to a minimum during the transplanting operation. Application may also be made post-directed to transplants after the first settling rain or irrigation. When an application is made post-directed, apply in a minimum of 20 gallons of water per acre and minimize contact with tomato plants. AX S-MET will not control emerged weeds. In bedded transplanted tomatoes, apply AX S-MET preplant non-incorporated to the top of the pressed bed, as the last step, prior to laying plastic. AX S-MET may also be used to treat row-middles in bedded tomatoes, as long as the total amount of AX S-MET does not exceed the maximum allowed per crop.

**Seeded**

AX S-MET may be applied post-directed to direct seeded tomatoes. Tomato plants must be at least 4 inches tall at the time of application and the product must be applied in a minimum of 20 gallons of water per acre. Minimize spray contact with tomato plants. AX S-MET will not control emerged weeds.

**Tomato Use Rates:** On coarse soils, apply 1.0-1.33 pts./A of AX S-MET if organic matter content is less than 3% or 1.33 pts./A if organic matter is 3% or greater. On medium soils, apply 1.33-1.67

pts./A of AX S-MET. On fine soils, apply 1.33-1.67 pts./A of AX S-MET if organic matter content is less than 3% or 1.67-2.0 pts./A if organic matter content is 3% or greater.

60/  
61

**Precautions:** TO AVOID CROP INJURY (1) Do not apply to varieties or cultivars with unknown tolerance to AX S-MET. (2) AX S-MET may damage transplants that have been weakened by any cause. To prevent damage only plant healthy transplants. Do not plant when wet, cool, or unfavorable growing conditions exist. (3) In transplanted tomatoes, if AX S-MET is applied preplant incorporated, incorporate to a depth less than the depth of transplanting, and use the lower end of the rate range for the given soil type, or damage may occur. (4) For row middle applications where tomatoes are grown on sandy soils and where high soil moisture conditions can exist (i.e. low binding and high evaporation conditions), as may be found in the States of Florida, Georgia, Maryland, and Virginia, there is potential for crop injury in the form of leaf epinasty. The risk of this type of injury can be reduced by: a) incorporating the AX S-MET immediately following application, b) applying the AX S-MET seven or more days before transplanting (but only after the beds have been formed), c) minimizing the application of AX S-MET onto the plastic of the bed, or d) any combination of the above.

**Restriction:** (1) Do not apply AX S-MET within 30 days of tomato harvest. (2) Do not exceed the maximum label rate for the soil texture per year. (3) Apply only by ground application.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

#### **Pesticide Storage**

This product may be stored at temperatures down to 30 degrees below 0°F.

#### **Pesticide Disposal**

Open dumping is prohibited. Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to federal, state, or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

#### **Container handling [less than 5 gallons]**

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

#### **Container Handling [greater than 5 gallons]**

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned stay out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

61/  
61

## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of AXION AG PRODUCTS LLC or Seller, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold AXION AG PRODUCTS LLC and Seller harmless for any claims relating to such factors.

AXION AG PRODUCTS LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or AXION AG PRODUCTS LLC, and TO THE EXTENT CONSISTENT WITH APPLICABLE LAW Buyer and User assume the risk of any such use. AXION AG PRODUCTS LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither AXION AG PRODUCTS LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF AXION AG PRODUCTS LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF AXION AG PRODUCTS, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

AXION AG PRODUCTS LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of AXION AG PRODUCTS LLC.

---

Mtrex®, AAtrex Nine-O®, Beacon®, Bicep II Magnum®, Caparol®, Concep®, AX S-MET®, Dual II Magnum®, Dual IIG Magnum®, Eptam®, Eradicane®, Evik®, Exceed®, Gramoxone Inteon®, Princep®, Princep® Caliber 90®, Touchdown®, the Syngenta logo and CP FRAME [icon] are trademarks of a Syngenta Group Company  
Accent®, Assure®, Canopy®, Lorox®, Upbeet®, Viton® trademarks of E. I. duPont de Nemours and Company, Inc.  
Ag-Chem RoGator® trademark of Ag-Chem Equipment Company  
Agsorb® trademark of Oil-Dri Corporation  
Balan®, Sonalan®, Stinger®, and Treflan® trademarks of Dow AgroSciences  
Balance® Pro, Betamix®, Betanex®, Buctril®, Butyrac®, Ignite® 280 SL, Liberty®, LibertyLink®, Progress®, Sencor® trademarks of Bayer CropScience  
Banvel®, Banvel SGF®, Basagran®, Clarity®, Marksman®, Poast®, Prowl®, Pursuit®, Scepter®, and Storm® trademarks of BASF Ag Products  
Butoxone® trademark of Vertac Chemical Corporation  
Celatom MP-79® trademark of Eagle-Picher Industries, Inc.  
Command® trademark of FMC Corporation  
Cotoran® trademark of Makhteshim Agan of North America, Inc.  
Hi-Cycle™ trademark of John Deere Company  
Landmaster®, Roundup®, Roundup Ultra™, Roundup Ready®, trademarks of Monsanto Company  
Select® trademark of Valent USA  
Tyler Patriot™ trademark of Tyler Ltd. Partnership  
Weedone® trademark of NuFarm, Inc.  
Willmar Air Ride® trademark of Willmar Manufacturing

All trademarks are the property of their respective owners.