



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

89167-62

Date of Issuance:

4/1/20

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

AX Clom 5 MEG

Name and Address of Registrant (include ZIP Code):

Axion Ag Products, LLC
1880 Fall River Dr., Suite 100
Loveland, CO 80538

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 89167-62."
3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Signature of Approving Official:

Erik Kraft, Product Manager 24
Fungicide Herbicide Branch, Registration Division (7505P)

Date:

4/1/20

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 02/12/2020

If you have any questions, please contact BeWanda Alexander by phone at (703)347-0313, or via email at alexander.bewanda@epa.gov.

Enclosure

AX CLOM 5 MEG

**HERBICIDE
INTENDED FOR AGRICULTURAL OR COMMERCIAL USE ONLY**

ACTIVE INGREDIENT:	% BY WT.
Clomazone: 2-(2-chlorophenyl)methyl-4,4-dimethyl-3-isoxazolidnone	5.0%
OTHER INGREDIENTS:	<u>95.0%</u>
TOTAL:	100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION / PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail).

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300


**[See inside label booklet for First Aid, Precautionary Statements and Directions for Use.]
[SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS.]**

EPA Reg. No.: 89167-AE

EPA Est. No.: _____

NET CONTENTS: ____ Lbs.

Manufactured For:
AXION AG PRODUCTS, LLC
1880 Fall River Drive, Suite 100
Loveland, CO 80538



033020

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
HOTLINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 or your poison control center at 1-800-222-1222. For Chemical Spill, Leak, Fire or Exposure, call CHEMTREC 800-424-9300.</p>	

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Causes moderate eye irritation. Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with skin, eyes or clothing. Avoid breathing vapor. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as Barrier Laminate or Viton
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

User should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Except as directed on this label, **DO NOT** apply directly to water, to areas where surface water is present or to intertidal areas below the mean highwater mark. **DO NOT** apply when weather conditions favor drift from the area treated. **DO NOT** apply where runoff is likely to occur. **DO NOT** contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

Non-Target Organism Advisory

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, chemical-resistant gloves such as Barrier Laminate or Viton and shoes plus socks.

PRODUCT INFORMATION

AX CLOM 5 MEG is a pre-emergence and post-emergence herbicide for the control of watergrass, barnyard grass (*Echinochloa* spp.) and sprangletop (*Leptochloa fascicularis*). AX CLOM 5 MEG controls weeds through foliar and root uptake as a chlorophyll-production inhibitor. AX CLOM 5 MEG is formulated as a 5% Micro encapsulated, uniform granule.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, this product is a Group 13 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 13 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

Weed Management

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 13 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in the field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical

method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.

- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact AXION AG PRODUCTS, LLC at [855-466-8428 or 844-425-8488 or other appropriate telephone number].

Management of Resistant Biotypes

Since the occurrence of resistant weeds cannot be determined until after product use and scientific confirmation, to the extent consistent with applicable law, manufacturer is not responsible for any losses that may result from the failure of this product to control resistant weed biotypes.

The following good agronomic practices are recommended to reduce the spread of resistant biotypes:

- If a naturally occurring resistant biotype is present in your application site, this product should be tank mixed or applied sequentially with an appropriately labeled herbicide with a different mode of action to achieve control.
- Cultural and mechanical control practices (e.g. crop rotation or tillage) may also be used as appropriate.
- Scout treated application site after herbicide applications and control escaping weeds including resistant biotypes before they set seed.
- Thoroughly clean equipment before leaving fields known to contain resistant biotypes.
- Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to this Mode of Actions have been found in your region. Do not assume that each listed weed is being controlled by multiple mechanisms of action. Co-formulated active ingredients are intended to broaden the spectrum of weeds that are controlled. Some weeds may be controlled only one of the active ingredients in this product.

Integrated Pest (Weed) Management

This product may be integrated into an overall weed pest management strategy whenever the use of an herbicide is required. Practices known to reduce weed development (tillage, crop competition) and herbicide use (weed scouting, proper application timing, banding) should be followed wherever possible. Consult local agricultural and weed authorities for additional IPM strategies established for your area.

MANDATORY SPRAY DRIFT

Aerial Applications

- **DO NOT** release spray at a height greater than 10 feet above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a coarse or coarser droplet size (ASABE S572.1).
- Applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- **DO NOT** apply during temperature inversions.

Ground Boom Applications

- User must only apply with the nozzle height recommended by the manufacturer, but no more than 4 feet above the ground or canopy.
- Applicators are required to use a coarse or coarser droplet size (ASABE S572.1)
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size - Aircraft

- **Adjust Nozzles** - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT - Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

ROTATIONAL CROP RESTRICTIONS

After application of AX CLOM 5 MEG, rotational crops may not be planted prior to the intervals specified below.

Note: When using AX CLOM 5 MEG with other registered herbicides always refer to rotational restrictions and precautions on the other product label. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Crop	Crop Rotation Interval (Months)
Cotton*, Peas, Peppers, Pumpkins, (processing) Rice, Soybeans, Squash, Sweet Potatoes, Tobacco, Tuberous & Corm Vegetables	No Restriction

(Arracacha, Cassava, Tanager, Yam beans and True Yams)	
Cotton, Corn, (field, pop, seed, sweet), Cucurbits, Dry Beans, Peanuts, Potatoes, Snap Beans, Sorghum, Sugar Beets, Tomatoes, (Transplanted)	9
All crops not listed	12
<p>Cover crops, however, may be planted anytime but stand reductions may occur in some areas. DO NOT graze or harvest for food or feed cover crops planted less than 9 months after AX CLOM 5 MEG treatment.</p> <p>*Requirements for Planting Time Applications: *If cotton is the rotational crop and is planted within 9 months after the application of this product, an organophosphate insecticide either disulfoton or phorate must be applied in-furrow with the seed at planting at a minimum of 0.75 pound per acre of active ingredient. Failure to apply either disulfoton or phorate in accordance with in-furrow label use direction can result in crop phytotoxicity (bleaching) and/or stand reduction. Diuron is not advised at this planting time application as plant injury may result.</p> <p>Refer to the insecticide product labels for appropriate in-furrow application directions and maximum use rates. Monitor application equipment to insure accurate and uniform placement of the insecticide.</p>	

CROP USE DIRECTIONS

RICE

Crop	Pests	Rate of Application (Pounds per Acre)
Rice	Barnyard Grass Watergrass Spp. Sprangletop	12 (0.6 lb ai/acre)
<p>AX CLOM 5 MEG may be used at reduced rates (no less than 8 pounds (0.4 lb ai) per acre) for the control of sprangletop or for use on lighter textured soils, decomposed granitic soils or soils subject to excess drainage. When used at rates less than 12 pounds (0.6 lb ai) per acre additional use of labeled post-emergence herbicide applications may be required for control of watergrass species.</p>		

Precautions

- Apply this product to rice only as specified on this label.
- Applications of this product to fields that have been terraced or precision leveled with deep cuts may result in rice crop injury including stand loss. This precaution does not apply to normal maintenance leveling.
- Consult with Agricultural Extension Specialist or County Farm Advisors before applying this product to newly developed rice varieties.
- If temperature is in excess of 98°F for prolonged periods of time, (3 to 4 days), bleaching of rice and stand loss may occur.
- Maintain a static water flood with no exposed soil for a minimum of 5 days after a post-flood application of this product.
- Lowering of water to allow exposed soil during the first 14 days following application of this product can cause bleaching of rice, possibly resulting in rice stand loss.
- To facilitate green-up of bleached rice fields, fields need to receive a rapid flush of fresh water. For proper water circulation, functional water control devices at each end of each levee within a field are advised.

Restrictions

- **DO NOT** apply more than 12 pounds (0.6 lb ai) per acre per application.
- **DO NOT** apply more than 12 pounds (0.6 lb ai) per acre per year.
- **DO NOT** make more than 1 application per year.
- **Preharvest Interval (PHI): DO NOT** apply later than 120 days before harvesting rice.

- **DO NOT** apply where runoff or irrigation water may flow directly onto agricultural land other than rice fields.
- **DO NOT** make applications of this product within 50 feet of any sensitive crops.
- **DO NOT** apply within 300 feet of downwind crops and other desirable non-target plants.
- **DO NOT** apply within 1,200 feet of Towns and Housing Developments, Commercial Fruit, Nut or Vegetable Production¹, Commercial Greenhouses or Nurseries.
¹ Except for vegetable crops registered for use on this label.
- **Chemigation: DO NOT** apply this product through any type of irrigation system.
- **DO NOT** fish or commercially grow fish, shellfish or crustaceans on treated acres during the year of treatment.
- **DO NOT** overlap or double apply ends of fields.
- **DO NOT** apply sequential applications of other herbicides if rice is bleached or under stress from the use of this product.
- **DO NOT** apply this product to stressed rice.
- **DO NOT** drain fields for a minimum of 14 days after application, unless the water is contained within a tailwater recovery system, or other system appropriate for preventing discharge from rice fields.
- **DO NOT** allow to any other crops or sites not specified on this label.
- **DO NOT** allow this product to come into contact with adjacent crops.
- **DO NOT** apply to ditches used to transport irrigation water.
- **DO NOT** allow exposed soil in bleached fields.
- **DO NOT** use this product to manufacture or reformulate other products.

POST-FLOODED, PREPLANT USE ON RICE (WATER- SEEDED RICE AT PRE-SEEDING)

For control of watergrass and barnyard grass, (*Echinochloa* spp.) and sprangletop (*Leptochloa fascicularis*), apply 12 pounds (0.6 lb ai) of AX CLOM 5 MEG per acre to a well-prepared seedbed free of large clods. See rate exception above.

Make application immediately after flooding by air or ground and follow with the seeding of rice within 3 days after the AX CLOM 5 MEG application. Maintain water level in the checks with no exposed soil until the watergrass, barnyard grass or sprangletop die, or for a minimum of 14 days after application.

Allowing water to recede or a prolonged soil exposure will result in loss of weed control and bleaching of rice. Excessively deep water will result in plant injury. Rice injury and/or stand thinning may be evident, especially when germinating rice is subject to stress conditions. Supplemental herbicide applications may be needed for season-long weed control.

POST-FLOODED, POST-PLANT USE ON RICE (WATER- SEEDED RICE)

For control of watergrass and barnyard grass (*Echinochloa* spp.) 1-leaf or less and sprangletop (*Leptochloa fascicularis*) 1-leaf or less apply 12 pounds (0.6 lb ai) of AX CLOM 5 MEG per acre to a well-prepared seedbed free of large clods. See rate exception above. Maintain a shallow flood in the checks after application with no exposed soil, or until watergrass, barnyard grass and sprangletop die, or for a minimum of 14 days.

Allowing water to recede or a prolonged soil exposure will result in loss of weed control and bleaching of rice. Excessively deep floodwater will result in plant injury. Rice injury and/or stand thinning may be evident, especially when germinating rice is subject to stress conditions. Supplemental herbicide applications may be needed for season-long weed control.

REPLANTING INSTRUCTIONS

If initial planting of rice fails to produce a uniform stand, rice may be replanted in fields treated with AX CLOM 5 MEG.

Replanting Restrictions

- **DO NOT** retreat fields with a second application of this product.
- **DO NOT** replant treated fields with any crop at intervals that are inconsistent with the ROTATIONAL CROP RESTRICTIONS on this label.

Dealer Must Sell in Original Package Only.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage and disposal.

Pesticide Storage: Keep out of reach of children and animals. Store in original container only. Store in a cool, dry place, preferably in a locked storage area and avoid excess heat. Carefully open containers. After partial use, close tightly. **DO NOT** put concentrate or dilute material into food or drink containers. In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call CHEMTREC (transportation and Spills): **(800) 424-9300**.

To confine spill: Cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Non-refillable container. **DO NOT** reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of AXION AG PRODUCTS, LLC or Seller, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold AXION AG PRODUCTS, LLC and Seller harmless for any claims relating to such factors.

AXION AG PRODUCTS, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or AXION AG PRODUCTS, LLC, and TO THE EXTENT CONSISTENT WITH APPLICABLE LAW Buyer and User assume the risk of any such use. AXION AG PRODUCTS, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither AXION AG PRODUCTS, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF AXION AG PRODUCTS, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF AXION AG PRODUCTS, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

AXION AG PRODUCTS, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of AXION AG PRODUCTS, LLC.

All trademarks are the property of their respective owners.