



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number: **89167-139**

Date of Issuance: **1/29/26**

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

AX FLUTRIAFOL 22.7

Name and Address of Registrant (include ZIP Code):

Axion AG Products, LLC
1880 Fall River Drive, Suite 100
Loveland, CO 80538

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Continues page 2

Signature of Approving Official:

Date:

1/29/26

Manjula Unnikrishnan, Product Manager 21
Fungicide Branch, Registration Division (7505T)

2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 89167-139."
3. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 05/25/2025

If you have any questions, please contact Raven Crosby at 202-566-0829 or at crosby.raven@epa.gov.

Enclosure

{Note to reviewer: [Text] in brackets denotes optional or explanatory language}
{Note to reviewer: {Text} in braces denotes where in the final label text will appear}
{BOOKLET FRONT PANEL LANGUAGE}

FLUTRIAFOL GROUP 3 FUNGICIDE

AX FLUTRIAFOL 22.7

[For use on alfalfa[*]; almond; barley[*]; walnut[*]; celery, brassica leafy vegetables crop group 5; grapes (fresh, juice, table, wine, raisin); leafy vegetables, except brassica, crop group 4; stone fruit[*]; strawberry; tomatoes, corn (field corn, field corn grown for seed, popcorn, and sweet corn)[*]; cotton; vegetables, cucurbit, group 9[*]; vegetables, fruiting, group 8-10; hops[*], peanut[*]; pome fruit (apple, crabapple, loquat, mayhaw, pear, oriental pear, quince)[*]; grain sorghum[*]; soybean[*]; sugar beet[*]; pecan and other tree nuts[*]; triticale[*]; wheat (spring and winter)[*].
[*Not Registered for Use By California]]

ACTIVE INGREDIENT:	(% by weight)
*Flutriafol.....	22.7%
OTHER INGREDIENTS:	77.3%
TOTAL	100.0%

*Contains 2.08 pounds flutriafol per gallon.

Suspension concentrate.

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

[See inside label booklet for [additional] Precautionary Statements, and Directions for Use.]

[See below additional Precautionary Statements] [See First Aid statement on back panel] [of booklet].

[Read the entire label before using this product.]

EPA Reg. No.: 89167-RGO

EPA Est. No.:

Net Contents: ___ Gal

A C C E P T E D

01/29/2026

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No.

89167-139

Manufactured For:

AXION AG PRODUCTS, LLC
1880 Fall River Drive, Suite 100
Loveland, CO 80538

RD120825

{LANGUAGE INSIDE BOOKLET}

FIRST AID	
If in eyes:	<ul style="list-style-type: none">Hold eye open and rinse slowly and gently with water for 15-20 minutes.Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">Call a poison control center or doctor immediately for treatment advice.Have person sip a glass of water if able to swallow.DO NOT induce vomiting unless told to do so by the poison control center or doctor.DO NOT give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none">Take off contaminated clothing.Rinse skin immediately with plenty of water for 15-20 minutes.Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact CHEMTRAC Day or Night at 1-800-424-9300 for emergency medical treatment information.	
Note to Physician: No specific antidote. Treat symptomatically.	

For Chemical Emergency:
Spill, Leak, Fire, Exposure, or Accident,
Call CHEMTRAC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes substantial but temporary eye injury. Harmful if swallowed. **DO NOT** get in eyes, or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks,
- Chemical-resistant gloves (barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinyl chloride \geq 14 mils, or Viton[®] \geq 14 mils),
- Protective eyewear

Mixers and Loaders must wear:

- Long-sleeved shirt and long pants, Shoes plus socks,
- Chemical-resistant gloves (barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinyl chloride \geq 14 mils, or Viton[®] \geq 14 mils)
- Protective eyewear

Follow manufacturer's instructions for cleaning / maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This formulated product is toxic to fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Flutriafol has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in any manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

{Note to Reviewer: Below is optional state-specific language.}

[Optional language for CA: In California, this product may only be applied to fields that are not irrigated, or are irrigated using pressurized irrigation methods such as macro-sprinkler, micro-sprinkler, or drip systems.]

[Optional language for NY: Not for sale, distribution, or use in Nassau or Suffolk Counties of New York State.]

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the Restricted Entry Interval (REI). The REI for each crop is located in the use directions for each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- coveralls,
- chemical resistant gloves,
- shoes plus socks, and
- protective eyewear.

PRODUCT DIRECTIONS

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH RESTRICTIONS, PRECAUTIONARY STATEMENTS, AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Not for use in greenhouses.

Product can be applied for all crops either by foliar application or chemigation unless otherwise specified in this label.

Chemigation:

Application through Irrigation Systems (Chemigation):

Apply **AX FLUTRIAFOL 22.7** through irrigation to crops at rates and timings specified in this label. Chemical tank and injector system must be thoroughly cleaned before and after use. Flush system with clean water.

Crop injury or lack of effectiveness can result from non-uniform distribution of **AX FLUTRIAFOL 22.7** across the treated area. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Drip Irrigation: Use **AX FLUTRIAFOL 22.7** for control of diseases at rates and timing as specified in this label. Ensure adequate soil moisture prior to utilizing **AX FLUTRIAFOL 22.7** in a drip irrigation system. **DO NOT** apply **AX FLUTRIAFOL 22.7** through drip irrigation that is deeper than 12" below the soil surface.

After application of **AX FLUTRIAFOL 22.7**, continue to drip irrigate the area with water to flush the irrigation system. **DO NOT** allow **AX FLUTRIAFOL 22.7** to remain in the irrigation system after the application is complete. The total volume of water, including the amount used for flushing the irrigation system, must be adequate to completely remove the product from the lines, but should be less than the amount that could over-saturate the beds (bed collapse can occur from over-saturation) in bedded production systems. Delay subsequent irrigation (water only) for at least 24 hours following drip application for best results.

AX FLUTRIAFOL 22.7 Rate (fl oz /A)	Fluid Ounces A2123.02/1000 Row Feet Based on Planted Row Spacing												
	30" Rows	34" Rows	36" Rows	38" Rows	40" Rows	44" Rows	48" Rows	60" Rows	66" Rows	72" Rows	80" Rows	84" Rows	120" Rows
5 (0.81 lb ai)	0.29	0.33	0.34	0.36	0.38	0.42	0.46	0.57	0.63	0.69	0.77	0.80	1.15
7 (0.114 lb ai)	0.40	0.46	0.48	0.51	0.54	0.59	0.64	0.80	0.88	0.96	1.07	1.12	1.61

Sprinkler Irrigation: Apply this product through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, micro and macro sprinkler or hand move irrigation systems. **DO NOT** apply this product through any other type of irrigation system except as specified on this label.

With Center Pivot irrigation equipment, use **AX FLUTRIAFOL 22.7** only with equipment with drive systems that provide uniform water distribution. **DO NOT** use end guns for chemigation due to the non-uniform application pattern.

Add **AX FLUTRIAFOL 22.7** to the pesticide supply tank containing sufficient water to maintain a continuous flow by the injection equipment. Maintain agitation during the entire application period.

For continuous-move irrigation systems, apply the labeled rate for that crop in ½ acre-inch or less per acre. For stationary or non-continuous moving systems, inject **AX FLUTRIAFOL 22.7** spray mixture during the last 20-30 minutes of the set.

DO NOT apply when winds are greater than 10-15 mph to avoid drift or wind skips. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. Thorough coverage of foliage is required for good control.

Operating Requirements for Application through Irrigation Systems:

1. **DO NOT** use chemigation when conditions are favorable for drift to non-target areas.
2. To prevent water-source contamination from backflow, a functional check valve, vacuum relief valve, and low-pressure drain should be located on the irrigation pipeline.
3. To prevent backflow back toward the injection pump, the pesticide injection pipeline must be equipped with a functional, automatic, quick-closing check valve.
4. To prevent fluid from being withdrawn from the supply tank when the irrigation system is shut down, the pesticide injection pipeline should also be equipped with a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock.
5. The system must also contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops and a functional pressure switch to stop the pump motor when water pressure decreases to a point at which pesticide distribution is adversely affected.
6. A metering pump, constructed of materials compatible with pesticides and capable of being fitted with a system interlock, such as a positive displacement injection pump (e.g., a diaphragm pump), must be included in the system.
7. A knowledgeable person responsible for the chemigation system should shut the system down and turn the irrigation water off, ensuring enough time for the pesticide to be flushed through all lines and nozzles.
8. No irrigation system, including those in greenhouses, used to distribute pesticides can be connected to a public water source unless safety measures and devices prescribed in the pesticide label for such connection are in place.

Specific Instructions for Public Water Systems:

1. Public water system means a system that provides piped water for human consumption if the system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. All measures and devices listed in the above section, "Operating Requirements for Application through Irrigation Systems," must be operational for connection to a public water system.
3. Additionally, chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There must be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

Rainfastness: **AX FLUTRIAFOL 22.7** is rainfast 2 hours after application. Disease control will be reduced if rainfall occurs within 2 hours of application.

Crop Rotation: Fields treated with **AX FLUTRIAFOL 22.7** may be planted to the following crops and plant back intervals:

Crop	Plant back Interval
Alfalfa, Barley, Field Corn; Field Corn Grown for Seed; Popcorn; Sweet Corn; Cotton; Cucurbits; Fruiting Vegetables; Grapes; Hops; Brassica Leafy Vegetables Crop Group 5; Leafy Vegetables, Except Brassica, Crop Group 4; Peanut; Pome Fruit; Grain Sorghum; Soybean; Stone Fruit; Strawberry; Sugar Beet; Tomato; Tree Nuts; Triticale; or Wheat	Immediately
Rice	30 days
All other crops	Prohibited

RATE EQUIVALENCY TABLE

Product Rate (Fl oz/A)	Equivalent Active Ingredient Rate (lb ai/A)
3.5	0.057
4	0.065
5	0.081
6	0.098
6.5	0.106
7	0.114
8	0.130
12	0.20
15	0.24
16	0.26
24	0.39

Use of Adjuvants: An adjuvant may be used with **AX FLUTRIAFOL 22.7** unless otherwise specified in the crop use directions. **AX FLUTRIAFOL 22.7** is compatible with most products, however not all have been tested. Use the compatibility jar test to ensure physical compatibility.

Tank Mixing: **AX FLUTRIAFOL 22.7** may be tank mixed with other fungicides, herbicides, insecticides, and/or other additives unless prohibited on the label of the tank mix partner. Follow more restrictive labeling of any tank mix partner. Although **AX FLUTRIAFOL 22.7** is compatible with most products, not all combinations have been tested. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Use the compatibility jar test to ensure physical compatibility. Before applying any tank mixture not specifically recommended on this label, the crop safety of the target crop should be confirmed by applying the mixture to a small area of the target crop according to the label instructions.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Compatibility Jar Test: **AX FLUTRIAFOL 22.7** is compatible with most products, however not all have been tested. Use the following compatibility test to ensure physical compatibility.

Using a quart jar, add the proportionate amounts of the products to approximately one quart of water with agitation. Add wettable powders and water dispersible granular products first, next liquid flowables, then emulsifiable concentrates, and last liquid soluble products. After thorough mixing, allow this mixture to stand for 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible. Once compatibility has been proven, use the same procedure for adding products to the spray tank. When tank mixing, follow more restrictive labeling of any tank mix partner. **DO NOT** tank mix with any product that contains a prohibition on tank mixing.

Spray Equipment/Volume: Apply **AX FLUTRIAFOL 22.7** in sufficient water volume to ensure thorough coverage as a broadcast foliar application for good disease control.

For ground application, use a minimum of 30 gallons of spray solution per acre by ground sprayer for hops, and 10 gallons of spray solution per acre by ground sprayer for all other crops. For aerial application, use aircraft spray equipment in a minimum of 10 gallons of spray solution per acre for tree nut crops and hops, 2 gallons of spray solution per acre for corn[*], cotton, grain sorghum[*], soybean[*], barley[*], wheat[*] and triticale[*], and 5 gallons of spray solution per acre for all other crops.

[*Not Registered for Use By California]

High Disease Pressure: Where a range is given, use the higher specified rate of **AX FLUTRIAFOL 22.7** and/or reduced spray interval when disease pressure is high and/or conditions are favorable for disease development.

Resistance Management:

For resistance management, **AX FLUTRIAFOL 22.7** contains a Group 3 fungicide (Flutriafol). Any fungal population may contain individuals naturally resistant to **AX FLUTRIAFOL 22.7** and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies must be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of **AX FLUTRIAFOL 22.7** or other Group 3 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM specifications for specific crops and pathogens.
- For further information or to report suspected resistance contact your pesticide distributor or university extension specialist to report resistance.

{**NOTE TO REVIEWER:** Registrant may add or remove the following state restriction statement as required throughout. (e.g., NOT REGISTERED FOR USE BY CALIFORNIA)}

[Not Registered For Use By _____]

CROP USE DIRECTIONS

Use Directions for ALFALFA[*]

AX FLUTRIAFOL 22.7 works by forming a protective barrier around the alfalfa root at the point of disease infection. Precipitation in the form of rainfall or irrigation is required to move the product into the infection zone. The amount of water required will vary with soil type, soil moisture, and rainfall or irrigation intensity.

[*Not Registered for Use By California]

Application Method:

Apply **AX FLUTRIAFOL 22.7** as a broadcast spray application at a minimum spray volume of 10 gallons per acre or 5 gallons by air, or through sprinkler irrigation systems designed as described in the Chemigation section of this label. For continuous-move irrigation systems, apply the labeled rate in $\frac{1}{2}$ acre-inch or more per acre. For stationary or non-continuous moving systems, inject **AX FLUTRIAFOL 22.7** spray mixture during the first 20-30 minutes of the set to ensure fungicide reaches the soil surface. Make applications postemergence to alfalfa at the timing(s) outlined below.

Disease	Product Rate (fl oz/A)	Application Directions
Texas Root Rot / Cotton Root Rot (<i>Phymatotrichopsis omnivora</i>)	15 – 24 (0.24 – 0.39 lb ai/A)	<p>Single Application: Apply in early spring prior to three (3) inches of new alfalfa growth OR after first cut of alfalfa when alfalfa is between two (2) and three (3) inches regrowth.</p> <p>Use the highest specified product rate when field is known to have high levels of cotton root rot disease pressure.</p>
	First application: 15 – 24 (0.24 – 0.39 lb ai/A)	<p>Split Application: Apply first application in early spring prior to three (3) inches of new alfalfa growth.</p> <p>Apply second application after second or third cutting of alfalfa when alfalfa is between two (2) and three (3) inches regrowth.</p>
	Second application: 12 - 15 (0.2 – 0.24 lb ai/A)	<p>Use split application under the following situations:</p> <ul style="list-style-type: none"> • High intensity production practices • 2nd year of established plantings and older <p>Use the highest specified product rate when field is known to have high levels of cotton root rot disease pressure.</p>

Restrictions:

- No single application may exceed 24 fl oz of product/A (0.39 lb ai/A).
- Minimum Retreatment Interval (RTI): 28 days
- Make only one application per cutting.
- **DO NOT** apply more than two applications per when applied at reduced rates.
- **DO NOT** exceed 39 fl oz of product (0.63 lb ai)/A per year.
- Preharvest Interval (PHI): **DO NOT** apply within 14 days of harvest.
- Restricted Entry Interval (REI) is 12 hours

Use Directions for Almond

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Brown Rot Blossom Blight (<i>Monilinia</i> spp.)	7 (0.114 lb ai/A)	For Brown Rot Blossom Blight start applications at 5 - 10% bloom followed by an application at 50 - 100% bloom.	7
Alternaria Leaf and Fruit Spot (<i>Alternaria alternata</i>)			
Anthracnose (<i>Colletotrichum acutatum</i>)		For all other diseases apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	
Scab (<i>Cladosporium carpophilum</i>)			
Shot Hole (<i>Wilsonomyces carpophilus</i>)			
Leaf Rust (<i>Tranzschelia discolor</i>)			

Restrictions:

- No single application may exceed 7 fl oz of product (0.114 lb ai)/A.
- The minimum retreatment interval is 7 days.
- **DO NOT** apply more than 4 applications per year.
- **DO NOT** apply more than 28 fl oz of product (0.455 lb ai)/A per year.
- **DO NOT** apply more than 0.455 lb ai of flutriafol/A per year.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Interval (PHI): **DO NOT** apply within 14 days of harvest.

Use Directions for Almond[*]

[*Not Registered for Use By California]

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Botryosphaeria Canker (<i>Botryosphaeria</i> spp.)	7 (0.114 lb ai/A)	Apply (via drip irrigation or foliar application) preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development.	7

Restrictions:

- No single application may exceed 7 fl oz of product (0.114 lb ai)/A.
- The minimum retreatment interval is 7 days.
- **DO NOT** apply more than 6 applications per year.
- **DO NOT** apply more than 42 fl oz of product (0.684 lb ai)/A per year.
- **DO NOT** apply more than 0.684 lb ai of flutriafol/A per year.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Interval (PHI): **DO NOT** apply within 14 days of harvest.

Use Directions for Walnut [[including] black and English (Persian) walnuts] [*]

[*Not Registered for Use By California]

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Botryosphaeria Panicle and Shoot Blight Phomopsis Canker	7 (0.114 lb ai/A)	Apply preventatively and repeat as necessary on a 7-day interval when conditions are favorable for disease development.	7

Restrictions:

- No single application may exceed 7 fl oz of product (0.114 lb ai)/A.
- The minimum retreatment interval is 7 days.
- **DO NOT** apply more than 4 applications per year.
- **DO NOT** apply more than 28 fl oz of product (0.455 lb ai)/A per year.
- **DO NOT** apply more than 0.455 lb ai of flutriafol/A per year.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Interval (PHI): **DO NOT** apply within 14 days of harvest.

Use Directions for Brassica Leafy Vegetables Crop Group 5. See below for a list of crops.

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Black Spot (<i>Alternaria</i> spp.) Cercospora Leaf Spot (<i>Cercospora</i> spp.) Powdery Mildew (<i>Erysiphe polygoni</i>)	5 – 7 (0.081- 0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development.	7
List of Brassica Leafy Vegetables Crop Group 5: broccoli; broccoli Chinese (gai lon); broccoli raab (rapini); Brussels sprouts; cabbage; cabbage, Chinese (bok choy); cabbage, Chinese (napa); cabbage, Chinese mustard (gai choy); cauliflower; cavolo broccolo; collards; kale; kohlrabi; mizuna; mustard greens; mustard spinach; rape greens and cultivars and/or hybrids of these crops.			
<p>Restrictions:</p> <ul style="list-style-type: none"> • No single application may exceed 7 fl oz of product (0.114 lb ai)/A. • The minimum retreatment interval is 7 days. • DO NOT apply more than 4 applications per year. • DO NOT apply more than 28 fl oz of product (0.455 lb ai)/A per year. • DO NOT apply more than 0.455 lb ai of flutriafol/A per year. • Restricted Entry Interval (REI) is 12 hours. • Preharvest Interval (PHI): DO NOT apply within 7 days of harvest. 			

Use Directions for Celery and Chinese Celery

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Alternaria Leaf Spot (<i>Alternaria sonchi</i> , A. spp.) Anthracnose (<i>Colletotrichum dematium</i>) Early Blight (<i>Cercospora apii</i>) Late Blight (<i>Septoria apicola</i>)	5 – 7 (0.081- 0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	7
Fusarium Yellows /Fusarium Wilt [*] (Suppression only) (<i>Fusarium oxysporum</i> f.sp <i>apii</i>) [*Not Registered for Use By California]	5 – 7 (0.081- 0.114 lb ai/A)	Product is to be applied through drip irrigation. It is advised to make initial application at transplanting. For spring and fall planting, a second application is required, approximately 30 days later. For fall planting, additional applications may be required at 14 day intervals, for a maximum of 4 applications per year.	14

Restrictions:

- No single application may exceed 7 fl oz of product (0.114 lb ai)/A.
- The minimum retreatment interval is 7 days.
- **DO NOT** apply more than 4 applications per year.
- **DO NOT** apply more than 28 fl oz of product (0.455 lb ai)/A per year.
- **DO NOT** apply more than 0.455 lb ai of flutriafol/A per year.
- Restricted Entry Interval (REI): The REI is 12 hours.
- Preharvest Interval (PHI): **DO NOT** apply within 7 days of harvest.

Use Directions for Grapes [(fresh, juice, table, wine, raisin)]

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Powdery Mildew (<i>Uncinula necator</i>)	4 – 5 (0.065- 0.081 lb ai/A)	[*] Foliar Application: Begin application at 6 to 10 inch shoot growth. [*Not Registered for Use By California]	14 – 21
	5 (0.081 lb ai/A)	[*] Drip Irrigation Application: Apply preventively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development and to provide extended foliar protection. Best results have been obtained when AX FLUTRIAFOL 22.7 is applied preventively, as early as bud break, to ensure time for movement of flutriafol into the foliage. Once fruit are present, foliar applications are needed for fruit protection. [*Not Registered for Use By California]	14
Black Rot (<i>Guignardia bidwellii</i>)	4 – 5 (0.065- 0.081 lb ai/A)	[*] Foliar Application: Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development. [*Not Registered for Use By California]	14 – 21
Restrictions:			
<ul style="list-style-type: none"> • No single application may exceed 5 fl oz of product (0.081 lb ai)/A. • The minimum retreatment interval is 14 days. • DO NOT apply more than 6 applications per year. • DO NOT apply more than 30 fl oz of product (0.488 lb ai)/A per year. • DO NOT apply more than 0.488 lb ai of flutriafol/A per year. • Restricted Entry Interval (REI): <ul style="list-style-type: none"> ○ For foliar applications: The REI is 5 days for girdling or turning of grapes. ○ For soil-applied applications[*]: The REI is 12 hours for girdling or turning of grapes. ○ The REI for all other activities is 12 hours. • Preharvest Interval (PHI): DO NOT apply within 14 days of harvest. <p>[*Not Registered for Use By California]</p>			

Use Directions for Leafy Vegetables, except Brassica, Crop Group 4. See below for a list of crops.

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Alternaria Leaf Spot (<i>Alternaria sonchi</i> , A. spp.)	5 – 7 (0.081- 0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development.	7
Anthracnose (<i>Microdochium panattonianum</i> , <i>Colletotrichum dematium</i>)			
Cercospora Leaf Spot (<i>Cercospora</i> spp.)			
Powdery Mildew (<i>Erysiphe cichoracearum</i> , <i>Leveillula taurica</i>)			
Septoria Leaf Spot (<i>Septoria</i> spp.)			

List of Leafy Vegetables, except Brassica, Crop Group 4: amaranth (Chinese spinach); amaranth (leafy); amaranth (tampala); arugula (roquette); cardoon; celtuce; chervil; chrysanthemum (edible-leaved); chrysanthemum (garland); corn salad; cress (garden); cress (upland); dandelion (leaves); dock (sorrel); endive (escarole); fennel, Florence; lettuce (head and leaf); orach; parsley; purslane (garden); purslane (winter); radicchio (red chicory); rhubarb; spinach; spinach (New Zealand); spinach (vine); Swiss chard and cultivars and/or hybrids of these crops.

Restrictions:

- No single application may exceed 7 fl oz of product (0.114 lb ai)/A.
- The minimum retreatment interval is 7 days.
- **DO NOT** apply more than 4 applications per year.
- **DO NOT** apply more than 28 fl oz of product (0.455 lb ai)/A per year.
- **DO NOT** apply more than 0.455 lb ai of flutriafol/A per year.
- Restricted Entry Interval (REI): 12 hours.
- Preharvest Interval (PHI): **DO NOT** apply within 7 days of harvest.

Use Directions for Stone Fruit [*]. See below for a list of crops.

[*Not Registered for Use By California]

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Powdery Mildew (<i>Sphaerotheca pannosa</i>)	7 (0.114 lb ai/A)	Start applications at petal fall. Repeat as necessary if disease conditions are favorable.	7
Rusty spot [*] (<i>Podosphaera leucotricha</i>)	7 (0.114 lb ai/A)	Start applications at petal fall and continue through 3rd Cover (3C) or until terminal shoot growth ceases and shoots harden off or conditions are no longer favorable for disease development.	7
Brown Rot Blossom Blight (<i>Monilinia</i> spp.)	7 (0.114 lb ai/A)	Start applications at 1-5% bloom followed by an application at 50-100% bloom.	7
Fruit Brown Rot (<i>Monilinia</i> spp.)	7 (0.114 lb ai/A)	Apply one to two weeks prior to harvest.	7
Shot Hole (<i>Wilsonomyces carpophilus</i>)	7 (0.114 lb ai/A)	Apply at 50-100% bloom. Repeat as necessary if disease conditions are favorable for disease development.	7
Cherry Leaf Spot (<i>Blumeriella jaapii</i>)	7 (0.114 lb ai/A)	Start applications at petal fall. Repeat as necessary if disease conditions are favorable for disease development.	7

List of Stone Fruit: Apricot, Sweet Cherry, Tart Cherry, Nectarine, Peach, Plum, Chickasaw Plum, Damson Plum, Japanese Plum, Plumcot, Prune.

Restrictions:

- No single application may exceed 7 fl oz of product (0.114 lb ai)/A.
- The minimum retreatment interval is 7 days.
- **DO NOT** apply more than 4 applications per year.
- **DO NOT** apply more than 28 fl oz of product (0.455 lb ai)/A per year.
- **DO NOT** apply more than 0.455 lb ai of flutriafol/A per year.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Interval (PHI): **DO NOT** apply within 7 days of harvest.

Use Directions for Strawberry

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Powdery Mildew (<i>Podosphaera aphanis</i>) Leaf Rust[*] (<i>Phragmidium potentillae</i>) Leaf Spot[*] (<i>Mycosphaerella fragariae</i>) [*Not Registered for Use By California]	5 – 7 (0.081- 0.114 lb ai/A)	<p>Foliar Application: Apply preventatively or when conditions are favorable for disease development.</p> <p>Repeat as necessary if conditions are favorable for disease development.</p>	7
	7 (0.114 lb ai/A)	<p>[*]Drip Irrigation: Apply preventatively or when conditions are favorable for disease development.</p> <p>Repeat as necessary if conditions are favorable for disease development. Best results for both foliar and soilborne pathogens have been obtained when AX FLUTRIAFOL 22.7 is applied preventively, shortly after transplanting, to provide protection of the roots from soilborne pathogens and to ensure time for movement of flutriafol into the foliage for protection against foliar pathogens. Once fruit are present, foliar applications are needed for fruit protection.</p> <p>[*Not Registered for Use By California]</p>	14
Anthracnose[*] (<i>Colletotrichum acutatum</i>) [*Not Registered for Use By California]	7 (0.114 lb ai/A)	<p>[*]Drip Irrigation: Apply preventatively or when conditions are favorable for disease development.</p> <p>Repeat as necessary if conditions are favorable for disease development. Best results for both foliar and soilborne pathogens have been obtained when AX FLUTRIAFOL 22.7 is applied preventively, shortly after transplanting, to provide protection of the roots from soilborne pathogens and to ensure time for movement of flutriafol into the foliage for protection against foliar pathogens. Once fruit are present, foliar applications are needed for fruit protection.</p> <p>[*Not Registered for Use By California]</p>	14
Charcoal Rot[*] (<i>Macrophomina phaseolina</i>) [*Not Registered for Use By California]	7 (0.114 lb ai/A)	<p>Product is to be applied through drip irrigation (see Chemigation for directions). It is advised to make initial application at transplanting. For spring and fall planting sequential applications are recommended on a 14 day interval.</p> <p>[In CA: For spring and fall planting, a second application is required approximately 30 days later. For fall planting, additional applications may be required at 30-day intervals.]</p>	14 [In CA: 30]

Restrictions:

- No single application may exceed 7 fl oz of product (0.114 lb ai)/A.
- The minimum retreatment interval is 7 days.
- **DO NOT** apply more than 4 applications per year.
- **DO NOT** apply more than 28 fl oz of product (0.228 lb ai)/A per year.
- **DO NOT** apply more than 0.455 lb ai of flutriafol/A per year.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Interval (PHI): May be applied the day of harvest (0 day PHI).

Use Directions for Tomatoes. See below for a list of crops.

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Powdery Mildew (<i>Leveillula Taurica</i> , <i>Oidium neolycorensis</i>) Anthracnose (<i>Collectotrichum coccodes</i>) Target Spot (<i>Corynespora cassiicola</i>) Septoria Leaf Spot[*] (<i>Septoria lycopersici</i>) Cercospora Leaf Spot[*] (<i>Cercospora capsici</i>) [*Not Registered for Use By California]	Foliar: 5 – 7 (0.081- 0.114 lb ai/A) [*] Drip: 7 (0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	Foliar – 7 [*] Drip - 14
Black Mold (<i>Alternaria alternata</i>) Early Blight (<i>Alternaria solani</i>)	3.5 – 7 (0.057- 0.114 lb ai/A)	Tank mix with mancozeb for improved disease control. Lower rate can be used when tank mixed with mancozeb or other effective fungicides.	7
Charcoal Rot [*] (<i>Macrophomina phaseolina</i>) [*Not Registered for Use By California]	7 (0.114 lb ai/A)	Apply through drip irrigation (see Chemigation for directions). It is advised to make initial application at transplanting. A second application is required approximately 14 days later. Two additional applications may be applied if necessary. [In CA: A second application is required approximately 30 days later.]	14 [In CA: 30]
Fusarium Wilt [*] (Suppression) (<i>Fusarium oxysporum f. sp lycopersici</i>) [*Not Registered for Use By California]	7 (0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development. Best results for soilborne pathogens has been obtained when AX FLUTRIAFOL 22.7 is applied preventively, shortly after transplanting, followed by subsequent applications at 14 day intervals to provide	14

		protection of the roots from soilborne pathogens.	
List of Tomatoes: Bush Tomato, Cocona, Currant Tomato, Garden Huckleberry, Goji Berry, Groundcherry, Narnajilla, Sunberry, Tomatillo, and Tree Tomato and cultivars, varieties and/or hybrids of these crops.			
Restrictions:			
<ul style="list-style-type: none"> • No single application may exceed 7 fl oz of product (0.114 lb ai)/A. • The minimum retreatment interval is 7 days. • DO NOT apply more than 4 applications per year. • DO NOT apply more than 28 fl oz of product/A per year. • DO NOT apply more than 0.455 lb ai of flutriafol/A per year. • Restricted Entry Interval (REI) is 12 hours. • Preharvest Interval (PHI): May be applied the day of harvest (0 day PHI). When tank mixed with mancozeb fungicide, the PHI is 5 days. 			

Use Directions for Corn (Field Corn, Field Corn Grown for Seed, Sweet Corn, and Popcorn) [*] (Foliar Applications)

[*Not Registered for Use By California]

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Gray Leaf Spot (<i>Cercospora zeae-maydis</i>)	3.5 - 7 (0.057- 0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. One repeat application may be made.	7
Southern Corn Leaf Blight (<i>Cochliobolus heterostrophus</i>)		Apply no later than growth stage R4 (early dough stage).	
Northern Corn Leaf Blight (<i>Setosphaeria turcica</i>)		For aerial application, use a minimum of 2 gallons of spray solution per acre.	
Common rust (<i>Puccinia sorghi</i>)		An adjuvant may be used with AX FLUTRIAFOL 22.7 prior to the V8 corn growth stage and after the VT corn growth stage.	
Southern Rust (<i>Puccinia polysora</i>)			
Eyespot (<i>Aureobasidium zeae</i>)			
Anthracnose Leaf Blight (<i>Colletotrichum graminicola</i>)			
Northern Corn Leaf Spot (<i>Cochliobolus carbonum</i>)			
Tar Spot (<i>Phyllachora maydis</i>)			

Restrictions:

- No single application may exceed 7 fl oz of product (0.114 lb ai)/A.
- The minimum retreatment interval is 7 days.
- **DO NOT** apply more than 2 applications per year.
- **DO NOT** apply more than 14 fl oz of product (0.228 lb ai)/A per year.
- **DO NOT** apply more than 0.228 lb ai of flutriafol/A per year.
- Restricted Entry Interval (REI):
 - **For detasselling field corn and popcorn grown for seed** is 5 days.
 - **For sweet corn** is 3 days.
 - **For all other activities** is 12 hours.
- Preharvest Interval (PHI):
 - **Harvest for forage:** Zero (0) days
 - **Stover and grain of field corn and popcorn and sweet corn:** 7 days

Use Directions for Cotton [*] (Foliar Application Only)

[*Not Registered for Use By California]

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Alternaria Leaf Spot, Boll Rot (<i>Alternaria</i> spp.) Anthracnose, Boll Rot (<i>Glomerella</i> spp.) Ascochyta Blight, Boll Rot (<i>Ascochyta gossypii</i>) Cercospora Blight and Leaf Spot (<i>Cercospora</i> spp.) Diplodia Boll Rot (<i>Diplodia</i> spp.) Fusarium Boll Rot (<i>Fusarium</i> spp.) Phoma Blight, Boll Rot (<i>Phoma</i> spp.) Rust (<i>Puccinia</i> spp., <i>Phyropsora</i> spp.) Stemphylium Leaf Spot (<i>Stemphylium</i> spp.) Target Spot (<i>Corynespora cassiicola</i>)	5 -7 (0.081- 0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	7

Restrictions:

- No single application may exceed 7 fl oz of product (0.114 lb ai)/A.
- The minimum retreatment interval is 7 days.
- **DO NOT** apply more than 2 foliar applications of product (0.228 lb ai) per year.
- **DO NOT** apply more than 14 fl oz (0.576 lb ai of flutriafol)/acre per year for foliar applications.
- **DO NOT** apply more than 30 fl oz of product (0.488 lb ai of flutriafol)/acre per year (including foliar and soil applications.)
- Restricted Entry Interval (REI): The REI is 12 hours.
- Preharvest Interval (PHI): **DO NOT** apply within 30 days of harvest.

Use Directions for Cotton[*] (Soil Applications)

[*Not Registered for Use By California]

Diseases	Product Rate (fl oz/A)
Cotton Root Rot (<i>Phymatotrichopsis omnivora</i>)	8-16 (0.13-0.26 lb ai/A) (See cotton row spacing table below for rate chart depending on row width)
Application Directions	
Application Methods:	
<p>T-Band Application: Apply AX FLUTRIAFOL 22.7 in a concentrated 3-4 inch wide band at planting perpendicular to row direction after furrow opening and seed placement but prior to furrow closure with press wheel.</p> <p>Modified In-Furrow Application: Apply using a splitter/Y-shaped application mechanism that directs the product on the sides of seed furrow and not in direct contact with the seed.</p> <p>Drip Irrigation: This product may be applied through drip irrigation systems where drip tape is not deeper than 12 inches below the soil surface. Apply only through drip irrigation systems configured with drip tape directly below the row and running in the same direction as the rows.</p>	

Application Methods to use by Field Irrigation Type

Overhead or Sprinkler Irrigation Fields:

- T-Band is the preferred application method under these cropping systems.
- Modified In-Furrow can be used. Avoid applying product in direct contact with seed.

Dryland Fields:

- Modified In-Furrow application technique may provide more consistent control under low rainfall conditions.
- Application using a T-Band method requires rainfall to move the product into the disease infection zone below the soil surface.

Furrow and Drip Irrigated Fields:

- Apply in T-Band, Modified In-Furrow, or through drip tape.
- When using the Modified In-Furrow application method sufficient irrigation must be applied to thoroughly wet the **AX FLUTRIAFOL 22.7** treated zone after cotton has emerged.
- For T-Band applications, the top of the bed must be thoroughly wetted after the cotton has emerged.
- Follow instructions in chemigation section of this label when applying through drip irrigation systems.

NOTICE for All Applications Methods and Field Conditions: Heavy rainfall or irrigation within 3 days after planting may delay emergence.

Restrictions:

- No single application may exceed 16 fl oz of product (0.26 lb ai)/A.
- **DO NOT** apply more than one soil application per year.
- **DO NOT** apply more than 16 fl oz (0.26 lb ai of flutriafol)/acre per year for soil applications.
- **DO NOT** apply more than 30 fl oz of product (0.488 lb ai flutriafol)/ acre per year (including soil and foliar applications).
- **DO NOT** apply with liquid fertilizer.
- Restricted Entry Interval (REI) for all activities is 12 hours.
- Preharvest Interval (PHI): **DO NOT** apply within 30 days of harvest.

Cotton Row Spacing Table

Cotton Row Spacing (Inches)	Fluid Ounces of Product to be Applied / 1000 Row Feet	
	8 Fl oz/A Product Rate (0.13 lb ai)	16 Fl oz/A Product Rate (0.26 lb ai)
30	0.46 (0.007 lb ai)	0.92 (0.015 lb ai)
32	0.49 (0.008 lb ai)	0.98 (0.016 lb ai)
36	0.55 (0.0089 lb ai)	1.10 (0.018 lb. ai)
38	0.58 (0.0094 lb ai)	1.16 (0.019 lb ai)
40	0.61 (0.010 lb ai)	1.22 (0.020 lb ai)

30" = 17,424 row ft per acre, 32" = 16,335 row ft per acre,

36" = 14,520 row ft per acre, 38" = 13,754 row ft per acre, 40" = 13,068 row ft per acre.

Formula:

Selected Use Rate (fl oz/A) = fl oz of product to be applied per 1000 row ft.
[Row ft. for row spacing / 1000 ft]

Example for 30" row spacing at 8 fl oz/A:

8 fl oz/A = 0.46 fl oz of product to be applied per 1000 row ft
[17,424 / 1000 ft]

Use Directions for Vegetables, Cucurbit, Group 9[*]. See below for a list of crops.

[*Not Registered for Use By California]

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
<p>Anthracnose[*] (<i>Colletotrichum</i> spp.)</p> <p>Powdery Mildew (<i>Sphaerotheca fuliginea</i>, <i>Erysiphe cichoracearum</i>)</p> <p>Cercospora Leaf Spot[*] (<i>Cercospora citrulline</i>)</p> <p>Gummy Stem Blight (<i>Didymella bryoniae</i>)</p> <p>Target Spot (<i>Corynespora cassiicola</i>)</p> <p>[*Not Registered for Use By California]</p>	<p>Foliar: 5 - 7 (0.081- 0.114 lb ai/A)</p> <p>[*]Drip: 7 (0.114 lb ai/A)</p>	<p>Apply preventively or when conditions are favorable for disease development.</p> <p>Repeat as necessary if conditions remain favorable for disease development and to provide extended foliar protection. Best results for both foliar and soilborne pathogens have been obtained when AX FLUTRIAFOL 22.7 is applied preventively, as early as 14 days after transplanting, to provide protection of the roots from soilborne pathogens and to ensure time for movement of flutriafol into the foliage for protection against foliar pathogens. Once fruit are present, foliar applications are needed for fruit protection.</p> <p>For gummy stem blight and target spot, tank mix with mancozeb for resistance management and in fields with known triazole resistance, apply AX FLUTRIAFOL 22.7 with mancozeb or alternative effective fungicide mode of action for disease control.</p>	<p>Foliar – 7</p> <p>[*]Drip - 14</p>
<p>Charcoal Rot[*] (<i>Macrophomina phaseolina</i>)</p> <p>[*Not Registered for Use By California]</p>	7 (0.114 lb ai/A)	<p>Apply through drip irrigation (see Chemigation section). Make initial application at transplanting. Make a second application 14 days later. Additional applications may be made on a 14 day interval, as needed.</p> <p>[In CA: Make a second application 30 days later. Additional applications may be made on a 30-day interval as needed.]</p>	<p>14</p> <p>[In CA: 30]</p>
<p>Suppression of: Fusarium Wilt[*] (<i>Fusarium oxysporum</i>)</p> <p>[*Not Registered for Use By California]</p>	7 (0.114 lb ai/A)	<p>Apply preventively or when conditions are favorable for disease development and repeat application every 14 days to provide extended in-season suppression. Best results for soilborne disease suppression have been obtained when AX FLUTRIAFOL 22.7 is applied preventively, as early as 14 days after transplanting, to provide protection of the roots from soilborne pathogens. AX FLUTRIAFOL 22.7 may be alternated with other non-Group 3 mode of action fungicides to improve Fusarium management.</p>	14

List of Cucurbits: Chayote (fruit); Chinese waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); *Momordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); muskmelon (includes true cantelope, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon); pumpkin; squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon¹ and hybrids and/or varieties of (*Citrullus* spp., *Cucumis melo*, *Cucurbita pepo*[*], *Lagenaria* spp., *Luffa* spp.[*], *Momordica* spp., *Sechium edule*[*]).

Restrictions for all cucurbit crop group 9 uses:

- No single application may exceed 7 fl oz of product (0.114 lb ai)/A.
- The minimum retreatment interval is 7 days.
- **DO NOT** apply more than 4 applications per year.
- **DO NOT** apply more than 28 fl oz of product (0.455 lb ai)/A per year.
- **DO NOT** apply more than 0.455 lb ai of flutriafol/A per year.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Interval (PHI): May be applied the day of harvest (0 day PHI).

Additional Watermelon only Restrictions:

- **DO NOT** apply more than 2 sequential applications, including soil and foliar applications of any Group 3 fungicide, before rotating with a fungicide from a different FRAC group.
- Use nonionic surfactants (NIS) only.
- **DO NOT** use organo silicone adjuvants, or crop oil concentrate (COC), or methylated seed oil (MSO) surfactants.

¹**Precaution for watermelon use:** Under certain environmental conditions, applications of **AX FLUTRIAFOL 22.7** may result in phytotoxicity symptoms that may reduce watermelon yields. Axion Ag Products recommends the grower and/or user test this product to determine its suitability for use in watermelon. Axion Ag Products provides this product to the grower and/or user solely to the extent that the benefit and utility, in the sole opinion of the grower and/or user, outweighs the potential of crop phytotoxicity following applications of **AX FLUTRIAFOL 22.7** to watermelon. The decision to use or not use **AX FLUTRIAFOL 22.7** in watermelon must be made by each individual grower and/or user on the basis of possible crop phytotoxicity, the severity of disease incidence, the cost of alternative disease control options, and other factors.

Use Directions for Hops[*]

[*Not Registered for Use By California]

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Powdery Mildew (<i>Podosphaera macularis</i> , <i>Sphaerotheca macularis</i>)	5- 7 (0.081- 0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development. Apply a minimum of 30 gallons per acre spray volume for ground applications and 10 gallons per acre spray volume for aerial applications. An adjuvant may be used with AX FLUTRIAFOL 22.7 .	14

Restrictions:

- No single application may exceed 7 fl oz product (0.114 lb ai)/A.
- The minimum retreatment interval is 14 days.
- **DO NOT** apply more than 4 applications per year.

- **DO NOT** apply more than 28 fl oz (0.455 lb ai)/A per year.
- **DO NOT** apply more than 0.455 lb ai of flutriafol/A per year.
- An adjuvant may be used with **AX FLUTRIAFOL 22.7**.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Interval (PHI): **DO NOT** apply within 7 days of harvest

Use Directions for Peanut[*]

[*Not Registered for Use By California]

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Early Leaf Spot (<i>Cercospora arachidicola</i>) Late Leaf Spot (<i>Cercosporidium personatum</i>) Peanut Rust (<i>Puccinia arachidis</i>) Leaf Scorch/ Pepper Spot (<i>Leptosphaerulina crassiasca</i>) Web Blotch (<i>Phoma arachidicola</i>)	5 - 7 (0.081- 0.114 lb ai/A) [In CA: 3.5 - 7 (0.057 - 0.114 lb ai/A)]	Apply on a preventive spray schedule for effective disease control in peanuts. AX FLUTRIAFOL 22.7 can be tank mixed with other fungicides for improved disease control, broader spectrum control and resistance management.	14
Suppression of Soil-Borne Diseases: White Mold / Southern Stem Rot (<i>Sclerotium rolfsii</i>)	5 - 7 (0.081- 0.114 lb ai/A) [In CA: 3.5 - 7 (0.057 - 0.114 lb ai/A)]	For soil-borne diseases, rainfall or irrigation will optimize activity of AX FLUTRIAFOL 22.7 . Tank-mix with other labeled fungicides for improved activity on soil-borne diseases.	14

Restrictions:

- No single application may exceed 7 fl oz of product (0.114 lb ai)/A.
- The minimum retreatment interval is 14 days.
- **DO NOT** apply more than 4 applications per year.
- **DO NOT** apply more than 28 fl oz of product (0.455 lb ai)/A per year.
- **DO NOT** apply more than 0.455 lb ai of flutriafol/A per year.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Interval (PHI): **DO NOT** apply within 7 days of harvest.

Use Directions for Pecan and Other Tree Nuts[*]. See below for list of crops.

[*Not Registered for Use By California]

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Pecan Scab (<i>Fusicladium effusum</i>) Powdery Mildew (<i>Microsphaera penicillata</i>) Alternaria Leaf and Fruit Spot (<i>Alternaria alternata</i>) Anthracnose (<i>Colletotrichum acutatum</i>) Downy Spot (<i>Mycosphaerella caryigena</i>) Zonate Leaf Spot (<i>Cristulariella moricola</i>)	5 - 7 (0.081- 0.114 lb ai/A) [In CA: 3.5 - 7 (0.057 - 0.114 lb ai/A)]	For Pecan Scab control, start application at bud break. For all other disease apply preventatively or when conditions remain favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	7

Eastern Filbert Blight (<i>Anisogramma anomalea</i>)	7 [In CA: 3.5 – 7 (0.057 - 0.114 lb ai/A)]	For Hazelnut only apply preventatively or when conditions remain favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	7
List of Tree Nuts: African Tree Nut, Brazil Nut, Burr Oak, Butternut, Cajou, Cashew, Castanha-Do-Maranhao, Coconut, Coquito Nut, Dika nut, Guiana Chestnut, Hazelnut, Heartnut, Hickory Nut, Japanese Horse-Chestnut, Macadamia Nut, Mongongo Nut, Monkeypot, Pachira Nut, Pecan, Sapucaia Nut.			
<p>Restrictions:</p> <ul style="list-style-type: none"> • No single application may exceed 7 fl oz of product (0.114 lb ai)/A. • The minimum retreatment interval is 7 days. • DO NOT apply more than 4 applications per year. • DO NOT apply more than 28 fl oz of product (0.455 lb ai)/A per year. • DO NOT apply more than 0.455 lb ai of flutriafol/A per year. • Restricted Entry Interval (REI) is 12 hours. • Preharvest Interval (PHI): DO NOT apply within 14 days of harvest. 			

Use Directions for Vegetables, Fruiting, Group 8-10. See below for list of crops.

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Powdery Mildew (<i>Leveillula taurica</i>) Anthracnose (<i>Colletotrichum cecedes</i> , <i>C. acutatum</i>) Cercospora Leaf Spot (<i>Cercospora capsici</i>) Black Mold[*] (<i>Alternaria alternata</i>) Early Blight[*] (<i>Alternaria solani</i>) [*Not Registered for Use By California]	7 (0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development. Best results for both foliar and soilborne pathogens have been obtained when AX FLUTRIAFOL 22.7 is applied preventively, shortly after transplanting, to provide protection of the roots from soilborne pathogens and to ensure time for movement of flutriafol into the foliage for protection against foliar pathogens. Once fruit are present, foliar applications are needed for fruit protection.	Foliar – 7 [*]Drip Irrigation 14
Target Spot[*] (<i>Corynespora cassiicola</i>) [*Not Registered for Use By California]	7 (0.114 lb ai/A)	Only apply through drip irrigation. Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development. Best results for both foliar and soilborne pathogens have been obtained when AX FLUTRIAFOL 22.7 is applied preventively, shortly after transplanting, to provide protection of the roots from soilborne pathogens and to ensure time for movement of flutriafol into the foliage for protection against foliar pathogens. Once fruit are present, foliar applications are needed for fruit protection.	14

Charcoal Rot [*] (<i>Macrophomina phaseolina</i>) Fusarium Wilt[*] (<i>Fusarium oxysporum</i>) [*Not Registered for Use By California]	7 (0.114 lb ai/A)	Product is to be applied through drip irrigation (see Chemigation section). Make initial application at transplanting. Make a second application 14 days later. Additional applications may be made on a 14 day interval, as needed. [In CA: Make a second application 30 days later. Additional applications may be made on a 30-day interval; as needed]	14 [In CA: 30]
List of Fruiting Vegetables: African Eggplant, Bell Pepper, Eggplant, Martynia, Non-Bell Pepper, Okra, Pea Eggplant, Pepino, Roselle, Scarlet Eggplant, and cultivars, varieties and/or hybrids of these crops.			
<p>Restrictions:</p> <ul style="list-style-type: none"> • No single application may exceed 7 fl oz of product (0.114 lb ai)/A. • The minimum retreatment interval is 7 days. • DO NOT apply more than 4 applications per year. • DO NOT apply more than 28 fl oz of product (0.455 lb ai)/A per year. • DO NOT apply more than 0.455 lb ai of flutriafol/A per year. • Restricted Entry Interval (REI) is 12 hours. • Preharvest Interval (PHI): May be applied the day of harvest (0 day PHI). 			

Use Directions for Pome Fruit (Apple, Crabapple, Loquat, Mayhaw, Pear, Oriental Pear, and Quince)[*]

[*Not Registered for Use By California]

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Powdery Mildew (<i>Podosphaera leucotricha</i>) Quince Rust (<i>Gymnosporangium clavipes</i>) Cedar Apple Rust (<i>Gymnosporangium juniperi-virginianae</i>) Brooks Fruit Spot (<i>Mycosphaerella pomi</i>) Flyspeck (<i>Zygophiala jamaicicensis</i>) (formerly <i>Schizothyrium pomi</i>)	4 - 6 (0.065 – 0.098 lb ai/A)	Initiate applications at green tip and continue through cover sprays.	10
Apple Scab (<i>Venturia inaequalis</i>) Sooty Blotch (<i>Gloeodes pomigena</i>) Bot rot or White rot (<i>Botryosphaeria dothidea</i>) Black Rot (<i>Botryosphaeria obtusa</i>) Bitter Rot (<i>Colletotrichum gloeosporioides</i> , <i>C. acutatum</i>) Alternaria Blotch (<i>Alternaria mali</i>)	6.5 (0.106 lb ai/A)	<p>Initiate applications at green tip or when environmental conditions are favorable for primary scab development.</p> <p>Continue applications through the duration of primary scab.</p> <p>Mix AX FLUTRIAFOL 22.7 with a protectant such as mancozeb or captan fungicide for scab control.</p> <p>Tank mix applications containing AX FLUTRIAFOL 22.7 with a protectant such as mancozeb or captan fungicide will also effectively control listed summer rots when applied at cover spray timings.</p>	7 - 10

Pear Scab (<i>Venturia pirina</i>)	6.5 (0.106 lb ai/A)	Start application at pink bud and continue through the duration of the primary scab period. Mix AX FLUTRIAFOL 22.7 with a protectant such as mancozeb for improved scab control.	7 - 10
---	------------------------	--	--------

Restrictions:

- No single application may exceed 6.5 fl oz of product (0.106 lb ai)/A.
- The minimum retreatment interval is 7 days.
- **DO NOT** apply more than 4 applications per year.
- **DO NOT** apply more than 26 fl oz of product (0.423 lb ai)/A per year.
- **DO NOT** apply more than 0.423 lb ai of flutriafol/A per year.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Interval (PHI): **DO NOT** apply within 14 days of harvest.

Use Directions for Grain Sorghum [*]

[*Not Registered for Use By California]

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Anthracnose Leaf Blight (<i>Colletotrichum graminicola</i>) Ergot (<i>Claviceps sorghi</i>) or <i>C. africana</i>) Gray Leaf Spot (<i>Cercospora sorghi</i>) Leaf Blight (<i>Exserohilum turcicum</i> or <i>Setosphaeria turcica</i>) Common Rust (<i>Puccinia purpuria</i>) Ladder Leaf Spot (<i>Cercospora fusimaculans</i>) Rough Spot (<i>Ascochyta sorghi</i>) Sooty Stripe (<i>Ramulispora sorghi</i>) Target Leaf Spot (<i>Bipolaris cookei</i>) Zonate Leaf Spot and Sheath Blight (<i>Gloecercospora sorghi</i>)	5 - 7 (0.081- 0.114 lb ai/A)	For Ergot make first application at, or just prior to, flowering. For all other diseases apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development.	7

Restrictions:

- No single application may exceed 7 fl oz of product (0.114 lb ai)/A.
- The minimum retreatment interval is 7 days.
- **DO NOT** apply more than 2 applications per year.
- **DO NOT** apply more than 14 fl oz of product (0.228 lb ai)/A per year.
- **DO NOT** apply more than 0.228 lb ai of flutriafol/A per year.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Interval (PHI): **DO NOT** apply within 30 days of harvest for stover, forage or grain.

Use Directions for Soybean[*] (Foliar Applications)

[*Not Registered for Use By California]

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Cercospora Leaf Blight and Purple Seed Stain (<i>Cercospora kikuchii</i>) Frogeye Leaf Spot (<i>Cercospora sojina</i>) Brown Spot (<i>Septoria glycines</i>) Powdery Mildew (<i>Microsphaera diffusa</i>) Target Spot (<i>Corynespora cassiicola</i>)	5 - 7 (0.081- 0.114 lb ai/A)	For Cercospora Leaf Blight begin applications at first flower (R1). For other diseases apply beginning at first visible flower or when conditions are favorable for disease development. Apply additional applications if environmental conditions are favorable for continued disease development.	14 – 35
Suppression: White Mold / Stem Rot (<i>Sclerotinia sclerotiorum</i>) Southern Blight (<i>Sclerotium rolfsii</i>)			
Rust (<i>Phakopsora pachyrhizi</i>)	5 - 7 (0.081- 0.114 lb ai/A)	Apply beginning at first visible flower or when conditions are favorable for disease development. Apply additional applications if environmental conditions are favorable for continued disease development.	21 – 35
Restrictions:			
<ul style="list-style-type: none"> No single foliar application may exceed 7 fl oz of product (0.114 lb ai)/A. The minimum retreatment interval is 14 days. DO NOT apply more than 3 foliar applications per year. DO NOT apply more than 21 fl oz of product (0.341 lb ai)/A per year For foliar applications: <ul style="list-style-type: none"> Two foliar applications can be made at 7 fl oz of product (0.114 lb ai)/A – OR – One foliar application at 7 fl oz of product (0.114 lb ai)/A followed by two additional foliar applications at 3.5 fl oz of product/A. DO NOT apply more than 30 fl oz of product (0.488 lb ai)/A per year. DO NOT apply more than 0.488 lb ai of flutriafol/A per year (including soil and foliar applications) Restricted Entry Interval (REI) is 12 hours. DO NOT feed forage or hay to animals or permit animals to graze. Preharvest Interval (PHI): DO NOT apply within 21 days of harvest. 			

Use Directions for Sugar Beet[*]

[*Not Registered for Use By California]

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Cercospora Leaf Spot (<i>Cercospora beticola</i>)	5 - 7 (0.081- 0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	14

		Tank mix with mancozeb for resistance management.	
Powdery Mildew (<i>Erysiphe beta</i> ae)	5 - 7 (0.081- 0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	14
Restrictions:			
<ul style="list-style-type: none"> • No single application may exceed 7 fl oz of product (0.114 lb ai)/A. • The minimum retreatment interval is 14 days. • DO NOT apply more than 2 applications per year. • DO NOT apply more than 14 fl oz of product (0.228 lb ai)/A per year. • DO NOT apply more than 0.228 lb ai of flutriafol/A per year. • Restricted Entry Interval (REI) is 12 hours. • Preharvest Interval (PHI): DO NOT apply within 21 days of harvest. 			

Use Directions for Barley, Wheat (Spring and Winter) and Triticale [*]

[*Not Registered for Use By California]

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Rust (<i>Puccinia</i> spp.) Powdery Mildew (<i>Blumeria graminis</i>) Leaf and Glume Blotch (<i>Septoria</i> spp.) Tan Spot (<i>P. Tritici-repentis</i>)	5 - 7 (0.081- 0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development. For aerial application, use a minimum of 2 gallons of spray solution per acre.	7
Suppression: Fusarium Head Blight (<i>Fusarium</i> spp.)	5 - 7 (0.081- 0.114 lb ai/A)	Apply at anthesis (Feekes Growth Stage 10.51). Properly set up spray equipment nozzles to ensure good coverage of the head. For aerial application, use a minimum of 2 gallons of spray solution per acre.	7

Restrictions:

- No single application may exceed 7 fl oz of product (0.114 lb ai)/A.
- The minimum retreatment interval is 7 days.
- **DO NOT** apply more than 2 applications per year.
- **DO NOT** apply more than 14 fl oz of product (0.228 lb ai)/A per year.
- **DO NOT** apply more than 0.228 lb ai of flutriafol/A per year.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Interval (PHI):
 - **DO NOT** apply within 30 days of harvest when grown for grain,
 - **DO NOT** apply within 15 days of harvest when grown for hay,
 - **DO NOT** apply within 3 days when grown for forage.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[For plastic containers ≤ 5 gallons: Nonrefillable Container: DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

[For plastic containers > 5 gallons: Nonrefillable container: DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

Note to PM, IBC's may be nonrefillable or refillable, the container handling section will state either Nonrefillable or Refillable.}

Nonrefillable container. {or}

Refillable container.

Bottom discharge IBC.

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of AXION AG PRODUCTS LLC or Seller, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold AXION AG PRODUCTS LLC and Seller harmless for any claims relating to such factors.

AXION AG PRODUCTS LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or AXION AG PRODUCTS LLC, and TO THE EXTENT CONSISTENT WITH APPLICABLE LAW Buyer and User assume the risk of any such use. AXION AG PRODUCTS LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither AXION AG PRODUCTS LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF AXION AG PRODUCTS LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE

OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF AXION AG PRODUCTS, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

AXION AG PRODUCTS LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of AXION AG PRODUCTS LLC.