

89168-3

12/6/2012

1/26



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D C 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Scott Baker  
Lighthouse Product Services  
3937 Cedarwood Lane  
Johnstown, CO 80534

12-6-12

Subject EPA Reg 89168-3 / Liberty Dicamba 4 DMA Notification

Dear Mr Baker

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 10-18-12 for the product EPA Reg 89168-3 / Liberty Dicamba 4 DMA. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records. The revised basic and alternate csf are acceptable.

If you have any questions please call Erik Kraft at 703-308-9358 or email at [Kraft.Erik@epa.gov](mailto:Kraft.Erik@epa.gov)

Sincerely,

A handwritten signature in black ink, appearing to read "K. Montague".

Kathryn Montague  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)

2/26



Please read instructions on reverse before completing form

Form Approved OMB No. 2070-0060, Approval expires 2 28 95

<p>United States <b>Environmental Protection Agency</b> Washington DC 20460</p>	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
---	--	-----------------------

### Application for Pesticide - Section I

<b>1 Company/Product Number</b> 89168 3	<b>2 EPA Product Manager</b> Montague	<b>3 Proposed Classification</b> <input type="checkbox"/> None <input checked="" type="checkbox"/> Restricted
<b>4 Company/Product (Name)</b> Liberty DICAMBA 4 DMA	<b>PM#</b> 23	
<b>5 Name and Address of Applicant (Include ZIP Code)</b> LIBERTY CROP PROTECTION, LLC 3937 CEDARWOOD LANE JOHNSTOWN, CO 80534 <input checked="" type="checkbox"/> Check if this is a new address		<b>6 Expedited Review</b> In accordance with FIFRA Section 3(c)(3) (b)(i) my product is similar or identical in composition and labeling to EPA Reg No _____ Product Name _____

### Section - II

<input type="checkbox"/> Amendment Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> Me Too Application
<input checked="" type="checkbox"/> Notification Explain below	<input type="checkbox"/> Other Explain below

**Explanation** Use additional page(s) if necessary (For section I and Section II )

Notification of new label and CSF following transfer from the previous registrant Change to Primary Brand Name under PR Notice 98-10

### Section - III

<b>1 Material This Product Will Be Packaged In</b>			
<b>Child Resistant Packaging</b> <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<b>Unit Packaging</b> <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<b>Water Soluble Packaging</b> <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<b>2 Type of Container</b> <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify _____)
<b>* Certification must be submitted</b> If "Yes" Unit Packaging wgt    No per container		If "Yes" Package wgt    No per container	
<b>3 Location of Net Contents Information</b> <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		<b>4 Size(s) Retail Container</b> 1 2 5 15 30 55 110 275 Gal Bulk	
<b>6 Manner in Which Label is Affixed to Product</b> <input type="checkbox"/> Lithograph Paper glued Stenciled		<input checked="" type="checkbox"/> Other self adhesive	
<b>5 Location of Label Directions</b> <input checked="" type="checkbox"/> Attached to container			

### Section - IV

<b>1 Contact Point</b> (Complete items directly below for identification of individual to be contacted if necessary to process this application)		
<b>Name</b> Scott Baker	<b>Title</b> Regulatory Agent	<b>Telephone No (Include Area Code)</b> 970 313 6654
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true accurate and complete I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law		<b>6 Date Application Received</b>  <b>(Stamped)</b>
<b>2 Signature</b> 	<b>3 Title</b> Regulatory Agent	
<b>4 Typed Name</b> Scott Baker	<b>5 Date</b> October 18, 2012	

3/26



**LIGHTHOUSE PRODUCT SERVICES**  
Regulatory Safety & Supply Chain Consulting

October 18, 2012

Document Processing Desk (REGIS)  
Office of Pesticide Programs  
U S Environmental Protection Agency  
One Potomac Yard Room S-4900  
2777 S Crystal Drive  
Arlington VA 22202

Subject LIBERTY DICAMBA 4 DMA EPA Reg No 89168-3

Lighthouse Product Services LLC Regulatory Agent for Liberty Crop Protection LLC is submitting the enclosed documentation to update the EPA files with new labeling and Confidential Statements of Formula for the subject product This product was recently transferred to Liberty Please find the following enclosed

- 1 Application Form 8570-1
- 2 1 copy of updated label
- 3 ~~1~~ copies of updated Basic CSF
- 4 ~~2~~ copies of updated Alternate 1 CSF

In addition to the product name, company identification EPA Registration number the date code and the Conditions of Sale and Limitation of Warranty and Liability being updated and some minor formatting

This notification is consistent with the provisions of PR Notice 98 10 and EPA regulations at 40 CFR 152.46 and no other changes have been made to the labeling or the confidential statement of formula of this product I understand that it is a violation of 18 U S C Sec 1001 to willfully make any false statement to EPA I further understand that if this notification is not consistent with the terms of PR Notice 98 10 and 40 CFR, 152 46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA

I also changed "recommended" rates to specified or "Listed as appropriate"

Please contact me at 970-313-6654 or by e-mail [sbaker@lhpsonline.com](mailto:sbaker@lhpsonline.com) if there are any questions or comments concerning this submission

Sincerely

Scott Baker  
Regulatory Agent for Liberty Crop Protection LLC

' SHINING THE LIGHT ON YOUR REGULATORY NEEDS '

( 11 Modification  
Acceptable "

4/26

# LIBERTY DICAMBA 4 DMA

HERBICIDE FOR WEED CONTROL IN CORN, COTTON SORGHUM, SOYBEAN SMALL GRAINS, PASTURE, HAY RANGELAND FARMSTEAD (NON CROPLAND), FALLOW SUGARCANE, ASPARAGUS, TURF AND GRASS

**ACTIVE INGREDIENT**

Dimethylamine salt of dicamba (3,6-dichloro-O-anisic acid)\* 50.2%

**OTHER INGREDIENTS**

49.8%

**TOTAL**

100.0%

\*This product contains 40.0% 3,6-dichloro-o-anisic acid (dicamba) or 4 pounds per gallon (480 g/L)

## KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle (If you do not understand the label find someone to explain it to you in detail)

FIRST AID	
<b>IF IN EYES</b>	Hold eye open and rinse slowly and gently with water for 15-20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eye Call poison control center or doctor for treatment advice
<b>IF SWALLOWED</b>	Call a poison control center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow Do not induce vomiting unless told to do so by a poison control center or doctor Do not give anything by mouth to an unconscious person
<b>IF ON SKIN OR CLOTHING</b>	Take off contaminated clothing Rinse skin immediately with plenty of water for 15-20 minutes Call a poison control center or doctor for treatment advice
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment You may also contact 1-800-222-1222	

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
WARNING / AVISO**

Causes substantial but temporary eye injury Do not get in eyes on skin or on clothing Harmful if swallowed Avoid breathing spray mist

EPA Reg No 89168-3

EPA Est No XXXXX XX XXX

101012

Net Contents \_\_\_\_\_

Manufactured For  
LIBERTY CROP PROTECTION LLC  
3937 CEDARWOOD LANE  
JOHNSTOWN CO 80534

**PERSONAL PROTECTIVE EQUIPMENT (PPE)** Some materials that are chemical resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.

All mixers, loaders, applicators, and other handlers must wear:

- Long sleeved shirt and long pants
- Shoes plus socks
- Goggles or faceshield and
- Chemical resistant gloves

See engineering controls for additional requirements.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENGINEERING CONTROLS STATEMENTS**

Pilots must use an enclosed cockpit in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)].

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Apply this product only as directed on label.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls worn over short sleeve shirt and short pants
- Chemical resistant footwear plus socks
- Chemical resistant gloves made of any waterproof material
- Chemical resistant headgear for overhead exposure
- Protective eyewear

**NON AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to the uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, in nurseries, in forests, or in greenhouses.

Do not enter or allow others to enter the treated areas until the spray has dried.

6/26

Before applying LIBERTY DICAMBA 4 DMA read all directions and precautions appearing on the container label and in this booklet. Failure to follow all directions and precautions may result in unsatisfactory weed control, crop injury or illegal residues.

### PRODUCT INFORMATION

The following directions apply to all uses of LIBERTY DICAMBA 4 DMA. Additional precautions and restrictions will be found in each specific use section.

Do not treat irrigation ditches or water used for crop irrigation or domestic uses.

Do not apply this product through any type of irrigation system.

Do not exceed the maximum single application rate of 2 pints (1.0 lb a.i.) LIBERTY DICAMBA 4 DMA per application with no more than 2 applications per year.

### MIXING AND APPLICATION

UNLESS OTHERWISE SPECIFIED UNDER THE INDIVIDUAL USE HEADINGS OF THIS BOOKLET, THE FOLLOWING DIRECTIONS APPLY TO ALL CROP AND NON CROP USES OF LIBERTY DICAMBA 4 DMA. REFER TO INDIVIDUAL USE SECTIONS FOR ADDITIONAL PRECAUTIONS, RESTRICTIONS, APPLICATION RATES AND TIMINGS.

LIBERTY DICAMBA 4 DMA is a water soluble formulation that can be applied using water or sprayable fluid fertilizer as the carrier. If a fluid fertilizer is to be used, a compatibility test (See COMPATIBILITY TEST) should be made prior to tank mixing.

Ground or aerial application equipment which will give good spray coverage of weed foliage should be used. HOWEVER, DO NOT USE AERIAL APPLICATION EQUIPMENT IF SPRAY PARTICLES CAN BE CARRIED BY WIND INTO AREAS WHERE SENSITIVE CROPS OR PLANTS ARE GROWING OR WHEN TEMPERATURE INVERSIONS EXIST.

Apply 3 to 50 gallons of diluted spray per treated acre when using ground application equipment or 1 to 10 gallons of diluted spray per treated acre (2 to 20 gallons of diluted spray per acre for preharvest uses) in a water based carrier when using aerial application equipment. Use the higher level of the listed spray volumes when treating dense or tall vegetation. Use coarse sprays.

Select nozzles designed to produce minimal amounts of fine spray particles. Spray with nozzles as close to the weeds as is practical for good weed coverage.

To avoid uneven spray coverage, LIBERTY DICAMBA 4 DMA should not be applied during periods of gusty wind or when wind is in excess of 15 mph.

Avoid disturbing (e.g. cultivating or mowing) treated areas for at least 7 days following application.

### BEST STEWARDSHIP PRACTICES

LIBERTY DICAMBA 4 DMA provides effective broadleaf weed and brush control when properly applied. Best stewardship practices in all mixing, loading, and application operations not only maximize weed control, but also protect ground and surface waters and minimize off target movement.

This chemical is known to leach through the soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

#### GROUND AND SURFACE WATERS PROTECTION

1) Point source contamination. To prevent point source contamination, do not mix, load, or apply this pesticide product within 50 feet of wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. Do not apply pesticide product within 50 feet of wells. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas as described below.

Mixing, loading, rinsing, or washing operations performed within 50 feet of a well are allowed only when conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be on or move across the pad. The pad must be self contained to prevent surface water flow over or from the pad. The pad capacity must be maintained at 110% that of the largest pesticide container or application equipment used on the pad and have sufficient capacity to contain all product spills, equipment or container leaks, equipment wash waters, and rainwater that may fall on the pad. The containment capacity does not apply to vehicles delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

Care must be taken when using this product to prevent a) back siphoning into wells, b) spills, or c) improper disposal of excess pesticide, spray mixtures, or rinsates. Check valves or anti siphoning devices must be used on all mixing equipment.

2) Movement by surface runoff or through soil. Do not apply under conditions which favor runoff. Do not apply to impervious substrates such as paved or highly compacted surfaces in areas with high potential for ground water contamination. Ground water contamination may occur in areas where soils are permeable or coarse and ground water is near the surface. Do not apply to soils classified as sand with less than 3% organic matter and where ground water depth is shallow (less than 8 feet in Arizona). To minimize the possibility of ground water contamination, carefully follow application rate recommendations as affected by soil type in the **Product Information** section of this label.

3) Movement by water erosion of treated soil. Do not apply or incorporate this product through any type of irrigation equipment nor by flood or furrow irrigation. Ensure treated areas have received at least one half inch rainfall (or irrigation) before using tailwater for subsequent irrigation of other fields.

**SENSITIVE CROP PRECAUTIONS**

LIBERTY DICAMBA 4 DMA may cause injury to desirable trees and plants particularly beans cotton flowers fruit trees grapes ornamentals peas potatoes soybeans sunflowers tobacco tomatoes and other broadleaf plants when contacting their roots stems or foliage These plants are most sensitive to LIBERTY DICAMBA 4 DMA during their development or growing stage FOLLOW THE PRECAUTIONS LISTED BELOW WHEN USING LIBERTY DICAMBA 4 DMA

- Do not treat areas where either possible downward movement into the soil or surface washing may cause contact of LIBERTY DICAMBA 4 DMA with the roots of desirable plants such as trees and shrubs
- Avoid making applications when air currents may carry spray particles to areas where sensitive crops and plants are growing or when temperature inversions exist Do not spray near sensitive plants if wind is gusty or in excess of 5 mph and moving in the direction of adjacent sensitive crops Leave an adequate buffer zone between area to be treated and sensitive plants Coarse sprays are less likely to drift out of the target area than fine sprays
- Use coarse sprays to avoid potential herbicide drift Select nozzles which are designed to produce minimal amounts of fine spray particles Examples of nozzles designed to produce coarse sprays via ground application are Delavan Rain drops Spraying Systems XR flat fans or large capacity flood nozzles such as D10 TK10 or greater capacity tips Keep the spray pressure at or below 20 psi and the spray volume at or above 20 GPA unless otherwise required by the manufacturer of drift reducing nozzles Consult your spray nozzle supplier concerning the choice of drift reducing nozzles
- Agriculturally approved drift reducing additives may be used
- Do not apply LIBERTY DICAMBA 4 DMA adjacent to sensitive crops when the temperature on the day of application is expected to exceed 85 F as drift is more likely to occur
- To avoid injury to desirable plants equipment used to apply LIBERTY DICAMBA 4 DMA should be thoroughly cleaned (See PROCEDURE FOR CLEANING SPRAY EQUIPMENT) before reusing to apply any other chemicals

All crop uses of LIBERTY DICAMBA 4 DMA are intended for a normal growing interval between planting and harvest No crop rotation restrictions exist if normal harvest of treated crop has occurred If this interval is shortened such as in cover crops that will be plowed under do not follow up with the planting of a sensitive crop

Crops growing under stress conditions such as drought poor fertility or foliar damage due to hail wind or insects can exhibit various injury symptoms that may be more pronounced if herbicides are applied

Consult your local or state authorities for possible application restrictions and advice concerning these and other special local use situations Tank mix recommendations are for use only in states where the tank mix product and application site are registered

**BAND TREATMENTS**

LIBERTY DICAMBA 4 DMA may be applied as a band treatment Use the formulas below to determine the appropriate rate and volume per treated acre

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast RATE} = \text{Band RATE}$$

per treated acre                      per treated acre

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast VOLUME} = \text{Band VOLUME}$$

per treated acre                      per treated acre

**COMPATIBILITY TEST**

Before mixing in the spray tank it is advisable to test compatibility by mixing all components in a small container in proportionate quantities (see following table)

Amount of Herbicide to Add to One Pint of Spray Carrier (Assuming Volume is 25 Gallons per Acre)

HERBICIDE FORMULATIONS	RATE PER ACRE	LEVEL TEASPOONS
Dry	1lb	1 1/2
Liquid	1 pt	1/2

If herbicide(s) do not ball up or form flakes sludge gels oily films or layers or other precipitates then the tested spray mix is compatible Usually incompatibility in any of the above described forms will occur with 5 minutes after mixing If components are incompatible the use of a compatibility agent is recommended Rerun the above COMPATIBILITY TEST with a suitable compatibility agent (1/4 teaspoon is equivalent to 2 pints per 100 gallons of fluid fertilizer)

**PROCEDURE FOR CLEANING SPRAY EQUIPMENT**

The steps listed below are suggested for thorough cleaning of spray equipment following applications of LIBERTY DICAMBA 4 DMA or tank mixes of LIBERTY DICAMBA 4 DMA or tank mixes of LIBERTY DICAMBA 4 DMA plus 2 4 D amine

- 1) Hose down thoroughly the inside as well as outside surfaces of equipment while filling the spray tank half full of water Flush by operating sprayer until the system is purged of the rinse water
- 2) Fill tank with water while adding 1 quart of household ammonia for every 25 gallons of water Operate the pump to circulate the ammonia solution through the sprayer system for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles Let the solution stand for several hours preferably overnight
- 3) Flush the solution out of the spray tank through the boom
- 4) Remove the nozzles and screens and flush the system with two full tanks of water

The steps listed below are suggested for thorough cleaning of spray equipment used to apply LIBERTY DICAMBA 4 DMA as a tank mix with wettable powders (WP) emulsifiable concentrates (EC) or other types of water dispersible formulations

LIBERTY DICAMBA 4 DMA tank mixes with water dispersible formulations require the use of a water/detergent rinse

5) Complete step 1

6) Fill tank with water while adding 2 lbs of detergent for every 40 gallons of water Operate the pump to circulate the detergent solution through the sprayer system for 5 to 10 minutes and discharge a small amount of the solution through the boom and nozzles Let the solution stand for several hours preferably overnight

7) Flush the detergent solution out of the spray tank through the boom

8) Repeat step 1 and follow with steps 2 3 and 4

**WEED LIST**

This is a list of weeds which may be treated with LIBERTY DICAMBA 4 DMA in accordance with this label as recommended under the rates and timing sections of the Individual Use headings Proper usage of this product will give control or growth suppression of many ANNUAL BIENNIAL and PERENNIAL broadleaf weeds and many WOODY brush and vine species including

<b>ANNUAL</b>			
Amaranth Spiny (Spiny Pigweed)	Evening Primrose Cutleaf	Pennycress Field (Fanweed)	Sicklepod
Aster Slender	Fleabane Annual	Frenchweed Stinkweed)	Sida Prickly (Teaweed)
Bedstraw	Goosefoot Nettleleaf	Pepperweed Virginia (Peppergrass)	Smartweed Green
Beggarweed Florida	Henbit	Pigweed Prostrate	Smartweed Pennsylvania
Broomweed Common	Jimsonweed	Pigweed Redroot (Carelessweed)	Sneezeweed Bitter
Buckwheat Wild	Knotweed	Pigweed Rough	Sowthistle Annual
Buffalobur	Kochia	Pigweed Smooth	Sowthistle Spiny
Burclover California	Ladythumb	Pigweed (triazine resistant)	Spikeweed Common
Burcucumber	Lambsquarters Common	Pigweed Tumble	Spurge Prostrate
Buttercup Roughseed	Lambsquarters (triazine resistant)	Poorjoe	Spurry Corn
Carpetweed	Lettuce Prickly	Puncturevine	Starbur Bristly
Catchfly Nightflowering	Mallow Common	Purslane Common	Sumpweed Rough
Chamomile Corn	Mallow Venice	Pusley Florida	Sunflower Common (Wild)
Chickweed Common	Mare s Tail (Horseweed)	Radish Wild	Sunflower Volunteer
Clovers (Annual)	Mayweed	Ragweed Common	Thistle Russian
Cockle Corn	Morning glory Ivyleaf	Ragweed Giant (Buffaloweed)	Velvetleaf
Cockle Cow	Morning glory Tall	Ragweed Lance Leaf	Waterhemp
Cocklebur Common	Mustard Tansy	Rubberweed bitter (Bitterweed)	Waterprimrose Winged
Croton Tropic	Mustard Wild	Sesbania Hemp	Wormwood Annual
Croton Woolly	Mustard (Yellowtops)	Shepherdspurse	
Daisy English	Nightshade Black		

<b>BIENNIALS</b>			
Burdock Common	Geranium Carolina	Plantain Bracted	Thistle Bull
Carrot Wild (Queen Anne s Lace)	Gromwell	Ragwort Tansy	Thistle Milk
Cockle White	Knapweed Diffuse	Starthistle Yellow	Thistle Musk
Evening Primrose Common	Knapweed Spotted	Sweetclover	Thistle Plumelless
	Mallow Dwarf	Teasel	

<b>PERENNIALS</b>			
Alfalfa	*Dock Broadleaf (Bitterdock)	Milkweed Western Whorled	Sundrop Halfshrub (Evening Primrose)
Artichoke Jerusalem	*Dock Curly	Nettle Stinging	Thistle Canada
Aster Spiny	Dogbane Hemp	Nightshade Silverleaf (White Horsenettle)	Toadflex Dalmation
Aster Whiteheath	*Dogfennel (Cypressweed)	Onion Wild	Tropical Soda Apple
Beadstraw Smooth	Fern Bracken	Plantain Broadleaf	Trumpetcreeper (Buckvine)
Bindweed Field	Garlic Wild	Plantain Buckhorn	Vetch
Bindweed Hedge	Goldenrod Canada	Pokeweed	Waterhemlock
Blueweed Texas	Goldenrod Missoun	Ragweed Western	Waterprimrose Creeping
*Bursage (Bur Ragweed Lakeweed Povertyweed)	Goldenweed Common	Redvine	*Woodsorrel Creeping
Buttercup Tall	Hawkweed	Sencia Lespedeza	Common Yellow
Campion Bladder	Henbane Black	Smartweed Swamp	Wormwood Common
Chickweed Field	Horsenettle Carolina	Snakeweed Broom	Wormwood Louisiana
Chickweed (Mouseear Canada)	Ironweed	*Sorrel Red (Sheep Sorrel)	*Yankeeweed
Chicory	Knapweed Black	Sowthistle	Yarrow Common
Clover Hop	Knapweed Russian	Sowthistle Perennial	
*Dandelion Common	Knapweed Climbing	Spurge Leafy	
	Milkweed Common		
	Milkweed Honeyvine		

\*Noted perennials may be controlled using LIBERTY DICAMBA 4 DMA at rates lower than those recommended for other listed perennial weeds (See application rates and timing sections in this label )



WOODY			
Alder	*Dewberry	Locust Black	Sagebrush Fringed
Ash	*Dogwood	Maple	Sassafras
Aspen	Elm	Mesquite	Serviceberry
Basswood	Grape	Oak	Spicebush
Beech	*Hawthorn (Thornapple)	Oak Poison	Spruce
Birch	Hemlock	Olive Russian	Sumac
Blackberry	Hickory	Persimmon Eastern	*Sweetgum
*Blackgum	Honeylocust	Pine	Sycamore
*Cedar	Honeysuckle	*Plum Sand (Wild Plum)	Tarbrush
Cherry	Hornbeam	Poplar	Willow
Chinquapin	Huckleberry	Rabbitbrush	Witchhazel
Cottonwood	Huisache	*Redcedar Eastern	*Yaupon
Creosotebush	Ivy Poison	*Rose McCartney	*Yucca
Cucumbertree	Kudzu	*Rose Multiflora	

Growth suppression

**FIELD SEED\* POPCORN\* AND SILAGE CORN**

Observe all precautions mixing and application instructions as well as the following

Do not apply LIBERTY DICAMBA 4 DMA to seed corn or popcorn without first verifying with your local seed corn company (supplier) the Dicamba selectivity on your inbred line or variety of popcorn. This precaution will help avoid potential injury of sensitive varieties.

LIBERTY DICAMBA 4 DMA is not registered for use on sweet corn.

Direct contact of LIBERTY DICAMBA 4 DMA with corn seed must be avoided. If corn seeds are less than 1 1/2 inches below the surface, delay application until corn has emerged.

Up to 2 applications of LIBERTY DICAMBA 4 DMA may be made during a growing season. Do not exceed a total of 1 1/2 pints of LIBERTY DICAMBA 4 DMA per treated acre per crop year. Allow two weeks or more between applications of LIBERTY DICAMBA 4 DMA. See appropriate section for rate information. For combination options or sequential treatments, refer to appropriate section.

Applications of LIBERTY DICAMBA 4 DMA to corn during periods of rapid growth may result in temporary leaning. Corn will usually become erect within 3 to 7 days. Cultivation should be delayed until after corn is growing normally to avoid breakage.

Agriculturally approved surfactants or sprayable fertilizers (1/2 to 1 gallon per acre of 28% 30% or 32% urea ammonium nitrate or 2.5 pounds per acre spray grade ammonium sulfate) may be added to the spray mixture to improve postemergence weed control, particularly in dry growing conditions.

Do not use adjuvants containing penetrants such as petroleum based oils after crop emergence or crop injury may result. Corn may be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity.

Several synthetic pyrethroid insecticides are labeled for tank mix applications of dicamba. Refer to their label for specific recommendations.

**WEEDS CONTROLLED**

LIBERTY DICAMBA 4 DMA will control many ANNUAL broadleaf weeds or give growth suppression of many PERENNIAL broadleaf weeds commonly found in corn. (Refer to the WEED LIST)

For best performance, make application when weeds have emerged and are actively growing.

Preemergence control of cocklebur, velvetleaf, and jimsonweed may be reduced if conditions such as low temperature or lack of soil moisture cause delayed or deep germination of weeds.

**PREPLANT/PREEMERGENCE IN NO TILLAGE CORN**

Applications of LIBERTY DICAMBA 4 DMA may be made before, during, or after planting to emerged and actively growing broadleaf weeds. Apply LIBERTY DICAMBA 4 DMA at 1 pint per treated acre on medium or fine textured soils containing 2% or greater organic matter. Use 1/2 pint per treated acre on coarse textured soils (sand, sandy loam, and loamy sand) or medium and fine textured soils with less than 2% organic matter.

When planting into a legume sod (e.g., alfalfa or clover), apply LIBERTY DICAMBA 4 DMA after 4 to 6 inches of regrowth has occurred.

**PREEMERGENCE IN CONVENTIONAL OR REDUCED TILLAGE CORN**

LIBERTY DICAMBA 4 DMA may be applied after planting and prior to corn emergence. Application at 1 pint per treated acre may be made to medium or fine textured soils which contain 2% or greater organic matter. DO NOT apply to coarse textured soils (sand, sandy loam, and loamy sand) until after crop emergence (see Early Postemergence uses below).

Preemergence application of LIBERTY DICAMBA 4 DMA does not require mechanical incorporation to become active. A shallow mechanical incorporation is recommended if application is not followed by adequate rainfall or sprinkler irrigation. Avoid tillage equipment (e.g., drags, harrows) which concentrates treated soil over seed furrow.

**EARLY POSTEMERGENCE (ALL TILLAGE SYSTEMS)**

**(Spike through 8 inch tall corn)**

LIBERTY DICAMBA 4 DMA at 1 pint per treated acre may be applied during the period from corn emergence through the five leaf stage or 8 inches tall, whichever comes first. Reduce the rate to 1/2 pint per treated acre if corn is growing on coarse textured soils (sand, sandy loam, and loamy sand). See LATE POSTEMERGENCE APPLICATIONS given below if the 6th true leaf is emerging from whorl or corn is greater than 8 inches tall.

**LATE POSTEMERGENCE (ALL TILLAGE SYSTEMS)  
(8 to 36 inch tall corn)**

Application of LIBERTY DICAMBA 4 DMA at 1/2 pint per treated acre may be made from 8 to 36 inch tall corn or 15 days before tassel emergence whichever comes first For best performance make applications when weeds are less than 3 inches tall

Make directed spray application when (1) corn leaves prevent proper spray coverage (2) sensitive crops are growing nearby (3) tank mixing with 2 4 D  
DO NOT apply LIBERTY DICAMBA 4 DMA when soybeans are growing nearby if any of these conditions exist

- corn is more than 24 inches tall
- soybeans are more than 10 inches tall
- soybeans have begun to bloom

**OVERLAY (SEQUENTIAL)TREATMENTS**

LIBERTY DICAMBA 4 DMA may be applied to ground previously treated with one or more of the following herbicides registered for use in corn

acetochlor alachlor (Lasso® Lasso MT® Partner®) atrazine Broadstrike® butylate (Sutan®) dimethenamid (Frontier®) EPTC	glyphosate halosulfuron (Battalion® Permit®) Lariat® metolachlor paraquat pendimethalin propachlor (Ramrod®) simazine
---	---

Apply LIBERTY DICAMBA 4 DMA at 1/2 pint per treated acre to ground previously treated with full rates of Clarity or Marksman herbicides Allow at least 2 weeks between applications

READ AND FOLLOW LABEL DIRECTIONS FOR EACH OF THE ABOVE PRODUCTS

**TANK MIX TREATMENTS FOR CORN**

LIBERTY DICAMBA 4 DMA may be tank mixed with one or more of the following herbicides for control of grasses or additional broadleaf weeds Read and follow the label of each tank mix product used for precautionary statements directions for use rates and timings, and other restrictions

RATES AND TIMINGS					
LIBERTY DICAMBA 4 DMA Plus	Preplant/ Preemergent (No Tillage Corn)	Pre-emergent (Conventional or Reduced Tillage Corn)	Early Post Emergent (All Tillage Systems)	Late Post Emergent (All Tillage Systems)	Additional Directions
Accent® (nicosulfuron)			1/2 1 oz a i /A	1/2 1 oz a i /A (To improve spray coverage of weeds and reduce risk of corn injury use drop pipes to direct spray beneath corn leaves when corn is greater than 8 inches tall)	Application may be made to emerged weeds before corn is greater than 24 inches tall Use non ionic surfactant at 25% (v/v) with this tank mixture
Atrazine	1 1/4 2 lbs a i /A	1 1/4 2 lbs a i /A	1 1/4 2 lbs a i /A Crop oil concentrates may be used with this mixture if corn is 5 inches or less in height	1 1/4 2 lbs a i /A Do not apply if corn is greater than 12 inches tall	Application may be made before grasses are 1 1/2 tall Follow all state and Federal restrictions pertaining to atrazine applications
Beacon® (primisulfuron)			0 31 0 62 oz a i /A	0 31 0 62 oz a i /A (To improve spray coverage of weeds and reduce risk of corn injury use drop pipes to direct spray beneath corn leaves when corn is greater than 8 inches tall)	Application may be made to emerged weeds when corn is 4 to 24 inches tall Use non ionic surfactant at 25% (v/v) with this tank mixture
Metolachlor	1 1/2 3 lbs a i /A	1 1/2 3 lbs a i /A (Use only on fine or medium textured soils with 2 1/2% or greater organic matter )	1 1/2 3 lbs a i /A		Application may be made before grasses reach the 2 leaf stage and before corn is greater than 3 inches tall

Frontier® (dimethenamid)	13.25 fl oz/A	13.25 fl oz/A (Use only on fine or medium textured soils with 2.5% or greater organic matter)	13.25 fl oz/A		Application may be made up to 8 inch tall corn. This treatment must be combined with a herbicide that provides post emergence control of grass weeds if they are greater than 1 inch tall at the time of application.
Frontier® 6.0 (dimethenamid)	16.32 fl oz/A	16.32 fl oz/A (Use only on fine or medium textured soils with 2.5% or greater organic matter)			Application may be made up to 8 inch tall corn. This treatment must be combined with a herbicide that provides post emergence control of grass weeds if they are greater than 1 inch tall at the time of application.
Paraquat	1/4 1 lb a.i./A	1/4 1 lb a.i./A			Application may be made to emerged weeds but prior to corn emergence.
Acetochlor	1 1/2 3 lbs a.i./A	1 1/2 3 lbs a.i./A (Use only on fine textured soils with greater than 2.5% organic matter)			Application should be made prior to corn emergence.
Lasso® (alachlor)	1 1/2 4 lbs a.i./A	1 1/2 4 lbs a.i./A (Use only on fine textured soils with greater than 2.5% organic matter)	1 1/2 4 lbs a.i./A		Application may be made before grasses reach the 2 leaf stage and before corn is greater than 3 inches tall. If microencapsulated forms of alachlor are used (Lasso MT Partner) applications must be made prior to grass emergence.
Simazine	2.0 3.0 lbs a.i./A	2.0 3.0 lbs a.i./A			Application may be made prior to corn or weed emergence.
Pendimethalin		3/4 1 1/2 lbs a.i./A (Use only on fine or medium textured soils with 2 1/2% or greater organic matter)	3/4 1 1/2 lbs a.i./A		Application may be made immediately after planting but prior to weed emergence. Corn should not be beyond the 2 leaf stage of growth.
Glyphosate	1.0 3.0 lbs a.i./A	1.0 3.0 lbs a.i./A			Application may be made to emerged weeds but prior to corn emergence.
Clopyralid			0.035 0.07 lb a.i./A	0.035 0.07 lb a.i./A	Application may be made any time after corn emergence through 24 inch tall corn. Use drop nozzles to direct spray after corn exceeds the 8 inch stage. Apply when the majority of the thistle plants have emerged and are at least 4 inches in height but before bud stage. Use higher rates listed for stand reduction or larger thistle plants or heavier infestations. Lower rates listed may provide seasonal thistle suppression only.

12/26

Tough® (pyridate)			0.47 lb a.i./A	0.47 lb a.i./A	Application may be made to emerged actively growing weeds. Directed applications are recommended when corn is large enough to prevent proper spray coverage.
2.4 D	1/4 1/2 lb a.i./A	1/4 1/2 lb a.i./A	Not recommended	1/8 lb a.i./A	Drop pipes are to be used when corn height is 8 inches or greater. Keeping the spray off the corn leaves and out of the whorl will reduce the likelihood of crop injury and improve spray coverage of weed foliage.

**COTTON  
EXCEPT CALIFORNIA**

**PREPLANT APPLICATION** Apply up to 8 fluid ounces of LIBERTY DICAMBA 4 DMA per acre to control emerged broadleaf weeds prior to planting cotton in conventional or conservation tillage systems. For best performance, apply LIBERTY DICAMBA 4 DMA when weeds are in the 2-4 leaf stage and rosettes are less than 2" across. Following application of LIBERTY DICAMBA 4 DMA and a minimum accumulation of 1" of rainfall or overhead irrigation, a waiting interval of 21 days is required per 8 fluid ounces per acre or less. These intervals must be observed prior to planting cotton. Do not apply preplant to cotton west of the Rockies. Do not make LIBERTY DICAMBA 4 DMA preplant applications to geographic areas with average annual rainfall less than 25". If applying a spring preplant treatment following application of a fall preplant (postharvest) treatment, then the combination of both treatments may not exceed 2 pounds acid equivalent per acre.

**COTTON TANK MIXES**

For control of grasses or additional broadleaf weeds, LIBERTY DICAMBA 4 DMA may be tank mixed with prometryn, paraquat, and glyphosate herbicides.

**SORGHUM (MILO)**

Observe all precautions, including the reference to crops growing under stress. Read and follow mixing and application instructions. Applications of LIBERTY DICAMBA 4 DMA to sorghum during periods of rapid growth may result in temporary leaning of plants or rolling of leaves. These effects are usually outgrown within 10 to 14 days. Do not graze or feed treated sorghum forage or silage prior to mature grain stage. If sorghum is grown for pasture or hay, refer to the pasture use section of this label. Do not apply LIBERTY DICAMBA 4 DMA to sorghum grown for seed production. Make no more than one application per growing season.

**WEEDS CONTROLLED**

LIBERTY DICAMBA 4 DMA, when applied at the recommended rate for sorghum, will control many actively growing ANNUAL broadleaf weeds and will reduce competition from established PERENNIAL broadleaf weeds as well as control their seedlings. (Refer to WEED LIST)

**RATES AND TIMINGS**

LIBERTY DICAMBA 4 DMA may be applied to emerged and actively growing weeds at least 15 days prior to planting. Postemergence application of LIBERTY DICAMBA 4 DMA must be made after sorghum is in the spike stage (all sorghum emerged) but before sorghum is 15 inches tall. For best performance, make applications when sorghum is in the 3 to 5 leaf stage and weeds are small (less than 3 inches tall). Use drop pipes (drop nozzles) if sorghum is taller than 8 inches. Keeping the spray off the sorghum leaves and out of the whorl will reduce the likelihood of crop injury and improve spray coverage of weed foliage.

**Broadcast rate per treated acre:** 1/2 pint (1/4 lb a.i.)

**TANK MIX TREATMENTS**

**LIBERTY DICAMBA 4 DMA plus Atrazine**

For improved control of emerged, actively growing broadleaf weeds including triazine resistant species and added suppression of perennial broadleaf weeds, tank mix 1/2 pint LIBERTY DICAMBA 4 DMA with 0.5 to 1.25 lbs a.i. atrazine per treated acre. For control of grasses (less than 1.5 inches tall), tank mix 1/2 pint LIBERTY DICAMBA 4 DMA with 2 lbs a.i. atrazine per treated acre. For best performance and minimal crop injury, make application when sorghum is 3-8 inches tall and when broadleaf weeds are small (less than 6 inches tall). Application of atrazine must be made before sorghum is beyond 12 inches tall. The atrazine rate will depend upon soil texture and length of residual weed control desired. Follow all state and Federal restrictions pertaining to atrazine applications.

**LIBERTY DICAMBA 4 DMA plus bromoxynil**

For improved control of broadleaf weeds, tank mix 1/2 pint LIBERTY DICAMBA 4 DMA with 1-1 1/2 pint bromoxynil herbicide.

per treated acre Make application at 4 leaf to 15 inch tall sorghum Use drop nozzles to direct spray beneath sorghum leaves when sorghum is greater than 8 inches tall

READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS DIRECTIONS FOR USE APPLICATION RATES AND TIMINGS AND OTHER RESTRICTIONS

**OVERLAY (SEQUENTIAL)TREATMENTS**

LIBERTY DICAMBA 4 DMA may be applied to ground previously treated with one or more of the following herbicides

Herbicide	Maximum Rate Per Treated Acre (lbs a i )
alachlor (Lasso®)	4
(Screen® treated seed)	2.5
atrazine <sup>1</sup>	2.5
metolachlor	5
propachlor (Ramrod®)	

<sup>1</sup> Maximum use rate for atrazine is determined by soil type tillage practices used surface residue and state or local restrictions Follow the more restrictive requirements when determining the maximum use rate for atrazine

**PREHARVEST USES**

FOR USE ONLY IN THE STATES OF TEXAS AND OKLAHOMA

LIBERTY DICAMBA 4 DMA may be applied for weed suppression any time after the sorghum has reached the soft dough stage An agriculturally approved surfactant may be used to improve performance For aerial applications use at least 2 gallons of water based carrier per treated acre

Delay harvest until 30 days after treatment

**Broadcast rate per treated acre** 1/2 pint (1/4 lb a i )

**SMALL GRAINS (WHEAT BARLEY AND OATS) NOT UNDERSEEDED TO LEGUMES**

**IMPORTANT**

Observe all precautions Read and follow cleaning mixing and application instructions

If small grains are used for pasture or hay the following restrictions apply

- Animals cannot be removed from treated area for slaughter prior to 30 days after last application
- There is no waiting period between treatment and grazing for non lactating dairy animals
- Treated areas may not be grazed by lactating dairy animals before 7 days after treatment
- Do not harvest hay from treated areas before 37 days after treatment

**NOTE** Observe all precautions and restrictions on the labels of products used in tank mix treatments

**WEEDS CONTROLLED**

LIBERTY DICAMBA 4 DMA or combinations with listed tank mix partners will provide control or suppression of the annual broadleaf weeds listed below For improved control of listed weeds it is recommended that LIBERTY DICAMBA 4 DMA be applied in a tank mix with other herbicides Refer to specific crop for tank mix options

Alkanet <sup>1</sup>	Knawel (German Moss)	Pigweed Tumble
Bedstraw Catchweed <sup>1</sup>	Knotweed Prostrate	Pineappleweed <sup>1</sup>
Bindweed Field <sup>2</sup>	Kochia	Plantain Broadleaf <sup>2</sup>
Buckwheat Tartary	Ladysthumb	Poppy Red Horned <sup>1</sup>
Buckwheat Wild	Lambsquarters Common	Puncturevine <sup>1</sup>
Carpetweed <sup>1</sup>	Lettuce Miners <sup>1</sup>	Purslane Common
Chamomile Corn	Lettuce Prickly	Radish Wild <sup>1</sup>
Chervil Bur <sup>1</sup>	Mallow Common	Ragweed Common
Chickweed Common <sup>1</sup>	Mayweed Chamomile	Ragweed Giant
Cockle Corn	(Dogfennel) <sup>1</sup>	(Buffaloweed) <sup>1</sup>
Cockle Cow	Mustard Blue	Rocket London <sup>1</sup>
Cocklebur Common	(Purple) <sup>1</sup>	Rocket Yellow <sup>1</sup>
Cornflower	Mustard Tansy	Salsify (Goatsbeard) <sup>1</sup>
(Bachelorbutton) <sup>1</sup>	Mustard Treacle <sup>1</sup>	Shepherdspurse <sup>1</sup>
Dandelion Common <sup>2</sup>	Mustard Tumble	Smartweed Green
Dock Curly <sup>2</sup>	(Jim Hill) <sup>1</sup>	Smartweed Pennsylvania
Dragonhead American <sup>1</sup>	Mustard Wild <sup>1</sup>	Sorrel Red
Evening Primrose	Nightshade Black	(Sheep Sorrel) <sup>1</sup>
Cutleaf <sup>1</sup>	Nightshade Cutleaf <sup>1</sup>	Sowthistle Annual
Falseflax Smallseeded <sup>1</sup>	Nightshade Silverleaf <sup>2</sup>	Starthistle Yellow <sup>1</sup>
Fiddleneck (Tarweed) <sup>1</sup>	(White Horsenettle)	Sunflower Common (Wild)
Flixweed <sup>1</sup>	Pennycress Field	Thistle Canada <sup>2</sup>
Fumitory <sup>1</sup>	(Fanweed Frenchweed	Thistle Russian
Gromwell Corn <sup>1</sup>	Stinkweed)	Velvetleaf
Groundsel Common <sup>1</sup>	Pepperweed Peppergrass <sup>1</sup>	Vetch <sup>1</sup>

Hempnettle <sup>1</sup>	Pigweed Redroot (Carelessweed)	Yarrow Common <sup>2</sup>
Henbit		
Jacobs Ladder <sup>1</sup>	Pigweed Rough	

<sup>1</sup> These weeds will be controlled with LIBERTY DICAMBA 4 DMA tank mixtures Refer to tank mix label for specific weeds controlled

<sup>2</sup> LIBERTY DICAMBA 4 DMA tank mixes will provide suppression of established broadleaf weeds and control their seedlings

**RATES AND TIMINGS**

Application of LIBERTY DICAMBA 4 DMA may be made before during or after planting small grains For best performance make applications when weeds are in the 2 3 leaf stage and rosettes are less than 2 inches across Application of LIBERTY DICAMBA 4 DMA to small grains during periods of rapid growth may result in crop leaning This condition is temporary and will not reduce crop yields

Use LIBERTY DICAMBA 4 DMA at 2 to 4 fluid ounces per treated acre in wheat fall seeded barley and oats and at 2 to 3 fluid ounces per treated acre in spring seeded barley Use the higher level of listed rate ranges when treating difficult to control weeds such as kochia wild buckwheat cow cockle prostrate knotweed Russian thistle and prickly lettuce or when dense vegetative growth occurs

LIBERTY DICAMBA 4 DMA used in a tank mix with other herbicides offers the best spectrum of weed control and herbicide tolerant or resistant weed management Refer to specific crop for LIBERTY DICAMBA 4 DMA rate and application timing

For applications prior to the emergence of weeds or when sulfonylurea resistant weeds are present or suspected use a minimum of 3 fluid ounces per treated acre of LIBERTY DICAMBA 4 DMA with a tank mix herbicide Non sulfonylurea herbicides such as 2 4 D or MCPA tank mixed with LIBERTY DICAMBA 4 DMA will offer more consistent control of sulfonylurea resistant weeds

When tank mixing with sulfonylurea herbicides such as Ally® Amber® Express® Finesse® Glean® and Harmony® Extra use an agriculturally approved surfactant of at least 80% active ingredient at the rate of 1-4 pints/100 gallons of spray or not more than 0 25 0 5% by volume Use the highest rate of surfactant when using the lower rate ranges of the tank mix and/or when treating more mature and difficult to control weeds or dense vegetative growth

**FALL AND SPRING SEEDED WHEAT**

LIBERTY DICAMBA 4 DMA MUST BE APPLIED TO FALL SEEDED WHEAT PRIOR TO THE JOINTING STAGE APPLICATIONS TO SPRING SEEDED WHEAT MUST BE MADE BEFORE WHEAT REACHES THE 6 LEAF STAGE

**NOTE** Early developing wheat varieties such as TAM 107 MADISON or WAKEFIELD must receive application between early tillering and the jointing stage Care should be taken in staging these varieties to be certain that the application occurs prior to the jointing stage

**TANK MIX TREATMENTS**

LIBERTY DICAMBA 4 DMA may be tank mixed with one or more of the following herbicides Read and follow the label of each tank mix product used for precautionary statements directions for use weeds controlled geographic and other restrictions

**Broadcast rate per treated acre**

Apply 2 4 fluid ounces LIBERTY DICAMBA 4 DMA with

Product	Active Ingredient	Formulation	Amount of Product Per Acre
2 4 D Amine or Ester	2 4 D	4 lb/gal	8 12 fluid oz ( 25 375 lb a i /A) <sup>1</sup>
MCPA Amine or Ester	MCPA	4 lb/gal	8 12 fluid oz ( 25 375 lb a i /A) <sup>1</sup>
Ally®	metsulfuron methyl	60% DF	1/10 oz
Amber®	triasulfuron	75% DF	0 28 oz
Express®	thifensulfuron + tribenuron methyl	75% DF	1/6 oz
Finesse®	chlorsulfuron + metsulfuron methyl	75% DF	1/3 oz
Glean®	chlorsulfuron	75% DF	1/6 oz
Harmony® Extra	thifensulfuron + tribenuron methyl	75% DF	1/3 oz
bromoxynil	bromoxynil	2 lb/gal	1 1 5 pts
Bronate®	bromoxynil + MCPA	4 lb/gal	1 2 pts
Curtail®	clopyralid+ 2 4 D	2 38 lb/gal	2 2 2/3pts
clopyralid	clopyralid	3 lb/gal	1/4 1/3 pt
diuron <sup>2</sup>	diuron	80% DF	1/2 1 5 lbs
metribuzin <sup>2</sup>	metribuzin	75% DF	1 10 oz
Dakota <sup>3</sup>	fenoxaprop ethyl+MCPA	3 1 lb/gal	16 fluid oz
Tiller® <sup>3</sup>	fenoxaprop ethyl + MCPA + 2 4D	2 7 lb/gal	1 1 7 pts

<sup>1</sup> When using formulations other than 4 lbs/gal use pounds active/acre listed

<sup>2</sup> Tank mixtures for fall seeded wheat only

<sup>3</sup> Use 2 fluid ounces of LIBERTY DICAMBA 4 DMA only Do not use if wild oats is the target weed Do not use LIBERTY DICAMBA 4 DMA as a tank mix treatment with Dakota® or Tiller® on Durum wheat

**SPECIAL USE TANK MIXES FOR SPRING AND FALL SEEDED WHEAT**  
(See Footnotes for Applicable Uses)

**Broadcast rate per treated acre**

Apply 3 4<sup>1</sup> fluid ounces LIBERTY DICAMBA 4 DMA with

Product <sup>2</sup>	Active Ingredient	Formulation	Amount of Product Per Acre
2 4 D or MCPA Amine	2 4 D or MCPA	4 lb/gal	1 2 pts <sup>3</sup> ( 5 1 0 lb a i /A ) <sup>4</sup>
2 4 D or MCPA Ester	2 4 D or MCPA	4 lb/gal	1 1 5 pts <sup>3</sup> ( 5 7 5 lb a i /A ) <sup>4</sup>
Allv®	metsulfuron methyl	60% DF	1/20 1/10 oz
Amber®	triasulfuron	75% DF	0.14 0.28 oz
Express®	thifensulfuron + tribenuron methyl	75% DF	1/12 1/6 oz
Finesse®	chlorsulfuron + metsulfuron methyl	75% DF	1/6 1/3 oz
Glean®	chlorsulfuron	75% DF	1/6 oz
Harmony® Extra	thifensulfuron + tribenuron methyl	75% DF	1/6 1/3 oz
Metsulfuron methyl + 2 4 D Amine or Ester <sup>5</sup>	Metsulfuron methyl + 2 4 D	60% DF + 4 lb/gal	1/20 1/10 oz + 8 fl oz
Amber® + 2 4 D Amine or Ester <sup>5</sup>	triasulfuron + 2 4 D	75% DF + 4 lb/gal	0.14 0.28 oz + 8 fl oz
Express® + 2 4 D Amine or Ester <sup>5</sup>	(thifensulfuron + tribenuron methyl) + 2 4 D	75% DF + 4 lb/gal	1/12 1/6 oz + 8 fl oz
Finesse® + 2 4 D Amine or Ester <sup>5</sup>	(chlorsulfuron + metsulfuron methyl) + 2 4 D	75% DF + 4 lb/gal	1/6 1/3 oz + 8 fl oz
Glean® + 2 4 D Amine or Ester <sup>5</sup>	chlorsulfuron + 2 4 D	75% DF + 4 lb/gal	1/6 + 8 fl oz
Harmony® Extra + 2 4 D Amine or Ester <sup>5</sup>	(thifensulfuron + tribenuron methyl) + 2 4 D	75% DF + 4 lb/gal	1/6 1/3 oz + 8 fl oz
glyphosate <sup>6</sup>	glyphosate	3 0 lb/gal	12 16 fl oz

LIBERTY DICAMBA 4 DMA may be used at 6 fluid ounces on fall seeded wheat in Western Oregon as a spring application only In CO KS NM OK and TX up to 8 fluid ounces of LIBERTY DICAMBA 4 DMA may be applied on fall seeded wheat after it exceeds the 3 leaf stage for suppression of perennial weeds such as field bindweed Applications may be made in the fall following a frost but before a killing freeze LIBERTY DICAMBA 4 DMA may be tank mixed with 2 4 D amine at 8 fluid ounces after wheat begins to tiller Periods of extended stress such as cold and wet weather may enhance the possibility of crop injury For fall applications only do not use if the potential for crop injury is not acceptable

<sup>2</sup> Do not use low rates of sulfonylurea herbicides such as Metsulfuron methyl Amber Express Finesse Glean and Harmony Extra on more mature weeds and/or on dense vegetative growth

<sup>3</sup> NOTE For use on Fall Seeded Wheat only Do not use unless potential crop injury will be acceptable

<sup>4</sup> When using formulations other than 4 lb/gal use pounds active/acre listed

<sup>5</sup> Use for improved control of Russian thistle flixweed gromwell mayweed and fiddleneck

<sup>6</sup> LIBERTY DICAMBA 4 DMA may be applied at 2 fluid ounces with any glyphosate formulation labeled for use as a preplant application to small grains with no waiting period prior to planting Read and follow label directions of the tank mix product for adjuvant use recommendations

**FALL SEEDED BARLEY**

LIBERTY DICAMBA 4 DMA MUST BE APPLIED TO FALL SEEDED BARLEY PRIOR TO THE JOINTING STAGE

**NOTE** For spring barley varieties that are seeded during the winter months or later follow the rates and timings given for Spring Seeded Barley

**TANK MIX TREATMENTS**

LIBERTY DICAMBA 4 DMA may be tank mixed with one or more but not limited to the following herbicides Read and follow the label of each tank mix product used for precautionary statements directions for use weeds controlled and geographic and other restrictions

**Broadcast rate per treated acre**

Apply 2 4 fluid ounces LIBERTY DICAMBA 4 DMA with

Product <sup>1</sup>	Active Ingredient	Formulation	Amount of Product Per Acre
2 4 D Amine or Ester	2 4 D	4 lb/gal	8 fluid oz ( 2 5 lb a i /A ) <sup>2</sup>
MCPA Amine or Ester	MCPA	4 lb/gal	8 12 fluid oz ( 2 5 3 7 5 lb a i /A )
metsulfuron methyl	metsulfuron methyl	60% DF	1/20 1/10 oz
Amber®	triasulfuron	75% DF	0 14 0 28 oz
Express®	thifensulfuron + tribenuron methyl	75% DF	1/12 1/6 oz
Finesse®	chlorsulfuron + metsulfuron methyl	75% DF	1/6 1/3 oz
Glean®	chlorsulfuron	75% DF	1/6 oz
Harmony® Extra	thifensulfuron + tribenuron methyl	75% DF	1/6 1/3 oz
metribuzin	metribuzin	75% DF	1 10 oz
bromoxynil	bromoxynil	2 lb/gal	1 1 1/2 pts
Bronate®	bromoxynil + MCPA	4 lb/gal	3/4 1 1/2 pts

<sup>1</sup> Do not use low rates of sulfonylureas (metsulfuron methyl Amber® Express® Finesse Glean® and Harmony® Extra) on more mature weeds and/or on dense vegetative growth

<sup>2</sup> When using formulations other than 4 lb/gal use pounds active/acre listed

**SPRING SEEDED BARLEY**

LIBERTY DICAMBA 4 DMA MUST BE APPLIED BEFORE SPRING SEEDED BARLEY EXCEEDS THE 4 LEAF STAGE  
**TANK MIX TREATMENTS**

LIBERTY DICAMBA 4 DMA may be mixed with one or more of the following herbicides Read and follow the label of each tank mix product used for precautionary statements directions for use weeds controlled and geographic and other restrictions

**Broadcast rate per treated acre**

Apply 2.4 fluid ounces LIBERTY DICAMBA 4 DMA with

Product <sup>1</sup>	Active Ingredient	Formulation	Amount of Product Per Acre
MCPA Amine or Ester	MCPA	4 lb/gal	8.12 fluid oz (25.375 lb a.i./A) <sup>2</sup>
metsulfuron methyl	metsulfuron methyl	60% DF	1/20 1/10 oz
Amber®	triasulfuron	75% DF	0.14 0.28 oz
Express®	thifensulfuron + tribenuron methyl	75% DF	1/12 1/6 oz
Finesse®	chlorsulfuron + metsulfuron methyl	75% DF	1/6 1/3 oz
Glean®	chlorsulfuron	75% DF	1/6 oz
Harmony® Extra	thifensulfuron + tribenuron methyl	75% DF	1/6 1/3 oz
metribuzin	metribuzin	75% DF	1.10 oz
bromoxynil	bromoxynil	2 lb/gal	1.1 1/2 pts
Bronate®	bromoxynil + MCPA	4 lb/gal	3/4 1 1/2 pts

<sup>1</sup> Do not use low rates of sulfonylureas (metsulfuron methyl Amber® Express® Finesse® Glean® and Harmony® Extra) on more mature weeds and/or on dense vegetative growth

<sup>2</sup> When using formulations other than 4 lb/gal use pounds active/acre listed

**FALL AND SPRING SEEDED OATS**

LIBERTY DICAMBA 4 DMA MUST BE APPLIED BEFORE SPRING SEEDED OATS EXCEED THE 5 LEAF STAGE  
 APPLICATIONS TO FALL SEEDED OATS MUST BE MADE PRIOR TO THE JOINTING STAGE

**TANK MIX TREATMENTS**

LIBERTY DICAMBA 4 DMA may be tank mixed with one or more of the following herbicides Read and follow the label of each tank mix product used for precautionary statements directions for use weeds controlled geographic and other restrictions

**Broadcast rate per treated acre**

Apply 2.4 fluid ounces LIBERTY DICAMBA 4 DMA with

Product	Active Ingredient	Formulation	Amount of Product Per Acre
MCPA Amine or Ester	MCPA	4 lb/gal	8.12 fluid oz (25.375 lb a.i./A) <sup>1</sup>

<sup>1</sup> When using formulations other than 4 lb/gal use pounds active/acre listed

**FALL AND SPRING SEEDED TRITICALE  
 EXCEPT CALIFORNIA**

**EARLY SEASON APPLICATIONS**

Apply 2.4 fluid ounces of LIBERTY DICAMBA 4 DMA to triticale

Early season applications to fall seeded triticale must be made prior to jointing stage

Early season applications to spring seeded triticale must be made before triticale reaches the 6 leaf stage

**TANK MIXES**

For best performance should be used in tank mix combination with bromoxynil

**SUGARCANE**

Observe all precautions Read and follow mixing and application instructions

Consult your local or state authorities for possible application restrictions especially concerning aerial applications and advice concerning special local use situations

**WEEDS CONTROLLED**

LIBERTY DICAMBA 4 DMA when applied at specified rates will control many ANNUAL BIENNIAL and PERENNIAL broadleaf weeds commonly found in sugarcane (Refer to WEED LIST)

**RATES AND TIMINGS**

Application of LIBERTY DICAMBA 4 DMA may be made any time after weeds have emerged and are actively growing but before the close in stage of sugarcane Application rates and timing of LIBERTY DICAMBA 4 DMA are given below Use the higher level of listed rate ranges when treating dense vegetative growth



17/26

**NOTE** Retreatments may be made as needed however, do not exceed a total of 2 pints (1 lb a i) of LIBERTY DICAMBA 4 DMA per treatment with a maximum of 2 treatments per year

Weed Stage & Type	Broadcast Rate Per Treated Acre		PHI
	Amount of Formulated LIBERTY DICAMBA 4 DMA (pints)	Equivalent Lbs a i	
Annual Small actively growing Established weed growth	1/2 1	1/4 1/2	87 days
	1 1 1/2	1/2 3/4	
Biennial	1 2	1/2 1	
Perennial	2	1 <sup>1</sup>	

<sup>1</sup> Application made over the top of actively growing sugarcane may result in crop injury  
When possible direct the spray beneath the sugarcane canopy in order to minimize the likelihood of crop injury The use of directed sprays will also aid in maximizing spray coverage of weed foliage

**TANK MIX TREATMENTS**

LIBERTY DICAMBA 4 DMA may be tank mixed with one or more of the following herbicides for control of grasses or additional broadleaf weeds Read and follow the label of each tank mix product used for precautionary statements directions for use rates and timings weeds controlled geographic and other restrictions

Herbicide	Rate Per Treated Acre (lbs a i)
ametryn	2/5 8
asulam	2 3 1/3
atrazine	2/5 4
2 4 D	1 /2 3*

Application of LIBERTY DICAMBA 4 DMA plus 2 4 D tank mix at the higher listed rate ranges may result in crop injury

**PASTURE, HAY RANGELAND AND FARMSTEAD (Non Cropland)**

LIBERTY DICAMBA 4 DMA is recommended for use for pasture hay rangeland farmstead (non cropland) (including fence rows and non irrigation ditchbanks) for broadleaf weed and brush control LIBERTY DICAMBA 4 DMA may also be applied to non cropland areas for the control of broadleaf weeds in Noxious Weed Control Programs Districts or Areas including broadcast or spot treatment of roadsides and highways utilities railroad and pipeline rights of way Noxious weeds must be recognized at the state level but programs may be administered at state county or other level

Observe all precautions Read and follow mixing and application instructions

LIBERTY DICAMBA 4 DMA uses described in this section also pertain to small grains (such as barley forage sorghum oats rye sudangrass or wheat) grown for pasture use only

**NEWLY SEEDED AREAS** including small grains grown for pasture may be severely injured if rates of LIBERTY DICAMBA 4 DMA greater than 1 pint/A are applied

**ESTABLISHED GRASS CROPS** growing under stress can exhibit various injury symptoms that may be more pronounced if herbicides are applied

Bentgrass carpetgrass buffalograss and St Augustine grass may be injured at rates exceeding 1 pint LIBERTY DICAMBA 4 DMA (1/2 lb a i) per treated acre Usually colonial bentgrasses are more tolerant than creeping types Velvetgrasses are most easily injured Treatments will kill or injure alfalfa clovers lespedeza wild winter peas vetch and other legumes

**ANIMALS CANNOT BE REMOVED FROM TREATED AREA FOR SLAUGHTER PRIOR TO 30 DAYS AFTER LAST APPLICATION**

**THERE IS NO WAITING PERIOD BETWEEN TREATMENT AND GRAZING FOR NON LACTATING ANIMALS**

**TIMING RESTRICTIONS FOR LACTATING DAIRY ANIMALS FOLLOWING TREATMENT**

LIBERTY DICAMBA 4 DMA Rate Per Treated Acre	Days Before Grazing	Days Before Hay Harvest
Up to 1 pint (1/2 lb a i)	7 days	37 days
Up to 2 pints (1 lb a i)	21 days	51 days

**NOTE** Observe all precautions and restrictions on labels of products used in tank mixtures

**MIXING AND APPLICATION**

LIBERTY DICAMBA 4 DMA can be applied using water oil in water emulsions including invert systems or sprayable fluid fertilizer as a carrier A COMPATIBILITY TEST (see COMPATIBILITY TEST section) should be made prior to tank mixing To prepare oil in water emulsions half fill spray tank with water then add appropriate amount of emulsifier With continuous agitation slowly add the herbicide and then the oil (such as diesel oil or fuel oil) or a premix of oil plus additional emulsifier to spray tank Complete filling of spray tank with water Maintain vigorous agitation during spray operation to prevent oil and water from forming separate layers

LIBERTY DICAMBA 4 DMA may be applied broadcast using either ground or aerial application equipment When using ground equipment apply 3 to 600 gallons of diluted spray per treated acre Volume of spray applied will depend on the height density and type of weeds or brush being treated and on the type of equipment being used When using aerial equipment apply 2 to 40 gallons of diluted spray per treated acre in a water based carrier

LIBERTY DICAMBA 4 DMA may be applied to individual clumps or small areas (SPOT TREATMENT) of undesirable vegetation using handgun or similar types of application equipment Apply diluted sprays to allow complete wetting (up to

run off) of foliage and stems

Herbicide adjuvants or other spray additives (emulsifiers surfactants wetting agents drift control agents or penetrants) may be used for wetting penetration or drift control Spray additives must be agriculturally approved when used in pasture applications If spray additives are used read and follow all use recommendations and precautions on product label

**WEEDS CONTROLLED**

LIBERTY DICAMBA 4 DMA when applied at specified rates will give control many ANNUAL BIENNIAL and PERENNIAL broadleaf weeds and many WOODY brush and vine species commonly found in pasture hay rangeland and farmstead (non cropland) areas (Refer to WEED LIST) Noted (\*)PERENNIAL weeds may be controlled with lower rates of either LIBERTY DICAMBA 4 DMA or LIBERTY DICAMBA 4 DMA plus 2 4 D See the following RATES AND TIMINGS section

**RATES AND TIMINGS**

Application rates and timing of LIBERTY DICAMBA 4 DMA are given below Use the higher level of listed rate ranges when treating dense or tall vegetative growth

**NOTE** Retreatments may be made as needed however, do not exceed a total of 2 pints (1 lb a i ) of LIBERTY DICAMBA 4 DMA per treatment with a maximum of 2 treatments per year

Weed Stage & Type	Broadcast Rate Per Treated Acre	
	Amount of Formulated LIBERTY DICAMBA 4 DMA (pints)	Equivalent Lbs a i
<b>Annual</b>		
Small actively growing	1/2 1	1/4 1/2
Established weed growth	1 1 1/2	1/2 3/4
<b>Biennial<sup>1</sup></b>		
Rosette diameter		
Less than 3 inches	1/2 1	1/4 1/2
3 inches or more	2	1
Bolting	2	1
<b>Perennial</b>		
Suppression or top growth control	1 2	1/2 1
Noted (*) Perennials	2	1
Other Perennials	2	1
<b>Woody Brush &amp; Vines</b>		
Top Growth Suppression	1 2	1/2 1
Top Growth Control <sup>2</sup>	2	1
Stems and Stem Suppression	2	1

<sup>1</sup> For best performance make application when BIENNIAL WEEDS are in the rosette stage

<sup>2</sup> Species noted in WEED LIST section will require tank mixtures for adequate control Rates above 1 0 lb a i /A are spot treatments only

**TANK MIX TREATMENTS**

READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS DIRECTIONS FOR USE APPLICATION RATES AND OTHER RESTRICTIONS

LIBERTY DICAMBA 4 DMA may be tank mixed with one or more of the following herbicides for control of grasses additional broadleaf weeds and woody brush and vines

Herbicide	Rate Per Treated Acre (lbs a i )
Pasture hay rangeland and farmstead (non cropland) use	
glyphosate	3/4 3 3/4
metsulfuron methyl	0 0038 0 011
paraquat	1/2 1
picloram	1/8 3
triclopyr	3/4 9
2,4 D	1/4 6

Due to the variations that may occur in formulated products and specific use ingredients (e g water supplies) a COMPATIBILITY TEST is recommended prior to actual tank mixing

**CUT SURFACE TREE TREATMENTS**

LIBERTY DICAMBA 4 DMA may be applied as a cut surface treatment for control of unwanted trees and prevention of sprouts of cut trees A mix of 1 part LIBERTY DICAMBA 4 DMA with 1 to 3 parts water should be used in application Use the lower dilution when treating difficult to control species

FRILL OR GIRDLE TREATMENTS Make a continuous cut or a series of overlapping cuts using an axe to girdle tree trunk Spray or paint cut surface with the LIBERTY DICAMBA 4 DMA/water mix

19/26

**STUMP TREATMENTS** Spray or paint freshly cut surface with the water mix. The area adjacent to the bark should be thoroughly wet.  
 Note: For more rapid foliar effects 2.4 D may be added to the LIBERTY DICAMBA 4 DMA/water mix

**DORMANT APPLICATIONS FOR CONTROL OF MULTIFLORA ROSE**

LIBERTY DICAMBA 4 DMA can be applied when plants are dormant as an undiluted SPOT CONCENTRATE directly to the soil or as a LO OIL BASAL BARK treatment using an oil water emulsion solution.  
 SPOT CONCENTRATE applications of LIBERTY DICAMBA 4 DMA should be applied directly to the soil as close as possible to the root crown but within 6-8 inches of the crown. On sloping terrain application should be made to the uphill side of the crown. Do not make application when snow or water prevents applying LIBERTY DICAMBA 4 DMA directly to the soil. The use rate of LIBERTY DICAMBA 4 DMA is dependent on the canopy diameter of the multiflora rose. Examples: Use LIBERTY DICAMBA 4 DMA at 1/4, 1 or 2 1/4 fluid ounces of product respectively for 5, 10 or 15 feet canopy diameters. Do not exceed a total of 2 quarts LIBERTY DICAMBA 4 DMA per acre per year.  
 LO OIL BASAL BARK applications of LIBERTY DICAMBA 4 DMA should be applied to the basal stem region from the ground line up to a height of 12 to 18 inches. Spray until runoff with special emphasis on covering the root crown. For best results make application when plants are dormant. Do not make application after bud break or when plants are showing signs of active growth. Do not make application when snow or water prevents applying LIBERTY DICAMBA 4 DMA to the ground line. Refer to Mixing and Applications above in this section for method of preparing oil in water emulsion. Example for making approximately 2 gallons of a LO OIL spray solution mixture: combine 1 1/2 gallons water plus 1 ounce emulsifier plus 1 pint LIBERTY DICAMBA 4 DMA plus 2 1/2 pints of No. 2 diesel fuel. Adjust amounts of materials used proportionately to the amount of final spray solution desired. Do not exceed 8 gallons of spray solution mix applied per acre per year.

**CONSERVATION RESERVE PROGRAM (CRP) ACRES**

LIBERTY DICAMBA 4 DMA can be used on both newly seeded and established grasses grown in Conservation Reserve or Federal Set Aside Programs. For program lands such as Conservation Reserve Program consult program rules to determine whether grass or hay may be used. The more restrictive requirements of the program rules or this label must be followed.  
 Observe all precautions, mixing and application directions.  
 LIBERTY DICAMBA 4 DMA treatment will injure or may kill alfalfa, clovers, lespedeza, wild winter peas, vetch, and other legumes.  
 Agriculturally approved surfactants may be added to the spray mixture to improve postemergence weed control, particularly in dry growing conditions.

Do not use adjuvants containing penetrants such as petroleum based oils after grass emergence on newly seeded grasses.

**NEWLY SEEDER AREAS**

LIBERTY DICAMBA 4 DMA may be applied either preplant or postemergence to newly seeded grasses or small grains such as barley, oats, rye, sudangrass, wheat, or other grain species grown as a cover crop. Postemergence applications may be made after seedling grasses exceed the 3 leaf stage. Rates of LIBERTY DICAMBA 4 DMA greater than 1 pint per treated acre may severely injure newly seeded grasses. Preplant applications: injury to new seedlings may occur if intervals between application and grass planting is less than 45 days per pint of LIBERTY DICAMBA 4 DMA per treated acre West of the Mississippi River or 20 days per pint East of the Mississippi River.

**ESTABLISHED GRASS STANDS**

Established grass stands are perennial grasses planted one or more seasons prior to treatment. Certain species bentgrass, carpetgrass, smooth brome, buffalograss, or St. Augustine grass may be injured when treated with LIBERTY DICAMBA 4 DMA at rates exceeding 1 pint per treated acre.

**WEEDS CONTROLLED**

LIBERTY DICAMBA 4 DMA when applied at specified rates will control many annual and biennial weeds and provide control or suppression of many perennial weeds. (Refer to WEED LIST)

**RATES AND TIMINGS**

Application rates and timing of LIBERTY DICAMBA 4 DMA treatment are given below. Use the higher rate of the rate range when vegetation is either dense or tall, or when weeds are growing under stressed conditions such as drought or cool temperature.

**NOTE:** Retreatments may be made as needed, however, do not exceed a total of 4 pints (2 lbs a.i.) of LIBERTY DICAMBA 4 DMA per treated acre during a growing season applied at a rate of 2 pints (1 lb a.i.) LIBERTY DICAMBA 4 DMA per treatment.

Weed Stage & Type	Broadcast Rate Per Treated Acre		PHI
	Amount of Formulated LIBERTY DICAMBA 4 DMA (pints)	Equivalent lbs a.i.	
<b>Annual</b>			
Small actively growing	1/4, 1	1/8, 1/2	
Established weed growth	1	1/2	

<b>Biennial</b> <sup>1 2</sup>			For grass forage 0 days
Rosette diameter			
Less than 3 inches	1/2 1	1/4 1/2	For grass hay 7 days
3 inches or greater	1 2	1/2 1	
Bolting biennial	2	1	
<b>Perennial</b> <sup>2</sup>			
Suppression/Control	2	1	

<sup>1</sup> For best results treat Biennial weeds with LIBERTY DICAMBA 4 DMA when they are in the rosette stage of growth

<sup>2</sup> Biennial and Perennial weeds will require follow up (sequential) treatments for seedling control and escapes

**TANK MIX TREATMENTS**

To control grasses and additional broadleaf weeds LIBERTY DICAMBA 4 DMA may be tank mixed with other herbicides registered for use in Conservation Reserve Programs such as 2 4 D glyphosate paraquat metsulfuron and others  
**READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS DIRECTIONS FOR USE APPLICATION RATES AND OTHER RESTRICTIONS**

**ASPARAGUS**

FOR USE ONLY IN THE STATES OF CALIFORNIA OREGON AND WASHINGTON

Observe all precautions Read and follow mixing and application instructions

**NOTE**

If spray contacts emerged spears crooking (twisting) of some spears may result If such crooking occurs discard affected spears

Do not harvest prior to 24 hours after treatment

Do not use in the Coachella Valley of California

Multiple applications may be made per growing season however **DO NOT EXCEED** a total of 1 pint (1/2 lb a i ) of LIBERTY DICAMBA 4 DMA per treated acre per crop year

**RATES AND TIMINGS**

Apply LIBERTY DICAMBA 4 DMA to emerged and actively growing weeds in 40 to 60 gallons of diluted spray per treated acre immediately after cutting the field but at least 24 hours before the next cutting

LIBERTY DICAMBA 4 DMA may be applied in a tank mixture with either 2 4 D or glyphosate herbicide for improved control of noted (\*) weeds **READ AND FOLLOW 2 4 D OR GLYPHOSATE PRODUCT LABELING FOR PRECAUTIONARY STATEMENTS DIRECTIONS FOR USE APPLICATION RATES AND TIMINGS AND OTHER RESTRICTIONS**

Weeds	Rate Per Treated Acre
Mustard Black Pigweed Redroot (Carelessweed) Sowthistle Annual *Thistle Canada Thistle Russian	1/2 1 pt (1/4 1/2 lb a i )
Bindweed Field Chickweed Common Goosefoot Nettleleaf Radish Wild Thistle Milk	1 pt (1/2 lb a i )

**TURF AND LAWNS**

FOR USE IN FARMSTEAD (NON CROPLAND) AND SOD FARMS

**IMPORTANT**

Observe all precautions Read and follow mixing and application instructions

To avoid injury to newly seeded grasses application of LIBERTY DICAMBA 4 DMA should be delayed until after the second mowing Further more application rates in excess of 1 pint (1/2 lb a i ) per treated acre may cause noticeable stunting or discoloration of sensitive grass species such as bentgrass carpetgrass buffalograss and St Augustine grass In areas where roots of sensitive plants extend do not apply in excess of 1/4 pint (1/8 lb a i ) of LIBERTY DICAMBA 4 DMA per treated acre on coarse textured (sandy type) soils or in excess of 1/2 pint (1/4 lb a i ) per treated acre on fine textured (clayey type) soils Do not make repeat applications in these areas for 30 days and until previous applications of LIBERTY DICAMBA 4 DMA have been activated in the soil by rain or irrigation

**WEEDS CONTROLLED**

LIBERTY DICAMBA 4 DMA when applied at specified rates will give control of many ANNUAL BIENNIAL and noted (\*) PERENNIAL broadleaf weeds commonly found in turf LIBERTY DICAMBA 4 DMA will also give growth suppression of many other listed PERENNIAL broadleaf weeds and WOODY brush and vine species (Refer to WEED LIST)

**MIXING AND APPLICATION**

Apply 30 to 200 gallons of diluted spray per treated acre (3 qts to 4 1 /4 gals per 1 000 sq ft ) depending on density or height of weeds treated and on the type of equipment used

21/26

**RATES AND TIMINGS**

Use the higher level of listed rate ranges when treating dense vegetative growth For best performance apply when weeds are emerged and actively growing

**NOTE** Retreatments may be made as needed however do not exceed a total of 2 pints (1 lb a i ) LIBERTY DICAMBA 4 DMA per treated acre with a maximum of 2 treatments per year

Weed Stage & Type	LIBERTY DICAMBA 4 DMA Herbicide		
	Pints per treated acre	Lbs a i per treated acre	Teaspoons per 1 000 sq ft
<b>Annual</b>			
Small actively growing	1/4 1	1/4 1/2	1 2 1/4
Established weed growth	1 1 1/2	1/2 3/4	2 1/4 3 1/4
<b>Biennial Rosette diameter</b>			
Less than 3 inches	1/2 1	1/4 1/2	1 2 1/4
3 inches or more	1 2	1/2 1	2 1/4 4 1/2
<b>Perennial and Woody</b>			
Brush and Vines	1 2	1/2 1	2 1/4 4 1/2

**TANK MIX TREATMENTS**

READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS DIRECTIONS FOR USE APPLICATION RATES AND TIMINGS AND OTHER RESTRICTIONS

Tank mix treatments of LIBERTY DICAMBA 4 DMA may be made with 2 4 D MCPA MCPP or bromoxynil for control of additional weeds listed on the tank mix product label Apply 1/5 to 1/2 pint (1/10 to 1/4 lb a i ) of LIBERTY DICAMBA 4 DMA per treated acre with 1/2 to 1 1/2 lbs acid equivalent of 2 4 D MCPA or MCPP or with 3/8 to 1/2 lb a i of bromoxynil Use the higher level of the listed rate ranges when treating established weeds Repeat treatments may be made as needed however do not exceed 2 pints (1 lb a i ) of LIBERTY DICAMBA 4 DMA per treated acre during the growing season

**GRASS SEED CROPS**

GRASSES GROWN FOR SEED SUCH AS BERMUDA GRASS BLUEGRASS FESCUE AND RYEGRASS

**IMPORTANT**

Observe all precautions Read and follow mixing and application instructions

Refer to the PASTURE HAY RANGELAND AND FARMSTEAD (NONCROPLAND AREAS) section for possible grazing and feeding restrictions

Do not use on bentgrass unless possible crop injury can be tolerated

**WEEDS CONTROLLED**

LIBERTY DICAMBA 4 DMA will provide control or suppression of annual broadleaf weeds listed below For improved control of listed weeds plus additional weeds it is recommended that LIBERTY DICAMBA 4 DMA be applied in a tank mix with other herbicides

Alfalfa <sup>1</sup>	Clover	Ladysthumb
Bedstraw Catchweed	Cockle White	Lambsquarters Common
Bindweed Field	Dock Broadleaf	Lettuce Prickly
Buttercup Corn	Dock Curly	Mayweed (Dogfennel)
Buttercup Creeping	Hemlock Poison	Ragwort Tansy
Buttercup Western Field	Knapweed Russian <sup>1</sup>	Sorrel Red (Sheep Sorrel)
Catchfly Nightflowering	Knawel	Sowthistle Annual
Chamomile Corn	Kochia	Starwort Little
Chickweed Common	Knotweed Prostrate	Thistle Canada <sup>1</sup>
Chickweed Mouseear		

<sup>1</sup> Top growth control only

**RATES AND TIMINGS**

Apply 1/2 to 1 pint of LIBERTY DICAMBA 4 DMA per treated acre on SEEDLING GRASS after the crop reaches the 3 5 leaf stage Apply up to 2 pints of LIBERTY DICAMBA 4 DMA on well established Perennial grass DO NOT APPLY AFTER THE GRASS SEED CROP BEGINS TO JOINT For best performance make applications when weeds are in the 2 4 leaf stage and rosettes are less than 2 inches across Use the higher level of listed rate ranges when treating more mature weeds or dense vegetative growth

**TANK MIX TREATMENTS**

For control of grasses or additional broadleaf weeds LIBERTY DICAMBA 4 DMA may be tank mixed with all broadleaf herbicides registered for use in Grass Seed Production Read and follow the label of each tank mix product used for precautionary statements directions for use weeds controlled and geographic and other restrictions

**Broadcast Rate Per Treated Acre**

Apply 1/2 to 2 pints LIBERTY DICAMBA 4 DMA with

Product	Active Ingredient	Formulation	Amount of Product Per Acre
2 4 D Amine or Ester	2 4 D	4 lb/gal	1 4 pts ( 5 2 0 lb a i /A ) <sup>1</sup>
MCPA Amine	MCPA	4 lb/gal	1 2 pts ( 5 1 0 lb a i /A ) <sup>1</sup>
bromoxynil	bromoxynil	2 lb/gal	1 2 pts
Curtail®	clopyralid + 2 4 D	2 38 lb/gal	1 3/44 pts
diuron	diuron	80% DF	2 4 lbs
clopyralid	clopyralid	3lb/qal	1/4 1 pt

<sup>1</sup> When using formulations other than 4 lb/gal use pounds active/acre listed

**ANNUAL GRASS CONTROL**

For suppression of ANNUAL GRASS WEEDS such as

- Brome Downy (Cheatgrass)
- Brome Ripgut
- Fescue Rattail
- Windgrass

Apply up to 2 pints (1lb a i ) of LIBERTY DICAMBA 4 DMA per treated acre in the fall or late summer after harvest and burning of established grass seed crops (maximum of 2 treatments per year) Applications should be made immediately following first irrigation when the soil is moist and before weeds have more than 2 leaves

**PREPLANT DIRECTIONS (POST HARVEST/FALLOW/CROP STUBBLE/SET A SIDE)  
FOR BROADLEAF WEED CONTROL BEFORE WHEAT CORN SORGHUM SOYBEANS**

**IMPORTANT**

Observe all precautions Read and follow mixing and application instructions

**WEEDS CONTROLLED**

LIBERTY DICAMBA 4 DMA may be applied alone or in tank mix combinations with other herbicides registered for this use LIBERTY DICAMBA 4 DMA can be applied either POST HARVEST in the fall spring or summer during the FALLOW period or to CROP STUBBLE/ SET A SIDE acres LIBERTY DICAMBA 4 DMA when applied at the specified rates will control many ANNUAL broadleaf weeds see the WEEDS CONTROLLED section under small grains In addition LIBERTY DICAMBA 4 DMA will control or suppress the following BIENNIAL and PERENNIAL broadleaf weeds

Alfalfa <sup>1</sup>	Dock Curly <sup>1</sup>	Sowthistle perennial <sup>1</sup>
Artichoke Jerusalem	Dogbane Hemp	Spurge leafy
Bindweed Field	Garlic Wild <sup>2</sup>	Thistle Bull
Bindweed Hedge	Horsenettle Carolina	Thistle Canada <sup>2</sup>
Blueweed Texas	Knapweed Diffuse	Thistle Milk
Bursage	Knapweed Spotted	Thistle Musk
(Bur Ragweed)	Nightshade Silverleaf	Thistle Plumless
(Povertyweed)	Redvine	Thistle Scotch
(Lakeweed) <sup>1</sup>	Smartweed Swamp	Trumpetcreeper (Buckvine)
Dandelion Common <sup>1</sup>		

<sup>1</sup> Perennials may be controlled using LIBERTY DICAMBA 4 DMA at rates lower than those recommended for other listed perennial weeds (See RATES AND TIMINGS under this heading )

<sup>2</sup> See the SPECIAL TANK MIX TREATMENTS section under this heading for specific control programs for these weeds

**RATES AND TIMINGS**

Apply LIBERTY DICAMBA 4 DMA as a broadcast or spot treatment to emerged and actively growing weeds after crop harvest (post harvest) and before a killing frost or in the fallow cropland or crop stubble the following spring or summer Agriculturally approved spray additives such as surfactants or oils may be used to enhance spray coverage and the herbicides penetration of weed foliage See Cropping restrictions for recommended interval between application and planting to prevent crop injury

For best performance make application when ANNUAL weeds are less than 6 inches tall when BIENNIAL weeds are in the rosette stage and to PERENNIAL weed regrowth in late summer or fall following a mowing or tillage treatment Most effective control of upright perennial broadleaf weeds such as Canada thistle and Jerusalem artichoke occurs if application is made when the majority of weeds such as field bindweed and hedge bindweed are best controlled when weeds are in or beyond the full bloom stage

Avoid disturbing treated areas following application Treatments may not kill weeds which develop from seed or underground plant parts such as rhizomes or bulblets after the effective period for LIBERTY DICAMBA 4 DMA For seedling control a follow up program or other cultural practices could be instituted For small grain in crop uses of LIBERTY DICAMBA 4 DMA see the RATE AND TIMINGS section under the SMALL GRAINS heading for details

**LIBERTY DICAMBA 4 DMA RATES PER TREATED ACRE**

**NOTE** Retreatments may be made as needed however do not exceed a total of 2 pints (1 lb a i ) per treatment of LIBERTY DICAMBA 4 DMA with a maximum of 2 treatments per year

WEED TYPE	AMOUNT OF PRODUCT PER ACRE PER APPLICATION
Annual	1/2 1 pt (8 16 fl oz )
Biennial	1 2 pts (16 32 fl oz )
Perennial	1 4* pts (16 64 fl oz )
Perennial suppression	1 2 pts (16 32 fl oz )
Noted (1) perennials	2pts (32 fl oz )
Other perennials	2 pts (32 fl oz )

**TANK MIX TREATMENTS**

LIBERTY DICAMBA 4 DMA may be tank mixed with one or more of the following herbicides for control of grasses or additional broadleaf weeds Read and follow the label of each tank mix product used for precautionary statements directions for use rates and timings weeds controlled geographic and other restrictions

**LIBERTY DICAMBA 4 DMA BROADCAST RATE PER TREATED ACRE FOR ANNUAL WEED CONTROL**

Apply 1/4 to 1 pint LIBERTY DICAMBA 4 DMA with

Product	Active Ingredient	Formulation	Amount of Product Per Acre
Atrazine <sup>1</sup>	atrazine	4 lb/gal	1 2 6 pts
		90% DF	1/2 3 3 lbs
metsulfuron methyl <sup>2</sup>	metsulfuron methyl	75% DF	0 1 oz
Amber® <sup>2</sup>	triasulfuron	75% DF	0 28 0 35oz
paraquat	paraquat	2 lb/gal	1 2 pts
		2 5 lb/gal	1 5 pts
Finesse® <sup>2</sup>	chlorsulfuron + metsulfuron methyl	75% DF	0 2 oz
pronamide <sup>1</sup>	pronamide	50 W	1/2 1 0 lb
Fallow Master®	glyphosate + dicamba	1 6 lb/gal	22 44 fluid oz
Landmaster® BW	glyphosate + 2 4 D	2 4 lb/gal	27 54 fluid oz
		glyphosate	glyphosate
metribuzin <sup>1</sup>	metribuzin	3 lb/gal	8 48 fluid oz
		75% DF	1/2 1 lb
2 4 D	2 4 D	4 lb/gal	3/4 1 1/2 pts
		4 lb/gal	1 2 pts (0 5 1 lb a i /A) <sup>3</sup>

<sup>1</sup> Tank mixes of LIBERTY DICAMBA 4 DMA with these products may be subject to special restrictions See the Product Label of the tank mix partner for intended use rates restrictions and other precautions

<sup>2</sup> When tank mixing with sulfonylurea herbicides refer to the product label for rates and restrictions Use a surfactant of at least 80% active ingredient at the rate of 1 2 quarts/100 gallons of spray or not more than 0 25 0 5% by volume Use the highest rate of surfactant when using the lower rate ranges of the tank mix and/or when treating more mature weeds or dense vegetative growth Sulfonylurea resistant weeds may not be controlled by tank mixes of LIBERTY DICAMBA 4 DMA and a sulfonylurea Refer to the LIBERTY DICAMBA 4 DMA tank mix section for alternative tank mixes

<sup>3</sup> When using formulations other than 4 lb/gal use pounds active/acre listed

**LIBERTY DICAMBA 4 DMA BROADCAST RATE PER TREATED ACRE FOR BIENNIAL AND PERENNIAL WEED CONTROL**

Apply 1 to 2 pints (0 5 1 0 lb a i ) of LIBERTY DICAMBA 4 DMA with

Product	Active Ingredient	Formulation	Amount of Product Per Acre
Curtail®	clopyralid + 2 4 D	2 38 lb/gal	2 4 pts
		4 lb/gal	2 6 pts (1 0 3 lb a i /A) <sup>1</sup>
Landmaster® BW	glyphosate +2 4 D	2 4 lb/gal	54 fluid oz
glyphosate	glyphosate	3 0 lb/gal	1 5 qts
picloram	picloram	2 lb/gal	1/2 1 pt

<sup>1</sup>When using formulation other than 4 lb/gal use pounds active/acre listed

**SPECIAL TANK MIX TREATMENTS**

For suppression of perennial weeds apply 1/2 1 pint of LIBERTY DICAMBA 4 DMA with 8 16 fluid ounces of glyphosate herbicide per treated acre

For wild garlic control apply 1 pint LIBERTY DICAMBA 4 DMA with 3 pints of 2 4 D LV Ester (4 lb/gal) per treated acre

Apply when wild garlic is 4 to 8 inches tall

For Canada thistle control use LIBERTY DICAMBA 4 DMA or LIBERTY DICAMBA 4 DMA plus Curtail® or LIBERTY DICAMBA 4 DMA plus glyphosate herbicide or glyphosate tank mix treatments Application may be made during fallow periods for control of volunteer barley bulbous bluegrass downy brome jointed goatgrass common rye and volunteer wheat when they are actively growing Use 1 pint LIBERTY DICAMBA 4 DMA with 1/2 1 lb pronamide 50W Fall seeded wheat may be planted 9 months or more after application For best performance make application between mid October and mid December prior to soil freeze up

During fallow periods apply LIBERTY DICAMBA 4 DMA plus Landmaster® BW or Fallow Master® herbicide to give improved control of kochia wild buckwheat prickly lettuce field bindweed and Canada thistle Use 1/8 1/4 pint of LIBERTY DICAMBA 4 DMA plus 22 to 54 fluid ounces of Landmaster® BW or Fallow Master® herbicide for annual weed control or 1/4 to 1/2 pint LIBERTY DICAMBA 4 DMA plus 22 to 54 fluid ounces of Landmaster® BW or Fallow Master® herbicide for perennial weed suppression

**CROPPING RESTRICTIONS**

The following use directions are based on a maximum single application rate of 1 0 lbs ae per acre and a maximum annual rate of 2 0 lbs ae per acre per year

CORN SORGHUM and SOYBEANS may be planted in the spring following applications made during the previous year If less than 1 inch of rainfall occurs between application and first killing frost treated areas should be cultivated to allow herbicide to come in contact with moist soil Cultivation may take place before or immediately after ground thaw

Soybean injury may occur if the interval between application and planting is less than specified In areas with greater than 30 inches of rainfall delay planting for 30 days per pint of LIBERTY DICAMBA 4 DMA per treated acre In areas with less than 30 inches of rainfall delay planting for 45 days per pint of LIBERTY DICAMBA 4 DMA per treated acre Exclude days when ground is frozen

WHEAT may be planted in the fall or spring following applications Also spot applications may be made any time prior to crop emergence if crop injury can be tolerated in treated areas Wheat injury may occur if the interval between application and planting is less than specified

East of the Mississippi River the interval is 20 days per pint of LIBERTY DICAMBA 4 DMA per treated acre or 1 25 days per 1 ounce Moisture is essential for LIBERTY DICAMBA 4 DMA degradation Exclude days when ground is frozen

West of the Mississippi River the interval is 45 days per pint of LIBERTY DICAMBA 4 DMA per treated acre or 3 days per ounce Moisture is essential for LIBERTY DICAMBA 4 DMA degradation Exclude days when ground is frozen

Following a normal harvest of barley oats or wheat any rotation crop may be planted If the interval before harvest is shortened such as when cover crops will be plowed under do not follow up with the planting of a sensitive crop

**CONTROL OF PERENNIAL BROADLEAF WEEDS IN CROPLAND (SPOT APPLICATION ONLY)**

FOR USE ONLY IN THE STATES OF IDAHO MONTANA NEVADA OREGON UTAH AND WASHINGTON

**IMPORTANT**

Observe all precautions Read and follow mixing and application instructions

Do not treat subirrigated cropland or areas where the soil remains saturated with water throughout the year

Make only one application on LIBERTY DICAMBA 4 DMA per year

**WEEDS CONTROLLED**

LIBERTY DICAMBA 4 DMA when applied at specified rates will control many broadleaf weeds including

- |                             |                  |
|-----------------------------|------------------|
| Bindweed Field              | Knapweed Russian |
| Dock Broadleaf (Bitterdock) | Ragwort Tansy    |
| Dock Curly                  | Spurge Leafy     |
| Knapweed Black              | Thistle Canada   |

**RATES AND TIMINGS**

LIBERTY DICAMBA 4 DMA may be applied at any time following a crop harvest to stubble fallow or other cropland

Application should be made when weeds are actively growing and prior to a killing frost

Apply a maximum of 2 pints (1 0 lb a i ) of LIBERTY DICAMBA 4 DMA per treated acre per application with a maximum of 2 applications per year Application may be made up to one month prior to the planting of wheat

NOTE Do not use unless injury to wheat or rotated barley will be acceptable

Barley oats corn sorghum (milo) annual or perennial grass crops may be planted into treated areas one year after application Crops grown for seed (other than perennial grass seed) should not be planted into treated areas until three years after application Do not plant broadleaf crops such as alfalfa beans peas potatoes or sugarbeets into treated areas until two years after application

In most cases treatments will not kill perennial weed seedlings which germinate from seed one or two years after treatment Once the effect of the chemical has been lost a follow up program for seedling control or other cultural practices should be instituted

**WIPER APPLICATION USES IMPORTANT** Observe all precautions

LIBERTY DICAMBA 4 DMA may be applied through wiper application equipment to control or suppress actively growing broadleaf weeds brush and vines Use a solution containing 1 part LIBERTY DICAMBA 4 DMA to 1 part water Do not contact desirable vegetation with herbicide solution Wiper application should only be made to crops (including pastures) and non cropland areas described in this label with the exception of Grain Sorghum (Milo)



**STORAGE AND DISPOSAL**

Do not contaminate water food or feed by storage or disposal *[Optional instructions for bulk storage and disposal Open dumping is prohibited This product may not be mixed loaded or used within 50 feet of all wells including abandoned wells drainage wells and sinkholes*

**AGITATE BEFORE USE]**

**PESTICIDE STORAGE**

Store in original containers in a well-ventilated area separately from fertilizer feed and foodstuffs Avoid cross contamination with other pesticides Spillage or leakage should be contained and absorbed with clay granules sawdust or equivalent material for disposal *[Optional instructions for bulk storage Ground water contamination may be reduced by diking and flooring of permanent liquid storage sites with an impermeable material ]*

**PESTICIDE DISPOSAL**

Triple rinse pesticide from containers and use rinsates in the pesticide application Wastes which cannot be used according to label instructions may be disposed of on site or at an approved waste disposal facility *[Optional instructions for bulk storage and disposal Pesticide spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal and local procedures under Subtitle C of the Resource Conservation and Recovery Act ]*

**CONTAINER DISPOSAL Non-refillable containers Plastic or Metal Do not reuse or refill this container**

Offer for recycling if available or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities such as burning of plastic containers If burned stay out of smoke Triple rinse or pressure rinse container (or equivalent) promptly after emptying Triple rinse as follows

**Non refillable container less than or equal to 5 gallons** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container ¼ full with water and recap Shake for 10 seconds Pour rinsate into formulation equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times

**Non-refillable container greater than 5 gallons** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container ¼ full with water Replace and tighten closures Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds Stand the container on its end and tip back and forth several times Turn the container over onto its other end and tip it back and forth several times Empty the rinsate into formulation equipment or a mix tank or store rinsate for later use or disposal Repeat this procedure two more times

**Pressure rinse as follows (all sizes)** Empty the remaining contents into formulation equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip Hold container upside down over formulation equipment or mix tank or collect rinsate for later use or disposal Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds Drain for 10 seconds after the flow begins to drip

**Refillable container (250 gallon & bulk)** Refill this container with pesticide only Do not reuse this container for any other purpose Empty containers retain vapor and product residue Observe all labeled safeguards until container is cleaned reconditioned or destroyed Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller

To clean the container before final disposal empty the remaining contents from the container into formulation equipment or mix tank Fill the container about 10 percent full with water Agitate vigorously or recirculate water with pump for 2 minutes Pour or pump rinsate into formulation equipment or rinsate collection system Repeat this rinsing process two more times

**CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product If the terms are not acceptable return the product at once unopened and the purchase price will be refunded The Directions for Use of this product must be followed carefully It is impossible to eliminate all risks inherently associated with the use of this product Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application weather presence of other materials or other influencing factors in the use of the product which are beyond the control of LIBERTY CROP PROTECTION LLC or Seller To the extent consistent with applicable law all such risks shall be assumed by Buyer and User and Buyer and User agree to hold LIBERTY CROP PROTECTION LLC and Seller harmless for any claims relating to such factors LIBERTY CROP PROTECTION LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use subject to the inherent risks referred to above when used in accordance with directions under normal use conditions This warranty does not extend to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or LIBERTY CROP PROTECTION LLC and Buyer and User assume the risk of any such use TO THE EXTENT CONSISTENT WITH APPLICABLE LAW LIBERTY CROP PROTECTION LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE

To the extent consistent with applicable law neither LIBERTY CROP PROTECTION LLC nor Seller shall be liable for any incidental

26/26

consequential or special damages resulting from the use or handling of this product **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF LIBERTY CROP PROTECTION LLC AND SELLER FOR ANY AND ALL CLAIMS LOSSES INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY CONTRACT NEGLIGENCE TORT STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR AT THE ELECTION OF LIBERTY CROP PROTECTION LLC OR SELLER THE REPLACEMENT OF THE PRODUCT** LIBERTY CROP PROTECTION LLC and Seller offer this product and Buyer and User accept it subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of LIBERTY CROP PROTECTION LLC

**REGISTERED TRADEMARKS**

Amber Beacon and Tough are registered trademarks of Syngenta  
Accent Express Finesse Glean and Harmony are registered trademarks of E I duPont de Nemours & Co Inc  
Bronate Dakota and Tiller are registered trademarks of Bayer Cropscience  
Battalion Bronco Bullet Harness Landmaster Lariat Lasso Partner Permit Ramrod and Screen are registered trademarks of Monsanto Company  
Broadstrike and Curtail are registered trademarks of Dow AgroSciences  
Clarity Fallow Master and Frontier are registered trademarks of BASF Corporation