



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

89168-71

Date of Issuance:

11/6/17

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Liberty Clop

Name and Address of Registrant (include ZIP Code):

MaryBeth Endres
Liberty Crop Protection, LLC
4850 Hahns Peak Dr., Suite 200
Loveland, CO 80538

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(C). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Kathryn Montague, Product Manager 23
Herbicide Branch, Registration Division (7505P)

Date:

11/6/17

2. You are required to comply with the data requirements described in the DCI or EDSP Order identified below:

- a. Clopyralid GDCI-117403-1454

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI or EDSP Order listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division:

<http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 89168-71.”
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 09/13/2017

If you have any questions, please contact Helen Hull-Sanders by phone at 703-347-0243, or via email at hull-sanders.helen@epa.gov.

Enclosure

ACCEPTED

11/6/2017

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 89168-71

LIBERTY CLOP

CLOPYRALID GROUP **4** HERBICIDE

For Selective Postemergence Control of Broadleaf Weeds in Apple, Asparagus, Barley, Oats and Wheat Not Underseeded with Legume, *Brassica*, Canola (Rapeseed) and Crambe, Christmas Tree Plantations, Corn (Field, Pop, Sweet), Cottonwood/Poplar and Eucalyptus Tree Plantations, Fallow Cropland, Garden Beet, Grasses Grown For Seed, Peppermint and Spearmint, Southern Pine Seedbeds in Forest Nurseries, Spinach, Stone Fruits, Sugar Beet, Turnip, Rangeland, Permanent Grass Pastures, Conservation Reserve Program (CRP) Acres, and Non-Cropland (Including Fencerows, Around Farm Buildings and Equipment Pathways)

ACTIVE INGREDIENT:	% BY WT.
Clopyralid: 3,6-dichloro-2- pyridinecarboxylic acid, monoethanolamine salt	40.9%
OTHER INGREDIENTS:	<u>59.1%</u>
TOTAL:	100.0%

Acid Equivalent: clopyralid: 3,6-dichloro-2-pyridinecarboxylic acid – 31% (3 lb/gal)

KEEP OUT OF REACH OF CHILDREN
CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you don't understand the label, find someone to explain it to you in detail).

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS.

EPA Reg. No.: 89168-TR

EPA Est. No.: _____

NET CONTENTS: ____ GAL (____ L)

Manufactured for:
LIBERTY CROP PROTECTION, LLC
4850 Hahns Peak Drive, Suite 200
Loveland, CO 80538

110517

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 or your poison control center at 1-800-222-1222. For Chemical Spill, Leak, Fire or Exposure, call CHEMTREC 800-424-9300.</p>	

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS
<p>Users should:</p> <ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Do not contaminate water used for irrigation or domestic purposes.

Ground Water Advisory

Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate groundwater which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an underlying aquifer is shallow, or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a medium potential for reaching surface water via runoff for several weeks after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of clopyralid from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Not for Sale, Sale Into, Distribution and/or Use in Nassau and Suffolk Counties of New York State.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: For applications to fallow cropland, rangeland, pasture, and non-crop areas, do not enter treated areas until sprays have dried. For early entry to treated areas, wear eye protection, chemical-resistant gloves made of any waterproof material, long-sleeved shirt, long pants, shoes and socks.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, this product a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

Weed Management

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 4 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in the field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will

control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.

- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact LIBERTY CROP PROTECTION, LLC at [855-466-8428 or 844-425-8488 or other appropriate telephone number].

Management of Resistant Biotypes

Since the occurrence of resistant weeds cannot be determined until after product use and scientific confirmation, manufacturer is not responsible for any losses that may result from the failure of this product to control resistant weed biotypes.

The following good agronomic practices are recommended to reduce the spread of resistant biotypes:

- If a naturally occurring resistant biotype is present in your application site, this product should be tank mixed or applied sequentially with an appropriately labeled herbicide with a different mode of action to achieve control.
- Cultural and mechanical control practices (e.g. crop rotation or tillage) may also be used as appropriate.
- Scout treated application site after herbicide applications and control escaping weeds including resistant biotypes before they set seed.
- Thoroughly clean equipment before leaving fields known to contain resistant biotypes.
- Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to this Mode of Actions have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of actions for each target weed.

Integrated Pest (Weed) Management

This product may be integrated into an overall weed pest management strategy whenever the use of an herbicide is required. Practices known to reduce weed development (tillage, crop competition) and herbicide use (weed scouting, proper application timing, banding) should be followed wherever possible. Consult local agricultural and weed authorities for additional IPM strategies established for your area.

PRODUCT INFORMATION

LIBERTY CLOP is a selective, postemergence herbicide for control of broadleaf weeds in apple, asparagus, barley, oats and wheat not underseeded with a legume, canola (rapeseed), Christmas tree plantations, conservation reserve program (CRP) acres, cottonwood/poplar and eucalyptus tree plantations, crambe, fallow cropland, field corn, garden beet, grasses grown for seed, *Brassica*, peppermint, popcorn, rangeland and permanent grass pastures, southern pine seedbeds in forest

nurseries, spearmint, spinach, stone fruits, sugar beet, sweet corn, turnip, and non-cropland areas including fence rows, around farm buildings, and equipment pathways.

LIBERTY CLOP may be applied by aircraft on the following crops only: canola (rapeseed), crambe, spinach and sugar beet.

Re-treatment is allowed, but do not apply more than the maximum allowable rate per crop growing season. An application to fallow cropland preceding or following an application to dryland small grains (wheat, barley or oats) is allowed, but is not allowed preceding or following an application to irrigated small grains.

RESTRICTIONS

- Not for Sale, Sale into, Distribution and/or Use in Nassau and Suffolk Counties of New York State.
- In California and New York, the maximum application rate for LIBERTY CLOP is 2/3 pint per acre per growing season. Do not exceed a cumulative amount of 2/3 pint of clopyralid (0.25 lb acid equivalent) per acre per crop year, unless specifically allowed.
- Do not contaminate irrigation ditches or water used for irrigation or domestic purposes.
- Do not use in greenhouses.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- Do not spray pastures containing desirable forbs, especially legumes, unless injury can be tolerated.
- Do not apply this product by aircraft to other labeled crops unless otherwise permitted by supplemental labeling.
- Do not transfer livestock from treated grazing areas (or feeding of treated hay) to sensitive broadleaf crop areas without first allowing 7 days of grazing on an untreated pasture (or feeding of treated hay). If livestock are transferred within less than 7 days of grazing untreated pasture or eating untreated hay, urine and manure may contain enough clopyralid to cause injury to sensitive broadleaf plants.

Field Bioassay Instructions: In fields previously treated with this product, plant short test rows of the intended rotational crop across the original direction of application in a manner to sample field conditions, such as soil texture, soil pH, drainage, and any other variable that could affect the seed bed of the new crop. Field bioassay at any time prior to the planting of the intended rotational crop. Observe the test crop for herbicidal activity, such as poor stand (effect on seed germination) chlorosis (yellowing), necrosis (dead leaves or shoots), or stunting (reduced growth). If herbicidal symptoms do not occur, the test crop can be grown. If there is apparent herbicidal activity, wait one year before repeating bioassay or plant only a labeled crop or crop listed in the table below for which the rotational interval has clearly been met.

Crop Rotation intervals

Residues of this product in treated plant tissues, including the treated crop or weeds, which have not completely decayed may affect succeeding susceptible crops.

Crop Rotation Intervals for Florida Only

Rotation Crops¹	Rotation Interval⁴ (Soils less than 2% organic matter AND rainfall greater than 15 inches during 12 months following application)
Barley, Canola (Rapeseed), Cole Crops (includes <i>Brassica</i> species grown for seed), Field Corn, Flax, Garden Beet, Grasses, Oats, Popcorn, Spinach, Sugar Beet, Sweet Corn, Turnip, Wheat	Anytime
Alfalfa, Asparagus, Grain Sorghum, Onions, Peppermint, Safflower, Spearmint, Strawberry	10.5 months
Dry Beans, Soybean, Sunflower	18 months ²
Lentils, Peas, Potatoes (including Potatoes Grown for Seed), and Broadleaf Crops grown for seed (excluding <i>Brassica</i> species)	18 months ^{2, 3}

- 1 For best results, conduct a field bioassay prior to planting any broadleaf crops that are not listed. Do not rotate to unlisted crops prior to 10.5 months following application.
- 2 Follow an 18-month crop rotation due to the potential for crop injury unless previous experience has shown no crop injury with the minimum 10.5-month rotation interval. **Precaution:** For these crops, a minimum 10.5-month rotation interval must be observed to avoid illegal residues in the harvested crop.
- 3 For best results, conduct a field bioassay prior to planting these sensitive crops.
- 4 **Precaution:** The above intervals are based upon average annual precipitation regardless of irrigation practices. Observance of listed crop rotation intervals should result in adequate safety to rotational crops. However, this product is dissipated in the soil by microbial activity and the rate of microbial activity is dependent upon several interrelating factors including soil moisture, temperature and organic matter. Therefore, accurate prediction of rotational crop safety is not possible. In areas of low organic matter (<2%) and less than 15 inches average annual precipitation, potential for crop injury may be reduced by burning or removal of plant residues, supplemental fall irrigation and deep moldboard plowing prior to planting the sensitive crop.

Crop Rotation Intervals for All States Except California, Florida, Idaho, Nevada, Oregon, Utah and Washington

Rotation Crops¹	Rotation Interval⁴ (Soils greater than 2% organic matter AND rainfall more than 15 inches during 12 months following application)	Rotation Interval⁴ (Soils less than 2% organic matter AND rainfall less than 15 inches during 12 months following application)
Barley, Canola (Rapeseed), Cole Crops (includes <i>Brassica</i> species grown for seed), Field Corn, Flax, Garden Beet, Grasses, Oats, Popcorn, Spinach, Sugar Beet, Sweet Corn, Turnip, Wheat	anytime	anytime
Alfalfa, Asparagus, Grain Sorghum, Onions, Peppermint, Safflower, Spearmint, Strawberry	10.5 months	10.5 months
Dry Beans, Soybean, Sunflower	10.5 months	18 months ²
Lentils, Peas, Potatoes (including Potatoes grown for seed), and Broadleaf crops grown for seed (excluding <i>Brassica</i> species)	18 months ²	18 months ^{2, 3}

- 1 For best results, conduct a field bioassay prior to planting any broadleaf crops that are not listed. Do not rotate to unlisted crops prior to 10.5 months following application.
- 2 Follow an 18-month crop rotation due to the potential for crop injury unless previous experience has shown no crop injury with the minimum 10.5-month rotation interval. **Precaution:** For these crops, a minimum 10.5-month rotation interval must be observed to avoid illegal residues in the harvested crop.
- 3 For best results, conduct a field bioassay prior to planting these sensitive crops.
- 4 **Precaution:** The above intervals are based upon average annual precipitation regardless of irrigation practices. Observance of listed crop rotation intervals should result in adequate safety to rotational crops. However, this product is dissipated in the soil by microbial activity and the rate of microbial activity is dependent upon several interrelating factors including soil moisture, temperature and organic matter. Therefore, accurate prediction of rotational crop safety is not possible. In areas of low organic matter (<2%) and less than 15 inches average annual precipitation, potential for crop injury may be reduced by burning or removal of plant residues, supplemental fall irrigation and deep moldboard plowing prior to planting the sensitive crop.

Crop Rotation Intervals for California, Idaho, Nevada, Oregon, Utah and Washington Only

Rotation Crops ¹	Rotation Interval ⁴ (Areas receiving greater than 18 inches of rainfall – not including irrigation)	Rotation Interval ⁴ (Areas receiving less than 18 inches of rainfall – not including irrigation)
Barley, Canola (Rapeseed), Cole Crops (includes <i>Brassica</i> species grown for seed), Field Corn, Flax, Garden Beet, Grasses, Oats, Popcorn, Spinach, Sugar Beet, Sweet Corn, Turnip, Wheat	anytime	anytime
Asparagus, Grain Sorghum, Onions, Peppermint, Safflower, Spearmint, Strawberry	12 months	12 months
Alfalfa, Dry Beans, Soybean, Sunflower	12 months	18 months ^{2, 3}
Broadleaf Crops grown for seed (excluding <i>Brassica</i> species), Carrot ² , Celery ² , Cotton ² , Lentils, Lettuce ² , Melons ² , Peas, Potatoes (including Potatoes grown for seed), Safflower and Tomato ²	18 months ²	18 months ^{2, 3}

1 For best results, conduct a field bioassay prior to planting any broadleaf crops that are not listed. Do not rotate to unlisted crops prior to 12 months following application.

2 Follow an 18-month crop rotation due to the potential for crop injury unless previous experience has shown no crop injury with the minimum 12-month rotation interval. **Precaution:** For these crops, a minimum 12-month rotation interval must be observed to avoid illegal residues in the harvested crop.

3 Crop injury and/or yield loss may occur up to 4 years after application. For best results, conduct a field bioassay prior to planting these sensitive crops. See instructions above.

4 **Precaution:** The above intervals are based upon average annual precipitation regardless of irrigation practices. Observance of listed crop rotation intervals should result in adequate safety to rotational crops. However, this product is dissipated in the soil by microbial activity and the rate of microbial activity is dependent upon several interrelating factors including soil moisture, temperature and organic matter. Therefore, accurate prediction of rotational crop safety is not possible. In areas of low organic matter (<2%) and less than 15 inches average annual precipitation, potential for crop injury may be reduced by burning or removal of plant residues, supplemental fall irrigation and deep moldboard plowing prior to planting the sensitive crop.

Avoid Injury to Non-Target Plants

This product can affect susceptible broadleaf plants directly through foliage and indirectly by root uptake from treated soil. Therefore, do not apply LIBERTY CLOP directly to, or allow spray drift to come in contact with, vegetables, flowers, tomatoes, potatoes, beans, lentils, peas, alfalfa, sunflowers, soybeans, safflower, or other desirable broadleaf crops or ornamental plants or soil where sensitive crops will be planted the same season. (See Crop Rotation Intervals.)

Residues in Plants or Manure: Do not use plant residues, including hay or straw from treated areas, or manure or bedding straw from animals that have grazed or consumed forage from treated areas, for composting or mulching where susceptible plants may be grown the following season. Do not spread manure from animals that have grazed or consumed forage or hay from treated areas on land used for growing susceptible broadleaf plants or apply such materials to land used for growing broadleaf crops, ornamentals, orchards, or other susceptible desirable plants. Plant materials or manure may contain enough clopyralid to cause injury to susceptible plant species. To promote herbicidal decomposition, plant residues should be evenly incorporated or burned. Breakdown of clopyralid in crop residues or manure is more rapid under warm, moist soil conditions and may be enhanced by supplemental irrigation.

Avoid Movement of Treated Soil

Avoid conditions under which soil from treated areas may be moved or blown to areas containing susceptible plants. Wind-blown dust containing clopyralid may produce visible symptoms, such as epinasty (downward curving or twisting of leaf petioles or stems), when deposited on susceptible plants; however, serious injury is unlikely. To minimize potential movement of clopyralid on wind-blown dust, avoid treatment of powdery dry or light sandy soils until soil is settled by rainfall or irrigation or irrigate the treated soil shortly after application.

Avoiding Spray Drift

Avoid spray drift since very small quantities of the spray, which may not be visible, may severely injure susceptible crops during active growth or dormant periods. Use coarse spray to minimize drift. To aid in further reducing drift, a drift control or deposition agent suitable for agricultural use may be used with this product. If used, follow all use instructions, restrictions and precautions on the product label.

Ground Application: With ground equipment, minimize spray drift by keeping the spray boom as low as possible, by applying 10 gallons or more of spray per acre, by keeping the operating spray pressure at the manufacturer's minimum specified pressures for the specified nozzle type used (low pressure nozzles are available from spray equipment manufacturers) and by spraying when the wind velocity is low (follow state regulations. Avoid application under completely calm conditions which may be conducive to air inversion. In hand-gun applications, select the minimum pressure required to obtain adequate plant coverage without forming a mist. Do not apply with a mist blower.

Aerial Application: Drift can be lessened by using straight stream nozzles directed straight back; by using drift control systems or use of drift control additives; and by keeping spray pressures low enough to provide coarse spray droplets. Do not use a thickening agent with the Microfoil or Thru-Value booms, or other systems that cannot accommodate thick sprays. Spray only when wind velocity is low (follow state regulations).

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses, or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward, parallel with the air stream and never be pointed downward more than 45 degrees.
3. Where states have more stringent regulations, they must be observed.

The applicator should be familiar with and take into account the information covered in the "Aerial Drift Reduction Advisory Information" section below.

Aerial Drift Reduction Advisory Information Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly or under unfavorable environmental conditions (See Wind, Temperature and Humidity, and Temperature Inversions sections of this label).

Controlling Droplet Size – Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

- **Pressure** – Do not exceed the nozzle manufacturer’s recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released backwards, parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

Boom Length

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height

Applications should not be made at a height greater than 10 feet above the top of the largest plants, unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

When making applications in low relative humidity, set up the equipment to produce larger droplets to reduce effects of evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Application should not occur during a temperature inversion because of potential drift. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. This cloud can move in unpredictable directions due to the light, variable winds common during inversions. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

This pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., non-target crops, bodies of water, residential areas, known habitat for threatened or endangered species) is minimal (e.g., when wind is blowing away from the sensitive areas).

Sprayer Clean-Out

To avoid injury to desirable plants, thoroughly clean equipment used to apply LIBERTY CLOP before re-using it to apply any other chemicals.

- Rinse and flush application equipment thoroughly at least three times with water after use. Dispose of rinse water by applying to treatment area or to non-cropland area away from water supplies.
- During the second rinse, add 1 quart of household ammonia for every 25 gallons of water. Circulate the solution through the entire system so that all internal surfaces are contacted (15 to 20 minutes). Let the solution stand for several hours, preferably overnight.
- Flush the solution out of the spray tank through the boom.
- Rinse the system twice with clean water, recirculating and draining each time.

- Remove nozzles and screens and clean separately.

Mixing Directions

LIBERTY CLOP - Alone

1. Add 3/4 of the required spray volume to the spray tank and start agitation.
2. Add the required amount of LIBERTY CLOP.
3. Add any surfactants, adjuvants or drift control agents according to manufacturer's label.
4. Agitate during final filling of the spray tank and maintain sufficient agitation during application to ensure uniformity of the spray mixture.

Precaution: Allow time for thorough mixing of each spray ingredient before adding the next. If allowed to stand after mixing, agitate spray mixture before use.

LIBERTY CLOP - Tank Mix

This product may be applied in tank mix combination with labeled rates of other products provided (1) the tank mix product is labeled for the timing and method of application for the use site to be treated; and (2) tank mixing is not prohibited by the label of the tank mix product. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Tank Mixing Precautions:

- Always perform a jar test to ensure the compatibility of products to be used in tank mixture.

Tank Mixing Restrictions:

- Do not exceed specified application rates.
- Do not tank mix with another pesticide product that contains the same active ingredient as this product unless the label of either tank mix partner specifies the maximum dosages that may be used.
- For products packaged in water soluble packaging, do not tank mix with products containing boron or mix in equipment previously used to apply a product mixture containing boron unless the tank and spray equipment have been adequately cleaned. (See Sprayer Clean-Out.)

Tank Mix Compatibility Testing: A jar test is recommended prior to tank mixing to ensure compatibility of LIBERTY CLOP and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in the required order and their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, gels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

APPLICATION DIRECTIONS

Application Timing

Apply to actively growing weeds. Extreme growing conditions, such as drought or near freezing temperatures prior to, at, or following application, may reduce weed control and increase the risk of crop injury at all stages of growth. Only weeds that have emerged at the time of application will be affected. If foliage is wet at the time of application, control may be decreased. Applications of LIBERTY CLOP are rainfast within 6 hours after application.

Application Rates

Generally, application rates at the lower end of the rate range will be satisfactory for young, succulent growth of susceptible weed species. For less sensitive species, perennials, and under conditions where control is more difficult (plant stress conditions, such as, drought or extreme temperatures, dense weed stands and/or larger weeds), use a higher rate within the rate range. Weeds in fallow land or other areas where competition from crops is not present will generally require higher rates for control or suppression.

Crop or Use Site	Rate Range (pt/acre)	Maximum Use Rate¹ (pt/acre /growing season)
Spinach	1/6 - 1/3 (0.062 – 0.125 lb a.e./acre)	1/2 (0.187 lb a.e./acre)
Barley, Oats, Wheat	1/4 - 1/3 (0.094 – 0.125 lb a.e./acre)	1/3 (0.125 lb a.e./acre)
Christmas Tree and Cottonwood/Poplar and Eucalyptus Tree Plantations, Fallow Cropland, Field Corn, Grasses grown for seed, Sugar Beet	1/4 - 2/3 (0.094 – 0.25 lb a.e./acre)	2/3 (0.25 lb a.e./acre)
Canola (Rapeseed), Cole Crops (<i>Brassica</i> species), Crambe, Garden Beet, Southern Pine Seedbeds	1/4-1/2 (0.094 – 0.187 lb a.e./acre)	1/2 (0.187 lb a.e./acre)
Apple, Popcorn, Stone Fruits, Sweet Corn	1/3 - 2/3 (0.0.125 – 0.25 lb a.e./acre)	2/3 (0.0.25 lb a.e./acre)
Turnip	1/3 - 1/2 (0.125 – 0.187 lb a.e./acre)	1/2 (0.187 lb a.e./acre)
Peppermint, Spearmint	1/3 – 1 (0.0.125 – 0.375 lb a.e./acre)	1 (0.375lb a.e./acre)
Noncropland, Non-Leguminous Trees, Permanent Grasses on CRP Land, Rangeland and Permanent Grass Pastures	1/3 – 1 1/3 (0.125 – 0.50 lb a.e./acre)	1 1/3 (0.50 lb a.e./acre)
Asparagus	1/2 - 2/3 (0.187 – 0.25 lb a.e./acre)	2/3 (0.25 lb a.e./acre)

¹ Do not exceed maximum rate in rate range per growing season,

Spot Treatments

To prevent misapplication, apply spot treatments only with a calibrated boom or with hand sprayers according to directions provided below,

Hand Held Sprayers: Hand held sprayers may be used for spot applications, Care should be taken to apply the spray uniformly and at a rate equivalent to a broadcast application. Application rates in the table are based upon an area of 1000 sq ft. Mix the amount of LIBERTY CLOP (fl oz or ml) corresponding to the desired broadcast rate in 1 gallon or more of spray. To calculate the amount of LIBERTY CLOP required for larger areas, multiply the table value (fl oz or ml) by the area to be treated in "thousands" of square feet, e.g., if the area to be treated is 3500 sq ft, multiply the table value by 3.5 (calc. 3500 ÷ 1000 = 3.5). An area of 1000 sq ft is approximately 10.5 x 10.5 yards (strides) in size.

Amount of Liberty Clop per Gallon of Spray to Equal Specified Broadcast Rate					
1/4 pt/acre	1/3 pt/acre	1/2 pt/acre	2/3 pt/acre	1 pt/acre	1 1/3 pt/acre
1/10 fl oz (2.7 ml)	1/8 fl oz (3.6 ml)	1/5 fl oz (5.4 ml)	1/4 fl oz (7.3 ml)	3/8 fl oz (11 ml)	0.5 fl oz (15 ml)

Use the following table for converting pints to fluid ounces.

Conversion Chart – Pints of Fluid Ounces	
Pints	Fluid Ounces
1/3	5
1/4	4
1/2	8
2/3	11

Band Application

LIBERTY CLOP may be applied as a band treatment. Use the formulas below to determine the appropriate rate and volume per treated acre.

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast rate per treated acre} = \text{Band rate per treated acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast volume per treated acre} = \text{Band volume per treated acre}$$

Use of Adjuvants

Addition of surfactants, crop oils, or other adjuvants is not usually necessary when using LIBERTY CLOP. Adding a surfactant to the spray mixture may increase effectiveness on weeds but may reduce selectivity to the crop, particularly under conditions of plant stress. If an adjuvant is added to the spray solution, follow all manufacturer use guidelines.

Spray Coverage

Use sufficient spray volume to provide thorough coverage and a uniform spray pattern. Do not broadcast apply in less than 2 gallons total spray volume per acre. For best results, and to minimize spray drift, apply in a spray volume of 10 gallons or more per acre. As vegetative canopy and weed density increase, increase spray volume to obtain equivalent weed control. Use only nozzle types and spray equipment designed for herbicide application. To reduce spray drift, follow precautions under Avoid Injury to Non-Target Plants.

Broadleaf Weeds Controlled and Guidelines for Control¹

Note: Letter in parentheses (-) after listed weed indicates if life cycle is annual (a), biennial (b), or perennial (p).

Weed Species	Stage of Growth	Rate for Control ² (pt/acre)
Biennial wormwood (a, b) ³	Up to 5 leaf	1/4 – 1/2
Black medic clover (a)		
Bull thistle (b)		
Clover (a)		
Cocklebur (a)		
Coffeeweed (a)		
Common burdock (b)		
Common cocklebur (a)		
Common groundsel (b)		
Common ragweed (a)		
Common teasel (b)		
Cornflower (bachelor button) (a)		
Curly dock (p)		
Dandelion (p)		
False chamomile (scentless) (a)		
Galinsoga (a)		
Giant ragweed (a)		
Hop clover (a)		
Horseweed (a)		
Jerusalem artichoke (p)		
Jimsonweed (a)		
Ladysthumb (a) ⁴		
Lambert locoweed (p)		
Marshelder (a)		
Mayweed chamomile (dogfennel) (a)		
Meadow salsify (goatsbeard) (b)		

Weed Species	Stage of Growth	Rate for Control² (pt/acre)
Musk thistle (b)		
Narrowleaf hawksbeard (a)		
Orange hawkweed (p)		
Oxeye daisy (p)		
Pineappleweed (a)		
Prickly lettuce (a)		
Ragweeds (a)		
Red clover (p)		
Red sorrel (p)		
Sicklepod (a)		
Sunflower (a)		
Sweet clover (b)		
Vetch (a)		
Volunteer alfalfa (p) (from seed only)		
Volunteer beans (a)		
Volunteer lentils (a)		
Volunteer peas (a)		
Volunteer soybean (a)		
White clover (p)		
White locoweed (p)		
Yellow hawkweed (p)		
Yellow starthistle (a)		
Wild buckwheat (a)	1-3 leaf stage, but before vining	1/2
Black nightshade (a)	2 – 4 leaf stage	
Buffalobur (a) ³		
Cutleaf nightshade (a)		
Eastern black nightshade (a)		
Hairy nightshade (a)		
Nightshade spp. (a)		
Green smartweed (a) ⁴	2 -3 leaf	
Smartweeds (suppression)		
Annual sowthistle (a) (suppression)	rosette up to bud stage	Degree of infestation: Light – 1/3 Moderate to heavy – 1/2 - 2/3
Canada thistle (p)		
Perennial sowthistle (p) ⁴		
Sowthistle (a) (suppression)		
Spotted/diffuse knapweeds (b)	up to bud stage	1/2 – 2/3
Russian knapweed (p) ⁴		2/3 – 1 1/3

1 This table is provided as a general reference only. Refer to use directions for specific crop or use site for application rates.

2 Where a rate range is provided, use a lower rate in the rate range for light to moderate infestations under good growing conditions and a higher rate in the rate range for dense infestations or under less favorable growing conditions such as drought.

3 Not registered for use in California.

4 These weeds may only be suppressed. Suppression is a visual reduction in weed competition (reduced population or vigor) as compared to untreated areas. The degree and duration of weed control will vary with weed size and density, application rate and coverage, and growing conditions before, during, and after treatment. For perennial weeds, this product will control the top growth and inhibit regrowth during the season of application (season-long control). At higher use rates shown on this label, this product may cause a reduction in shoot regrowth in the season following application; however, plant response may be inconsistent due to inherent variability in shoot regrowth from perennial root systems.

CROP USES

Agricultural Use Requirements for Crops: For the following crop uses, follow PPE and Reentry instructions in the Agricultural Use Requirements section of this label.

APPLE

(Not for Sale, Sale Into, Distribution and/or Use in Nassau and Suffolk Counties of New York State)

Use LIBERTY CLOP for postemergence control of broadleaf weeds listed below that are infesting apples.

Target Broadleaf Weeds	Application Rate (pt/acre)
Aster	1/3 – 2/3
Burdock	
Clover, red	
Clover, white	
Curly dock	
Dandelion	
Goldenrod	
Horseweed (marestail)	
Nightshade, black	
Nightshade, hairy	
Pineappleweed	
Sowthistle, annual	
Thistle, Canada	
Thistle, musk	
Vetch	
Volunteer alfalfa	

Application Timing

Apply LIBERTY CLOP to clover and vetch from weed emergence up to the 5-leaf stage of growth. Apply LIBERTY CLOP to nightshade (black and hairy) at the 2- to 4-leaf stage of growth. For control of Canada thistle and annual sowthistle, apply this product from rosette up to bud stage.

Application Rate

Apply 1/3 to 2/3 pint of LIBERTY CLOP per acre with ground equipment in 10 gallons or more total spray volume per acre. Use a higher rate in the rate range for heavy infestations or when greater residual control is desired.

Tank Mixtures: LIBERTY CLOP may be tank mixed with other herbicides labeled for use on apple. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Restrictions

- **Preharvest Interval:** Do not apply within 30 days of harvest.
- Make one to two broadcast applications per crop year, not to exceed a total of 2/3 pint per acre.
- Apply this product to non-bearing (well established trees, 1 year or older) and bearing trees.
- East of the Rocky Mountains, do not apply this product during bloom.
- Avoid direct contact with foliage, fruit or tree trunks.

ASPARAGUS
(Not Registered for Use in Florida)

Use LIBERTY CLOP for selective postemergence control of specific annual and perennial broadleaf weeds infesting asparagus.

Application Timing

Apply before or during the asparagus cutting season, or after harvest is complete, but prior to fern growth. Treat annual weeds before they send up a flower stalk. For best results on perennial weeds, such as Canada thistle, apply this product after the majority of basal leaves have emerged up to bud stage. Following application, wait at least two weeks before cultivating.

Application Rate

Apply LIBERTY CLOP at a rate of 1/2 to 2/3 pint per acre in a total spray volume of 10 to 40 gallons per acre. Use a higher rate in the rate range for more effective control of perennial weeds. A second application may be made as long as the total amount applied does not exceed 2/3 pint of this product per acre during the growing season.

Precautions

- Make postharvest (layby) applications as soon as possible after cutting provided weeds are in the proper stage of growth for treatment. Malformed ferns may result from application when spears are longer than 3 inches or have open seed heads.
- When this product is applied during the cutting season, some crooking (twisting) of asparagus spears may occur. Do not apply during the cutting season if crooking cannot be tolerated. Clear-cutting of spears just before applying this product may reduce the occurrence of crooking.

Restrictions

- **Preharvest Interval:** Do not apply within 48 hours of harvest.
- Make one to two applications per crop year, not to exceed a total of 2/3 pint per acre per growing season.

BARLEY, OATS AND WHEAT NOT UNDERSEEDED WITH LEGUME
(Not Registered for Use in Florida)

Application Rate

Apply 1/4 to 1/3 pint of LIBERTY CLOP per acre when crop is from the 3-leaf stage up to early boot stage of growth. For control of perennial weeds, such as Canada thistle, apply 1/3 pint of LIBERTY CLOP per acre. Russian knapweed will only be suppressed at this rate.

Restrictions

- Do not permit lactating dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 1 week after treatment.
- Do not harvest hay from treated grain fields.
- Do not apply more than 1/3 pint per acre per growing season.

BRASSICA (COLE) LEAFY VEGETABLES (CROP GROUP 5)¹

**For use and distribution only in the states of Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Maine, Maryland, Massachusetts, Michigan, Missouri, New Hampshire, New Jersey, New Mexico, New York, North Carolina, Ohio, Oklahoma, Pennsylvania, Rhode Island, South Carolina, Texas, Vermont, Virginia, West Virginia, and Wisconsin
(Not for Sale, Sale Into, Distribution and/or Use in Nassau and Suffolk Counties of New York State)**

Brassica (cole) leafy vegetables (crop group 5) including broccoli, broccoli raab (rapini), Brussels sprouts, cabbage, cauliflower, cavalo broccolo, Chinese broccoli (gai ion), Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens

Target Broadleaf Weeds	LIBERTY CLOP (pt./acre)
Chamomile	1/4 – 1/2
Clover	
Common cocklebur	
Dandelion	
Galinsoga	
Pineappleweed	
Prickly lettuce	
Ragweed	
Smartweed	
Wild buckwheat	
Annual sowthistle ¹	
Canada thistle ¹	

¹ Suppression only

Broadcast Application Rates

Apply uniformly with ground equipment in a minimum of 10 to 40 gallons of water per acre. For suppression of Canada thistle, apply after the majority of basal leaves have emerged, but prior to bud stage, and at least 30 days prior to harvest.

Tank Mixes

This product may be tank mixed with other herbicides labeled for use on mustard greens. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. See Tank Mixing section under Mixing Directions.

Restrictions

- **Preharvest Interval:** Do not apply within 30 days of harvest.
- Make one to two broadcast applications per crop per year, not to exceed a total of 1/2 pint per acre per year.
- **In New York and California,** the maximum application rate is 2/3 pint per acre per growing season. Do not exceed the cumulative amount of 2/3 pint per acre per crop year.
- **In Florida,** this product may be used only on cabbage, Chinese cabbage (napa) and Chinese mustard cabbage (gai choy).

CANOLA (RAPESEED) AND CRAMBE (Not Registered for Use in California and Florida)

Application Timing

Apply to canola or crambe in the 2- to 6-leaf stage of crop growth at rates shown in the following table. Consult the table entitled Broadleaf Weeds Controlled and Guidelines for Control for additional information. Apply this product uniformly with ground or aerial equipment in 10 to 20 gallons total spray volume per acre (minimum of 5 gallons per acre by air).

Target Broadleaf Weeds	LIBERTY CLOP (pt/acre)
Canada thistle	1/3 For top growth suppression
Canada thistle	1/2 For season long control
Perennial sowthistle	
Annual sowthistle	1/4 – 1/2
Biennial wormwood	
Dandelion	
Dock, curly	

False chamomile	
Green smartweed	
Mayweed chamomile	
Nightshade species	
Sunflower	
Wild buckwheat	

Restrictions

- **Preharvest Interval:** Do not apply within 50 days of harvest.
- Make one broadcast application per crop per year, not to exceed 1/2 pint per acre.

CHRISTMAS TREE PLANTATIONS (Not Registered for Use in Florida)

Application Timing

Use LIBERTY CLOP for over the top application to actively growing balsam fir, blue spruce, Douglas fir, fraser fir, grand fir, lodgepole pine, noble fir, ponderosa pine, and white pine. In the Pacific Northwest, do not apply in the first year of transplanting because some needle curling has been observed on first year transplants. For control of annual weeds, apply this product from weed emergence up to the 5-leaf stage of growth. For control of wild buckwheat, apply at 3- to 5-leaf stage of growth, but before vining. For control of weeds, such as Canada thistle and knapweeds, apply after the majority of the basal leaves have emerged up to bud stage. Later application may result in less consistent control.

Application Rate

Apply 1/4 to 1/2 pint of LIBERTY CLOP per acre for control of annual weeds. Apply 1/2 to 2/3 pint of LIBERTY CLOP per acre for difficult to control weeds, such as Canada thistle and knapweeds. Apply as a broadcast or band application in a minimum of 10 gallons per acre by ground application. Use the formulas under Band Application to determine the rate and volume per treated acre.

LIBERTY CLOP may be applied as a spot treatment using a hand held sprayer at an equivalent broadcast rate of 1/2 to 2/3 pint per acre. Refer to Hand Held Sprayers under Spot Treatment in the Application Directions section.

Precautions

- Tree injury may occur with the addition of a surfactant or crop oil with this product. Do not use unless previous experience shows injury is tolerable.

Restrictions

- Re-treat as necessary, but do not exceed 2/3 pint per acre per annual growing season.
- **Blue spruce:** Do not exceed 1/2 pint per acre per annual growing season.
- Do not apply with an air blast sprayer.

CORN (FIELD, POP, SWEET) (Not Registered for Use in Florida)

Use LIBERTY CLOP for postemergence control of annual sowthistle, Canada thistle, common cocklebur, common sunflower, giant and common ragweed, Jerusalem artichoke, jimsonweed and other broadleaf weeds infesting field corn. Apply LIBERTY CLOP at specified timing and rates for field, pop and sweet corn as indicated below.

Weed Control

For control of common cocklebur, common ragweed, giant ragweed, sunflower, other annual weeds and Jerusalem artichoke, apply 1/4 to 1/2 pint of LIBERTY CLOP per acre from weed emergence up to the 5-leaf stage of growth. Use a higher rate in the rate range for heavy infestations or when greater residual control is desired. Consult the table entitled Broadleaf Weeds Controlled and Guidelines for Control for additional information.

Control of Canada Thistle

For effective control of Canada thistle, apply 1/3 to 2/3 pint of LIBERTY CLOP per acre as a broadcast treatment to the entire infested area. Apply when the majority of thistle plants have emerged and thistles are at least 6 to 8 inches in diameter or height up to bud stage. Cultivation can disrupt translocation to the roots of Canada thistle. For best long-term control, do not cultivate before or after application. If cultivation is necessary, wait 14 to 20 days after application before cultivating to allow for thorough translocation.

Control of Canada thistle is influenced by growing conditions, density and size of thistle plant at application, tillage practices used, etc. Light infestations (less than 10 plants per square yard) will generally be adequately controlled with a rate of 1/3 pint per acre. For medium to heavy infestations (more than 10 plants per square yard), rates of 1/2 to 2/3 pint per acre are generally more effective since these Canada thistle stands involve an extensive rhizome system.

The following are general descriptions of control to be expected from each application rate given a medium to heavy population of Canada thistle. Control of lighter infestations may be better than that described.

- A rate of 1/3 pint per acre will suppress top growth of Canada thistle for 6 to 8 weeks. Some regrowth may occur by the end of the season, but this will not interfere with harvesting of the crop.
- A rate of 1/2 pint per acre will generally provide season-long control of Canada thistle. Not all rhizomes will be killed and some regrowth may occur by the end of the growing season.
- A rate of 2/3 pint per acre will provide season-long control of Canada thistle plus suppression into the following season, resulting in a reduction of the total number of Canada thistle plants in the treated area.

FIELD CORN

Application Timing

Apply LIBERTY CLOP to actively growing broadleaf weeds any time after corn emergence through 24 inch tall corn. Apply with ground equipment as a postemergence broadcast or directed spray in 10 gallons or more of spray volume per acre to ensure uniform and thorough spray coverage of the weed foliage. Use only spray nozzles designed for herbicide application. Using flat fan nozzles provides the best coverage and distribution of chemical on the plant foliage. Use spray pressures (at the boom) specified by nozzle manufacturers to obtain desired spray volume. Use higher spray volumes when weed foliage is dense.

Tank Mixes or Sequential Applications

See Tank Mixing section under Mixing Directions. If LIBERTY CLOP is applied sequentially or in combination with Clopyralid + Flumetsulam to the current corn crop, apply 4.0 – 8.1 fl oz/acre of this product with labeled rate of Clopyralid + Flumetsulam. When the high rate, 8.1 fl oz, of LIBERTY CLOP is used sequentially or in combination with Clopyralid + Flumetsulam; the Clopyralid + Flumetsulam will be used at the low rate. When LIBERTY CLOP is applied using the low rate, 4.0 fl oz, Clopyralid + Flumetsulam will be mixed or used sequentially using its high rate. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Precautions

- Maximum use rate for clopyralid is 0.25 lb a.e. per acre of all clopyralid containing products.
 - One fluid ounce of this product contains 0.023 lb of clopyralid.

Corn Inbred Lines or Breeding Stock

Susceptibility of corn to injury from LIBERTY CLOP is highly related to varietal response. Inbred lines or any breeding stock may be injured by this product. Contact your seed production agronomist for advice before applying this product to inbred lines or breeding stock.

Hand Held Sprayers

LIBERTY CLOP may be applied as a spot treatment using a hand held sprayer at an equivalent broadcast rate of 2/3 pint per acre. Refer to Hand Held Sprayers under Spot Treatment in the Application Directions

section. Make applications on a spray-to-wet basis with spray coverage uniform and complete. Do not spray to the point of runoff.

Restrictions

- Retreat as necessary, but do not apply more than 2/3 pint per acre per year.
- Do not apply to field corn more than 24 inches tall.
- Do not allow livestock to graze treated areas or harvest treated corn silage as feed within 40 days after last treatment.

POPCORN AND SWEET CORN (Not Registered for Use in California)

Application Timing

For popcorn, apply LIBERTY CLOP any time after popcorn emergence through 24-inch tall popcorn. For sweet corn, apply this product any time after sweet corn emergence through 18-inch tall sweet corn.

Application Rate

Apply 1/3 to 2/3 pint of LIBERTY CLOP per acre uniformly with ground equipment as a broadcast or directed spray in 10 to 20 gallons total spray volume per acre. For control of Canada thistle, apply this product when the majority of thistle plants have emerged and thistles are at least 6 to 8 inches in diameter or height, but before bud stage. For control of annual sowthistle, common cocklebur, Jerusalem artichoke, jimsonweed, ragweed (common and giant), and sunflower, apply this product from weed emergence up to the 5-leaf stage of growth. Use a higher rate in the rate range for heavy infestations or when greater residual control is desired. Consult the table entitled Broadleaf Weeds Controlled and Guidelines for Control for additional information.

Restrictions

- **Preharvest Interval:** Do not apply within 30 days of harvest for ears and forage and within 60 days of harvest for stover.
- Make one to two broadcast applications per crop per year, not to exceed a total of 2/3 pint per acre.
- **Re-Treatment Interval:** 21 days.
- Do not apply to popcorn more than 24 inches tall or sweet corn more than 18 inches tall.
- Apply only to sweet corn or popcorn that is to be used for processing.

COTTONWOOD/POPLAR AND EUCALYPTUS TREE PLANTATIONS (Not Registered for Use in Florida)

Application Timing

LIBERTY CLOP may be used for selective postemergence control of labeled broadleaf weeds in new and established plantings of cottonwood/poplar and eucalyptus tree plantations.

Application Rate

Apply as a broadcast foliar spray over trees or as a banded or directed spray at a rate of 1/3 to 2/3 pint per acre. Apply in 10 gallons or more total spray volume per acre using ground equipment only. Multiple applications of LIBERTY CLOP may be made as long as the total rate per growing season does not exceed 1 1/3 pints per acre. Apply to new plantings only after they are well established as indicated by several inches of new healthy growth.

See Broadleaf Weeds Controlled and Guidelines for Control for specified rates and timing for specific susceptible annual, biennial, and perennial weeds.

Hand Held Sprayers

Spot applications using hand held equipment are also allowed, but avoid contact with tree foliage or limit it to lower branches. Apply to weeds on a spray-to-wet basis with uniform and complete spray coverage. Do not spray to the point of runoff. Prepare a spray solution by adding 1/4 fl oz of LIBERTY CLOP per gallon

of water. When applied at 1 gallon of spray per 1000 sq ft, this spray concentration is equivalent to a broadcast rate of 2/3 pint per acre.

Precautions

- This product will not control certain broadleaf weeds including mustards, henbit, chickweed, kochia, lambsquarters, pigweed, Russian thistle and bindweed.

Restrictions

- Do not tank mix this product with other herbicides labeled for this use unless spray avoids all contact with tree foliage.
- Retreat as necessary, but do not apply more than 1 1/3 pints per acre per year per growing season.

FALLOW CROPLAND (Not Registered for Use in Florida)

Application Timing

LIBERTY CLOP can be applied either postharvest, in the spring/summer (during fallow period), or to set aside acres to control or suppress listed weeds (refer to rotation restrictions). Apply to young, emerged weeds under conditions that promote active growth. For best results on perennial weeds, such as Canada thistle, apply after the majority of the basal leaves have emerged up to bud stage. Later applications may result in less consistent control.

For best results, wait 14 to 20 days after application before cultivating or fertilizing with shank-type applicators to allow for thorough translocation.

Application Rate

Apply 1/4 to 2/3 pint of LIBERTY CLOP per acre. Use a higher rate in the rate range on perennial weeds or when the condition of weeds at treatment may prevent optimum control.

Tank Mixes

To improve control of certain broadleaf weeds, LIBERTY CLOP may be applied with 0.5 to 2 lb a.e. of 2,4-D per acre. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. See Tank Mixing section under Mixing Directions.

Restriction

- Retreat as necessary, but do not apply more than 2/3 pint per acre per year.

GARDEN BEET (Not Registered for Use in California and Florida)

Use LIBERTY CLOP for postemergence control of common ragweed, galinsoga, nightshade (black cutleaf, Eastern black and hairy), prickly lettuce, sowthistle, sweet clover, and wild buckwheat infesting garden beet.

Application Timing

Apply to garden beet in the 2- to 8-leaf stage of crop growth when weeds are young and actively growing. Apply LIBERTY CLOP to wild buckwheat at the 1- to 3-leaf stage of growth before vining begins. Apply this product to common ragweed and sweet clover from weed emergence up to the 5-leaf stage of growth. Apply this product to all species of nightshade at the 2- to 4-leaf stage of growth. Apply this product to sowthistle from rosette up to bud stage. Apply in 10 gallons or more total spray volume per acre with ground equipment.

Application Rate

Apply 1/4 to 1/2 pint of LIBERTY CLOP per acre with ground equipment in 10 gallons or more total spray volume per acre. Use a higher rate in the rate range for heavy infestations or when greater residual control is desired.

Restrictions

- **Preharvest Interval:** Do not apply within 30 days of harvest.
- Make one to two broadcast applications per crop per year, not to exceed a total of 1/2 pint per acre.

**GRASSES GROWN FOR SEED
(Not Registered for Use in Florida)**

Application Timing

Apply only to established grasses before the boot stage of growth. Applications in the boot stage and beyond can result in increased potential for injury. Do not apply to bentgrass unless injury can be tolerated. For control of late emerging Canada thistle, a preharvest treatment may be made after grass seed is fully developed. Treatment of Canada thistle at the bud stage or later may result in less consistent control. Postharvest fall treatments may be made to actively growing Canada thistle after the majority of basal leaves have emerged.

Application Rate

Use 1/4 to 2/3 pint of LIBERTY CLOP per acre for control of annual weeds and Canada thistle.

Tank Mixes

LIBERTY CLOP may be tank mixed with 2,4-D, MCPA, dicamba, or bromoxynil to control additional broadleaf weeds. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. See Tank Mixing section under Mixing Directions.

Precautions

- Dicamba or bromoxynil tank mixes may be useful in broadening the annual weed control spectrum, but may reduce long-term control of perennials, such as Canada thistle. Do not tank mix LIBERTY CLOP with 2,4-D, MCPA, or dicamba unless the risk to crop injury is acceptable.

Restrictions

- Re-treat as necessary, but do not exceed 2/3 pint per acre per season.

**PEPPERMINT AND SPEARMINT
(Not Registered for Use in Florida)**

LIBERTY CLOP may be used for selective postemergence control of specific annual and perennial broadleaf weeds infesting peppermint and spearmint.

Application Timing

Treat annual weeds when they are small and actively growing before they send up a flower stalk. For Canada thistle, apply LIBERTY CLOP after the majority of basal leaves have emerged, but prior to bud stage.

Application Rate

Apply as a broadcast foliar spray in 10 gallons or more total spray volume per acre using ground equipment only. A nonionic surfactant of at least 80% active ingredient may be added at a rate of 1 pint per 100 gallons of spray solution.

Application Timing and Weeds Controlled	LIBERTY CLOP (pt/acre)
Fall treatment only (Sept. 15 to first frost)	
Annuals	1/2
Perennials	1/3
Hard-to-kill perennials (Canada thistle, dandelion)	1
Spring treatment only	
Annuals	1/3
Perennials	1/2

Fall plus spring treatment	Maximum of 2/3 in fall + 1/3 in spring
----------------------------	-------------------------------------------

Precautions

- Discoloration or malformation of peppermint and spearmint leaves may occur following treatment. This effect is generally temporary and does not reduce oil yields.
- This product will not control many broadleaf weeds, such as chickweed, field bindweed, henbit, kochia, lambsquarters, mustards, pigweed, and Russian thistle.

Restrictions

- **Preharvest Interval:** Do not apply within 45 days of harvest.
- Retreat as necessary, but do not apply more than 1 pint per acre per growing season.
- Treated peppermint and spearmint may be used for distillation (oil extraction) only.
- Do not feed spent peppermint and spearmint hay slugs to livestock.
- Peppermint and spearmint straw, hay or spent hay (slugs) from treated areas cannot be used for composting or mulching. If hay slugs are disposed of on cropland, distribute in a thin layer and incorporate. Do not dispose of hay slugs on land to be rotated to a susceptible crop. (See Residues in Plants or Manure section.)

SOUTHERN PINE SEEDBEDS IN FOREST NURSERIES

**Registered for Use in Alabama, Arkansas, Georgia, Louisiana, Mississippi, North Carolina, South Carolina, Tennessee, Texas and Virginia
(Not Registered for Use in Florida)**

LIBERTY CLOP may be applied over the top of loblolly pine, slash pine, and longleaf pine to control sicklepod and other susceptible broadleaf weeds in southern pine seedbeds in forest nurseries. Apply as a broadcast or spot treatment from May through July when weeds are actively growing.

Application Timing

For best results, apply when weeds are small and actively growing. For control of sicklepod, apply after the majority of basal leaves have emerged.

Application Rate

Apply at a broadcast rate of 1/4 to 1/2 pint per acre in a spray volume of 20 gallons or more per acre. Application may be made any time after May 1, but some needle curling may occur if applied during active conifer growth. When making spot applications, use a calibrated boom, or if a hand held sprayer is used, care should be taken to apply the spray uniformly and at a rate equivalent to a broadcast application. Otherwise, do not use more than 1/5 fl oz (1 tsp.) of LIBERTY CLOP per gallon of spray and direct spray onto weeds. Avoid spraying pine seedlings whenever possible.

Precautions

- Application of this product during active growth of conifers may cause some needle curling.
- Do not use surfactants or crop oils in spray mixtures as the potential for tree injury in the form of needle curling may be increased.

Restriction

- Do not apply more than 1/2 pint per acre per growing season.

SPINACH

(Not Registered for Use in California and Florida)

Use LIBERTY CLOP for postemergence control of annual sowthistle, black nightshade, Canada thistle, clover, common cocklebur, common groundsel, hairy nightshade, jimsonweed, pineappleweed, prickly lettuce, and ragweed infesting spinach.

Application Timing

Apply to spinach in the 2- to 5-leaf stage of crop growth. Apply LIBERTY CLOP to clover, common cocklebur, common groundsel, jimsonweed, prickly lettuce, pineappleweed and ragweed from weed

emergence up to the 5-leaf stage of growth. For top growth suppression of annual sowthistle and Canada thistle, apply this product from rosette up to bud stage. For control of Canada thistle, apply after the majority of basal leaves have emerged but prior to bud stage and at least 21 days prior to harvest.

Application Rate

Apply 1/4 to 1/2 pint per acre of LIBERTY CLOP uniformly with ground or aerial equipment in 10 to 20 gallons total spray volume per acre (minimum of 5 gallons per acre by air). Use a higher rate in the rate range for heavy infestations or when greater residual control is desired.

Precautions

- Some leaf curling may be observed on smaller spinach, particularly at higher use rates. Crop tolerance may be optimized by selecting the lower application rate necessary for weed control, especially where non-uniform emergence has caused variable plant sizes

Restrictions

- **Preharvest Interval:** Do not apply within 21 days of harvest.
- Make one to two broadcast applications per crop per year, not to exceed a total of 1/2 pint per acre.

STONE FRUITS (CROP GROUP 12)¹ (Not Registered for Use in California and Florida)

¹ Stonefruits (Crop Group 12) including apricot, chickasaw plum, damson plum, fresh prune, Japanese plum, nectarine, peach, plum, plumcot, sweet cherry, tart cherry.

Use LIBERTY CLOP for postemergence control of annual sowthistle, Canada thistle, clover, dandelion, horseweed, musk thistle, nightshade (black and hairy), and vetch infesting stone fruits.

Application Timing

Apply LIBERTY CLOP to clover and vetch from weed emergence up to the 5-leaf stage of growth. Apply this product to nightshade (black and hairy) at the 2- to 4-leaf stage of growth. For control of Canada thistle and annual sowthistle, apply this product from rosette up to bud stage.

Application Rate

Apply 1/3 to 2/3 pint of LIBERTY CLOP per acre with ground equipment in 10 gallons or more total spray volume per acre. Use a higher rate in the rate range for heavy infestations or when greater residual control is desired.

Restrictions

- **Preharvest Interval:** Do not apply within 30 days of harvest.
- Make one to two broadcast applications per crop per year, not to exceed a total of 2/3 pint per acre.

SUGAR BEET (Not Registered for Use in Florida)

Use LIBERTY CLOP for the control of various annual and perennial broad leaf weeds infesting sugar beet.

Application Rate

Apply 1/4 to 2/3 pint of LIBERTY CLOP per acre with ground equipment as a broadcast foliar spray or band treatment or with aerial equipment in 5 gallons or more total spray volume per acre. See instructions for band application under Application Directions in the Product Information section. Apply in 10 gallons or more total spray volume per acre when the sugar beets are in the cotyledon to 8-leaf stage of growth and the weeds are young and actively growing

For annual weed control apply 1/4 to 1/2 pint of LIBERTY CLOP per acre from weed emergence up to the 5-leaf stage of growth. Apply to wild buckwheat at the 1- to 3-leaf stage of growth before vining begins.

For the most effective control of perennials, such as Canada thistle and sowthistle, apply 1/2 to 2/3 pint of LIBERTY CLOP per acre as a broadcast treatment to the entire infested area. Apply when the majority of basal leaves have emerged up to the bud stage. Cultivation can disrupt translocation to the roots of perennials, such as Canada thistle. For best results, do not cultivate thistle patches.

To promote herbicidal efficacy, wait a minimum of 7 days after application before flood or furrow irrigation.

Tank Mixes

To control additional broadleaf weeds and provide consistent control of difficult to control weeds, such as wild buckwheat, LIBERTY CLOP may be applied in combination with labeled rates of a product containing phenmedipham/desmedipham, esmedipham, triflurosulfuron, or other products registered for postemergence application in sugar beets. For best results, tank mix 1/4 pint of this product per acre with a product containing phenmedipham/desmedipham or desmedipham followed one to two weeks later by a second application of 1/4 to 1/3 pint of this product per acre tank mixed with a product containing phenmedipham/desmedipham or desmedipham. This product may also be tank mixed with a grass herbicide containing sethoxydim. Crop oil or Dash surfactant may be added to the tank mixture to optimize grass weed control. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. See Tank Mixing section under Mixing Directions.

Restrictions

- **Preharvest Interval:** Do not apply within 45 days of harvest.
- Re-treat as necessary, but do not exceed 2/3 pint per acre per season.
- Aerial application of this product in sugar beet is allowed only in the states of Colorado, Idaho, Michigan, Minnesota, Montana, Nebraska, North Dakota, Oregon, Washington, and Wyoming.

TURNIP

(Not Registered for Use in California and Florida)

Use LIBERTY CLOP for postemergence control of common ragweed, galinsoga, prickly lettuce, sweet clover, and wild buckwheat and postemergence suppression of sowthistle infesting turnip harvested for roots and tops.

Application Timing

Apply LIBERTY CLOP to wild buckwheat at the 1- to 3-leaf stage of growth before vining begins. Apply LIBERTY CLOP to common ragweed and sweet clover from weed emergence up to the 5-leaf stage of growth. For suppression of sowthistle, apply LIBERTY CLOP from rosette up to bud stage.

Application Rate

Apply 1/3 to 1/2 pint of LIBERTY CLOP per acre with ground equipment in 10 gallons or more total spray volume per acre. Use a higher rate in the rate range for heavy infestations or when greater residual control is desired.

Restrictions

- **Preharvest Interval:** Do not apply within 30 days of harvest of turnip roots or within 15 days of harvest of turnip tops.
- Make one broadcast application per crop per year, not to exceed a total of 1/2 pint per acre.

RANGELAND, PASTURE, CRP AND NON-CROP USES

(Not Registered for Use in Florida)

Rotation to Broadleaf Crops: Do not plant broadleaf crops in treated areas until an adequately sensitive bioassay shows that no detectable clopyralid is present in the soil. (See Crop Rotation Restrictions in Product Information section.)

Rangeland and Permanent Grass Pastures

Apply 1/2 to 1 1/3 pints of LIBERTY CLOP per acre when weeds are young and actively growing. Established grasses are tolerant to LIBERTY CLOP, but new grass seedlings may be injured to varying degrees until the grass has become well established as indicated by vigorous growth and development of tillers and secondary roots.

Precaution

- Some forbs (desirable broadleaf forage plants) are susceptible to this product. However, the stand and growth of established perennial grasses is usually improved after spraying, especially when rainfall is adequate and grazing is deferred.

Restrictions

- Do not use hay or straw from treated areas for composting or mulching on susceptible broadleaf crops. (See Residues in Plants or Manure section.)
- There are no further restrictions on grazing or hay harvest following application of this product at labeled rates.
- Re-treat as necessary, but do not exceed 1 1/3 pint per acre per season.

Conservation Reserve Program (CRP) for Seeding to Permanent Grasses Only

Application Timing: Apply LIBERTY CLOP when perennial grasses are well established as indicated by vigorous growth and development of tillers and secondary roots. At this stage, most perennial grasses have shown adequate tolerance to this product. For optimum results, apply prior to the flowering stage (still in the bud stage).

Application Rate: For control of actively growing weeds, such as Canada thistle, knapweed (spotted, diffuse, and Russian), and musk thistle, apply 2/3 to 1 1/3 pints of LIBERTY CLOP per acre after the majority of basal leaves have emerged up to bud stage. For control of musk thistle rosettes, volunteer sunflower, and wild buckwheat, apply 2/3 pint of this product per acre. For best results; use in 10 gallons or more of water per acre by ground. Increasing the application rate increases the risk of injury.

Tank Mixes: LIBERTY CLOP can also be tank mixed with 1/2 to 1 lb of 2,4-D per acre where species present are sensitive to 2,4-D. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. See Tank Mixing section under Mixing Directions.

Precautions

- Conditions of plant stress, such as drought, will increase potential for injury to grasses at all stages of growth.

Restrictions

- Do not use this product if legumes or bentgrass are a desired cover during CRP.
- Do not apply to newly seeded areas until grass is established.
- Do not apply more than 1 1/3 pint per acre per year.

Non-Cropland

LIBERTY CLOP may be applied in non-cropland areas, such as fencerows, around farm buildings and equipment pathways.

Application Rate: For control of broadleaf weeds, apply 1/4 to 1 1/3 pints of LIBERTY CLOP per acre. The lower rate of 1/4 pint per acre provides acceptable control of weeds only under highly favorable growing conditions and when plants are 1 to 3 inches tall. Apply 1/2 pint per acre when weeds are 3 to 6 inches tall or under dry conditions. Where Canada thistle or knapweeds are the primary pest, best results are obtained by applying 2/3 to 1 1/3 pints of this product per acre.

Tank Mixes: To improve spectrum of weed control, or to increase control of more mature weeds, LIBERTY CLOP may be tank mixed with 0.5 to 2 lb a.e. of 2,4-D amine per acre or low volatile ester herbicide or other herbicides registered for this use site. It is the pesticide user's responsibility to ensure that all products

in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. See Tank Mixing section under Mixing Directions.

Precaution

- This product is not registered for use in landscaping or on turfgrass or lawns.

Restriction

- Do not apply more than 1 1/3 pint per acre per year.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store above 28°F or warm to 40°F and agitate before use. Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling:

NONREFILLABLE CONTAINER (EQUAL TO OR LESS THAN 5 GALLONS): Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

NONREFILLABLE CONTAINER (GREATER THAN 5 GALLONS): Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration, or by other procedures allowed by state and local authorities.

REFILLABLE CONTAINER: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. After triple rinsing is complete, and the container is not suitable for refilling or reconditioning, offer the container for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of LIBERTY CROP PROTECTION LLC or Seller. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW all such risks shall be assumed by Buyer and User and Buyer and User agree to hold LIBERTY CROP PROTECTION LLC and Seller harmless for any claims relating to such factors.

LIBERTY CROP PROTECTION LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or LIBERTY CROP PROTECTION LLC, and TO THE EXTENT CONSISTENT WITH APPLICABLE LAW Buyer and User assume the risk of any such use. To the extent consistent with applicable law LIBERTY CROP PROTECTION LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither LIBERTY CROP PROTECTION LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF LIBERTY CROP PROTECTION LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LIBERTY CROP PROTECTION, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

LIBERTY CROP PROTECTION LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of LIBERTY CROP PROTECTION LLC.

All trademarks are the property of their respective owners.