



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Registration Division (7505P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

89168-75

Date of Issuance:

3/26/18

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Liberty Flumiox 44

Name and Address of Registrant (include ZIP Code):

Mary Beth Enders
 Liberty Crop Protection LLC
 4850 Hahns Peak Drive, Suite 200
 Loveland, CO 80538

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.
2. You are required to comply with the data requirements described in the DCI identified below:
 - a. Flumioxazin GDCI-129034-1236

Signature of Approving Official:

Shaja Joyner
 Shaja Joyner, Product Manager 20
 Fungicide Herbicide Branch, Registration Division (7505P)

Date:

3/26/18

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division: <http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 89168-75.”
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 8/4/2017
- Alternate CSF 1 dated 8/4/2017

If you have any questions, please contact Nathan Mellor by phone at 703-347-8562, or via email at mellor.nathan@epa.gov.

Enclosure

LIBERTY FLUMIOX 44

FOR CONTROL AND/OR SUPPRESSION OF CERTAIN WEEDS IN ALFALFA, ARTICHOKE, ASPARAGUS, HEAD AND STEM BRASSICA, BUSHBERRIES, CACTUS (PRICKLY PEAR), CANEBERRIES, CELERY, CITRUS FRUIT, CLOVER, COTTON, CUCURBIT VEGETABLES, DRY BEANS, FIELD CORN, FIELD PEAS, FLAX, FRUITING VEGETABLES (INCLUDING OKRA), GARLIC, GRAPE, HOPS, LENTILS, MINT, NUT TREES (INCLUDING PISTACHIO), ONION (DRY BULB), OLIVE, PEANUT, POME FRUIT, POMEGRANATE, POTATO, SOYBEAN, STONE FRUIT, STRAWBERRY, SUGARCANE, SUNFLOWER AND SAFFLOWER, SWEET POTATO, WHEAT, NON-BEARING FRUIT TREES, FALLOW LAND AND TO MAINTAIN BARE GROUND ON NON-CROP AREAS OF FARMS, ORCHARDS AND VINEYARDS.

ACTIVE INGREDIENT:	% BY WT.
Flumioxazin*	44.0%
OTHER INGREDIENTS:	<u>56.0%</u>
TOTAL:	100.0%


*2-[7-fluoro-3,4-dihydro-3-oxo-4-(2-propynyl)-2H-1,4-benzoxazin-6-yl]-4,5,6,7-tetrahydro-1H-isoindole-1,3(2H)-dione
 This product is a suspension concentrate containing 4 pounds flumioxazin per gallon.

Shake Well Before Use

KEEP OUT OF REACH OF CHILDREN
CAUTION / PRECAUCIÓN
 Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand the label, find someone to explain it to you in detail.)

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS.



EPA Reg. No.: 89168-TL

EPA Est. No.: _____

NET CONTENTS: ___ GAL (___ L)

Manufactured for:
 LIBERTY CROP PROTECTION, LLC
 1880 Fall River Drive, Suite 100
 Loveland, CO 80538

032218

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 or your poison control center at 1-800-222-1222. For Chemical Spill, Leak, Fire or Exposure, call CHEMTREC 800-424-9300.</p>	

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some of the materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material including polyethylene or polyvinyl chloride, shoes and socks.

For aerial application to sugarcane, mixer/loaders must also wear: coveralls, chemical resistant apron and chemical resistant boots.

For aerial application to artichoke; field peas; flax; lentils; safflower; sunflower and wheat, mixer/loaders must also wear: filtering face piece respirator (N95, R95 or P95).

For ground boom application to cactus (prickly pear); olive and pomegranate, mixer/loaders must also wear: filtering face piece respirator (N95, R95 or P95).

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS
<p>Users should:</p> <ul style="list-style-type: none"> • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to non-target plants and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff maybe hazardous to non-target plants and aquatic organisms in neighboring areas. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

This pesticide is toxic to plants and should be used strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposures.

Under some conditions this product may have a potential to run-off to surface water or adjacent land. Where possible, use methods which reduce soil erosion, including no till, limited till and contour plowing; these methods also reduce pesticide run-off. Use of vegetation filter strips along rivers, creeks, streams, wetlands or on the downhill side of fields where run-off could occur will minimize water run-off.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow coming in contact with oxidizing agent. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read the entire label before using this product. Use strictly in accordance with label precautionary statements and directions, and with applicable state and federal regulations.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statement of this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box only apply to users of this product that are covered by the WPS.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil or water is: coveralls, chemical resistant gloves made of waterproof material and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the WPS for agricultural pesticides (40CFR Part 170). The WPS applies when this product is used to produce agricultural crops on farms, forests, nurseries, or greenhouses.

Keep all unprotected persons out of operating areas, or vicinity where there may be drift. Do not enter or allow others to enter the treated area until sprays have dried.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product must be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of LIBERTY CROP PROTECTION LLC. The Buyer must be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are

required to take the crop to harvest. If the Buyer chooses not to accept these risks, then this product must not be applied. By applying this product buyer acknowledges and accepts these inherent unintended risks and to the fullest extent allowed by law, agrees that all such risks associated with the application and use are assumed by the buyer.

Liberty shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

See also **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY** sections of the label for additional information.

RESISTANCE MANAGEMENT

For resistance management, this product is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

Weed Management

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in the field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact LIBERTY CROP PROTECTION, LLC at [855-466-8428 or 844-425-8488 or other appropriate telephone number].

Management of Resistant Biotypes

The following good agronomic practices are recommended to reduce the spread of resistant biotypes:

- If a naturally occurring resistant biotype is present in your application site, this product should be tank mixed or applied sequentially with an appropriately labeled herbicide with a different mode of action to achieve control.
- Cultural and mechanical control practices (e.g. crop rotation or tillage) may also be used as appropriate.
- Scout treated application site after herbicide applications and control escaping weeds including resistant biotypes before they set seed.
- Thoroughly clean equipment before leaving fields known to contain resistant biotypes.
- Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to this Mode of Actions have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of actions for each target weed.

Integrated Pest (Weed) Management

This product may be integrated into an overall weed pest management strategy whenever the use of an herbicide is required. Practices known to reduce weed development (tillage, crop competition) and herbicide use (weed scouting, proper application timing, banding) should be followed wherever possible. Consult local agricultural and weed authorities for additional IPM strategies established for your area.

TANK MIXES

Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor, to the extent allowed by applicable law. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

PRODUCT INFORMATION

- This product can be used on farms, orchards and vineyards for non-selective vegetation control to maintain bare ground non-crop areas that must be kept weed free.
- This product provides residual control of susceptible weeds.
- This product provides additional burndown activity when used as part of a burndown program.
- This product can be applied as part of a fall burndown program for control of susceptible winter annuals.
- This product can be applied with a hooded or shielded sprayer, as well as part of a layby application, in selected crops for postemergence weed control as well as residual control of susceptible weeds.
- Read tank mix product label for rates and weeds controlled. Always read and follow label directions for all tank mix products before using. The most restrictive labeling of any tank mix product must be followed. When this product is applied according to label use directions, will control the weeds claimed in crop specific use directions. This label makes no claims concerning control of other weed species.

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.

The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making decisions. Where states have more stringent regulations, they must be observed.

RESTRICTIONS

- Do not apply this product when weather conditions favor spray drift from treated areas.
- Do not apply during low-level inversion conditions, including fog.
- Do not apply to frozen or snow covered soil.
- Do not apply to farm alleys or roads where traffic may result in treated dust settling onto crops or other desirable vegetation.
- Do not apply within 300 yards of non-dormant pears.
- Do not apply to powdery soils or soils that are susceptible to wind displacement unless irrigation can be applied immediately after application.

PRECAUTIONS

- When applying by air, observe drift management restrictions and precautions listed under "AERIAL APPLICATION".
- Mechanical incorporation into the soil will reduce residual weed control.

Note:

- Post directed and layby applications of this product should be applied only to healthy growing crops.

Do not use spray equipment used to apply this product to apply other materials to any crop foliage, unless the proper cleanout procedures are followed. See "SPRAYER CLEANUP" for more information.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL PERFORMANCE

Preemergence Application (Conventional Tillage)

Important: Crop injury may occur from applications made to poorly drained soils and/or applications made under cool, wet conditions. Risk of crop injury can be minimized by using on well drained soils, planting at least 1.5 inches deep, using high quality seed and completely covering seeds with soil prior to preemergence applications. Treated soil that is splashed onto newly emerged crops may result in temporary crop injury.

Moisture is necessary to activate this product in soil for residual weed control. Dry weather following applications of this product may reduce effectiveness. However, when adequate moisture is received after dry conditions, this product will control susceptible germinating weeds. This product may not control weeds that germinate after application but before an activating rainfall/irrigation or weeds that germinate through cracks resulting from dry soil.

When adequate moisture is not received after an application of this product, weed control may be improved by irrigation with at least 1/4 inch of water. If emerged weeds are controlled by cultivation, residual weed control will be reduced.

Burndown Application

For best results, apply this product as part of a burndown program to actively growing weeds. Applying this product under conditions that do not promote active weed growth will reduce herbicide effectiveness. Do not apply this product when weeds are under stress due to drought, excessive water, extremes in temperature, disease or low humidity. Weeds under stress tend to become less susceptible to herbicidal action. This product is most effective when applied under warm sunny conditions.

Reduced residual weed control may occur when burndown applications are made to fields where heavy crop and/or weed residue exist.

Postemergence Application

Only apply this product to healthy crops labeled for postemergence use. Do not apply this product to crops that have been weakened by disease, drought, flooding, excessive fertilization, soil salts, previously applied pesticides, nematodes, insects or winter injury.

Rainfast

This product is rainfast one hour after application. Postemergence efficacy may be reduced if rain is expected within one hour of application.

Soil Characteristics

Application of this product to soils with high organic matter and/or high clay content may require higher dosages than soils with low organic matter and/or low clay content. Application to cloddy seedbeds can result in reduced weed control.

HERBICIDE RATE

Residual Weed Control (Including Preemergence Applications or Applications as Part of a Fall or Spring Burndown and Fallow Seedbed Program)

Based upon soil characteristics (organic matter content and texture), the most difficult to control weed species being targeted, and the crop being grown, select the proper dosage of this product from the rate range tables contained in this label.

CARRIER VOLUME AND SPRAY PRESSURE (Ground Equipment only. See Information for Aerial Equipment under "AERIAL APPLICATION".)

Preemergence Application (Conventional Tillage)

To ensure uniform coverage, use 10 to 30 gallons of spray solution per acre for conventional tillage applications. Nozzle selection must meet manufacturer's gallonage and pressure specifications for preemergence herbicide application.

Burndown Application (Prior to Crop Emergence)

To ensure thorough coverage in burndown applications, use 15 to 60 gallons spray solution per acre. Use 20 to 60 gallons per acre if dense vegetation or heavy crop residue is present. Nozzle selection must meet manufacturer's gallonage and pressure specifications for postemergence herbicide application. Do not use flood jet nozzles.

Postemergence Application (Emerged Crop)

Check use directions for specific crops in which this product can be applied postemergence. To ensure thorough coverage in burndown applications, use a minimum of 15 gallons spray solution per acre. Use a minimum of 20 gallons per acre if dense vegetation or heavy crop residue is present. Nozzle selection must meet manufacturer's gallonage and pressure specifications for postemergence herbicide application.

ADDITIVES

Burndown Application (Prior to Crop Emergence)

Postemergence control of weeds from tank mixes of this product will require the addition of an agronomically approved adjuvant to the spray mixture. Either a crop oil concentrate or methylated seed oil which contains at least 15% emulsifiers and 80% oil or a non-ionic surfactant at 0.25% v/v, may be used when applying this product as part of a burndown program. Some tank mix partners, for example glyphosate, are formulated with sufficient adjuvants and do not require the addition of a crop oil concentrate, methylated seed oil or non-ionic surfactant when tank mixed with this product. The addition of a crop oil concentrate or methylated seed oil may increase the burndown activity on certain weeds including cutleaf eveningprimrose and Carolina geranium. Verify mixing compatibility qualities by a jar test.

A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 pounds per acre or a 28 to 32% nitrogen solution at 1 to 2 quarts per acre) may be added to the spray mixture along with either a crop oil concentrate, methylated seed oil or non-ionic surfactant to enhance weed control. The addition of a nitrogen source does not replace the need for a crop oil concentrate, a methylated seed oil or a non-ionic surfactant.

JAR TEST TO DETERMINE COMPATIBILITY OF ADJUVANTS AND LIBERTY FLUMIOX 44

When using this product and an adjuvant, including in stale seed bed, layby, hooded/shielded or reduced tillage situations, perform a jar test before mixing commercial quantities of this product, when using this product for the first time, when using new adjuvants or when a new water source is being used.

1. Add 1 pint of the water to a quart jar. Use water from the same source and temperature as which will be used in the spray tank mixing operation.
2. Add 1 milliliter of this product to the quart jar for every 3 fluid ounces of this product per acre being applied (4 milliliters if 12 fluid ounces per acre is the desired rate of this product), gently mix until product goes into suspension.
3. Add 60 milliliters (4 Tablespoons or 2 fluid ounces) of the crop oil or methylated seed oil to the quart jar or 1 milliliter of non-ionic surfactant if it is being used in place of oil, gently mix.

4. If nitrogen is being used, add 16 milliliters (1 Tablespoon or 0.5 ounce) of the 28 to 32% nitrogen source to the quart jar. If ammonium sulfate is being used, add 19 g AMS to the quart jar in place of the 28 to 32% nitrogen.
5. Place cap on jar, invert 10 times, let stand for 15 minutes, evaluate.
6. An ideal tank mix combination will be uniform. If any of the following conditions are observed question the choice of adjuvant:
 - a. Layer of oil or globules on the mixture's surface.
 - b. Flocculation: fine particles in suspension or as a layer on the bottom of the jar.
 - c. Clabbering: Thickening texture (coagulated) like gelatin.

SPRAYER PREPARATION

Before application of this product, start with clean, well maintained application equipment. The spray tank, as well as all hoses and booms, must be cleaned to ensure no residue from the previous spraying operation remains in the sprayer. Some pesticides, including but not limited to, the sulfonylurea and phenoxy herbicides, (i.e., chlorimuron ethyl and 2,4-D respectively) are active at very small amounts and can cause crop injury when applied to susceptible crops. The spray equipment must be cleaned according to the manufacturer's directions for the last product used before the equipment is used to apply this product. If two or more products were tank mixed prior to application of this product, follow the most restrictive cleanup procedure.

MIXING INSTRUCTIONS

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. If a drift retardant is to be used, add 10 pounds of spray grade ammonium sulfate per 100 gallons of spray solution.
3. While agitating, slowly add the LIBERTY FLIMIOX 44 to the spray tank. Agitation creates a rippling or rolling action on the water surface.
4. If tank mixing this product with other labeled herbicides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates and then solutions. Prepare no more spray mixture than is required for the immediate spray operation.
5. Add any required adjuvants.
6. Fill spray tank to desired level with water. **Continue agitation until all spray solution has been applied.**
7. Mix only the amount of spray solution that can be applied the day of mixing. Apply this product within 6 hours of mixing.

SPRAYER CLEANUP

Spray equipment, including mixing vessels and nurse tanks, must be cleaned each day following application of this product. After this product is applied, the following steps must be used to clean the spray equipment:

1. Completely drain the spray tank, rinse the sprayer thoroughly, including the inside and outside of the tank and all in-line screens.
2. Fill the spray tank with clean water and flush all hoses, booms, screens and nozzles.
3. Top off tank, add 1 gallon of 3% household ammonia (or equivalent) for every 100 gallons of water, circulate through sprayer for 5 minutes, and then flush all hoses, booms, screens and nozzles for a minimum of 15 minutes. If diaphragms are being used on the spray boom, loosen diaphragms before flushing the spray system, allowing cleaning solution to spray through the open diaphragm. If spray lines have any end caps, they must be loosened before flushing the system, allowing cleaning solution to spray through the loosened caps. To enhance removal of this product from the spray system, add a tank cleaner including [INNVICTIS PREMIUM TANK CLEANSER or ALTITUDE TANK CLEANER or other appropriate product name], in place of ammonia and allow the cleaning solution to remain in the pressurized spray system (spray tank, hoses and boom) overnight before flushing the system for a minimum of 15 minutes.
4. Drain tank completely.
5. Add enough clean water to the spray tank to allow all hoses, booms, screens and nozzles to be flushed for 2 minutes.
6. Remove all nozzles and screens and rinse them in clean water.

Spray equipment, including all tanks, hoses, booms, screens and nozzles, must be thoroughly cleaned before it is used to apply postemergence pesticides. Equipment with residue of this product remaining in the system may result in crop injury to the subsequently treated crop.

APPLICATION EQUIPMENT

Application equipment must be clean and in good repair. Nozzles must be uniformly spaced on boom and frequently checked for accuracy.

BROADCAST APPLICATION

Apply this product and tank mixes of this product, with ground equipment using standard commercial sprayers equipped with flat fan or flood nozzles (preemergence applications only) designed to deliver the desired spray pressure and spray volume.

BAND APPLICATION

When banding, use proportionately less water and this product per acre. The rate of this product required per acre, when applied as a banded application, can be calculated with the following formula:

$$\frac{\text{Band Width in Inches}}{\text{Row Width in Inches}} \times \text{Rate per Broadcast Acre} = \text{Amount Needed per Acre For Banded Application}$$

AERIAL APPLICATION

Spray drift away from the site of application may cause damage to non-target vegetation. To minimize drift, apply the largest droplet size consistent with uniform coverage and satisfactory weed control. To obtain satisfactory application and avoid drift, the following directions must be observed:

RESTRICTIONS

- Do not apply during low-level inversion conditions (including fog), when winds are gusty or under other conditions that favor drift. Do not spray when wind velocity is less than 2 mph or more than 10 mph.
- Do not apply this product by air within 40 feet of non-target plants including non-target crops.
- Do not apply this product by air within 100 feet of emerged cotton crops.
- Do not apply this product by air within 40 feet of streams, wetlands, marshes, ponds, lakes and reservoirs.
- **Carrier Volume and Spray Pressure:** When used as part of a burndown weed control program, apply this product in 7 to 10 gallons of water per acre. Application at less than 7 gallons per acre may provide inadequate control. When used for preemergence weed control, apply this product in 5 to 10 gallons of water per acre. The higher gallonage applications afford more consistent weed control. Do not exceed the nozzle manufacturer's specified pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Nozzle Selection and Orientation:** Formation of very small drops may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray pressure. Use nozzles that produce flat or hollow cone spray patterns. Use non-drip type nozzles, for example diaphragm type nozzles, to avoid unwanted discharge of spray solution. The nozzles must be directed toward the rear of the aircraft, at an angle between 0 and 15° downward. Do not place nozzles on the outer 25% of the wings or rotors.
- **Adjuvants and Drift Control Additives:** Refer to tank mix partner's label for adjuvant directions. Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

CHEMIGATION

Follow all label instructions for crops regarding rates, timing of application, special instructions and precautions.

Apply this product only through center pivot systems. End guns must be turned off due to uneven application. Do not apply this product through any other type of irrigation system.

Crop injury, lack of efficacy or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

The system must be properly calibrated (with water only) to ensure that the amount of this product applied corresponds to the labeled rate.

Apply this product in 1/2 to 3/4 inches of water during the first sprinkler set. Allow time for all lines to flush the herbicide through all nozzles before turning off irrigation water. To ensure the lines are flushed and free of remaining herbicide, a dye indicator may be injected into the lines to mark the end of the application period. Once chemigation has begun, the run must be completed to ensure no product is left in the system.

If you have any questions about calibration, contact your State Extension Service Specialist, equipment manufacturers or other experts.

Special Precautions for Chemigation

1. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
2. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments in the event the need arises.
3. The system must be free of leaks and clogged nozzles.
4. The pesticide must be supplied continuously for the duration of the aqueous application. An uneven application may cause injury to the crop or poor weed control.
5. Agitation must be maintained in the nurse tank.
6. The sprinkler chemigation system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.
7. The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.
8. The pesticide injection pipeline must contain a functional, normally closed, solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
9. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in the case where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
10. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
11. Systems must use a metering pump, for example a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with the pesticides and capable of being fitted with a system interlock.
12. Do not apply when wind speed favors drift beyond the area intended for treatment.

Chemigation Systems Connected to Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption, if such a system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to the public water system must contain a functional, reduced pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, discharge the water from the public water system into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

3. All Chemigation systems connected to the public water system must also follow restrictions listed in the preceding section titled "**Special Precautions for Chemigation**".

APPLICATION WITH DRY BULK FERTILIZERS

Dry bulk fertilizer may be impregnated or coated with this product. Application of dry bulk fertilizer with this product provides weed control equal to, or slightly below, the same rate of this product applied in liquid carriers, due to better coverage with application via spray equipment. Follow label instructions for this product regarding rates, special instructions, cautions and special precautions. Apply 400 to 700 pounds of the fertilizer/herbicide mixture per acre to obtain adequate soil coverage. Apply the mixture to the soil with properly calibrated equipment immediately after blending. Uniform application of the herbicide/fertilizer mixture is essential to prevent possible crop injury and to obtain uniform weed control.

Do not use ammonium nitrate and/or limestone as the sole source of fertilizer, as this product may not adhere to these materials.

Compliance with all Federal and State regulations relating to blending pesticide mixtures with dry bulk fertilizer, registrations, labeling and application are the responsibility of the individual and/or company offering the fertilizer and mixtures of this product for sale.

This product must be premixed with water to form a slurry prior to impregnation on dry bulk fertilizer. For best results, use a minimum of 1 pint of water for each 2 fluid ounces of this product. Use a minimum of 6 pints of slurry of this product to impregnate 2000 pounds of the fertilizer for uniform coverage of the fertilizer. Closed drum, belt, ribbon or other commonly used dry bulk blenders may be used. The amount of this product required can be calculated with the following formula:

$$\text{Fluid Ounces of This Product Per Acre} \times 2,000 \div \text{Pounds of Fertilizer Per Acre} = \text{Fluid Ounces of This Product Per Ton of Fertilizer}$$

Thoroughly clean dry fertilizer blending equipment after this product has been placed in the system to avoid injury to sensitive crops that may be treated with fertilizers blended after the equipment has been used for this product. Rinse the sides of the blender and the herbicide tank with water. Then impregnate the rinsate onto a load of dry fertilizer intended for an approved crop. Use a maximum rate of 1 gallon of rinsate per ton of fertilizer. Follow with 1 to 2 loads of unimpregnated fertilizer in the blender before switching herbicides.

ROTATIONAL RESTRICTIONS

The following rotational crops may be planted after applying this product at the listed rate.

- **Do not plant any crop, except corn (field), cotton, peanut, soybean, sugarcane and sweet potato earlier than 30 days after applying this product.**

APPLICATION RATE	CROPS	ROTATION INTERVALS
1 fl oz/A	Cotton (no-till or strip-till only)	14 days ¹
1.5 to 2 fl oz/A	Cotton (no-till or strip-till only)	21 days ¹
2 fl oz/A or less	Peanut, Soybean, Sugarcane and Sweet Potato	immediately
	Field Corn (minimum and no-till)	7 days
	Cotton and Field Corn (conventional tillage), Rice, Sorghum, Sunflower, Tobacco and Wheat	30 days ¹
	Barley, Dry and Snap Beans, Flax, Peas, Rye, Safflower and Sweet Corn	3 months
	Alfalfa, Canola, Clover, Oats, Potato, Sugar Beet and all other crops not listed ²	4 months if soil is tilled prior to planting 8 months if no tillage is performed

APPLICATION RATE	CROPS	ROTATION INTERVALS
	Lentil	6 months
Up to 3 fl oz/A	Peanut, Soybean, Sugarcane and Sweet Potato	immediately
	Field Corn (minimum and no-till)	14 days
	Field Corn (conventional tillage) and Sorghum	30 days ¹
	Cotton, Rice, Sunflower, Tobacco and Wheat	2 months ¹
	Barley, Dry and Snap Beans, Flax, Pea, Rye, Safflower and Sweet Corn	4 months
	Alfalfa, Clover, Oats, Potato, Sugar Beet	5 months if soil is tilled prior to planting 10 months if no tillage is performed
	Canola and all other crops not listed ²	6 months if soil is tilled prior to planting 12 months if no tillage is performed
	Lentil	7 months
Up to 4 fl oz/A	Sugarcane	immediately
	Cotton, Field Corn, Peanut, Rice, Sorghum, Soybean, Sunflower, Tobacco and Wheat	4 months
	Alfalfa, Canola, Potato, Sugar Beet and all other crops not listed ²	6 months if soil is tilled prior to planting 12 months if no tillage is performed
	Transplanted on raised beds only: melon, pepper and tomato ³	2 months (if the top 4 inches of the beds have been removed)
6 to 12 fl oz/A	Cotton, Field Corn, Peanut, Rice, Sorghum, Soybean, Sunflower, Tobacco and Wheat	9 months
	Alfalfa, Canola, Sugar Beet and all other crops not listed ² Trees can be transplanted 2 months after an application of this product ⁴	12 months if soil is tilled prior to planting 18 months if no tillage is performed

¹ At least one inch of rainfall/irrigation must occur between application and planting or crop injury may occur.

² Successful soil bioassay must be performed prior to planting crops.

³ **Arizona, California and Hawaii only:** see "DIRECTIONS FOR USE FOLLOWBED USE ON TRANSPLANTED MELON, PEPPER AND TOMATO BEDS" for use instructions.

⁴ Transplanted apple, apricot, avocado, bushberries (including blueberry), cherry, fig, grape, grapefruit, lemon, nectarine, nut trees (including pistachio), olive, orange, peach, pear, plum (including dried plum), and tangerine can be planted 2 months after application of 2 to 12 fluid ounces per acre of this product.

Table 1. Broadleaf Weeds Controlled by Residual Activity of LIBERTY FLUMIOX 44

BROADLEAF WEED SPECIES		ORGANIC MATTER	SOIL TYPE	APPLICATION RATE
SECTION A				
COMMON NAME	SCIENTIFIC NAME	Up to 5%	All Soil Types	2 fl oz/A
Carpetweed	<i>Mollugo verticillata</i>			
Chickweeds,				
Common	<i>Stellaria media</i>			
Mouseear	<i>Cerastium vulgatum</i>			
Dandelion	<i>Taraxacum officinale</i>			
Eclipta	<i>Eclipta prostrata</i>			
Eveningprimrose, Cutleaf	<i>Oenothera laciniata</i>			
Field Pennycress	<i>Thlaspi arvense</i>			
Florida Pusley	<i>Richardia scabra</i>			

Henbit	<i>Lamium amplexicaule</i>			
Lambsquarters, Common	<i>Chenopodium album</i>			
Little Mallow	<i>Malva parviflora</i>			
Marestail/Horseweed	<i>Conyza canadensis</i>			
Mayweed/False Chamomile	<i>Matricaria maritima</i>			
Nightshades,				
Black	<i>Solanum nigrum</i>			
Eastern Black	<i>Solanum ptycanthum</i>			
Hairy	<i>Solanum sarrachoides</i>			
Pigweeds,				
Redroot	<i>Amaranthus retroflexus</i>			
Smooth	<i>Amaranthus hybridus</i>			
Spiny Amaranth	<i>Amaranthus spinosus</i>			
Tumble	<i>Amaranthus albus</i>			
Prickly Lettuce	<i>Lactuca serriola</i>			
Prickly Sida (Teaweed)	<i>Sida spinosa</i>			
Puncturevine	<i>Tribulus terrestris</i>			
Purslane, Common	<i>Portulaca oleracea</i>			
Radish, Wild	<i>Raphanus raphanistrum</i>			
Redmaids	<i>Calandrinia ciliata</i> var. <i>menziessii</i>			
Shepherd's-purse	<i>Capsella bursa-pastoris</i>			
Smallflower Morningglory	<i>Jacquemontia tamnifolia</i>			
Sowthistle, Prickly	<i>Sonchus asper</i>			
Spotted Spurge	<i>Euphorbia maculata</i>			
Venice Mallow	<i>Hibiscus trionum</i>			

Table 1. Broadleaf Weeds Controlled by Residual Activity of LIBERTY FLUMIOX 44 (continued)

SECTION B – All weeds listed in Section A Plus:		ORGANIC MATTER	SOIL TYPE	APPLICATION RATE ²
COMMON NAME	SCIENTIFIC NAME			
Coffee Senna	<i>Cassia occidentalis</i>	Up to 3%	All Soil Types	2 fl oz/A Cotton and Dry Bean 2.5 fl oz/A Field Corn and Soybean 3 fl oz/A Peanut and all other labeled crops
Common Ragweed ¹	<i>Ambrosia artemisiifolia</i>			
False Chamomile	<i>Tripleurospermum maritima</i>			
Florida Beggarweed	<i>Desmodium tortuosum</i>			
Golden Crownbeard	<i>Verbesina encelioides</i>			
Hairy Indigo	<i>Indigofera hirsuta</i>			
Hemp Sesbania	<i>Sesbania exaltata</i>	3 to 5%	Coarse and Medium Soils (sandy loam, loamy sand, loamy, silt-loam, silt, sandy clay, sandy clay loam)	2 fl oz/A Cotton and Dry Bean 2.5 fl oz/A Field Corn and Soybean 3 fl oz/A Peanut and all other labeled crops
Jimsonweed	<i>Datura stramonium</i>			
Kochia	<i>Kochia scoparia</i>			
London Rocket	<i>Sisymbrium irio</i>			
Morningglories, ³				
Entireleaf	<i>Ipomoea hederacea</i> var. <i>integriuscula</i>			
Ivyleaf	<i>Ipomoea hederacea</i>			
Red/Scarlet	<i>Ipomoea coccinea</i>			
Tall	<i>Ipomoea purpurea</i>			
Mustard, Wild	<i>Brassica kaber</i>			
Palmer Amaranth	<i>Amaranthus palmeri</i>			
Spurred Anoda	<i>Anoda cristata</i>			
Tropic Croton	<i>Croton glandulosus</i>			
Waterhemp, ¹		Fine Soils: (silty clay, loam, clay, clay loam)	2 fl oz/A Cotton and Dry Bean 3 fl oz/A Field Corn, Peanut, Soybean, and all other labeled crops	
Common	<i>Amaranthus rudis</i>			
Tall	<i>Amaranthus tuberculatus</i>			
Wild Poinsettia	<i>Euphorbia heterophylla</i>			
Yellow Rocket	<i>Barbarea vulgaris</i>			

- ¹ A postemergence herbicide, including lactofen or glyphosate (Roundup soybeans only) may be needed following a preemergence application of this product to adequately control common ragweed or waterhemp in soybean fields with heavy pressure.
- ² Due to differences in crop canopy timing between peanuts and soybeans, use 3 fluid ounces per acre of this product in peanuts, regardless of soil type and organic matter content, except in the states of North Carolina, Oklahoma and Virginia where a maximum of 2 fluid ounces per acre can be applied in peanuts. This product will provide residual control of these weeds at 2 fluid ounces per acre when applied under a cotton canopy.
- ³ Morningglory species are not adequately controlled on fine soils or soils with greater than 3% organic matter.

Table 2. Weeds Suppressed by Residual Activity of LIBERTY FLUMIOX 44

COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	APPLICATION RATE
BROADLEAF WEED SPECIES		Up to 5%	2.0 to 3.0 fl oz/A
Bristly Starbur	<i>Acanthospermum hispidum</i>		
Copperleaf, Hophornbeam	<i>Acalypha ostryifolia</i>		
Ragweed, Giant	<i>Ambrosia trifida</i>		
Russian Thistle	<i>Salsola iberica</i>		
Smartweeds,			
Ladysthumb	<i>Polygonum persicaria</i>		
Pennsylvania	<i>Polygonum pennsylvanicum</i>		
Smellmelon ¹	<i>Cucumis melo</i>		
Velvetleaf	<i>Abutilon theophrasti</i>		
Wild Buckwheat	<i>Polygonum convolvulus</i>		
Wormwood, Biennial	<i>Artemisia biennis</i>		
GRASS WEED SPECIES			
Barnyardgrass	<i>Echinochloa crus-galli</i>		
Bluegrass, Annual	<i>Poa annua</i>		
Crabgrass, Large	<i>Digitaria sanguinalis</i>		
Foxtail, Giant	<i>Setaria faberi</i>		
Goosegrass	<i>Eleusine indica</i>		
Lovegrass, California	<i>Eragrostis diffusa</i>		
Panicums,			
Fall	<i>Panicum dichotomiflorum</i>		
Texas	<i>Panicum texanum</i>		
Ryegrass, Italian	<i>Lolium multiflorum</i>		
Signalgrass, Broadleaf	<i>Brachiaria platyphylla</i>		
Cheat	<i>Bromus secalinus</i>	Up to 5%	1.5 to 3 fl oz/A
Downy Brome ¹	<i>Bromus tectorum</i>		

¹ Not for use in California

DIRECTIONS FOR USE IN FALL AND SPRING PREPLANT BURNDOWN AND FALLOW SEEDBED PROGRAMS IN FIELD CORN, PEANUT AND SOYBEAN (Preemergence to Crop)
[For Use in the States of Arizona, California and Hawaii Only]

RESTRICTIONS

- Do not apply to frozen or snow covered soil.
- Do not perform any tillage operation after application or residual weed control will be reduced.
- Observe all rotational intervals prior to planting as listed in the "ROTATIONAL RESTRICTIONS" table.

FALL BURNDOWN AND FALLOW SEEDBED PROGRAMS

This product [, at 2 to 4 fluid ounces per acre] can be used in the fall to provide residual weed control in fields that will be planted the following spring with field corn, peanut or soybean (refer to "ROTATIONAL

RESTRICTIONS" table for rates and rotational intervals prior to planting). Weeds controlled by residual activity are listed in **Table 1 (sections A and B), Table 3** and **Table 7**. If weeds have emerged at the time of application, use this product in combination with a labeled burndown herbicide. [Application must be made no earlier than October 15 in Region 2 or November 15 in Region 1 or when soil temperature falls below 50°F at a 2 inch depth to maintain residual weed control into the spring (April 1 in Region 1 and May 1 in Region 2) or up until planting, whichever comes first.] This product can be used in a fall burndown or fallow seedbed program [outside of Regions 1 and 2], however the length of residual control may be variable.

Abnormally warm or wet winters will reduce the length of weed control observed in the spring.

Fall Application Regions:

Region 1: Alabama, Arkansas, Georgia, Kentucky, Mississippi, Oklahoma, Tennessee and Virginia
Region 2: Delaware, Kansas, Illinois, Indiana, Iowa, Maryland, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, Pennsylvania, South Dakota, West Virginia and Wisconsin]

Weeds controlled by postemergence or residual activity are listed in **Table 3**. Preplant burndown treatment tank mixes and rates are:

Herbicide	Application Rate
Program 1¹	
Liberty Flumiox 44 Plus	2 to 3 fl oz/A
Glyphosate Plus	0.5 to 1.0 lb ai/A
2,4-D LVE (2,4-D for use on preplant soybeans only) Plus	0.5 to 1.0 lb ai/A
NIS + AMS	0.5% v/v + 17 lbs/100 gals of water
or	
Program 2¹	
Liberty Flumiox 44 Plus	2 to 3 fl oz /A
Glyphosate Plus	0.5 to 1.0 lb ai/A
COC ² or NIS + AMS	1 pt/A or 0.5% v/v + 17 lbs/100 gals of water
or	
Program 3¹	
Liberty Flumiox 44 Plus	2 to 3 fl oz /A
2,4-D LVE (2,4-D for use on preplant soybeans only) Plus	0.5 to 1.0 lb ai/A
COC	1 pt/A

¹ Dicamba at 0.188 pounds AI per acre can be added to Programs 1, 2 & 3 to assist in the control of emerged broadleaves. Refer to dicamba label for rotational restrictions.

² Crop oil concentrate has been found to increase glyphosate burndown of emerged cutleaf eveningprimrose and Carolina geranium.

Table 3. Weeds Controlled by Fall and Spring Preplant Burndown Programs

WEEDS CONTROLLED ¹		POSTEMERGENCE			RESIDUAL
COMMON NAME	SCIENTIFIC NAME	Program 1	Program 2	Program 3	
Weeds 3 Inches or Less					
Chamomile, False	<i>Matricaria maritima</i>	Yes	Yes	No	Yes
Cheatgrass	<i>Bromus tectorum</i>	Yes	Yes	No	Yes
Chickweed, Common	<i>Stellaria media</i>	Yes	Yes	No	Yes
Chickweed, Mouseear	<i>Cerastium vulgatum</i>	Yes	Yes	No	Yes
Cockle, White	<i>Silene latifolia</i>	No	Yes	Yes	Yes
Dandelion	<i>Taraxacum officinale</i>	Yes	No	Yes ²	Yes
Deadnettle, Purple	<i>Lamium purpureum</i>	Yes	Yes	Yes	Yes
Groundsel, Cressleaf	<i>Senecio glabellus</i>	Yes	Yes	-	Yes
Henbit	<i>Lamium amplexicaule</i>	Yes	Yes	Yes	Yes
Kochia	<i>Kochia scoparia</i>	Yes	Yes	Yes	Yes
Marestail/Horseweed	<i>Conyza canadensis</i>	Yes	Yes ³	Yes	Yes
Mallow, Common	<i>Malva Neglecta</i>	Yes	Yes	No	Yes
Prickly Lettuce	<i>Lactuca serriola</i>	Yes	Yes	Yes	Yes
Wormwood, Biennial	<i>Artemisia biennis</i>	Yes	Yes	Yes	Yes
Weeds 12 Inches or Less					
Canola, Volunteer	<i>Brassica napus</i>	Yes	Yes	Yes	Yes
Carolina Geranium	<i>Geranium carolinianum</i>	Yes	Yes	Yes	-
Eveningprimrose, Cutleaf ⁴	<i>Oenothera laciniata</i>	Yes	Yes	Yes	Yes
Flixweed	<i>Descurainia sophia</i>	Yes	Yes	Yes	Yes
Mustard, Tansy	<i>Descurainia pinnata</i>	Yes	Yes	Yes	Yes
Mustard, Wild	<i>Brassica kaber</i>	Yes	Yes	Yes	Yes
Shepherd's-purse	<i>Capsella bursa-pastoris</i>	Yes	Yes	Yes	Yes

¹ Refer to glyphosate and/or 2,4-D labels for additional weeds controlled and rotational restrictions.

² Use 1 pound AI per acre of 2,4-D LVE for control of emerged dandelion.

³ Program 2 will not control emerged glyphosate resistant marestail/horseweed.

⁴ Use Program 1 to control cutleaf eveningprimrose that are nearing 12 inches in height or are past the rosette stage. Use Programs 2 or 3 to control cutleaf eveningprimrose that are 12 inches or less and in the rosette stage.

SPRING BURNDOWN PROGRAMS

This product may be used in combination with labeled preplant burndown herbicides to assist in the postemergence burndown of emerged weeds and provide residual weed control prior to crop emergence. Weeds controlled by residual activity are listed in **Table 1**.

No-till planters that incorporate the soil during planting may result in decreased weed control in the row. Apply this product after planting peanuts and soybeans when these types of planters are used (within 3 days after planting soybeans, within 2 days after planting peanuts and before the crop emerges). This product cannot be applied after planting field corn.

This product can be used [at 1 to 3 fluid ounces per acre] with labeled preplant burndown herbicides to enhance the speed of burndown and increase weed spectrum.

This product can be used [at 1 to 3 fluid ounces per acre] [1 to 2 fluid ounces per acre] in field corn, peanut and soybean burndown programs. See "DIRECTIONS FOR USE IN FIELD CORN", "DIRECTIONS FOR USE IN PEANUT", "DIRECTIONS FOR USE IN SOYBEAN" for more information.

**DIRECTIONS FOR USE IN FALL AND SPRING BURNDOWN PROGRAMS
IN COTTON AND SUGARCANE
[For Use in the States of Arizona, California and Hawaii Only]**

RESTRICTIONS

- Do not apply to frozen or snow covered soil.
- Do not perform any tillage operation after application or residual weed control will be reduced.
- A minimum of 30 days must pass, and 1 inch of rainfall/irrigation must occur, between application of this product and planting of conventionally tilled cotton.
- A minimum of 14 days must pass, and 1 inch of rainfall/irrigation must occur, between application of this product and planting of no-till or strip-till cotton when a rate of this product at 1 fluid ounce (0.031 lb a.i.) per acre is used and 21 days when a rate of this product at 1.5 to 2 fluid ounces (0.047 to 0.063 lb a.i.) per acre is used. The field must contain the stubble from the previous crop.
- Observe all rotational intervals prior to planting as listed in the "ROTATIONAL RESTRICTIONS" table.
- Refer to most restrictive label for minimum interval between application and planting.

Notes

- This product can be used [at 1 to 2 fluid ounces (0.031 to 0.063 lb a.i.) per acre] with labeled burndown herbicides to enhance the speed of burndown and increase weed spectrum.
- This product can be applied as part of a burndown application to sugarcane until cane emergence.

FALL BURNDOWN PROGRAMS

This product [, at 2 to 4 fluid ounces per acre,] can be used in the fall to provide residual weed control in fields that will be planted the following spring with cotton or sugarcane (refer to "ROTATIONAL RESTRICTIONS" table for rates and rotational intervals prior to planting). Weeds controlled by residual activity are listed in **Table 1** and **Table 7**. If weeds have emerged at the time of application, use this product in combination with a labeled burndown herbicide. [Application must be made no earlier than October 15 in Region 2 or November 15 in Region 1 or when soil temperature falls below 50°F at a 2 inch depth to maintain residual weed control into the spring (April 1 in Region 1 and May 1 in Region 2) or up until planting, whichever comes first.] [This product can be used in a fall burndown or fallow seedbed program outside of Regions 1 and 2.]

Abnormally warm or wet winters will reduce the length of weed control observed in the spring.

SPRING BURNDOWN PROGRAMS

This product [, at 1 to 2 fluid ounces per acre,] can be used in combination with labeled preplant burndown herbicides to assist in the postemergence burndown of emerged weeds and provide residual weed control prior to crop emergence in fields that will be planted with cotton or sugarcane. Weeds controlled by residual activity are listed in **Table 1**.

No-till planters that incorporate the soil during planting may result in decreased weed control in the row.

**DIRECTIONS FOR USE IN FALL AND SPRING BURNDOWN PROGRAMS IN
RICE, SORGHUM, SUNFLOWERS, TOBACCO AND WHEAT
(Preplant to Crop)
[For Use in the States of Arizona, California and Hawaii Only]**

RESTRICTIONS

- Do not apply to frozen or snow covered soil.
- Do not perform any tillage operation after application or residual weed control will be reduced.
- This product can be used [at 1 to 2 fluid ounces (0.031 to 0.063 lb a.i.) per acre] with labeled burndown herbicides to enhance the speed of burndown and increase weed spectrum. A minimum of 30 days must pass, and 1 inch of rainfall/irrigation must occur, between application of this product and planting of rice, sorghum, sugarcane, sunflowers, tobacco or wheat. Refer to most restrictive label for minimum interval between application and planting.
- Observe all rotational intervals prior to planting as listed in the "ROTATIONAL RESTRICTIONS" table.

FALL BURNDOWN PROGRAMS

This product can be used in combination with labeled burndown programs to control emerged weeds and provide residual weed control in fields that will be planted the following spring (refer to "ROTATIONAL RESTRICTIONS" table for rates and rotational intervals prior to planting). [Application must be made no earlier than October 15 in Region 2 or November 15 in region 1 or when soil temperature falls below 50°F at a 2 inch depth to maintain residual weed control into the spring.]

Abnormally warm winters may reduce the length of weed control observed in the spring.

SPRING BURNDOWN PROGRAMS

This product can be used in combination with labeled burndown programs to control emerged weeds and provide residual weed control prior to crop emergence. Weeds controlled by residual activity are listed in **Table 1, Section A**. Crops that will be planted following application must be in compliance with the rotational interval listed in the "ROTATIONAL RESTRICTIONS" table above.

No-till planters that incorporate the soil during planting may result in decreased weed control in the row.

DIRECTIONS FOR USE IN FALL BURNDOWN PROGRAMS IN FIELDS TO BE PLANTED TO BARLEY, FIELD PEA, FLAX, LENTIL, SAFFLOWER, SUNFLOWER AND SPRING WHEAT (Preplant to Crop)

[For Use in the States of Arizona, California and Hawaii Only]

RESTRICTIONS

- Do not apply to frozen or snow covered soil.
- Do not perform any tillage operation after application or residual weed control will be reduced.
- This product can be mixed with 2,4-D and/or glyphosate formulations labeled for burndown programs (preplant to crop) in accordance with the most restrictive label limitations and precautions. Labeled application rates must not be exceeded. Do not mix this product with any product containing a label prohibition against such mixing.
- Observe all rotational intervals prior to planting as listed in the "ROTATIONAL RESTRICTIONS" table.

FALL BURNDOWN PROGRAMS

This product can be used [at 2 to 4 fluid ounces per acre] with labeled burndown herbicides to enhance the speed of burndown, increase weed spectrum and provide residual weed control of the weeds listed in **Table 3** until the following spring. Rotational intervals must be followed for crop to be planted in the spring following the fall application of this product. Refer to most restrictive label for minimum interval between application and planting. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

DIRECTIONS FOR USE IN FALLOW LAND [For Use in the States of Arizona, California and Hawaii Only]

This product may be used as a preemergence fallow treatment. Weeds controlled by residual activity are listed in **Table 1**. This product [, at 2 to 4 fluid ounces per acre,] can be used in the fall to provide residual weed control in fallow fields (refer to "ROTATIONAL RESTRICTIONS" table for rates and rotational intervals prior to planting). If weeds have emerged at the time of application, use this product in combination with a labeled fallow herbicide. [Application must be made no earlier than October 15 in Region 2 or November 15 in Region 1 or when soil temperature falls below 50°F at a 2 inch depth to maintain residual weed control into the spring (April 1 in Region 1 and May 1 in Region 2).] Abnormally warm or wet winters will reduce the length of weed control observed in the spring.

This product [, at 1 to 4 fluid ounces per acre,] can be used in spring in combination with labeled burndown herbicides to control emerged weeds and provide residual weed control.

**DIRECTIONS FOR USE IN FALLOWBED USE ON TRANSPLANTED
MELON, PEPPER AND TOMATO BEDS
[For Use in the States of Arizona, California and Hawaii Only]**

RESTRICTIONS

- Do not apply more than 4 fluid ounces (0.125 lb a.i.) of this product per acre per application.
- Do not apply more than 4 fluid ounces (0.125 lb a.i.) of this product per acre per year.
- Do not make more than 1 application per year.

Many weather related factors, including high wind or heavy rains or cool conditions at or near crop transplanting, may result in crop injury in fields treated with this product. On occasion this has resulted in a delay in maturity. User assumes these risks before using this product.

TIMING TO CROP


LIBERTY FLUMIOX 44 Fallowbed Use Prior To Transplanting

APPLICATION RATE	ADJUVANT	GPA	TRANSPLANTING INTERVAL
4 fl oz/A	Required by burndown tank mix partner	Ground — 20 to 40	2 Months


Application Method: Apply with a burndown herbicide labeled for the control of emerged weeds. When using this product alone satisfactory control of emerged weeds will not be attained.

USE RESTRICTIONS FOR PREEMERGENCE FALLOWBED WEED CONTROL PRIOR TO TRANSPLANTING


1. Always read and follow all label directions when using any pesticide alone or in tank mix combinations.
2. The top 4 inches of the bed, from a horizontal and vertical perspective, where the crop will be transplanted, must be removed prior to transplanting.
3. Use only healthy transplants. Do not use on direct seeded crops.
4. [On flat beds (tomato only), the soil must be incorporated to a depth of at least 4 inches, twice, prior to transplanting. Failure to incorporate may result in stand reduction and/or crop injury.]
5. This use pattern makes no claim for in-season weed control after the beds have been disturbed.
6. Do not apply when weather conditions favor spray drift.



Beds are formed and this product is applied with a burndown herbicide.



A minimum of 2 months after application of this product, the tops of the beds are removed and the soil from the tops of the beds is placed in the area between the beds.



Crops are transplanted into beds.

DIRECTIONS FOR USE IN ESTABLISHED ALFALFA

RESTRICTIONS

- Do not apply more than 4 fluid ounces (0.125 lb a.i.) of this product per acre per application.
- Do not apply more than 8 fluid ounces (0.250 lb a.i.) of this product per acre per year.
- Do not make more than 2 applications per year.
- Do not make a sequential application of this product within 60 days of the first application of this product.
- Do not apply to alfalfa with greater than 6 inches of growth. Application will result in burning of treated leaves and stems. Users must understand and accept this risk before using this product on alfalfa.
- Do not apply within 25 days of harvest or grazing.
- Do not use on alfalfa grown for seed unless approved by a State authority to support a Special Local Need (SLN) under FIFRA section 24(c).
- Do not use on intended mixed alfalfa-grass stands.

Notes

- Application with paraquat can be used to burndown winter annuals prior to winter dormant period.
- Only apply with an adjuvant or tank mix with products formulated as an emulsifiable concentrate "EC" when targeting control of emerged weeds (crop burn and/or stunting should be expected and accepted if this product is used with an adjuvant, a tank mix partner formulated as an emulsifiable concentrate (EC) or a tank mix partner formulated with an adjuvant.)

TIMING TO ALFALFA

This product may be applied to established alfalfa with a maximum amount of growth of 6 inches or less for the preemergence control of the weeds listed in **Table 7**. Established alfalfa is defined as alfalfa planted in the fall or spring which has gone through a first cutting/mowing. Application to alfalfa with greater than 6 inches of growth may result in unacceptable crop injury.

For control of winter annual weeds: the best timing for preemergence control is in the fall immediately after the last cutting or sheepling-off has occurred.

For control of summer annual weeds: the best timing for preemergence control is in the spring prior to alfalfa growth and before 6 inches of growth.

TIMING TO WEEDS

Preemergence – Preemergence to Weeds

Apply this product before alfalfa growth exceeds 6 inches in height for the preemergence control of weeds listed in **Table 7**. Make applications as soon as possible after cutting and removing alfalfa to minimize injury to alfalfa growth.

Postemergence Dodder¹ Suppression

Apply this product at 4 fluid ounces per acre with an adjuvant for postemergence suppression of dodder¹. Tank mixes with imazamox or imazethapyr will increase control.

¹ Not for use in California.

DIRECTIONS FOR USE IN ARTICHOKE

RESTRICTIONS

- Do not apply more than 4 fluid ounces (0.125 lb a.i.) of this product per acre per application on annual or perennial artichoke varieties after new planting.
- Do not apply more than 6 fluid ounces (0.188 lb a.i.) of this product per acre per application on perennial artichoke varieties after cutback.
- Do not apply more than 6 fluid ounces (0.188 lb a.i.) of this product per acre per year.
- Do not make more than 2 applications per year.

PRECAUTIONS

- Application to artichoke foliage may result in unacceptable crop injury.

TIMING TO ARTICHOKE

Annual Varieties: This product may be applied to artichoke beds prior to transplanting. Application of this product must be made to the beds no later than 2 days prior to transplanting. Irrigation or rainfall after transplanting is necessary to activate this product. Do not irrigate this product before transplanting. Heavy irrigation or rainfall may result in crop injury. The injury is usually transitory and the plants will quickly grow out of the crop damage. Take care to minimize soil disturbance during transplanting, as preemergence weed control will decrease as soil disturbance increases.

Perennial Varieties: This product may be applied to artichokes after planting of crown pieces or "cut back" of mature plants. Applications of this product must be made within 2 days after planting or cut back and prior to artichoke emergence. Application after the artichokes have begun to crack, or are emerged, will result in crop injury. Do not apply when artichokes have begun to emerge (cracking).

TIMING TO WEEDS

Pre-plant (annual)/Preemergence (perennial) to Artichokes - Preemergence to Weeds

Apply this product pre-plant to annual artichokes for preemergence control of the weeds. For perennial artichokes apply before cracking for preemergence control the weeds. Apply prior to weed emergence. A post-emergence herbicide may be necessary to control emerged weeds. This product may be applied to annual or perennial artichokes as specified above for preemergence control of weeds listed in **Table 7**.

DIRECTIONS FOR USE IN ESTABLISHED ASPARAGUS

RESTRICTIONS

- Do not apply more than 6 fluid ounces (0.188 lb a.i.) of this product per acre per application.
- Do not apply more than 6 fluid ounces (0.188 lb a.i.) of this product per acre per year.
- Do not make more than 1 application per year.
- Apply only to dormant asparagus no less than 14 days before spears emerge. Application to non-dormant asparagus may result in unacceptable crop injury.

Note

- Do not work soil within 60 days prior to application in the spring. Soil can be worked after spear harvest in preparation for applications of this product prior to fern emergence. Treated soil that is splashed onto the ferns may result in spotting.

TIMING TO ASPARAGUS – Dormant

This product may be applied to dormant asparagus for preemergence control of the weeds listed in **Table 10**. Application to non-dormant asparagus will result in unacceptable crop injury. Apply no less than two weeks prior to spear emergence and must be sprinkler or rainfall incorporated with 0.5 to 0.75 inches of water or some scoring may result.

TIMING TO ASPARAGUS - Post Harvest

Apply this product after the final harvest of the season, but prior to fern emergence, for preemergence control of the weeds listed in **Table 10**. Application after fern emergence will result in unacceptable crop injury. Apply no less than two weeks prior to fern emergence and must be sprinkler or rainfall incorporated with 0.5 to 0.75 inches of water. Add a burndown tank mix partner for the control of emerged weeds labeled for asparagus in accordance with the most restrictive labeled limitations and precautions.

TIMING TO WEEDS

Burndown - Dormant Asparagus, Postemergence to Weeds

This product may be used for residual weed control, as well as to assist in postemergence burndown of many annual and perennial weeds where asparagus is dormant. For control of emerged weeds, tank mix this product with paraquat. Refer to paraquat label for directed rate and application parameters. To ensure thorough coverage, use a minimum of 15 gallons of spray solution per acre. Tank mixes of this product applied to assist in the control of emerged weeds must be applied with a non-ionic surfactant at 0.25% v/v. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 pounds per acre or 28 to 32% nitrogen solution at 1 to 2 quarts per acre) may be added to increase herbicidal activity.

Burndown - After Last Harvest of Season, Postemergence to Weeds

Use this product for residual weed control and to assist in postemergence burndown for many annual and perennial weeds where asparagus harvest has been completed for the year. For control of emerged weeds, use a labeled tank mix partner with activity on the emerged weeds.

Preemergence - Dormant Asparagus or After Last Harvest of Season, Preemergence to Weeds

Apply this product to dormant asparagus for the preemergence control of weeds listed in **Table 10**.

DIRECTIONS FOR USE ON BRASSICA HEAD AND STEM VEGETABLE GROUP 5-16 ROW MIDDLES

Includes: Broccoli; Brussels sprouts; Cabbage; Cabbage, Chinese, Napa; Cauliflower; cultivars, varieties and/or hybrids of these.

[FOR DISTRIBUTION AND USE ONLY WHERE THIRD PARTY INDEMNIFICATION IS IN EFFECT]

DIRECTIONS FOR USE IN ROW MIDDLES

RESTRICTIONS

- Do not apply after crops are transplanted.
- Do not apply more than 3 fluid ounces (0.093 lb a.i.) per acre per application. For Cabbage do not apply more than 4 fluid ounces (0.125 lb a.i.) per acre per application.
- Do not apply more than 6 fluid ounces (0.188 lb a.i.) per acre per year. For Cabbage do not apply more than 8 fluid ounces (0.250 lb a.i.) per acre per year.
- Do not make more than 2 applications per year.

PRECAUTIONS

- Injury can occur if soil particles treated with this product contact the crop.

Notes

- This product can only be applied in row middles between raised plastic mulched beds that are at least **4 inches higher** than the treated row middle and the mulched bed must have **a minimum of a 24-inch bed width**.
- Spray must remain between raised beds and contact no more than the bottom 1 inch of the side of the raised bed.
- All applications must be made with shielded or hooded equipment.
- A rainfall after application but prior to transplanting is required.

APPLICATION RATE

Do not exceed more than 4 fluid ounces of this product per acre.

TIMING TO CROP

This product may be applied at 3 fluid ounces per acre (except cabbage may be applied at 4 fluid ounces per acre) as a shielded or hooded application to row middles after plastic is laid up to transplanting. Transplanting or seeding can take place any time after spray has dried. Spray must be directed to the row middle and contact no more than approximately the bottom 1 inch of the side of the raised bed. If the top of the mulch beds (where plants are to be transplanted) is contacted, severe injury can occur due to foliage contact with treated plastic.

WEED CONTROL AND TANK MIXING

This product provides preemergence residual control of the weeds listed in **Table 7**, as well as to assist in the postemergence control of emerged weeds. A registered preemergence grass herbicide may be added for control of additional grassy weeds. For control of emerged weeds, tank mix this product with carfentrazone, glyphosate, paraquat, or other registered burndown herbicide. Refer to tank mix partner label for directed rates. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

DIRECTIONS FOR USE ON CACTUS (PRICKLY PEAR)

RESTRICTIONS

- Do not apply more than 12 fluid ounces (0.375 lb a.i.) of this product per acre per application.
- Do not apply more than 12 fluid ounces (0.375 lb a.i.) of this product per acre per year.
- Do not make more than 2 applications per year.
- Use a maximum rate of 6 fluid ounces (0.188 lb a.i.) per acre per application of this product on any soil that has a sand plus gravel content over 80% if plants are less than 3 years of age. (Two applications of 6 fluid ounces per acre in a year can still be made as long as there have been 60 days between applications).
- Do not apply to farm alleys or roads where traffic may result in treated dust settling onto crops or other desirable vegetation.
- Do not mow treated areas. Dust created by mowing may drift onto desirable vegetation resulting in injury.
- Follow the most restrictive label limitations and precautions of the tank mix product(s) being used.
- Do not apply within 60 days prior to harvest.
- Do not apply to plants established less than one year.

PRECAUTIONS

- Raise mower height during all mowing to reduce dust. Dust created by mowing can drift onto desirable vegetation resulting in injury.
- Avoid direct or indirect spray contact to foliage or injury may occur.

Apply this product as a uniform broadcast application to the plantation floor or as a uniform band directed at the base of the cactus. The preferred application timing for this product is in the fall to maximize the potential for rainfall to activate and set the herbicide. Do not apply over the top of crop or allow spray to come in contact with crop as a result of application or drift.

Preemergence Application

Apply 6 to 12 fluid ounces of this product per broadcast acre as a preemergence application. Applications of this product must be made prior to weed emergence for control of weeds listed in **Table 10**. Make preemergence (to weed emergence) applications of this product to a weed-free soil surface. Preemergence applications of this product must be completed prior to weed emergence. Moisture is necessary to activate this product on soil for residual weed control. Dry weather following application of this product may reduce effectiveness. However, when adequate moisture is received after dry conditions, this product will control susceptible germinating weeds.

[Postemergence Application

Apply 6 to 12 fluid ounces of this product per broadcast acre plus an adjuvant (0.25% v/v non-ionic surfactant or 1 quart per acre crop oil concentrate). The addition of an adjuvant enhances the activity of this product on emerged weeds. Thorough spray coverage is necessary to maximize the postemergence activity of this product.

Refer to **Table 7** for weeds controlled by the residual activity of this product. Tank mix this product with a labeled burndown herbicide for control of the emerged weeds.

Residual weed control will be reduced if vegetation prevents this product from reaching the soil surface. If vegetation is heavy, use a burndown herbicide with this product and make sequential applications of this product prior to the emergence of new weeds.]

Carrier Volume and Spray Pressure

To ensure thorough coverage in burndown applications, use a minimum of 15 gallons of spray solution per acre. Use higher gallonage if dense vegetation or heavy crop residue is present. Nozzle selection must meet manufacturer's gallonage and pressure specifications.

Banded Application

Rates listed in **Table 13**, refer to a broadcast application covering the entire acre. Refer to the Band Application table in Use Information Section to calculate amount needed per acre when making a banded application.

DIRECTIONS FOR USE IN CELERY **[For Use in the States of [California], Michigan and Wisconsin Only]**

RESTRICTIONS

- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre during a pre-transplant application.
- [In the state of California, use as pre-transplant application only.]
- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre during a post-transplant application.
- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre per year.
- Do not make more than 1 application per year.
- Do not use with an adjuvant.
- Post transplant applications must be made between 3 to 7 days following transplanting.
- Do not apply as part of a tank mix.

TIMING TO CELERY

Apply this product at 3 fluid ounces per acre prior to transplanting, or between 3 and 7 days following transplanting, for preemergence control of the weeds listed in **Table 1**.

TIMING TO WEEDS

Use this product prior to weed emergence for residual control.

Refer to Product Information section for tank mix guidance. This product, when applied according to label use directions, will control the weeds listed in **Table 1**. This label makes no claims concerning control of other weed species.

DIRECTIONS FOR USE IN ESTABLISHED CLOVER **For Use in Idaho, Oregon and Washington Only**

RESTRICTIONS

- Do not apply more than 4 fluid ounces (0.125 lb a.i.) per acre per application.
- Do not apply more than 4 fluid ounces (0.125 lb a.i.) per acre per year.
- Do not make more than 1 application per year.
- Do not apply within 25 days of harvest or grazing.
- Do not use on intended mixed clover-grass stands.

PRECAUTIONS

- Do not apply to clover with greater than 6 inches of growth. Application will result in burning of treated leaves and stems. Users must understand and accept this risk before using this product on clover.
- Application to clover with greater than 6 inches of growth may result in unacceptable crop injury.

Notes

- Only apply with an adjuvant or tank mix with products formulated as an emulsifiable concentrate “EC” when targeting control of emerged weeds (expect and accept crop may be burned and/or stunting when applying tank mixes of this product with an adjuvant).
- Application with paraquat can be used to burndown winter annuals prior to winter dormant period.

TIMING TO CLOVER

This product may be applied to established clover with a maximum amount of growth of 6 inches or less for the preemergence control of the weeds listed in **Table 7**. Established Clover is defined as clover planted in the fall or spring which has gone through a first cutting/mowing.

For control of winter annual weeds: the best timing for preemergence control is in the fall immediately after the last cutting or sheepling-off has occurred.

For control of summer annual weeds: the best timing for preemergence control is in the spring prior to clover growth and before 6 inches of growth.

TIMING TO WEEDS

Preemergence – Preemergence to Weeds

Apply this product before clover growth exceeds 6 inches in height for the preemergence control of weeds listed in **Table 7**. Apply as soon as possible after cutting and removing clover to minimize injury to clover growth.

Postemergence Dodder Suppression

Apply this product at 4 fluid ounces per acre with an adjuvant for postemergence suppression of dodder.

Tank mixes with Imazethapyr or Imazamox will increase control. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Read and follow applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

DIRECTIONS FOR USE IN COTTON **[For Use in the States of Arizona, California and Hawaii Only]**

RESTRICTIONS

- Do not apply more than 2 fluid ounces (0.063 lb a.i.) of this product per acre per application.
- Do not apply more than 4 fluid ounces (0.125 lb a.i.) of this product per acre per year.
- Do not make more than 2 applications per year.
- Do not make a sequential application of this product within 30 days of the first application of this product.
- Do not apply within 60 days of harvest.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL PERFORMANCE

Hooded, Shielded and Layby Application

For best results, apply this product to actively growing weeds within the growth stages indicated in this label. Applying this product under conditions that do not promote active weed growth will reduce herbicide effectiveness. Do not apply this product when the crop or weeds are under stress due to drought, excessive water, extremes in temperature, disease or low humidity. Weeds under stress tend to become less susceptible to herbicidal action. This product is most effective when applied under sunny conditions at temperatures above 65°F.

This product is rainfast one hour after application. Do not apply if rain is expected within one hour of application or postemergence efficacy may be reduced. Rainfall within one hour of application will not adversely affect residual activity.

HERBICIDE RATE

Hooded, Shielded and Layby Application

For postemergence weed control, apply this product through a hooded or shielded sprayer or at layby, at 2 fluid ounces per acre, in combinations with MSMA or at 1 to 2 fluid ounces per acre in combination with glyphosate, to assist in the control of weeds listed in **Table 4**. Residual weed control can also be obtained through hooded, shielded and layby application of this product. Weeds that are controlled through residual activity of this product are listed in **Table 1**. Weeds that are suppressed by residual activity of this product are listed in **Table 2**.

Table 4. Emerged Broadleaf Weeds Controlled by Hooded, Shielded and Layby Application of Tank Mixes of This Product with Glyphosate or MSMA in Cotton

COMMON NAME	SCIENTIFIC NAME	WEED HEIGHT (inches) 2 fl oz/A
BROADLEAF WEED SPECIES		
Bindweed, Field ¹	<i>Convolvulus arvensis</i>	4
Carpetweed	<i>Mollugo verticillata</i>	4
Chickweed, Common	<i>Stellaria media</i>	4
Cocklebur, Common	<i>Xanthium strumarium</i>	4
Florida Beggarweed	<i>Desmodium tortuosum</i>	2
Hemp Sesbania	<i>Sesbania exaltata</i>	6
Jimsonweed	<i>Datura stramonium</i>	4
Lambsquarters, Common	<i>Chenopodium album</i>	4
Morningglories,		
Entireleaf	<i>Ipomoea hederacea</i> var. <i>integriuscula</i>	4
Ivyleaf	<i>Ipomoea hederacea</i>	4
Pitted	<i>Ipomoea lacunose</i>	4
Red	<i>Ipomoea coccinea</i>	4
Tall	<i>Ipomoea purpurea</i>	2
Mustard, Wild	<i>Brassica kaber</i>	6
Nightshades,		
Black	<i>Solanum nigrum</i>	4
Eastern Black	<i>Solanum ptycanthum</i>	4
Hairy	<i>Solanum sarrachoides</i>	4
Pigweeds,		
Palmer Amaranth	<i>Amaranthus palmeri</i>	4
Redroot	<i>Amaranthus retroflexus</i>	4
Smooth	<i>Amaranthus hybridus</i>	4
Plantain, Broadleaf	<i>Plantago major</i>	6
Prickly Sida (Teaweed)	<i>Sida spinosa</i>	4
Purslane, Common	<i>Portulaca oleracea</i>	2
Ragweeds,		
Common	<i>Ambrosia artemisiifolia</i>	2
Giant	<i>Ambrosia trifida</i>	4
Rice Flatsedge	<i>Cyperus iria</i>	2
Sicklepod	<i>Senna obtusifolia</i>	4
Smartweeds,		
Ladysthumb	<i>Polygonum persicaria</i>	4
Pale	<i>Polygonum lapathifolium</i>	4
Pennsylvania	<i>Polygonum pennsylvanicum</i>	4
Spotted Spurge	<i>Euphorbia maculata</i>	4
Velvetleaf	<i>Abutilon theophrasti</i>	4
Venice Mallow	<i>Hibiscus trionum</i>	2
Waterhemp,		
Common	<i>Amaranthus rudis</i>	2
Tall	<i>Amaranthus tuberculatus</i>	2

¹ Tank mixes of this product will control the above ground portion of field bindweed. Repeated applications will be needed to control regrowth.

**CARRIER VOLUME AND SPRAY PRESSURE
Hooded, Shielded and Layby Application**

To ensure thorough coverage in hooded, shielded and layby applications, use 15 to 30 gallons spray solution per treated acre. Use 20 to 30 gallons per treated acre under heavy weed pressure. Nozzle

selection must meet manufacturer's gallonage and pressure specifications for application method being used. Do not use "Flood Jet" nozzles, as they tend to increase the chance of crop injury.

ADDITIVES

Hooded, Shielded and Layby Application

Weed control from hooded, shielded or layby application of this product in cotton requires the addition of an agronomically approved non-ionic surfactant to the spray mixture. Non-ionic surfactant must contain at least 80% active ingredient. Verify mixing compatibility qualities by a jar test. **The use of crop oil concentrates, methylated seed oils, organo-silicant surfactants or products containing these ingredients may result in severe crop injury.**

APPLICATION EQUIPMENT

Apply tank mixes of this product, with ground equipment using standard commercial sprayers equipped with nozzles designed to deliver the desired spray pressure and spray volume. Application equipment must be clean and in good repair. Nozzles must meet manufacturer's specifications for spray pattern and placement on spray boom and must be checked frequently for accuracy.

TIMING TO COTTON

Hooded and Shielded Application

Tank mixes of this product may be applied with a hooded or shielded sprayer after cotton has reached a minimum of 6 inches in height. All nozzles must be under the hood or behind the shield to ensure no spray solution comes in contact with the cotton. **Care must be taken to ensure the spray solution or drift does not come in contact with the cotton or severe crop injury can occur.**

Layby Application

Layby application of tank mixes of this product may be made once cotton has reached a minimum of 16 inches in height. Cotton that is smaller than 16 inches in height may be injured by applications of this product. Application of this product must be directed to the lower 2 inches of the cotton stem to avoid crop injury.

TIMING TO WEEDS

Tank mix applications of this product must be made to weeds within the height range given in **Table 4**.

TANK MIXES

This product must be tank mixed with one of the herbicides listed in **Table 5** for postemergence control of the weeds listed in **Table 4**. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Table 5. Tank Mixes for Hooded, Shielded and/or Layby Use in Cotton

TANK MIX PARTNER	TARGET WEEDS	HOODED AND SHIELDED	LAYBY
Glyphosate	Perennial Grasses and Broadleaves	X	X ¹
MSMA	Annual Grasses Yellow Nutsedge	X	X

¹ For use only in cotton with the Roundup Ready gene.

DIRECTIONS FOR USE IN CUCURBIT VEGETABLES

Cucurbit Vegetables (Crop Group 9) including: chayote (fruit); Chinese Waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); *Momordica* spp. (includes balsam apple, balsam pear, bittermelon, Chinese cucumber); muskmelon (includes cantaloupe); pumpkin; squash, summer; squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon

Many weather related factors, including high wind or heavy rains or cool conditions at or near crop transplanting, may result in crop injury in fields treated with this product. On occasion this has resulted in a delay in maturity. User assume these risks before using this product.

Refer to Product Information section for tank mix guidance. This product, when applied according to label use directions, will control the weeds listed in **Table 7**. This label makes no claims concerning control of other weed species.

RESTRICTIONS

- Do not apply more than 4 fluid ounces (0.125 lb a.i.) of this product per acre per application.
- Do not apply more than 8 fluid ounces (0.250 lb a.i.) of this product per acre per year.
- Do not make more than 2 applications per year.
- Do not apply during or after bloom.

DIRECTIONS FOR USE IN ROW MIDDLES

[FOR DISTRIBUTION AND USE ONLY WHERE THIRD PARTY INDEMNIFICATION IS IN EFFECT]

RESTRICTIONS

- Do not use with an adjuvant.
- Spray must be directed to the row middle, away from the crop bed and with minimal contact with plastic, including the sides of the bed. If top of mulch beds (where plants are to be transplanted) is contacted, severe injury can occur due to foliage contact with treated plastic. In this scenario, a rainfall event of 1/2 inch (natural or irrigation) must occur prior to transplanting to reduce residues of this product.
- **[Arizona, California and Hawaii only:** For fallow bed application on transplanted peppers and tomato beds see “DIRECTIONS FOR USE FALLOWBED USE ON TRANSPLANTED MELON, PEPPER AND TOMATO BEDS” use instructions.]
- Irrigate treated field after application and prior to transplanting with minimum of 1/4 inch of water if rainfall does not occur between application and transplanting.
- All applications must be made with hooded or shielded equipment.

PRECAUTIONS

- Drift of treated soil particles onto plants may cause contact injury.

Note

- Grow plants on raised plastic mulched beds that are higher than the treated row middle.

TIMING TO CUCURBIT VEGETABLES

Apply this product at 4 fluid ounces per acre as a hooded or shielded application to row middles up to 14 days prior to transplanting or seeding for preemergence control of the weeds listed in **Table 7**, as well as to assist in the postemergence control of emerged weeds. A second application of this product at 4 fluid ounces per acre may be applied up to 21 days after transplanting or emergence if needed. Do not apply during or after bloom.

TIMING TO WEEDS

This product may be used for residual weed control, as well as to assist in postemergence burndown of many annual and perennial weeds in row middles. A registered preemergence grass herbicide may be added for control of additional grassy weeds. For assisting in the control of emerged weeds, tank mix this product with carfentrazone, paraquat, or other registered burndown herbicide. Do not tank mix with glyphosate after transplanting.

DIRECTIONS FOR USE IN DRY BEANS

[Dried cultivars of bean (*Lupinus*); bean (*Phaseolus*) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); bean (*Vigna*) (includes adzuki bean, blackeyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean); broad bean (dry); chickpea (garbanzo bean); guar; lablab bean and lentil]

WEED SUPPRESSION IN DRY BEANS AND WEED CONTROL IN CHICKPEAS (GARBANZO BEANS) [Arizona, California, Colorado, Hawaii, Idaho, Montana, Oregon and Washington only]

RESTRICTIONS

- For Chickpeas, do not apply more than 2 fluid ounces (0.063 lb a.i.) of this product per acre per application. For all other Dry Beans, do not apply more than 1.5 fluid ounces (0.047 lb of this product per acre application).
- For Chickpeas, do not apply more than 2 fluid ounces (0.063 lb a.i.) of this product per acre per year. For all other Dry Beans, do not apply more than 1.5 fluid ounces (0.047 lb a.i.) of this product per acre per year.
- Do not make more than 1 application per year.

Many weather related factors, including high wind, splashing or heavy rains or cool conditions at or near crop emergence, may result in dry bean injury in fields treated with this product. On occasion this has resulted in a delay in maturity. User assumes these risks before using this product.

TIMING TO DRY BEANS AND CHICKPEAS

This product may be applied to dry beans within 2 days after planting for the preemergence suppression of the weeds listed in **Table 1** or **Table 8**. Tank mix this product with other labeled herbicides for broad spectrum weed control.

TIMING TO WEEDS

This product may be applied to dry beans prior to planting or preemergence (after planting). Preemergence application of this product must be made within 2 days after planting and prior to dry bean emergence. To avoid severe crop injury, do not apply to dry beans after beans begin to crack or have emerged.

Preplant incorporation (PPI) applications may result in reduced weed control.

ADDITIONAL RESIDUAL GRASS CONTROL

This product can be tank mixed with pendimethalin for additional grass control. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

HARVEST AID RESTRICTIONS

- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre application.
- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre per year.
- Do not make more than 2 applications per year.
- Do not harvest within 5 days of application.

Desiccation from this product requires the addition of an agronomically approved adjuvant to the spray mixture. Use a methylated seed oil which contains at least 15% emulsifiers and 80% oil at 2% v/v. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 pounds per acre or a 28 to 32% nitrogen solution at 1 to 2 quarts per acre) may be added to the spray mixture along with either a crop oil concentrate or methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for a crop oil concentrate or a methylated seed oil. Tank mixing this product with glyphosate or paraquat will increase control of emerged weeds and aid in harvest. Add a burndown tank mix partner for the control of emerged weeds labeled for dry bean in accordance with the most restrictive labeled limitations and precautions.

TIMING TO DRY BEANS AND CHICKPEAS

Apply when crop is mature and at least 80% of the pods are yellowing and mostly ripe with no more than 40% (bush type beans) or 30% (vine type beans) of the leaves still green in color. Dry beans can be harvested 5 days after application. To ensure thorough coverage use 15 to 30 gallons spray solution per

acre. Nozzle selection must meet manufacturer's gallonage and pressure specifications for postemergence application.

DIRECTIONS FOR USE IN FIELD CORN
[For Use in the States of Arizona, California and Hawaii Only]

RESTRICTIONS

- Use only on no-till or minimum tillage fields where last year's crop residue has not been incorporated into the soil.
- Corn must be planted between 14 and 30 days after application unless the application is made as part of a Fall burndown program.
- Corn can be planted 7 days after an application of 2 fluid ounces (0.063 lb a.i.) per acre if a minimum of 25% of the soil surface is covered with the residue of the preceding crop and a minimum of 1/4 inch of rainfall has occurred between application and planting.
- Do not apply more than [2 fluid ounces (0.063 lb a.i.)] [3 fluid ounces (0.093 lb a.i.)] of this product per acre per year.
- Do not make more than 1 application per year.
- Do not irrigate between emergence and 2-leaf corn
- Do not use on popcorn, sweet corn or corn grown for seed.

TIMING TO FIELD CORN

- Apply this product, at 2 to 3 fluid ounces per acre, between 7 and 30 days prior to planting field corn, for the preemergence control of the weeds listed in **Table 1**.
- Apply this product at 2 fluid ounces per acre between 7 and 30 days prior to planting field corn if a minimum of 25% of the soil surface is covered with the residue of the preceding crop and a minimum of 1/4 inch of rainfall has occurred between application and planting.
- Apply this product at 3 fluid ounces per acre between 14 and 30 days prior to planting field corn.

Burndown Use Directions - For Preplant Applications in Field Corn

This product, applied as part of a burndown program, may be used for residual weed control, as well as to assist in postemergence burndown of many weeds where field corn will be planted directly into the residue of the previous year. See "DIRECTIONS FOR USE IN FALL AND SPRING PREPLANT BURNDOWN AND FALLOW SEEDBED PROGRAMS IN FIELD CORN, PEANUT AND SOYBEAN" for rates and timing of applications. For control of emerged weeds, this product must be applied with an appropriate burndown tank mix partner listed in **Table 6**. To ensure thorough coverage, use a minimum of 15 gallons of spray solution per acre. Refer to tank mix partner's label for specified application pressure and adjuvant systems.

INCREASING SPEED OF GLYPHOSATE BURNDOWN ACTIVITY

This product, at 1 fluid ounce per acre, may be tank mixed with glyphosate to increase the speed of burndown activity compared to glyphosate applied alone. Residual weed control will not be provided at rates lower than 2 fluid ounces per acre; however, suppression of the weeds in **Table 2** may occur at rates of this product as low as 1 fluid ounce per acre. Applications of this product at 1 fluid ounce per acre must be made a minimum of 14 days prior to planting field corn.

TANK MIXES

This product may be tank mixed with the herbicides listed in **Table 6** for pre-plant burndown applications. Refer to tank mix partner's label for adjuvant directions. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Table 6. Tank Mix Partners for Burndown and/or Residual Control of Weeds in Field Corn

TANK MIX PARTNERS¹	
2,4-D LVE	Metribuzin
Atrazine	Paraquat

Clopyralid + Flumetsulam	Rimsulfuron
Dicamba	Rimsulfuron + Thifensulfuron methyl
Dicamba + 2,4-D	Simazine
Flumetsulam	Tribenuron methyl
Glyphosate	

¹ Refer to tank mix product labels for directions.

TANK MIX RESTRICTIONS

Tank mixes with alachlor, acetochlor, dimethenamid/dimethenamid-p, flufenacet or metolachlor/s-metolachlor may result in injury to field corn when application is followed by prolonged periods of cool wet weather and must not be used with this product.

DIRECTIONS FOR USE IN FIELD PEAS [For use in Idaho, Montana, Oregon and Washington only.]

WEED CONTROL RESTRICTIONS

- Do not apply more than 2 fluid ounces (0.063 lb a.i.) of this product per acre per application.
- Do not apply more than 2 fluid ounces (0.063 lb a.i.) of this product per acre per year.
- Do not make more than 1 application per year.

Many weather related factors, including high wind, splashing or heavy rains or cool conditions at or near crop emergence, may result in pea injury in fields treated with this product. On occasion this has resulted in a delay in maturity. User assumes these risks before using this product.

TIMING TO FIELD PEAS

This product may be applied to field peas within 2 days after planting for the preemergence control of the weeds listed in **Table 1** or **Table 2**. Tank mix this product with other labeled herbicides for broad spectrum weed control. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

TIMING TO WEEDS

This product may be applied to field peas prior to planting or preemergence (after planting). Preemergence application of this product must be made within 2 days after planting and prior to field pea emergence. To avoid severe crop injury, do not apply to field peas after peas begin to crack or have emerged.

Preplant incorporation (PPI) applications may result in reduced weed control.

ADDITIONAL RESIDUAL GRASS CONTROL

This product can be tank mixed with pendimethalin for additional grass control. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

HARVEST AID RESTRICTIONS

- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre per application.
- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre per year.
- Do not make more than 1 harvest aid application per year.
- Do not harvest within 5 days of application.

Desiccation from this product requires the addition of an agronomically approved adjuvant to the spray mixture. Use a methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 quart per acre. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lbs per acre or a 28 to 32% nitrogen solution at 1 to 2 quarts per acre) may be added to the spray mixture along with methylated seed oil to

enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil. Tank mixing this product with glyphosate will increase control of emerged weeds and aid in harvest.

TIMING TO FIELD PEAS

Apply this product, at 1.5 to 2 fluid ounces per acre, when crop is physiologically mature and a minimum of 80% of the pods are yellow to tan in color and 20% are yellow in color. If field peas are treated too early, a reduction in seed quality may occur. Do not spray this product on any area of the field with a significant amount of plants with green color. Peas can be harvested 5 days after application.

To ensure thorough coverage, use 15 to 30 gallons of spray solution per acre and select nozzle type using manufacturer's gallonage and pressure specifications for postemergence application.

DIRECTIONS FOR USE IN FLAX

HARVEST AID RESTRICTIONS

- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre per application.
- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre per year.
- Do not make more than 1 application per year.
- Do not harvest within 5 days of application.

Desiccation from this product requires the addition of an agronomically approved adjuvant to the spray mixture. Use a methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 quart per acre. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 pounds per acre or 28 to 32% nitrogen solution at 1 to 2 quarts per acre) may be added to the spray mixture along with methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil.

TIMING TO FLAX

Apply this product, at 1.5 to 2 fluid ounces per acre, when crop is physiologically mature and at least 75% of the bolls are brown in color. Flax can be harvested 5 days after application.

To ensure thorough coverage, use 15 to 30 gallons of spray solution per acre and select nozzle type using manufacturer's gallonage and pressure specifications for postemergence application.

DIRECTIONS FOR USE IN FRUITING VEGETABLES (INCLUDING OKRA)

[FOR DISTRIBUTION AND USE ONLY WHERE THIRD PARTY INDEMNIFICATION IS IN EFFECT]

African eggplant; bush tomato; cocona; currant tomato; eggplant; garden huckleberry; goji berry; groundcherry; martynia; naranjilla; okra; pea eggplant; pepino; pepper, bell; pepper, non-bell; roselle; scarlet eggplant; sunberry; tomatillo; tomato; tree tomato; cultivars, varieties, and/or hybrids of these.

Many weather related factors, including high wind or heavy rains or cool conditions at or near crop transplanting, may result in crop injury in fields treated with this product. On occasion this has resulted in a delay in maturity. User assumes these risks before using this product.

RESTRICTIONS

- Do not apply more than 4 fl oz (0.125 lb a.i.) of this product per acre per application.
- Do not apply more than 8 fl oz (0.250 lb a.i.) of this product per acre per year.
- Do not make more than 2 applications per year.

DIRECTIONS FOR USE IN ROW MIDDLES

[FOR DISTRIBUTION AND USE ONLY WHERE THIRD PARTY INDEMNIFICATION IS IN EFFECT]

RESTRICTIONS

- All applications must be made with hooded or shielded equipment.
- Do not apply during or after bloom.

- Do not tank mix with glyphosate after transplanting or crop emergence.

PRECAUTIONS

- Drift of treated soil particles onto plants may cause contact injury.
- Spray must be directed to the row middle, away from the crop bed and with minimal contact with plastic, including the sides of the bed. If top of mulch beds (where plants are to be transplanted) is contacted, severe injury can occur due to foliage contact with treated plastic. In this scenario, a rainfall event of 1/2 inch (natural or irrigation) must occur prior to transplanting to reduce this product's residues.

Notes

- Grow plants on raised or plastic mulched beds that are higher than the treated row middle.
- Irrigate treated field after application and prior to transplanting with minimum of 1/4 inch of water if rainfall does not occur between application and transplanting.

TIMING TO FRUITING VEGETABLES

Apply this product at 4 fluid ounces per acre as a hooded or shielded application to row middles up to 14 days prior to transplanting or seeding for preemergence control of the weeds listed in **Table 7**, as well as to assist in the postemergence control of emerged weeds. A second application of this product at 4 fluid ounces per acre may be applied up to 21 days after transplanting or emergence if needed.

TIMING TO WEEDS

This product may be used for residual weed control, as well as to assist in postemergence burndown of many annual and perennial weeds in row middles. A registered preemergence grass herbicide may be added for control of additional grassy weeds. For assisting in the control of emerged weeds, tank mix this product with carfentrazone-ethyl, paraquat or other registered burndown herbicide. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

DIRECTIONS FOR USE IN GARLIC

RESTRICTIONS

- Do not apply more than 6 fluid ounces (0.188 lb a.i.) of this product per acre per application.
- Do not apply more than 6 fluid ounces (0.188 lb a.i.) of this product per acre per year.
- Do not make more than 1 application per year.

TIMING TO GARLIC

This product may be applied, at 6 fluid ounces per acre, to garlic prior to garlic emergence. Apply within 3 days after planting garlic.

TIMING TO WEEDS

Preemergence - Preemergence to Weeds

Apply this product to weed free garlic for preemergence control of the weeds listed in **Table 10**.

DIRECTIONS FOR USE IN HOPS [Not for Use in California or New York]

RESTRICTIONS

- Do not apply more than 6 fluid ounces (0.188 lb a.i.) of this product per acre per application.
- Do not apply more than 6 fluid ounces (0.188 lb a.i.) of this product per acre per year.
- Do not make more than 1 application per year.
- Do not apply within 30 days of harvest.
- Do not use with an adjuvant.

PRECAUTIONS

- Do not allow spray to contact green stem (unless used for sucker control), foliage, flowers or cones or unacceptable injury may occur.

This product can be used in hops for preemergence weed control as well as sucker control.

TIMING TO HOPS FOR SUCKER CONTROL

Apply this product at 6 fluid ounces per acre as a directed application after hops have reached a minimum of 6 feet in height for sucker control. Direct applications to the lower 2 feet of the hops.

TIMING TO HOPS FOR PREEMERGENCE WEED CONTROL

Apply this product at 6 fluid ounces per acre as a 1 to 1.5 foot band to each side of the hop row, to dormant hops January thru March to ensure time for rain incorporation and activation. If weeds are emerged at the time of application, tank mix this product with a labeled burndown herbicide including paraquat or glyphosate to assist with control of emerged weeds. Do not mow or rake over treated areas, as dust created by mowing may drift onto sensitive crops or vegetation resulting in injury.

TIMING TO WEEDS

Applications of this product must be made prior to weed emergence for control of weeds listed in **Table 10**.

Refer to Product Information section for tank mix guidance. This product, when applied according to label use directions, will control the weeds listed in **Table 10**. This label makes no claims concerning control of other weed species.

DIRECTIONS FOR USE IN LENTILS

HARVEST AID RESTRICTIONS

- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre per application.
- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre per year.
- Do not make more than 1 application per year.
- Do not harvest within 5 days of application.

Desiccation from this product requires the addition of an agronomically approved adjuvant to the spray mixture. Use a methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 quart per acre. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 pounds per acre or a 28 to 32% nitrogen solution at 1 to 2 quarts per acre) may be added to the spray mixture along with methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil. Tank mixing this product with glyphosate or paraquat will increase control of emerged weeds and aid in harvest.

TIMING TO LENTILS

Apply this product, at 1.5 to 2 fluid ounces per acre, when crop is physiologically mature and a minimum of 80% of the pods are yellow to tan in color and 20% are yellow in color. If lentils are treated to early, a reduction in seed quality may occur. Lentils can be harvested 5 days after application.

To ensure thorough coverage, use 15 to 30 gallons of spray solution per acre and select nozzle type using manufacturer's gallonage and pressure specifications for postemergence application.

DIRECTIONS FOR USE IN MINT (Peppermint and Spearmint)

RESTRICTIONS

- Do not apply more than 4 fluid ounces (0.125 lb a.i.) of this product per acre per application.
- Do not apply more than 8 fluid ounces (0.250 lb a.i.) of this product per acre per year.
- Do not make more than 2 applications per year.
- Do not make a sequential application of this product within 60 days of the first application of this product.
- Do not apply within 80 days of harvest.

PRECAUTIONS

To avoid crop injury:

- Do not apply to stands established longer than 3 years.
- Do not apply a Fall application if roots and rhizomes are weak, thin or damaged.
- Do not apply this product on mint in Southern Union County (south of Ladd Canyon) or Baker County in Oregon.
- Do not apply to row or baby mint, use only on established meadow mint.
- Do not apply to mint that has been weakened by diseases, insects (example mint root borer), nematodes, drought, soil salts, high soil pH, previous pesticides, winter injury or double cutting, as severe injury may occur. Apply only to healthy vigorous mint with undamaged rhizomes.
- Do not apply before November 25 or after March 1.
- Apply only to dormant mint. Application to non-dormant mint may result in unacceptable crop injury.

Many weather related factors, including high wind, splashing or heavy rains or cool conditions at or near mint emergence, may result in mint injury in fields treated with this product. User assumes these risks before using this product.

Tank mixes with labeled rates of paraquat are recommended to control emerged weeds and increase crop safety. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

TIMING TO MINT

As a spray, this product may be applied only to established, dormant mint for preemergence control of the weeds listed in **Table 7**, as well as to assist in the postemergence control of emerged weeds. Application to non-dormant mint or to baby (row) mint (time from planting of mint roots through the first cutting), may result in unacceptable crop injury. As a bulk fertilizer application, this product may be applied at least 80 days prior to harvest. Leaves must be dry at the time of applications or severe injury may occur.

TIMING TO WEEDS

Burndown - Dormant Mint, Postemergence to Weeds

This product may be used for residual weed control, as well as to assist in postemergence burndown of many annual and perennial weeds where established mint is dormant. For control of emerged weeds, tank mix this product with paraquat. Refer to paraquat label for rate and application parameters. To ensure thorough coverage, use a minimum of 15 gallons of spray solution per acre. Tank mixes of this product applied to assist in the control of emerged weeds must be applied with a non-ionic surfactant at 0.25% v/v. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 pound per acre, or 28 to 32% nitrogen solution at 1 to 2 quarts per acre) may be added to increase herbicidal activity.

Preemergence - Dormant Mint, Preemergence to Weeds

Apply this product to dormant mint for the preemergence control of weeds listed in **Table 7**. Fall applications of this product, followed by a sequential application in the Spring, have resulted in better Summer annual weed control than a single Fall or single Spring application.

Fall application is most effective for Fall germinating weeds, for example groundsel. Fields plowed or harrowed after an application of this product will result in less effective preemergence activity. In furrow irrigated fields, corrugating that is done after an application of this product will expose untreated soil and break the herbicide barrier resulting in poor weed control.

Table 7. Weeds Controlled by Residual Activity of LIBERTY FLUMIOX 44

COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	SOIL TYPE	APPLICATION RATE
BROADLEAF WEED SPECIES		Up to 5%	All Soil Types	4 fl oz/A
Bristly Starbur	<i>Acanthospermum hispidum</i>			
Carpetweed	<i>Mollugo verticillata</i>			

COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	SOIL TYPE	APPLICATION RATE
Chickweeds,				
Common	<i>Stellaria media</i>			
Mouseear	<i>Cerastium vulgatum</i>			
Coffee Senna	<i>Cassia occidentalis</i>			
Copperleaf, Hophornbeam	<i>Acalypha ostryifolia</i>			
Dandelion	<i>Taraxacum officinale</i>			
Dodder (suppression only) ^{1, 2}	<i>Cuscuta</i> spp.			
Eclipta	<i>Eclipta prostrata</i>			
Eveningprimrose, Cutleaf	<i>Oenothera laciniata</i>			
False Chamomile	<i>Tripleurospermum maritima</i>			
Fiddleneck, Coast ²	<i>Amsinckia menziesii</i>			
Field Pennycress ²	<i>Thlaspi arvense</i>			
Fleabane, Hairy ²	<i>Conyza bonariensis</i>			
Flixweed	<i>Descurainia spophia</i>			
Florida Beggarweed	<i>Desmodium tortuosum</i>			
Florida Pusley	<i>Richardia scabra</i>			
Golden Crownbeard	<i>Verbesina encelioides</i>			
Groundsel, Common	<i>Senecio vulgaris</i>			
Hairy Indigo	<i>Indigofera hirsute</i>			
Hemp Sesbania	<i>Sesbania exaltata</i>			
Henbit	<i>Lamium amplexicaule</i>			
Jimsonweed	<i>Datura stramonium</i>			
Kochia	<i>Kochia scoparia</i>			
Lambsquarters, Common	<i>Chenopodium album</i>			
Little Mallow	<i>Malva parviflora</i>			
London Rocket	<i>Sisymbrium irio</i>			
Marestail/Horseweed	<i>Conyza canadensis</i>			
Mayweed/False Chamomile	<i>Matricaria maritima</i>			
Morningglories,				
Entireleaf	<i>Ipomoea hederacea</i> var. <i>integrifolia</i>			
Ivyleaf	<i>Ipomoea hederacea</i>			
Red/Scarlet	<i>Ipomoea coccinea</i>			
Smallflower	<i>Jacquemontia tamnifolia</i>			
Tall	<i>Ipomoea purpurea</i>			
Mustard,				
Tansy	<i>Descurainia pinnata</i>			
Tumble	<i>Sisymbrium altissimum</i>			
Wild	<i>Brassica kaber</i>			
Nettle, Burning	<i>Urtica urens</i>			
Nightshades,				
Black	<i>Solanum nigrum</i>			
Eastern Black	<i>Solanum ptycanthum</i>			
Hairy	<i>Solanum sarrachoides</i>			
Pigweeds,				
Palmer Amaranth	<i>Amaranthus palmeri</i>			
Redroot	<i>Amaranthus retroflexus</i>			
Smooth	<i>Amaranthus hybridus</i>			
Spiny Amaranth	<i>Amaranthus spinosus</i>			
Tumble	<i>Amaranthus albus</i>			
Prickly Lettuce (China Lettuce)	<i>Lactuca serriola</i>			
Prickly Sida (Teaweed)	<i>Sida spinosa</i>			
Puncturevine	<i>Tribulus terrestris</i>			
Purslane,				
Common	<i>Portulaca oleracea</i>			
Horse	<i>Trianthema portulacastrum</i>			
Radish, Wild	<i>Raphanus raphanistrum</i>			

COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	SOIL TYPE	APPLICATION RATE
Ragweed, Common	<i>Ambrosia artemisiifolia</i>			
Redmaids	<i>Calandrinia ciliata</i> var. <i>menziesii</i>			
Russian Thistle	<i>Salsola iberica</i>			
Shepherd's-purse	<i>Capsella bursa-pastoris</i>			
Smartweeds,				
Ladysthumb	<i>Polygonum persicaria</i>			
Pennsylvania	<i>Polygonum pennsylvanicum</i>			
Smellmelon ²	<i>Cucumis melo</i>			
Sowthistle, Prickly ²	<i>Sonchus asper</i>			
Spotted Spurge	<i>Euphorbia maculata</i>			
Spurred Anoda	<i>Anoda cristata</i>			
Tropic Croton	<i>Croton glandulosus</i>			
Velvetleaf	<i>Abutilon theophrasti</i>			
Venice Mallow	<i>Hibiscus trionum</i>			
Waterhemp,				
Common	<i>Amaranthus rudis</i>			
Tall	<i>Amaranthus tuberculatus</i>			
White Cockle ²	<i>Silene latifolia</i>			
Wild Poinsettia	<i>Euphorbia heterophylla</i>			
Wormwood, Biennial	<i>Artemisia biennis</i>			
Yellow Rocket	<i>Barbarea vulgaris</i>			
GRASS WEED SPECIES				
Barnyardgrass	<i>Echinochloa crus-galli</i>			
Bluegrass, Annual	<i>Poa annua</i>			
Craggrass, Large	<i>Digitaria sanguinalis</i>			
Foxtail, Giant	<i>Setaria faberi</i>			
Goosegrass	<i>Eleusine indica</i>			
Lovegrass, California	<i>Eragrostis diffusa</i>			
Panicums,				
Fall	<i>Panicum dichotomiflorum</i>			
Texas	<i>Panicum texanum</i>			
Ryegrass, Italian	<i>Lolium multiflorum</i>			
Signalgrass, Broadleaf	<i>Brachiaria platyphylla</i>			

¹ This product at 4 fluid ounces per acre will provide postemergence dodder² suppression when applied in combination with imazethapyr or imazamox at labeled rates. The use of imazethapyr or imazamox require the use of a NIS, which will result in burn and stunting of alfalfa. Growers must expect and accept this prior to using this tank mix.

² Not for use in California.

DIRECTIONS FOR USE IN ONION (DRY BULB)
[For Use in the States of Michigan, New York and North Dakota Only]

[For chemigation applications on onion follow the "CHEMIGATION" section of this label.]

RESTRICTIONS

- Do not apply more than 2 fluid ounces (0.063 lb a.i.) of this product per acre per application.
- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre per year.
- Do not make more than 2 applications per year.
- Do not make sequential application within 14 days of the first application.
- Do not apply more than 1 fluid ounce (0.031 lb a.i.) of this product per year on soils that contain greater than 90% sand plus gravel.
- Do not apply with any type of adjuvant.
- Do not apply within 45 days of harvest.

PRECAUTION

- Do not apply as part of a tank mix, other than with Prowl® H₂O or unacceptable injury may result. **Note:** Other formulations of pendimethalin should not be tank mixed with this product for use in onions.

Use of this product may result in necrotic spotting of onion leaves that come in contact with the spray. User assumes this potential crop response before using this product.

[Microrate Application]

Sequential applications of this product may be applied to onions (dry bulb), between the 2-leaf and 6-leaf stage, at rates of 0.5 to 1 fluid ounce per acre, on a 7 day interval.]

TIMING TO ONIONS (Dry Bulb)

Apply this product to transplanted onions (dry bulb) between the 2-leaf and 6-leaf stage and on direct seed onions (dry bulb) between the 3- leaf and 6-leaf stage.

TIMING TO WEEDS

Preemergence - Emerged Onions (dry bulb), Preemergence to Weeds

Apply this product to weed free onions (dry bulb) for preemergence control of the weeds listed in **Table 1, Section A.**

DIRECTIONS FOR USE IN PEANUT

RESTRICTIONS

- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre per year.
- Do not make more than 1 application per year.
- Do not irrigate when peanuts are cracking.
- Do not graze treated fields or feed treated hay to livestock.

[PRECAUTION

- Do not apply more than 2 fluid ounces (0.063 lb a.i.) per acre in the states of North Carolina, Oklahoma or Virginia where climatic conditions may result in unacceptable injury to peanuts [or under conditions specified below under **PREEMERGENCE APPLICATION IN PEANUT**].

Many weather related factors, including high wind, splashing or heavy rains or cool conditions at or near peanut emergence, may result in peanut injury in fields treated with this product. On occasion this has resulted in a delay in maturity or even a slight decrease in yield.

WIND MANAGEMENT

In areas where shallow cultivation is used between rows to reduce wind-borne sand damage to peanuts, weed control from this product may be reduced.

TIMING TO PEANUTS

This product may be applied to peanuts prior to planting or preemergence (after planting). Preemergence applications of this product must be made within 2 days after planting and prior to peanut emergence. Application after the peanuts have begun to crack, or are emerged, will result in severe crop injury. Do not apply when peanuts have begun to crack. Select rate of this product from **Table 1**, according to anticipated weed spectrum.

TIMING TO WEEDS

Burndown - Preemergence to Peanuts, Postemergence to Weeds

This product, applied as part of a burndown program, may be used for residual weed control, as well as to assist in postemergence burndown of many annual and perennial weeds where peanuts will be planted directly into a stale seedbed, cover crop or in previous crop residues. Apply this product before planting, during planting or after planting, but before the crop emerges. For control of emerged weeds, tank mix this product with glyphosate. Refer to glyphosate label for rate and application pressure. To ensure thorough coverage, use a minimum of 15 gallons of spray solution per acre. Tank mixes of this product applied to assist in the control of emerged weeds must be applied with an adjuvant, including a non-ionic surfactant at 0.25% v/v or a crop oil concentrate or a methylated seed oil at 1 to 2 pints per acre. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 pounds per acre or 28 to 32% nitrogen solution at 1 to 2 quarts per acre) may be added to increase herbicidal activity. Preemergence (conventional tillage) applications of this product must be applied prior to weed emergence.

[PREEMERGENCE APPLICATION IN PEANUT (North Carolina, Oklahoma and Virginia Only)

This product, at 3 fluid ounces per acre, can be applied within 2 days of planting to control common ragweed, tropic croton and entireleaf, ivyleaf and tall/scarlet morningglories.

Cool temperatures near emergence (2 consecutive nighttime lows in the 50's F) in combination with heavy rainfall may result in severe crop injury. Only use this product, at 3 fluid ounces per acre, in these states when other alternatives are not available for adequate control of the weeds listed above and the user acknowledges the risks associated with this use rate under the adverse environmental conditions listed above.]

ADDITIONAL RESIDUAL GRASS CONTROL: SEQUENTIAL

This product may be applied sequentially following a preplant incorporated application of trifluralin (states of New Mexico, Oklahoma and Texas only), ethafluralin, metolachlor/S-metolachlor, pendimethalin or dimethenamid /dimethenamid-P.

ADDITIONAL RESIDUAL GRASS CONTROL: TANK MIXED

This product can be tank mixed with alachlor, metolachlor/S-metolachlor or dimethenamid/dimethenamid-P for additional grass and broadleaf weed control. This product can also be tank mixed with pendimethalin or ethafluralin in states where they are labeled, provided overhead irrigation guidelines on the pendimethalin and/or ethafluralin labels are followed. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

DIRECTIONS FOR USE IN POTATO

[Arizona, California, Colorado, Delaware, Florida, Hawaii, Idaho, Maryland, Minnesota, Montana, Nebraska, Nevada, New Jersey, New Mexico, North Carolina, North Dakota, Oregon, South Dakota, Texas, Utah, Virginia, Washington, Washington DC and Wyoming only.]

RESTRICTIONS

- Do not apply more than 1.5 fluid ounces (0.047 lb a.i.) of this product per acre per application.
- Do not apply more than 1.5 fluid ounces (0.047 lb a.i.) of this product per acre per year.
- Do not make more than 1 application per year.
- Do not apply to Rill (Furrow) irrigated potatoes.

Many weather related factors, including high wind, splashing or heavy rains or cool conditions at or near potato emergence, may result in potato injury in fields treated with this product. On occasion this has resulted in a delay in maturity. User assumes these risks before using this product.

TIMING TO POTATOES

This product may be applied to potatoes after hilling for the preemergence suppression of the weeds listed in **Table 8**. Tank mix this product with other labeled herbicides for broad spectrum weed control. A minimum of 2 inches of settled soil must cover the vegetative portion of the potato plant at the time of application of this product. Application to potatoes with less than 2 inches of soil covering the vegetative portion of the potato may result in crop injury. In areas with historically higher amounts of rainfall during the time of preemergence herbicide applications, including the Red River Valley, Minnesota and North Dakota, the requirement for 2 inches of settled soil is critical to avoid crop injury. Mechanical incorporation of this product will result in decreased weed control. In areas with sprinkler irrigation, incorporate this product with 0.5 to 0.75 inches of irrigation, after application and before any sprouts are within 2 inches of the settled soil surface if a rainfall event has not yet occurred.

TIMING TO WEEDS

Preemergence - Soil Covered Potatoes, Preemergence to Weeds

Apply this product to soil covered potatoes for the preemergence suppression of the weeds listed in **Table 8**. Harrowing, cultivation or corrugating after this product application will reduce weed control.

Read tank mix product label for rates and weeds controlled. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

[CHEMIGATION

This product may be applied through sprinkler systems in potatoes. For chemigation applications on potatoes follow the "CHEMIGATION" section of this label.]

Table 8. Weeds Suppressed by Residual Activity of LIBERTY FLUMIOX 44 at 1.5 Fluid Ounces per Acre

COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	APPLICATION RATE
Lambsquarters, Common	<i>Chenopodium album</i>	Up to 5%	1.5 fl oz/A
Mustard, Wild	<i>Brassica kaber</i>		
Nightshades,			
Black	<i>Solanum nigrum</i>		
Eastern Black	<i>Solanum ptycanthum</i>		
Hairy	<i>Solanum sarrachoides</i>		
Pigweeds,			
Palmer Amaranth	<i>Amaranthus palmeri</i>		
Redroot	<i>Amaranthus retroflexus</i>		
Smooth	<i>Amaranthus hybridus</i>		
Spiny Amaranth	<i>Amaranthus spinosus</i>		
Tumble	<i>Amaranthus albus</i>		
Prickly Lettuce (China Lettuce)	<i>Lactuca serriola</i>		
Radish, Wild	<i>Raphanus raphanistrum</i>		

DIRECTIONS FOR USE IN SOYBEAN

RESTRICTIONS

- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre per year.
- Do not make more than 1 application per year.
- Do not graze treated fields or feed treated hay to livestock.
- Do not irrigate when soybeans are cracking.
- Do not tank mix this product with acetochlor, alachlor, flufenacet, metolachlor/S-metolachlor, or dimethenamid/ dimethenamid-p within 14 days of planting soybeans, unless soybeans are planted under no-till or minimum tillage conditions on wheat stubble or no-till field corn stubble.

TIMING TO SOYBEANS

This product may be applied to soybeans prior to planting or preemergence (after planting). Preemergence application of this product must be made within 3 days after planting and prior to soybean emergence. Application after the soybeans have begun to crack, or are emerged, will result in severe crop injury. Do not apply when soybeans have begun to crack. Select rate of this product from **Table 1** according to anticipated weed spectrum.

TIMING TO WEEDS

Burndown - Preemergence to Soybeans, Postemergence to Weeds

This product, applied as part of a burndown program, may be used for residual weed control, as well as to assist in postemergence burndown of many annual and perennial weeds where soybeans will be planted directly into a stale seedbed, cover crop or in previous crop residues. For control of emerged weeds, choose the most appropriate tank mix partner from **Table 9**. Apply this product with ground equipment before planting, during planting or within 3 days after planting, **but before the crop emerges**. To ensure thorough coverage, use a minimum of 15 gallons of spray solution per acre. Refer to tank mix partner's label for specified application pressure. All tank mixes of this product applied to assist in the control of emerged

weeds must be applied with crop oil concentrate or methylated seed oil at 1 to 2 pints per acre or a non-ionic surfactant at 0.25% v/v.

INCREASING SPEED OF GLYPHOSATE BURNDOWN ACTIVITY

This product, at rates as low as 1 fluid ounce per acre, may be tanked mixed with glyphosate to increase the speed of burndown activity compared to glyphosate applied alone. Residual weed control will not be provided at rates lower than 2 fluid ounces per acre; however, suppression of the weeds in **Table 2**, may occur at rates of this product as low as 1 fluid ounce per acre.

TANK MIXES

This product may be tank mixed with the herbicides listed in **Table 9** for increased burndown activity, additional residual broadleaf and/or additional grass control. Refer to tank mix partner's label for adjuvant directions. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Table 9. Tank Mix Partners for Control of Emerged Weeds in Reduced Tillage Soybeans

TANK MIX PARTNERS	TARGET WEEDS¹
2,4-D LVE	Marestail Giant Ragweed Dandelion
Clethodim	Annual Grasses
Dicamba + 2,4-D	Marestail Giant Ragweed Dandelion
Glyphosate	General Burndown
Imazaquin	Cocklebur Common Sunflower
Paraquat	Annual Grasses Henbit

¹ Refer to tank mix product labels for specific directions for control of emerged weeds present.

ADDITIONAL RESIDUAL BROADLEAF CONTROL

This product can be tank mixed with cloransulam-methyl, flumetsulam, imazaquin, imazaquin + imazethapyr, imazethapyr + pendimethalin, linuron or metribuzin for additional broadleaf control.

ADDITIONAL RESIDUAL GRASS CONTROL

This product can be tank mixed with clomazone or pendimethalin for additional grass control. Tank mixes with alachlor, flufenacet, dimethenamid/ dimethenamid/-P or metolachlor/S-metolachlor, may result in severe injury to soybeans when application is followed by prolonged periods of cool wet weather.

ROUNDUP READY® PROGRAM

This product may be applied as part of a burndown program or preemergence in conventional tillage programs, at 2 to 3 fluid ounces per acre to reduce early season weed competition from waterhemp, velvetleaf, nightshade and morningglories as well as other weeds listed in Tables 2 and 3 in Roundup Ready programs. A sequential post emergence application of glyphosate will be required to control weeds not controlled by this product.

DIRECTIONS FOR USE IN STRAWBERRY

RESTRICTIONS

- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre per application.
- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre per year.
- Do not make more than 1 application per year.

TIMING TO STRAWBERRIES

- This product, at 3 fluid ounces (0.093 lb a.i.) per acre, can be applied to the soil a minimum of 30 days prior to transplanting strawberries provided the strawberries will be transplanted through a plastic mulch.
- This product at 3 fluid ounces (0.093 lb a.i.) per acre can be applied to dormant (established or newly planted) strawberries for the preemergence control of the weeds listed in **Table 1**.
- This product, at 3 fluid ounces (0.093 lb a.i.) per acre, can be applied in strawberry row middles with a shielded or hooded sprayer for the preemergence control of the weeds listed in **Table 1**.

Application Method	Minimum Time From Application to Harvest (PHI)	Use Rate Per Acre Per Application (fl oz)	Use Rate Per Acre Per Year (fl oz)	Special Use Instructions
Pre-transplant	Not applicable	3	3	Apply a minimum of 30 days prior to transplanting and prior to plastic mulch being laid. Apply as part of a tank mix to control emerged weeds.
Preemergence to dormant strawberries	Not applicable			Crop oil concentrate, at 1% v/v, or non-ionic surfactant, at 0.25% v/v, may be added to help control emerged broadleaf weeds.
Hooded or shielded sprayer application to row middles	Do not apply after fruit set			Apply only to row middles - do not apply over strawberries. Apply prior to weed emergence. Crop spotting may occur if an adjuvant is added. Application after fruit set may result in spotting of fruit. Do not allow spray drift to come in contact with fruit or foliage.

Table 10. Weeds Controlled by Preemergence Application of LIBERTY FLUMIOX 44

COMMON NAME	SCIENTIFIC NAME	ORGANIC	SOIL	APPLICATION	
BROADLEAF WEED SPECIES		Up to 10% ¹	All Soil Types ²	Asparagus, Caneberries, Garlic, Hops 6 fl oz/A	
Bristly Starbur	<i>Acanthospermum hispidum</i>			Sugarcane 6 to 8 fl oz/A	
Carpetweed	<i>Mollugo verticillata</i>				
Chickweeds					
Common	<i>Stellaria media</i>				
Mouseear	<i>Cerastium vulgatum</i>				
Coffee Senna	<i>Cassia occidentalis</i>				
Dandelion	<i>Taraxacum officinale</i>				
Eclipta	<i>Eclipta prostrata</i>				Bushberries, Cactus, Citrus Fruit, Grapes, Nut Trees, Olive, Pome Fruit, Pomegranate, Stone Fruit, and Non-Bearing Fruit Trees 6 to 12 fl oz/A ²
Eveningprimrose, Cutleaf	<i>Oenothera laciniata</i>				
False Chamomile	<i>Tripleurospermum maritima</i>				
Filaree					
Redstem	<i>Erodium cicutarium</i>				
Whitestem	<i>Erodium moschatum</i>				
Fiddleneck, Coast	<i>Amsinckia menziesii</i>				
Fleabane, Hairy	<i>Conyza bonariensis</i>				
Field Pennycress	<i>Thlaspi arvense</i>				
Florida Beggarweed	<i>Desmodium tortuosum</i>				
Florida Pusley	<i>Richardia scabra</i>				
Golden Crownbeard	<i>Verbesina encelioides</i>				

COMMON NAME	SCIENTIFIC NAME	ORGANIC	SOIL	APPLICATION
Groundsel, Common	<i>Senecio vulgaris</i>			To Maintain Bare Ground on Non-Crop Areas of Farms, Orchards & Vineyards 6 to 12 fl oz/A.
Hairy Indigo	<i>Indigofera hirsuta</i>			
Hemp Sesbania	<i>Sesbania exaltata</i>			
Henbit	<i>Lamium amplexicaule</i>			
Jimsonweed	<i>Datura stramonium</i>			
Kochia	<i>Kochia scoparia</i>			
Lambsquarters, Common	<i>Chenopodium album</i>			
Mallow				
Common (Cheeseweed)	<i>Malva neglecta</i>			
Little				
Horseweed/Marestail	<i>Conyza canadensis</i>			
Mayweed/False Chamomile	<i>Matricaria maritima</i>			
Morningglories				
Entireleaf	<i>Ipomoea hederacea</i> var.			
Ivyleaf				
Red/Scarlet	<i>Ipomoea coccinea</i>			
Smallflower	<i>Jacquemontia tamnifolia</i>			
Tall	<i>Ipomoea purpurea</i>			
Mustards				
London Rocket	<i>Sisymbrium irio</i>			
Tansey	<i>Desurainia pinnata</i>			
Tumble	<i>Sisymbrium altissimum</i>			
Wild	<i>Brassica kaber</i>			
Nettle, Burning	<i>Urtica urens</i>			
Nightshades				
Black	<i>Solanum nigrum</i>			
Eastern Black	<i>Solanum ptycanthum</i>			
Hairy	<i>Solanum sarrachoides</i>			
Pigweeds				
Palmer Amaranth	<i>Amaranthus palmeri</i>			
Redroot	<i>Amaranthus retroflexus</i>			
Smooth	<i>Amaranthus hybridus</i>			
Spiny Amaranth	<i>Amaranthus spinosus</i>			
Tumble	<i>Amaranthus albus</i>			
Prickly Lettuce	<i>Lactuca serriola</i>			
Prickly Sida (Teaweed)	<i>Sida spinosa</i>			
Puncturevine	<i>Tribulus terrestris</i>			
Purslane				
Common	<i>Portulaca oleracea</i>			
Horse	<i>Trianthema portulacastrum</i>			
Radish, Wild	<i>Raphanus raphanistrum</i>			
Ragweed, Common	<i>Ambrosia artemisiifolia</i>			
Redmaids	<i>Calandrinia ciliata</i> var. <i>menziessi</i> .			
Redweed	<i>Melochia corchorifolia</i>			
Shepherd's-purse	<i>Capsella bursa-pastoris</i>			
Smellmelon	<i>Cucumis melo</i>			
Sowthistle, Annual ³	<i>Sonchus oleraceus</i>			
Spotted Spurge	<i>Euphorbia maculata</i>			
Spurred Anoda	<i>Anoda cristata</i>			
Thistle, Russian	<i>Salsola iberica</i>			
Tropic Croton	<i>Croton glandulosus</i>			
Venice Mallow	<i>Hibiscus trionum</i>			
Waterhemp				
Common	<i>Amaranthus rudis</i>			
Tall	<i>Amaranthus tuberculatus</i>			
Wild Poinsettia	<i>Euphorbia heterophylla</i>			
White Cockle	<i>Silene latifolia</i>			
Wormwood, Biennial	<i>Artemisia biennis</i>			

COMMON NAME	SCIENTIFIC NAME	ORGANIC	SOIL	APPLICATION
Yellow Rocket	<i>Barbarea vulgaris</i>			
GRASS WEED SPECIES				
Barnyardgrass	<i>Echinochloa crus-galli</i>			
Bluegrass, Annual	<i>Poa annua</i>			
Crabgrass				
Large	<i>Digitaria sanguinalis</i>			
Smooth	<i>Digitaria ischaemum</i>			
Foxtails				
Bristly	<i>Setaria verticillata</i>			
Giant	<i>Setaria faberi</i>			
Green	<i>Setaria viridis</i>			
Yellow	<i>Setaria glauca</i>			
Goosegrass	<i>Eleusine indica</i>			
Guineagrass	<i>Panicum maximum</i>			
Johnsongrass, Seedling	<i>Sorghum halepense</i>			
Lovegrass, California	<i>Eragrostis diffusa</i>			
Panicum				
Fall	<i>Panicum dichotomiflorum</i>			
Texas	<i>Panicum texaum</i>			
Ryegrass, Italian	<i>Lolium multiflorum</i>			
Signalgrass, Broadleaf	<i>Brachiaria platyphylla</i>			

- ¹ This product can be used on soils with greater than 10%; however, length of residual control may be shorter than on soils with lower organic matter content.
- ² Use a maximum rate of 6 fluid ounces per acre per application on any soil that has a sand plus gravel content over 80% if bushes, trees or vines are under 3 years of age.
- ³ Not for use in California.

DIRECTIONS FOR USE IN SUGARCANE

RESTRICTIONS

- Do not apply more than 8 fluid ounces (0.250 lb a.i.) of this product per acre per application.
- Do not apply more than 12 fluid ounces (0.375 lb a.i.) of this product per acre per year.
- Do not make more than 2 applications per year.
- Do not make a sequential application within 14 days of the first application.
- Do not apply within 90 days of harvest.

TIMING TO SUGARCANE

This product may be applied from 2 weeks prior to planting to before the sugarcane emerges, post directed or at layby. Select the proper rate of this product from **Table 10** according to anticipated weed spectrum and soil organic matter content for preemergence applications. Select rate of this product from **Table 11** according to emerged weed spectrum and weed heights for post-directed and layby applications.

TIMING TO WEEDS

Burndown — Preemergence to Sugarcane, Postemergence to Weeds

This product may be used for preemergence control, and to assist in postemergence burndown, of many annual broadleaf weeds in sugarcane. For control of emerged weeds, choose the most appropriate tank mix partner from **Table 12**. Apply this product **before the crop emerges**. To ensure thorough coverage, use a minimum of 15 gallons of spray solution per acre. All tank mixes of this product applied to assist in the control of emerged weeds must be applied with crop oil concentrate or methylated seed oil at 1 quart per acre or a non-ionic surfactant at 0.25% v/v. Some tank mix products, including glyphosate, may be formulated with a suitable adjuvant and do not require additional adjuvant.

Preemergence — Preemergence to Sugarcane, Preemergence to Weeds

This product may be used for preemergence control of many annual broadleaf and grassy weeds in sugarcane. Select rate based on anticipated weed spectrum and soil organic matter content from **Table 10**. Apply this product **before the crop emerges**.

Post-Directed — Postemergence to Sugarcane, Postemergence to Weeds

Only make post-directed applications to upright sugarcane varieties after the sugarcane has exceeded 24 inches in height and has begun to joint. Post-directed applications to "PINEAPPLE" varieties. Do not apply post-directed applications to "PINEAPPLE" varieties or to upright varieties that have not exceeded 24 inches in height and have not begun to joint, may result in unacceptable crop injury. To ensure thorough coverage, use a minimum of 15 gallons of spray solution per acre. Post-directed applications of this product must include a crop oil concentrate or methylated seed oil at 1 quart per acre or a non-ionic surfactant at 0.25% v/v. Select the proper rate of this product based on weed spectrum and weed height from **Table 11**.

Layby — Postemergence to Sugarcane, Postemergence to Weeds

Layby applications can be made to upright and "PINEAPPLE" varieties after the sugarcane has exceeded 30 inches in height and the spray solution will not contact foliage above 6 inches from the base of the sugarcane. To ensure thorough coverage, use a minimum of 15 gallons of spray solution per acre. Layby applications of this product must be applied with crop oil concentrate or methylated seed oil at 1 quart per acre or a non-ionic surfactant at 0.25% v/v. Select the proper rate of this product based on weed spectrum and weed height from **Table 11**.

Table 11. Broadleaf Weeds Controlled by Post-Directed or Layby Application of LIBERTY FLUMIOX 44 in Sugarcane

COMMON NAME	SCIENTIFIC NAME	WEED HEIGHT (inches)	
		3 fl oz/A	4 fl oz/A
BROADLEAF WEED SPECIES			
Bindweed, Field ¹	<i>Convolvulus arvensis</i>	4	8
Carpetweed	<i>Mollugo verticillata</i>	4	4
Cocklebur, Common	<i>Xanthium strumarium</i>	4	4
Florida Beggarweed	<i>Desmodium tortuosum</i>	2	2
Hemp Sesbania	<i>Sesbania exaltata</i>	6	8
Jimsonweed	<i>Datura stramonium</i>	4	4
Lambsquarters, Common	<i>Chenopodium album</i>	4	4
Morningglories,			
Entireleaf	<i>Ipomoea hederacea</i> var. <i>integriuscula</i>	-	4
Ivyleaf	<i>Ipomoea hederacea</i>	4	4
Pitted	<i>Ipomoea lacunosa</i>	4	6
Red	<i>Ipomoea coccinea</i>	-	4
Tall	<i>Ipomoea purpurea</i>	2	4
Mustard, Wild	<i>Brassica kaber</i>	6	6
Pigweeds,			
Palmer Amaranth	<i>Amaranthus palmeri</i>	4	6
Redroot	<i>Amaranthus retroflexus</i>	4	6
Smooth	<i>Amaranthus hybridus</i>	4	6
Plantain, Broadleaf	<i>Plantago major</i>	6	6
Prickly Sida	<i>Sida spinosa</i>	4	6
Purslanes,			
Common	<i>Portulaca oleracea</i>	2	4
Rock	<i>Calandrinia</i> spp.	-	2
Ragweeds,			
Common	<i>Ambrosia artemisiifolia</i>	2	2
Giant	<i>Ambrosia trifida</i>	4	4
Rice Flatsedge	<i>Cyperus iria</i>	2	4
Sicklepod	<i>Senna obtusifolia</i>	4	4
Smartweeds,			
Ladysthumb	<i>Polygonum persicaria</i>	4	4
Pale	<i>Polygonum lapathifolium</i>	4	4
Pennsylvania	<i>Polygonum pennsylvanicum</i>	4	4

COMMON NAME	SCIENTIFIC NAME	WEED HEIGHT (inches)	
		3 fl oz/A	4 fl oz/A
Spotted Spurge	<i>Euphorbia maculata</i>	4	4
Velvetleaf	<i>Abutilon theophrasti</i>	4	6
Venice Mallow	<i>Hibiscus trionum</i>	2	2
Waterhemp, Common	<i>Amaranthus rudis</i>	2	2
Tall	<i>Amaranthus tuberculatus</i>	2	2

¹ Tank mixes of this product will only control the above ground portion of field bindweed. Repeated applications will be needed to control regrowth.

TANK MIXES

This product may be tank mixed with the herbicides listed in **Table 12** for additional weed control in burndown, preemergence, post-directed and layby applications. Refer to tank mix partner's label for adjuvant directions. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Read and follow the application restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Table 12. Tank Mixes with LIBERTY FLUMIOX 44 for Post-Directed or Layby Use in Sugarcane

TANK MIX PARTNER ¹	TARGET WEEDS	BURNDOWN	POST-DIRECTED ²	LAYBY
2,4-D amine	Annual and Perennial Broadleaf Weeds	X		
ametryn ³	Annual Grasses		X	X
asulam ⁴	Annual Grasses		X	X
atrazine	Pigweeds Cocklebur	X	X	X
dicamba + 2,4-D	Annual and Perennial Broadleaf Weeds	X		
glyphosate ⁵	Annual and Perennial Weeds	X		X
halosulfuron-methyl	Purple Nutsedge Yellow Nutsedge	X	X	X
metribuzin ⁶	Broadleaf Panicum Goosegrass		X	X

¹ Refer to tank mix product labels for specific directions for control of emerged weeds present not listed in **Table 11**.

² Only apply post-directed to upright sugarcane varieties after the sugarcane has exceeded 24 inches in height. Post-directed applications to "PINEAPPLE" varieties. Do not apply post-directed applications to "PINEAPPLE" varieties or to upright varieties that have not exceeded 24 inches in height may result in unacceptable crop injury.

³ Apply before weeds are greater than 6 inches tall.

⁴ Apply to sugarcane at least 24 inches tall.

⁵ Glyphosate applications must be made with a hooded sprayer. Sugarcane must be at least 3 feet tall. Contact with the sugarcane foliage by either the spray mixture or the treated weed foliage will result in sugarcane injury.

⁶ Refer to metribuzin label for restrictions based on soil type.

ADDITIONAL PREEMERGENCE BROADLEAF CONTROL

This product can be tank mixed with atrazine or diuron for additional preemergence broadleaf control.

ADDITIONAL PREEMERGENCE GRASS CONTROL

This product can be tank mixed with pendimethalin for additional preemergence grass control provided sugarcane has not emerged.

DIRECTIONS FOR USE IN SUNFLOWER AND SAFFLOWER

HARVEST AID

RESTRICTIONS

- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre per application.
- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre per year.
- Do not make more than 1 application per year.
- Do not harvest within 5 days of application.

Desiccation from this product requires the addition of an agronomically approved adjuvant to the spray mixture. Use a methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 quart per acre. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 pounds per acre or a 28 to 32% nitrogen solution at 1 to 2 quarts per acre) may be added to the spray mixture along with methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil. Tank mixing this product with glyphosate or paraquat will increase control of emerged weeds and aid in harvest for sunflowers. Tank mixing this product with glyphosate will increase control of emerged weeds and aid in harvest for safflower.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

TIMING TO SUNFLOWER AND SAFFLOWER

Apply this product, at 1.5 to 2 fluid ounces per acre, when crop is mature (when seed is 35% moisture or less). For many varieties, this is when the backs of the heads are turning yellow and the bracts are turning brown. Sunflower and safflower can be harvested 5 days after application.

To ensure thorough coverage, use 15 to 30 gallons of spray solution per acre and select nozzle type using manufacturer's gallonage and pressure specifications for postemergence application.

DIRECTIONS FOR USE IN SWEET POTATO

[For Use in the States of Arizona, California and Hawaii Only]

RESTRICTIONS

- Do not apply more than 3 fluid ounces (0.093 lb a.i.) of this product per acre per year.
- Do not make more than 1 application per year.
- Do not apply postemergence to sweet potatoes.
- Do not use greenhouse grown transplants.
- Do not use transplants harvested more than 2 days prior to transplanting.
- Do not apply as a part of any tank mix, except with labeled rates of clomazone, if tank mix is applied prior to transplanting.
- Do not use on any sweet potato variety other than "Beauregard", unless user has tested this product on other variety and has found crop tolerance to be acceptable.

TIMING TO SWEET POTATOES

This product must be applied prior to transplanting sweet potatoes.

TIMING TO WEEDS

Preemergence to Weeds

Apply this product to soil prior to transplanting sweet potato slips for the preemergence control of the weeds listed in **Table 1**.

DIRECTIONS FOR USE IN WHEAT

[For Use in the States of Delaware, Idaho, Kentucky, Maryland, Minnesota, Montana, North Carolina, North Dakota, New Jersey, Oregon, South Carolina, South Dakota, Tennessee, Virginia and Washington Only]

RESTRICTIONS

- Do not apply more than 2 fluid ounces (0.063 lb a.i.) of this product per acre per application.
- Do not apply more than 2 fluid ounces (0.063 lb a.i.) of this product per acre per year.
- Do not make more than 1 application per year.

PRE-PLANT APPLICATIONS. PRE-EMERGENCE WEED CONTROL

RESTRICTIONS

- For pre-plant weed control, use only on no-till or minimum tillage fields where the previous year's crop residue has not been incorporated into the soil.
- [Plant wheat no sooner than 7 days after application of this product in the states of DE, ID, KY, MD, MN, MT, NC, ND, NJ, OR, SC, SD, TN, VA or WA]
- [Plant wheat no sooner than 14 days after application of this product in the states of DE, ID, KY, MD, MN, MT, NC, ND, NJ, OR, SC, SD, TN, VA or WA]
- Do not use on Durum wheat.
- Do not irrigate between emergence and spike.
- Wheat must be planted a minimum of 1 inch deep.
- Do not graze until wheat has reached 5 inches in height.

Burndown Use Directions

This product, applied as part of a burndown program at 2 fluid ounces per acre, may be used for residual weed control, as well as to assist in postemergence burndown of many weeds where wheat will be planted directly into the residue of the previous crop. See "DIRECTIONS FOR USE IN FALL BURNDOWN PROGRAMS IN FIELDS TO BE PLANTED TO BARLEY, FIELD PEA, FLAX, LENTIL, SAFFLOWER, SUNFLOWER AND WHEAT" for rates and timing of applications. For control of emerged weeds, this product must be applied with an appropriate burndown tank mix partner. To ensure thorough coverage, use a minimum of 15 gallons of spray solution per acre. Refer to tank mix partner's label for specified application pressure and adjuvant systems.

[POST-PLANT PRE-EMERGENCE WEED CONTROL]

[RESTRICTIONS]

- For post-plant, pre-emergence weed control, use only on no-till or minimum tillage fields where the previous crop residue has not been incorporated into the soil.
- Apply this product up to 2 days after planting.
- Do not use on Durum wheat.
- Do not irrigate between emergence and spike.
- Wheat must be planted a minimum of 1 inch deep.
- Do not graze until wheat has reached 5 inches in height.

Use Directions

This product, applied at 2 fluid ounces per acre, may be used for residual weed control, where wheat has been planted directly into the residue of the previous year. Application must be made no later than 2 days after planting.]

HARVEST AID

RESTRICTIONS

- Do not harvest within 10 days of application.

Use Directions

This product, applied at 2 fluid ounces per acre for desiccation requires the addition of an agronomically approved adjuvant to the spray mixture. Use a methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 quart per acre. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 pounds per acre or a 28 to 32% nitrogen solution at 1 to 2 quarts per acre) may be added to the spray mixture along with methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil. Tank mixing this product with glyphosate will increase control of emerged weeds and aid in harvest.

To ensure thorough coverage, use a minimum of 10 gallons spray solution per acre by ground application and a minimum of 5 gallons per acre by aerial application. Nozzle selection must meet manufacturer's gallonage and pressure specifications for postemergence application.

TIMING TO WHEAT

Apply this product, at 1.5 to 2 fluid ounces per acre, after wheat reaches the hard dough stage and grain has no more than 30% moisture. Wheat can be harvested 10 days after application. Liberty recommends tank mixing with glyphosate.

DIRECTIONS FOR USE IN BUSHBERRY, CANEBERRY, CITRUS FRUIT, GRAPE, TREE NUT, OLIVE, POME FRUIT, POMEGRANATE, STONE FRUIT AND NON-BEARING FRUIT TREES

Bushberry (Subgroup 13-07B): Aronia Berry; Blueberry, Highbush; Blueberry, Lowbush; Buffalo Currant; Chilean Guava; Cranberry, Highbush; Currant, Black; Currant, Red; Elderberry, European; European Barberry; Gooseberry; Honeysuckle, edible; Huckleberry; Jostaberry; Juneberry (Saskatoon Berry); Lingonberry; Native Currant; Salal; Sea Buckthorn; cultivars, varieties, and/or hybrids of these.

Caneberry (Subgroup 13-07A): Blackberry, Loganberry, Black Raspberry, Red Raspberry, Wild Raspberry cultivars, varieties and/or hybrids of these.

Citrus Fruit (Crop Group 10-10): Australian Desert Lime; Australian Finger-lime; Australian Round Lime; Brown River Finger Lime; Calamondin; Citron; Citrus hybrids; Grapefruit; Japanese Summer Grapefruit; Kumquat; Lemon; Lime; Mediterranean Mandarin; Mount White Lime; New Guinea Wild Lime; Orange, Sour; Orange, Sweet; Pummelo; Russell River Lime; Satsuma Mandarin; Sweet Lime; Tachibana Orange; Tahiti Lime; Tangelo; Tangerine (mandarin); Tangor; Trifoliolate Orange; Uniq Fruit; cultivars, varieties and/or hybrids of these.

Tree Nut (Crop Group 14-12): African Nut-tree; Almond, Beechnut; Brazil Nut; Brazilian Pine; Bunya; Bur Oak; Butternut; Cajou Nut; Candenut; Cashew; Chestnut; Chinquapin; Coconut; Coquito Nut; Dika Nut; Ginkgo; Guiana Chestnut; Hazelnut (Filbert); Heartnut; Hickory Nut; Japanese Horse-chestnut; Macadamia Nut; Mongongo Nut; Monkey-pot; Monkey Puzzle Nut; Okari Nut; Pachira Nut; Peach Palm Nut; Pecan; Pequi; Pili Nut; Pine Nut; Pistachio; Sapucaia Nut; Tropical Almond; Walnut, Black; Walnut, English; Yellowhorn, cultivars, varieties and/or hybrids of these.

Pome Fruit (Crop Group 11-10): Apple; Azarole; Crabapple; Loquat; Mayhaw; Medlar; Pear; Pear, Asian; Quince; Quince, Chinese; Quince, Japanese; Tejocote; cultivars, varieties and/or hybrids of these.

Stone Fruit (Crop Group 12-12): Apricot; Apricot, Japanese; Capulin; Cherry, Black; Cherry, Nanking; Cherry, Sweet; Cherry, Tart; Jujube, Chinese; Nectarine; Peach; Plum; Plum, American; Plum, Beach; Plum, Canada; Plum, Cherry; Plum, Chickasaw; Plum, Damson; Plum, Japanese; Plum, Klamath; Plum, Prune; Plumcot; Sloe and cultivars, varieties and/or hybrids of these.

RESTRICTIONS

- Do not apply more than 12 fluid ounces (0.375 lb a.i.) of this product per acre per application, except Caneberries; for Caneberries do not apply more than 6 fluid ounces (0.188 lb a.i.) this product per acre per application.
- Do not apply more than 24 fluid ounces (0.750 lb a.i.) of this product per acre per year, except Bushberries; for Bushberries do not apply more than 12 fluid ounces (0.357 lb a.i.) of this product per acre per year.
- Do not make more than 2 applications per year.
- Do not make a sequential application within 30 days of the first application, except nut trees, do not make a sequential application within 60 days of the first application.
- Do not apply to farm alleys or roads where traffic may result in treated dust settling onto crops or other desirable vegetation.
- Do not apply within 300 yards of non-dormant pears.
- Do not apply to powdery soils or soils that are susceptible to wind displacement unless irrigation can be applied immediately after application.
- Follow the most restrictive label limitations and precautions of the tank mix product(s) being used.
- Do not apply to nut trees established less than one year, unless protected from spray contact by non-porous wraps, grow tubes, or waxed containers.

- Preharvest Interval (PHI)
 - Citrus Fruit: 3 days
 - Bushberries: 7 days
 - Caneberries: 7 days
 - Grape: 60 days
 - Nut Trees: 60 days
 - Olive: 60 days
 - Pome Fruit: 60 days
 - Pomegranate: 60 days
 - Stone Fruit: 60 days

PRECAUTIONS

- Raise mower height during all mowing to reduce dust. Dust created by mowing can drift onto desirable vegetation resulting in injury.
- Do not mow treated areas between bud break and final harvest. Dust created by mowing may drift onto desirable vegetation resulting in injury.

Notes

- Use a maximum rate of 6 fluid ounces (0.188 lb a.i.) per acre per application of this product on any soil that has a sand plus gravel content over 80% if bushes, trees or vines are less than 3 years of age. (Two applications of 6 fluid ounces (0.188 lb a.i.) per acre in a year can still be made as long as there have been 60 days between applications).
- Avoid direct or indirect spray contact to foliage and green bark or canes (non-barked trunk and non-barked vines with the exception of undesirable suckers).
- Irrigate after application with minimum of 1/4 inch of water to activate the herbicide and to reduce wind displacement of soil.
- Follow the most restrictive label limitations and precautions of the tank mix product(s) being used.

For bushberries, caneberries, citrus fruit, grape, nut trees, olive, pomegranate and non-bearing fruit trees, a uniform broadcast application of this product to the orchard or vineyard floor or as a uniform band directed at the base of the bush, cane, trunk or vine. For stone fruit and pear this product can only be applied as a uniform band directed at the base of the trunk prior to “bud break” in stone fruit and pear. For apple, this product can only be applied as a uniform band directed at the base of the trunk prior to “pink bud”. For other pome fruit, check with your LIBERTY CROP PROTECTION LLC representative for application timing. The preferred application timing for this product is in the fall to maximize the potential for rainfall to activate and set the herbicide. Do not apply over the top of crop or allow spray to come in contact with crop as a result of application or drift.

Preemergence Application

Apply 6 to 12 fluid ounces (maximum 6 fluid ounces per acre for caneberries) of this product per broadcast acre as a preemergence application. Make preemergence (to weed emergence) applications of this product to a weed-free soil surface. Preemergence applications of this product must be completed prior to weed emergence. Moisture is necessary to activate this product on soil for residual weed control. Dry weather following application of this product may reduce effectiveness. However, when adequate moisture is received after dry conditions, this product will control susceptible germinating weeds.

Postemergence Application

Apply 6 to 12 fluid ounces (maximum 6 fluid ounces per acre for caneberries) of this product per broadcast acre plus an adjuvant (0.25% v/v non-ionic surfactant or 1 quart per acre crop oil concentrate). The addition of an adjuvant enhances activity of this product on emerged weeds. Thorough spray coverage is necessary to maximize the postemergence activity of this product.

Refer to **Table 10** for weeds controlled by the residual activity of this product. Tank mix this product with a labeled burndown herbicide for control of the emerged weeds listed in **Table 13**. Refer to tank mix partner’s label for additional weed species and increased weed heights claimed. Refer to tank mix partner’s label for additional restrictions, including minimum carrier volume and crops in which tank mix partner may be used. Burndown tank mix partners include glyphosate, paraquat, 2,4-D and Glufosinate-ammonium. Tank mixes

with glyphosate or 2,4-D containing products are not recommended during the period after bloom through final harvest to ensure crop safety from drift.

Residual weed control will be reduced if vegetation prevents this product from reaching the soil surface. If vegetation is heavy, use a burndown herbicide with this product and make a sequential application of this product prior to the emergence of new weeds.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Carrier Volume and Spray Pressure

To ensure thorough coverage in burndown applications, use a minimum of 15 gallons of spray solution per acre. Use higher gallonage if dense vegetation or heavy crop residue is present.

Nozzle selection must meet manufacturer's gallonage and pressure specifications.

Banded Application

Rates listed in **Table 13**, refer to a broadcast application covering the entire acre. When making a banded application, use the following formula to calculate amount needed per acre:

$$\frac{\text{Band Width in Inches}}{\text{Row Width in Inches}} \times \text{Rate per Broadcast Acre} = \text{Amount Needed per Acre for Banded Application}$$

USE RESTRICTIONS FOR BUSHBERRIES

- Do not use in the states of Idaho, Oregon or Washington except west of the Cascade Mountains in the following counties:
 - Oregon:** Benton, Clackamas, Clatsop, Columbia, Coos, Curry, Douglas, Jackson, Josephine, Lane, Lincoln, Linn, Marion, Multnomah, Polk, Tillamook, Umatilla, Yamhill and Washington
 - Washington:** Benton, Clallam, Clark, Cowlitz, Franklin, Grant, Grays Harbor, King, Jefferson, Kitsap, Lewis, Pacific, Pierce, Skagit, Snohomish, Thurston, Wahkiakum, Walla Walla and Whatcom
- Do not apply to bushberries established less than 2 years unless they are protected from spray contact by nonporous wrap, grow tubes or waxed containers.
- Do not apply within 7 days of harvest.

USE RESTRICTIONS FOR GRAPES

- Do not apply within 60 days of harvest.
- Do not apply to grapes established less than 2 years unless they are trellised at least 3 feet from the soil surface or are protected from spray contact by non-porous wrap, grow tubes or waxed containers.
- Do not apply to grapes that are not trellised or staked unless they are free standing.

Juice, Raisin and Wine Grapes:

- Do not apply during the period after bud break through final harvest, unless using shielded application equipment and applicator can ensure spray drift will not come in contact with crop fruit or foliage. Shielded applications during this time period must not be made with glyphosate or products containing glyphosate.

Table Grapes

- Do not apply after bud break. **Note:** This product may be applied during the period following final harvest up to bud break.

NOTES FOR GRAPES

- Avoid direct or indirect spray contact to foliage and green bark (non-barked vines, with the exception of undesirable suckers).

- Plant new plantings of "own-rooted varieties", including Concord, so that all roots are a minimum 8 inches below the soil surface to be treated. In some situations, this may require hilling soil around newly planted vines so that the settled depth of the hill will be 4 to 5 inches above the vineyard floor.

USE RESTRICTIONS FOR CITRUS FRUIT, TREE NUTS, OLIVE, POME FRUIT, POMEGRANATE AND STONE FRUIT

- Do not apply to pears in the states of Oregon or Washington.
- For pome fruit and stone fruit, this product can only be applied as a uniform band directed at the base of the trunk prior to silver tip in apples and bud break in stone fruit.
- For pome fruit and stone fruit do not apply to row middles (area between berms)
- For nut trees, olive and pomegranate apply after bud break through final harvest using shielded application equipment if the applicator can ensure the spray drift will not come into contact with non-target vegetation, crop fruit and/or foliage. Shielded application equipment is not required if the following application parameters are followed:
 - Application pressure (at boom) < 30 PSI.
 - Application speed < 5 MPH.
 - Applicator can ensure the spray drift will not come into contact with non-target vegetation, crop fruit and/or foliage.
- Do not apply to trees established less than one year, unless protected from spray contact by non-porous wraps, grow tubes, paint or waxed containers.
- Do not use in the states of Oregon or Washington except in the following counties unless the additional restrictions listed below are followed:
 - Oregon:** Benton, Clackamas, Clatsop, Columbia, Coos, Curry, Douglas, Jackson, Josephine, Lane, Lincoln, Linn, Marion, Morrow, Multnomah, Polk, Tillamook, Umatilla, Yamhill and Washington
 - Washington:** Clallam, Cowlitz, Grays Harbor, King, Jefferson, Kitsap, Lewis, Pacific, Pierce, Skagit, Snohomish, Thurston, Wahkiakum and Whatcom
 - For apples east of the Cascade Mountains in Washington (counties not listed above), follow the restrictions above plus:
 - Apply between final harvest and January 1.
 - Apply only to apple blocks with an established (2 years or older) permanent cover crop that covers a minimum of 60% of the surface area in the block.
 - Application must be incorporated with a minimum of one half inch of water within 48 hours after application.
 - Do not apply to powdery soils or soils susceptible to wind displacement.
 - Apply only to orchard berms.
 - Do not mow the treated berm areas of the orchard

USE RESTRICTIONS FOR NON-BEARING FRUIT AND NUT TREES

Non-Bearing Avocado and Fig

- Do not apply more than 12 fluid ounces (0.375 lb a.i.) of this product per acre per application.
- Do not apply more than 24 fluid ounces (0750 lb a.i.) of this product per acre per year.
- Do not harvest fruit from treated trees within one year of application.
- Do not apply to trees established less than one year, unless protected from spray contact by non-porous wraps, grow tubes or waxed containers.
- Do not apply during the period after flowering through leaf drop, unless using shielded application equipment and the applicator can ensure spray drift will not come in contact with the crop foliage.

USE PRECAUTIONS FOR ALMOND AND STONE FRUIT IN DEFINED AREAS OF MERCED, SAN JOAQUIN AND STANISLAUS COUNTIES OF CALIFORNIA

The use of this product in soils common in parts of Merced, San Joaquin and Stanislaus counties in California is known to have resulted in injury to almonds under drought stress conditions. These soils are characterized by having been cut or filled, high sand content, low clay content and shallow profiles. Growers in the Defined Area must be aware and assume the risk of using this product on almond or stone fruit crops. The Defined Area can be seen on the Map or by the description that follows:

- Intersection of Highway 4 and Escalon Bellota Road at Farmington in San Joaquin County;
- Directly South on Escalon-Bellota to the Santa Fe Avenue and railroad tracks at Escalon
- Southeast on Santa Fe Avenue down to the Merced River;
- East following the Merced River to the Merced/Mariposa County line;
- Northwest following the Merced County line through the intersection of Merced and Stanislaus County line following the Stanislaus/Tuolumne County and Calaveras County line to Highway 4;
- West on Highway 4 back to the Farmington intersection of Escalon — Bellota Road.



Table 13. Weeds Controlled by Postemergence Activity of Tank Mixes of LIBERTY FLUMIOX 44

COMMON NAME	SCIENTIFIC NAME	WEED HEIGHT/ LENGTH (Inches)	APPLICATION RATE
BROADLEAF WEED SPECIES			
Bindweed, Field ¹	<i>Convolvulus arvensis</i>	8	6 to 12 fl oz/A
Carpetweed	<i>Mollugo verticillata</i>	4	
Chickweeds, Common	<i>Stellaria media</i>	4	
Mouseear	<i>Cerastium vulgatum</i>	4	
Cocklebur, Common	<i>Xanthium strumarium</i>	4	

COMMON NAME	SCIENTIFIC NAME	WEED HEIGHT/ LENGTH (Inches)	APPLICATION RATE
Eveningprimrose, Cutleaf ²	<i>Oenothera laciniata</i>	12	
Filaree,			
Broadleaf	<i>Erodium botrys</i>	4	
Redstem	<i>Erodium cicutarium</i>	4	
Florida Beggarweed	<i>Desmodium tortuosum</i>	2	
Hemp Sesbania	<i>Sesbania exaltata</i>	8	
Jimsonweed	<i>Datura stramonium</i>	4	
Lambsquarters, Common	<i>Chenopodium album</i>	4	
Morningglories,			
Entireleaf	<i>Ipomoea hederacea</i> var. <i>integriuscula</i>	4	
Ivyleaf	<i>Ipomoea hederacea</i>	4	
Pitted	<i>Ipomoea lacunosa</i>	6	
Red/Scarlet	<i>Ipomoea coccinea</i>	4	
Tall	<i>Ipomoea purpurea</i>	4	
Mustard, Wild	<i>Brassica kaber</i>	6	
Pigweeds,			
Palmer Amaranth	<i>Amaranthus palmeri</i>	6	
Redroot	<i>Amaranthus retroflexus</i>	6	
Smooth	<i>Amaranthus hybridus</i>	6	
Plantain, Broadleaf	<i>Plantago major</i>	6	
Prickly Sida (Teaweed)	<i>Sida spinosa</i>	6	
Purslanes,			
Common	<i>Portulaca oleracea</i>	4	
Rock	<i>Calandrinia</i> spp.	2	
Ragweeds,			
Common	<i>Ambrosia artemisiifolia</i>	2	
Giant	<i>Ambrosia trifida</i>	4	
Rice Flatsedge	<i>Cyperus iria</i>	4	
Sicklepod	<i>Senna obtusifolia</i>	4	
Smartweeds,			
Ladysthumb	<i>Polygonum persicaria</i>	4	
Pale	<i>Polygonum lapathifolium</i>	4	
Pennsylvania	<i>Polygonum pennsylvanicum</i>	4	
Spotted Spurge	<i>Euphorbia maculata</i>	4	
Velvetleaf	<i>Abutilon theophrasti</i>	4	
Venice Mallow	<i>Hibiscus trionum</i>	4	
Waterhemp,			
Common	<i>Amaranthus rudis</i>	2	
Tall	<i>Amaranthus tuberculatus</i>	2	

¹ This product will only provide control of the above ground portion of bindweed. Repeated applications will be needed to control regrowth.

² For acceptable control, cutleaf evening primrose should be 12 inches or less and in the rosette stage. Add crop oil concentrate, at 1 pint per acre, or non-ionic surfactant at 0.25% v/v, to glyphosate tank mixes for cutleaf evening primrose control, including glyphosate formulations that contain a built-in adjuvant system.

ADDITIONAL RESIDUAL WEED CONTROL

This product may be tank mixed with diuron, oryzalin or simazine for additional residual weed control. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

DIRECTIONS FOR USE TO MAINTAIN BARE GROUND ON NON-CROP AREAS OF FARMS, ORCHARDS AND VINEYARDS

RESTRICTIONS

- Do not apply to farm alleys or roads where traffic may result in treated dust settling onto crops or other desirable vegetation.
- Do not apply to ditch banks.

This product, when used as directed, can be used on farms, orchards and vineyards for non-selective vegetation control to maintain bare ground on non-crop areas that must be kept weed free. Follow all applicable directions as outlined above under "USE INFORMATION".

This product offers residual and postemergence control of susceptible broadleaf and grass weeds as well as an additional mode of action to assist in the control of ALS (acetolactate synthase) resistant weeds. This product can be tank mixed with the herbicides listed in **Table 14** for increased residual or postemergence control. The length of residual control is dependent on the rate applied as well as on rainfall and temperature conditions. Length of residual control will decrease as temperature and precipitation increase. Rates of this product of 6 to 12 fluid ounces per acre are required to provide residual control of the weeds listed in **Table 10**.

PREEMERGENCE APPLICATION

Apply 6 to 12 fluid ounces of this product per broadcast acre as a preemergence application. Make preemergence (to weed emergence) applications of this product to a weed-free soil surface. Preemergence applications of this product must be completed prior to weed emergence. Moisture is necessary to activate this product on soil for residual weed control. Dry weather following application of this product may reduce effectiveness. However, when adequate moisture is received after dry conditions, this product will control susceptible germinating weeds.

POSTEMERGENCE APPLICATION

Apply 6 to 12 fluid ounces of this product per broadcast acre plus an adjuvant (0.25% v/v non-ionic surfactant or 1 quart per acre crop oil concentrate). The addition of an adjuvant enhances activity of this product on emerged weeds. Thorough spray coverage is necessary to maximize the postemergence activity of this product. Emerged weeds are controlled postemergence with this product; however, translocation of this product within a weed is limited, and control is affected by spray coverage and by the addition of an adjuvant. The most effective postemergence weed control with this product occurs when applied in combination with a surfactant to weeds less than 2 inches in height. Use a tank mix partner in combination with this product for the postemergence control of weeds larger than 2 inches. Tank mix partners are listed in **Table 14**.

Important: Completely read and follow the label of any potential tank mix partner with this product. When using tank mixtures, use conditions must be in accordance with the most restrictive of the label limitations and precautions on either herbicide label.

Table 14. Tank Mix Combinations to Maintain Bare Ground on Non-Crop Areas

2,4-D	Glufosinate-ammonium	Glyphosate	Paraquat
-------	----------------------	------------	----------

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Pesticide Storage

Keep pesticide in original container. Store in a cool, dry, secure place. Do not put formulation or dilute spray solution into food or drink containers. Do not contaminate food or foodstuffs. Do not store or transport near feed or food. Not for use or storage in or around the home. For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC (800) 424-9300.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling

NONREFILLABLE CONTAINER (EQUAL TO OR LESS THAN 5 GALLONS): Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

NONREFILLABLE CONTAINER (GREATER THAN 5 GALLONS): Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

REFILLABLE CONTAINER: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. After triple rinsing is complete, and the container is not suitable for refilling or reconditioning, offer the container for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of LIBERTY CROP PROTECTION LLC or Seller. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW all such risks shall be assumed by Buyer and User and Buyer and User agree to hold LIBERTY CROP PROTECTION LLC and Seller harmless for any claims relating to such factors.

LIBERTY CROP PROTECTION LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or LIBERTY CROP PROTECTION LLC, and TO THE EXTENT CONSISTENT WITH APPLICABLE LAW Buyer and User assume the risk of any such use. To the extent

consistent with applicable law LIBERTY CROP PROTECTION LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither LIBERTY CROP PROTECTION LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF LIBERTY CROP PROTECTION LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LIBERTY CROP PROTECTION, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

LIBERTY CROP PROTECTION LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of LIBERTY CROP PROTECTION LLC.

All trademarks are the property of their respective owners.