



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505P)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

89168-116

Date of Issuance:

1/25/22

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

LIBERTY ST-SB

Name and Address of Registrant (include ZIP Code):

Liberty Crop Protection  
1880 Fall River Drive Suite 100  
Loveland, Colorado 80538

Karen Murphy  
Regulatory Manager  
Liberty Crop Protection

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

- 1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Cynthia Giles-Parker, Chief  
Fungicide Branch, Registration Division (7505P)

Date:

1/25/22

2. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, “EPA Reg. No. 89168-116.”
3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 10/31/2020
- Alternate CSF 1 dated 10/31/2020
- Alternate CSF 2 dated 10/31/2020
- Alternate CSF 3 dated 03/31/2021

If you have any questions, please contact Senedu Alemu via email at [alemu.senedu@epa.gov](mailto:alemu.senedu@epa.gov)

Enclosure -stamped “accepted” label

METALAXYL	GROUP	4	FUNGICIDE
THIOPHANATE-METHYL	GROUP	1	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE

# LIBERTY ST – SB

For use as a seed treatment for protection of seed and seedlings against listed diseases on soybeans, dry beans, and snap beans.

**ACTIVE INGREDIENTS:**

Metalaxyl.....14.26%  
 Thiophanate -methyl .....9.30%  
 Fludioxonil .....2.31%

**OTHER INGREDIENTS:** .....74.13%

**TOTAL:** ..... 100.00%

Contains 1.31 lb metalaxyl, 0.86 lb thiophanate-methyl, and 0.21 lb fludioxonil per gallon.

**KEEP OUT OF REACH OF CHILDREN**


## CAUTION

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, Exposure or Medical Emergencies Call CHEMTREC (800) 424-9300

EPA REG. NO. 89168-XX  
 EPA EST. NO. \_\_\_\_\_

NET CONTENTS: \_\_\_\_\_ GAL



Manufactured for:  
 LIBERTY CROP PROTECTION, LLC  
 1880 Fall River Drive  
 Suite 100  
 Loveland, CO 80537

011422

<b>FIRST AID</b>	
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
<p>Have the product container or label with you when calling a poison control center, doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents) call CHEM-TREC at 1-800-424-9300.</p>	

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Avoid contact with eyes or clothing. Wear long-sleeved shirt and long pants, socks and shoes.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks,
- Chemical-resistant gloves of Barrier Laminate, Butyl Rubber ≥14 mils, Nitrile Rubber ≥14 mils, Neoprene Rubber ≥14 mils, Polyvinyl Chloride (PVC) ≥14 mils, or Viton ≥14 mils for all mixers and loaders and for applicators using hand held equipment, and
- Chemical-resistant apron for mixers, loaders and other handlers exposed to the concentrate.

**Note:** Persons involved in bagging treated seed, sewing or moving bags of treated seed, or cleaning up bagging areas or seed treatment equipment are pesticide handlers and must wear the PPE required on this label for pesticide handlers.

**USER SAFETY REQUIREMENTS**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. For terrestrial use, **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wash water or rinsate. Cover or incorporate spilled treated seeds.

Groundwater Advisory: Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

### PHYSICAL AND CHEMICAL HAZARDS

**DO NOT** mix or allow coming in contact with oxidizing agents. Hazardous chemical reaction may occur.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Maximum usage when applying metalaxyl containing products to the same crop within the same season: **DO NOT** apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.

The maximum annual rate must not exceed 12.3 lb a.i./A

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

LIBERTY ST–SB is formulated for on-farm or commercial seed treatment using standard slurry or mist-type seed treatment equipment.

Mix thoroughly before use or use entire container at one time. Pre-test all tank mixes to determine physical compatibility between formulations. Observe the most restrictive use precautions and limitations on labeling of the products used in mixtures.

LIBERTY ST–SB contains an EPA approved dye/colorant that imparts an unnatural color to the seed.

Treated seed must not be used for or mixed with food or animal feed or processed for oil. Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the Federal Seed Act.

[NOTE TO REVIEWER: [One of the following statements will be used on the final printed labeling. Liberty Crop Protection, LLC reserves the option to choose the statement that will be used].

[STATEMENT 1: Use the entire contents of LIBERTY ST–SB in this container in the use season immediately following purchase from Liberty Crop Protection, LLC. Store at temperatures above 40 °F but below 80 °F.]

[STATEMENT 2: The LIBERTY ST–SB in this container has been manufactured for use in this use season. Use the entire contents of LIBERTY ST–SB in this container in the use season immediately following purchase from Liberty Crop Protection, LLC. Store at temperatures above 40 °F but below 80 °F. Remaining product is non-returnable.]

### **PRODUCT INFORMATION**

LIBERTY ST–SB is a seed treatment fungicide for use against damping-off, seed, and seedling rots due to Pythium, Fusarium, Rhizoctonia spp. In soybeans, dry beans, and snap beans. LIBERTY ST–SB also suppresses seed-borne Sclerotinia and Phomopsis spp. particularly in soybeans.

Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application to seed is necessary to ensure seed safety and best disease protection. Seed must be sound and well cured prior to treatment. Product must be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates for the crop to be treated with this product.

### **RESISTANCE-MANAGEMENT**

For resistance management, LIBERTY ST-SB contains Group 1/thiophanate-methyl, Group 4/metalaxyl and Group 12/fludioxonil fungicides. Any fungal population may contain individuals naturally resistant to LIBERTY ST-SB and other Group 1, 4 and 12 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Follow appropriate resistance-management strategies.

To delay fungicide resistance, take one or more of the following steps:

- Avoid application of more than labeled maximum number of applications and consecutive sprays of LIBERTY ST-SB or other fungicides in the same group in a season.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM guidelines for specific crops and pathogens.
- For further information or to report suspected resistance contact Liberty Crop Protection at 844-425-8488. You can also contact your pesticide distributor or university extension specialist to report resistance.

### **TANK MIXES**

Pre-test all tank mixes to determine physical compatibility between formulations.

**NOTICE:** Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor, to the extent consistent with applicable law.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

### **MIXING INSTRUCTIONS FOR SOYBEANS**

LIBERTY ST–SB is formulated for on-farm or commercial treatment to be used with liquid rhizobia products, using standard mechanical slurry or mist-type seed treatment equipment which accurately measures and mixes a flowable seed treatment.

The equipment must provide uniform coverage of LIBERTY ST–SB on the seed. Uneven seed coverage may not give the desired level of disease control. LIBERTY ST–SB must be diluted with either water or liquid inoculants prior to application to soybeans. Consult the manufacturer of the application equipment you plan to use for suitability for this application and for instructions on operation and calibration of the equipment. **DO NOT** mix LIBERTY ST–SB fungicide with other pesticides unless sufficient compatibility, loading rates, and seed and crop safety factors are determined. **DO NOT** mix with any product that contains a prohibition on mixing. Follow the more restrictive labeling of any tank-mix partner.

In a clean container large enough to contain the entire mixture add the specified amount of LIBERTY ST – SB. Add half of the required water or liquid inoculant and mix thoroughly. Add the remaining amount of water or liquid inoculant and mix again. The slurry is ready for application to the soybeans. Mix as needed and **DO NOT** store diluted mixture.

Additional or continuous agitation or mixing of the slurry mixture may be necessary to prevent the LIBERTY ST–SB from settling out of solution when using water or liquid inoculants. LIBERTY ST–SB has been found to be compatible with some liquid inoculant products. LIBERTY ST–SB may be mixed or applied sequentially with approved liquid inoculants. Consult the maker of the liquid inoculants and a Liberty Crop Protection representative for directions before applying LIBERTY ST–SB with liquid inoculants.

**NOTE:** LIBERTY ST–SB is formulated to be compatible with some liquid inoculants. If water is used to dilute LIBERTY ST–SB when rhizobia are present, use only nonchlorinated water. Chlorinated water will kill the live rhizobia contained in the inoculant.

### **MIXING INSTRUCTIONS FOR OTHER BEANS (DRY and SUCCULENT)**

Apply LIBERTY ST–SB as a water based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease protection. Thoroughly mix the labeled amount of LIBERTY ST–SB into the required amount of water for the slurry treater and dilution rate to be used.

Follow the manufacturer application instructions for the seed treatment equipment being used. Maintain constant agitation of the slurry during the treatment. Allow seed to dry before bagging.

### **USE RESTRICTIONS**

**DO NOT** use treated seed for feed, food, or oil.

**DO NOT** use for hopper box or planter box seed treatments.

### **Seed Bag Label Requirements**

The Federal Seed Act requires that containers containing treated seeds shall be labeled with the following statements:

- This seed has been treated with Liberty ST-SB a fungicide containing the active ingredients metalaxyl, thiophanate-methyl, fludioxonil.
- **DO NOT** use for food, feed, oil production or any other purpose except planting.

THE U.S. ENVIRONMENTAL PROTECTION AGENCY REQUIRES THE FOLLOWING STATEMENTS ON CONTAINERS CONTAINING TREATED SEED:

- Store treated seed away from food and feedstuffs.
- **DO NOT** allow children, pets, or livestock to have access to treated seeds.
- Wear long pants, long-sleeved shirt, shoes, socks, and chemical-resistant gloves when handling treated seed.
- Treated seeds exposed on the soil surface may be hazardous to birds and wildlife.
- Dispose of all excess treated seed by burying seed away from bodies of water. Cover or collect treated seeds spilled during loading and planting.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- **DO NOT** contaminate bodies of water when disposing of planting equipment wash water.
- **DO NOT** graze or feed livestock on treated areas for 30 days after planting.
- Dispose of seed packaging or containers in accordance with local requirements.
- Excess treated seed may be used for ethanol production if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.

**Groundwater Advisory:** Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

After the seeds have been planted, **DO NOT** enter or allow worker entry into treated areas during the **restricted-entry interval (REI) of 48 hours**. Exception: If the treated seed is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil or water is: coveralls, chemical-resistant gloves made of any waterproof material, and chemical-resistant footwear.

**CROP SPECIFIC USE DIRECTIONS**

<b>SOYBEANS</b>				
<b>DISEASE</b>	<b>USE RATE (based on 3,000 seeds / lb)</b>			
	fl.oz. / cwt. seed*	fl.oz. / 1,000 seed	fl.oz. / 140,000 seed	mg AI / seed
Protection against damping-off and seed and seedling rots caused by <i>Pythium</i> , <i>Fusarium</i> , and <i>Rhizoctonia</i> spp., and others	1.5	0.005	0.7	0.0232 metalaxyl 0.0152 thiophanate-methyl 0.0037 fludioxonil
Protection against <i>Phytophthora</i> (see application instructions below)				
Suppresses seed-borne <i>Sclerotinia</i> and <i>Phomopsis</i> spp.				*Contains 4.642 g of metalaxyl, 3.048 g of thiophanate-methyl, and 0.744 g of fludioxonil per fl.oz (100 kg seed contains 15.35 grams metalaxyl, 10.08 grams thiophanate-methyl, and 2.28 grams fludioxonil)
<b>Application Instructions</b>				
Apply Liberty ST-SB at the rate specified in the rate table, plus water to prepare a slurry of sufficient volume and viscosity to adequately cover the seed. As an alternative to water, you may use approved liquid rhizobia inoculants at rates specified by the manufacturer and may add the inoculants to the slurry to achieve sufficient total slurry volume and viscosity to adequately cover the seed.				
The addition of an application of another approved metalaxyl or mefenoxam containing seed treatment product may be used if targeting fields with high <i>Phytophthora</i> pressure. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.				



**RESTRICTIONS**

After the seeds have been planted, **DO NOT** enter or allow worker entry into treated areas during the **restricted-entry interval (REI) of 48 hours**. Exception: If the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil or water is: coveralls, chemical resistant gloves made of any waterproof material, and chemical-resistant footwear.

**DRY BEANS, SUCCULENT (SNAP) BEANS** (*Cicer arietinum* (chickpea, garbanzo bean); *Lupinus* spp. (sweet lupine, white sweet lupine, white lupine, and grain lupine). *Phaseolus* spp. (kidney bean, lima bean, mung bean, navy bean, pinto bean, snap bean, and waxbean); *Vicia faba* (broad bean, fava bean); *Vigna* spp. (asparagus bean, blackeyed pea and cowpea)

<b>DISEASE</b>	<b>USE RATE (fl.oz / cwt. seed)</b>
<p>For protection against damping-off and seed and seedling rots caused by <i>Pythium</i>, <i>Fusarium</i>, and <i>Rhizoctonia</i> spp. and others.</p> <p>Protection against <i>Phytophthora</i> (see application instructions below)</p> <p>Suppresses seed-borne <i>Sclerotinia</i> and <i>Phomopsis</i> spp.</p>	<p><b>1.5</b></p>

**Application Instructions**  
 The addition of an application of another approved metalaxyl or mefenoxam containing seed treatment product may be used if targeting fields with high *Phytophthora* pressure. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**RESTRICTIONS**

After the seeds have been planted, **DO NOT** enter or allow worker entry into treated areas during the **restricted-entry interval (REI) of 48 hours**. Exception: If the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil or water is: coveralls, chemical resistant gloves made of any waterproof material, and chemical-resistant footwear.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool place, **DO NOT** store in direct sunlight. Protect from freezing temperatures.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### [Nonrefillable Containers 5 Gallons or Less]

**Nonrefillable container. DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

### [Nonrefillable containers larger than 5 gallons]

**Nonrefillable container. DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

### [Refillable containers larger than 5 gallons]

**Refillable container.** Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of LIBERTY CROP PROTECTION LLC or Seller. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW all such risks shall be assumed by Buyer and User and Buyer and User agree to hold LIBERTY CROP PROTECTION LLC and Seller harmless for any claims relating to such factors.

LIBERTY CROP PROTECTION LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or LIBERTY CROP PROTECTION LLC, and TO THE EXTENT CONSISTENT WITH APPLICABLE LAW Buyer and User assume the risk of any such use. To the extent consistent with applicable law LIBERTY CROP PROTECTION LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither LIBERTY CROP PROTECTION LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF LIBERTY CROP PROTECTION LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LIBERTY CROP PROTECTION, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

LIBERTY CROP PROTECTION LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of LIBERTY CROP PROTECTION LLC.