



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

89168-117

Date of Issuance:

3/17/22

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Liberty ST-W

Name and Address of Registrant (include ZIP Code):

Karen Murphy
Liberty Crop Protections LLC.
1880 Fall River Drive suite 100
Loveland, CO 80538

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. A one year study is required to satisfy these data requirements. You have 18 months from the date of registration to provide these data.

Signature of Approving Official:

Nate Mellor, Product Manager 21
Fungicide Branch, Registration Division (7505P)

Date:

3/17/22

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 89168-117.”
4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 12/3/2020
- Alternate CSF 1 dated 12/3/2020
- Alternate CSF 2 dated 12/3/2020
- Alternate CSF 3 dated 12/3/2020
- Alternate CSF 4 dated 12/3/2020

If you have any questions, please contact Carmen Swinger by phone at 202-566-2923, or via email at swinger.carmen@epa.gov.

Enclosure

ACCEPTED

03/17/2022

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 89168-117

METALAXYL	GROUP	4	FUNGICIDE
TEBUCONAZOLE	GROUP	3	FUNGICIDE
THIOPHANTE-METHYL	GROUP	1	FUNGICIDE

LIBERTY ST – W

For use as a seed treatment for protection of seed and seedlings against listed diseases on soybeans, dry beans, and wheat.

ACTIVE INGREDIENTS:

- Metalaxyl..... 3.60%
- Thiophanate-methyl 3.00%
- Tebuconazole 1.20%

OTHER INGREDIENTS: 92.20%

TOTAL: 100.00%

Contains 0.31 lb metalaxyl, 0.26 lb thiophanate-methyl, and 0.10 lb tebuconazole per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, Exposure or Medical Emergencies Call CHEMTREC (800) 424-9300

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to by a poison control center or doctor. • DO NOT give anything to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call poison control center or doctor.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information on this pesticide product, including health concerns, medical emergencies, or pesticide incidents, call CHEM-TREC at 1-800-424-9300.	

EPA REG. NO. 89168-XX

EPA EST. NO. _____

NET CONTENTS: _____ GAL

Manufactured for:
LIBERTY CROP PROTECTION, LLC
1880 Fall River Drive
Suite 100
Loveland, CO 80537

031522

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

1. Long-sleeved shirt and long pants
2. Shoes plus socks,
3. Chemical resistant gloves made out of: nitrile rubber \geq 14 mils, butyl rubber \geq 14 mils, neoprene rubber \geq 14 mils, barrier laminate, polyvinyl chloride (PVC) \geq 14 mils or Viton \geq 14 mils.

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170-240 (d) (4)], the handler PPE may be reduced or modified as specified in the WPS.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

Exposed treated seed may be hazardous to birds and wildlife. Dispose of all excess treated seed and seed packaging or containers by burial away from bodies of water in accordance with any local requirements. Cover, incorporate, or collect treated seeds spilled during loading and planting. **DO NOT** contaminate bodies of water when disposing of planting equipment wash water.

Groundwater Advisory: Metalaxyl and Tebuconazole are known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Surface Water Advisory: Tebuconazole This product contains tebuconazole and may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications, when rainfall is forecasted within 48 hours.

DO NOT discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. **DO NOT** discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact you State Water Board or Regional office of the EPA.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT mix or allow coming in contact with oxidizing agent. Hazardous Chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application: For any requirements specific to your State or Tribal Agency, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

After seeds have been planted, **DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Exception: If the product is soil injected or soil incorporated, the WPS, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

1. Long-sleeved shirt and long pants
2. Chemical resistant gloves made out of: nitrile rubber \geq 4 mils, butyl rubber \geq 14 mils, neoprene Rubber \geq 14 mils, barrier laminate, polyvinyl chloride (PVC) \geq 14 mils or Viton \geq 14 mils
3. Shoes plus socks
4. Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170) The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

RESTRICTION:

When applying both metalaxyl-containing and mefenoxam-containing products to the same crop within the same season **DO NOT** apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.

All seed must be adequately dyed with a suitable colorant to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, part 2.25. Any colorant added to treated seed must be cleared for use under 40 CFR, part 153.155.

USE INFORMATION

LIBERTY ST-W is a systemic and contact broad-spectrum fungicide for seed treatment that protects against a wide variety of specified diseases. LIBERTY ST-W is effective against the diseases Fusarium, Cochliobolus sativus, Pythium, smuts, bunt and leaf stripe. It is also effective against seed rot fungi including Aspergillus and Penicillium.

DO NOT use LIBERTY ST-W in combination with other seed treatment products unless compatibility has been verified. Read and follow carefully all label directions of each combination product. When using combinations of products, the most restrictive of label limitations and precautions must be followed.

Allow seed to dry before bagging.

Manual seed treatments are prohibited.

ALWAYS SHAKE OR STIR WELL BEFORE USE

RESISTANCE-MANAGEMENT

For resistance management, LIBERTY ST-W contains Group 1/thiophanate-methyl, Group 4/metalaxyl and Group 3/tebuconazole fungicides. Any fungal population may contain individuals naturally resistant to LIBERTY ST-W and other Group 1, 4 and 12 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Follow appropriate resistance-management strategies.

To delay fungicide resistance, take one or more of the following steps:

- Avoid application of more than labeled maximum number of applications and consecutive sprays of LIBERTY ST-W or other fungicides in the same group in a season.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as Cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact LIBERTY CROP PROTECTION at 844-425-8488. You can also contact your pesticide distributor or university extension specialist to report resistance.

SEED TREATMENT EQUIPMENT

LIBERTY ST-W may be used both in commercial seed treatment facilities and in on-farm application. **DO NOT** apply this product as a planter box or hopper box treatment. Apply LIBERTY ST-W with mechanical, slurry, or mist-type seed treating equipment, provided that the equipment can be calibrated to accurately and uniformly apply the product to seed. Uniform application to seed is necessary to ensure best disease and insect protection and optimum performance. For best activity of this product, seed must be sound and well-cured before treatment. Refer to the APPLICATION INSTRUCTIONS below for label rates.

In most cases, LIBERTY ST-W is ready to use and may be applied to seed as is; however, dilution with water may be appropriate for some types of treaters and/or when treating under dry and/or hot conditions, to achieve more uniform product to seed coverage. Mix LIBERTY ST-W and water thoroughly into a slurry prior to treating seed. Recalibrate treating equipment to compensate for the addition of water. The amount of dilution water that is most suitable depends on the seed, treater type and temperature. Contact your local Liberty Crop Care, LLC representative or supplier for specific recommendations.

WHEAT (SPRING and WINTER) APPLICATION INSTRUCTIONS

100 lbs. seed	Symptoms Controlled	Symptoms
4.0* fl. oz.	<p>General seed rots (including those caused by <i>Penicillium</i> and <i>Aspergillus</i>)</p> <p>Damping off and seedling blight (seed- and soil-borne <i>Fusarium</i>, <i>Pythium</i>, <i>Rhizoctonia</i> and <i>Cochliobolus sativus</i>)</p> <p>Loose Smut (<i>Ustilago tritici</i>)</p> <p>Common Bunt (<i>Tilletia caries</i> & <i>foetida</i>)</p> <p>Early Season Root Rot (<i>Rhizoctonia</i>)</p> <p>Early season Stinking smut, Flag smut & Loose smut</p> <p>Early season Septoria disease complex</p>	<p>Common root rot (<i>Cochliobolus sativus</i>)</p> <p>Crown and Foot Rot (<i>Fusarium</i>)</p> <p>Early season suppression of powdery mildew and wheat leaf rust</p> <p>Yellow Dwarf Virus symptoms and spread.</p>

*4.0 fluid ounces of Liberty ST-W contains 0.01 lb of metalaxyl, 0.008 lb of thiophanate-methyl and 0.003 lb of tebuconazole.

SOYBEAN & DRY BEAN APPLICATION INSTRUCTIONS

100 lbs. seed	Symptoms Controlled
4.0* fl. oz.	<p><i>Pythium</i> damping-off and early season <i>Phytophthora</i> control</p> <p>For early season control of seed decay (<i>Phomopsis</i>, <i>Fusarium</i>, <i>Rhizoctonia</i>) and suppression of seedling blight (<i>Fusarium</i> and <i>Rhizoctonia</i>)</p>

*4.0 fluid ounces of Liberty ST-W contains 0.01 lb of metalaxyl, 0.008 lb of thiophanate-methyl and 0.003 lb of tebuconazole.

USE RESTRICTIONS:

- **DO NOT** use treated seed for food, feed or oil purposes.
- **DO NOT** apply as a planter box or hopper box treatment.

- Care must be exercised in the handling of treated seed. **DO NOT** use augers used for handling treated seed to move seed for feed, food or oil processing.
- **DO NOT** re-use bags from treated seed to handle food or feed products.
- Wheat: **DO NOT** graze or feed livestock on treated areas.
- **DO NOT** feed vines or hay grown from treated seed to livestock.
- Maximum of one application per season.
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- **WHEAT: DO NOT** apply more than the maximum annual rate for metalaxyl of 12.3 lb a.i./A per acre per year.
- **WHEAT: DO NOT** apply more than the maximum annual of 0.7 lb a.i./A per acre per year from all applications containing thiophanate-methyl.
- **WHEAT: DO NOT** apply more than the maximum annual rate of 0.11 lb a.i./A per acre per year applications containing tebuconazole.
- **Soybeans: DO NOT** apply more than the maximum annual rate of 0.625 lb a.i./A per acre per year from all applications containing metalaxyl. Do not apply by seed treatment more than 0.0166 pounds of metalaxyl
- **Soybeans: DO NOT** apply more than the maximum annual rate of 1.4 lb a.i./A per acre per year from all applications containing thiophanate-methyl.
- **Soybeans: DO NOT** apply more than the maximum annual rate of 0.34 lb a.i./A per acre per year applications containing tebuconazole

Seed Bag Label Requirements

The Federal Seed Act requires that containers containing treated seeds shall be labeled with the following statements:

- This seed has been treated with Liberty ST-W, a fungicide containing the active ingredients metalaxyl, thiophanate-methyl and tebuconazole.
- **DO NOT** use for food, feed, oil production or any other purpose except planting.

The U.S. Environmental Protection Agency requires the following statements on containers containing seed treated with metalaxyl, thiophanate-methyl and tebuconazole:

- Store treated seed away from food and feedstuffs.
- **DO NOT** allow children, pets, or livestock to have access to treated seeds.
- **DO NOT allow livestock to graze cotton plants grown from treated seed.**
- Wear long pants, long-sleeved shirt and protective gloves when handling treated seed.
- Treated seeds exposed on the soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Treated seed must be planted into the soil at a depth greater than one inch.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- **DO NOT** contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed packaging or containers in accordance with local requirements.
- Excess treated seed may be used for ethanol production if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- This soybean/ dry bean seed has been treated with 0.01 lb ai metalaxyl, 0.008 lb ai thiophanate-methyl, 0.003 lb ai tebuconazole per 100 lb seed.
- This Wheat seed has been treated 0.01 lb ai metalaxyl, 0.008 lb ai thiophanate-methyl, 0.003 lb ai tebuconazole per 100 lb seed.

[Rotational Crop Plant-Back Intervals from the product label must be applied to seed bags]

User is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act and EPA regulations.

ROTATIONAL CROPS:

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application. The following plant-back intervals are required for listed crops:

Rotational Plant-Back Intervals*		
Immediate plant-back		30-DAY PLANT BACK
Artichoke	Oats	Cereals, including: Buckwheat Rice Millet
Barley	Okra	
Brassica (cole) leafy veg.	Pepinos	
Borage	Pepper	
Bulb Veg	Popcorn	
Canola	Potato	
Cilantro	Rapeseed	
Corn, Field	Rye	
Com, Sweet	Safflower	
Cotton	Sorghum	
Cranberry	Soybean	
Crambe	Strawberry	
Cucurbits	Sugarbeet	
Eggplant	Sunflower	
Flax	Tomatillo	
Ground cherry	Tomato	
Leaf petiole veg.	Triticale	
Leafy veg.	Veg. Root and Tuber	
Legume veg. (succulent Or dried, including soybean)	Watercress	
Mustard Seed	Wheat	
<p>* Cover crops for soil building or erosion control may be planted any time, but DO NOT graze or harvest for food or feed.</p> <p>* For all other crops not listed on an Imidacloprid label, or for crops for which no tolerance for the active ingredient has been established, a 12-month plant-back interval is required.</p>		

Notification of the crop rotational restriction must be conveyed to the grower by appropriate seed tag labeling or bag printing on all seed units.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry secure area at temperatures above freezing and in a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out the reach of children.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[For containers ≤ 5 gallons:] Nonrefillable container: DO NOT reuse or refill this container.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

[Nonrefillable Containers Larger than 5 Gallons:] Nonrefillable container: DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of

the product, which are beyond the control of LIBERTY CROP PROTECTION LLC or Seller. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW all such risks shall be assumed by Buyer and User and Buyer and User agree to hold LIBERTY CROP PROTECTION LLC and Seller harmless for any claims relating to such factors.

LIBERTY CROP PROTECTION LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or LIBERTY CROP PROTECTION LLC, and TO THE EXTENT CONSISTENT WITH APPLICABLE LAW Buyer and User assume the risk of any such use. To the extent consistent with applicable law LIBERTY CROP PROTECTION LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither LIBERTY CROP PROTECTION LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF LIBERTY CROP PROTECTION LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LIBERTY CROP PROTECTION, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

LIBERTY CROP PROTECTION LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of LIBERTY CROP PROTECTION LLC.