



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460-0001

EPA Reg. Number:

89168-163

Date of Issuance:

2/15/24

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

LIBERTY INDOXACARB EC

Name and Address of Registrant (include ZIP Code):

Daniel Sandoval
Regulatory Manager
Liberty Crop Protection, LLC
1880 Fall River Drive, Suite 100
Loveland, CO 80538

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 89168-163."

Continues page 2

Signature of Approving Official:

Tamica L. Cain, Product Manager 10
Invertebrate-Vertebrate Branch 2
Registration Division (7505T)

Date:

2/15/24

3. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSFs:

- Basic CSF, dated 6/12/23
- Alternate CSF #1, dated 6/12/23
- Alternate CSF #2, dated 6/12/23

If you have any questions, please contact Carmen Rodia via e-mail at Rodia.Carmen@epa.gov.

*Enclosures: Stamped "Accepted" Master Label, dated 2/15/24
 Acute Toxicity Review, dated 1/12/24
 Product Chemistry Review, dated 10/18/23*

{Note to reviewer: [Text] in brackets denotes optional or explanatory language}
 {Note to reviewer: {Text} in braces denotes where in the final label text will appear}
 {BOOKLET FRONT PANEL LANGUAGE}

INDOXACARB	GROUP	22A	INSECTICIDE
------------	-------	-----	-------------

LIBERTY INDOXACARB EC

ACTIVE INGREDIENT:

% By WT.

Indoxacarb:

(S)-methyl 7-chloro-2,5-dihydro-2-[[[(methoxycarbonyl)[4-(trifluoromethoxy)phenyl]amino]

carbonyl]indeno[1,2-e][1,3,4]oxadiazine-4a-(3H)-carboxylate.....15.84%

OTHER INGREDIENTS:84.16%

TOTAL100.00%

Contains 1.25 pounds of active ingredient per gallon of formulated product.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you DO NOT understand the label, find someone to explain it to you in detail.)

[See inside label booklet for First Aid Statements, [additional] Precautionary Statements, and Directions
 for Use.]

[See below additional First Aid Statements and Precautionary Statements]


EPA Reg. No. 89168-163

EPA Est. No. _____

Net Contents: _____ gal

Manufactured For:

LIBERTY CROP PROTECTION, LLC
 1880 Fall River Drive, Suite 100
 Loveland, CO 80538



021324

{LANGUAGE INSIDE BOOKLET}

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by the poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBERS Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergencies call the poison control center at 1-800-222-1222 . For non-emergency resource information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 Monday - Friday 8 am - Noon Pacific Time, (NPIC Web site: www.npic.orst.edu). For Chemical Spill, Leak, Fire or Exposure, call CHEMTREC 800-424-9300 .	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Avoid contact with skin or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks, and
- Chemical-resistant gloves such as: Barrier Laminate, Butyl Rubber ≥ 14 mils, Nitrile Rubber ≥ 14 mils, Neoprene Rubber ≥ 14 mils, Natural Rubber ≥ 14 mils, Polyethylene, Polyvinyl Chloride (PVC) ≥ 14 mils, and Viton ≥ 14 mils.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS. **IMPORTANT:** when reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "**Applicators and other handlers**" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

USER SAFETY RECOMMENDATIONS

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, birds, fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding application when rainfall is forecasted to occur within 24 hours. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems. **DO NOT** apply to any impervious surfaces which may contact or lead directly to surface water, storm drains, or urban runoff conveyance systems (gutters).

Pollinator Advisory: This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. **DO NOT** apply this product or allow it to drift to blooming crops or weeds while bees are foraging the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT mix or allow to come in contact with any oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of federal law to use this produce in a manner inconsistent with its labeling.

USE RESTRICTIONS

- **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- **DO NOT** formulate this product into any other end-use products without written permission of Liberty Crop Protection, LLC.
- **DO NOT** use **Liberty Indoxacarb EC** in greenhouses.
- **DO NOT** use **Liberty Indoxacarb EC** on ornamental plants or plants being grown for ornamental purposes.
- **DO NOT** use **Liberty Indoxacarb EC** in residential areas.
- For fields to which applications of **Liberty Indoxacarb EC** will be made, construct a vegetative filter strip if one does not already exist. Existing and new filter strips must be, at a minimum, 10-foot wide and composed of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds). Vegetative filter strips must be maintained to optimize their utility. Only apply products containing indoxacarb onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.
- For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Liberty Indoxacarb EC must be used only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Coveralls over long-sleeved shirt and long pants,
- Socks plus chemical resistant footwear, and
- Chemical-resistant gloves such as: Barrier Laminate, Butyl Rubber \geq 14 mils, Nitrile Rubber \geq 14 mils, Neoprene Rubber \geq 14 mils, Natural Rubber \geq 14 mils, Polyethylene, Polyvinyl Chloride (PVC) \geq 14 mils, and Viton \geq 14 mils.

PRODUCT INFORMATION

Liberty Indoxacarb EC is an emulsifiable concentrate that can be applied as a foliar spray to control many important insects. This product is diluted with water for application. Always shake well before use.

CHEMIGATION: DO NOT apply this product through any type of irrigation system except for application to alfalfa, corn (field), corn, (pop), corn (grown for seed), cotton, peanut, and soybean, and as allowed by Federal Supplemental and Special Local Need (SLN) labeling. (See “**APPLICATION BY CHEMIGATION**” section of the label.)

INTEGRATED PEST MANAGEMENT

Liberty Crop Protection, LLC supports the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an IPM program which can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other pest detection methods, correct target pest identification, population monitoring, rotation of insecticides with different modes-of-action and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants, product manufacturer or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop or site systems in your area.

SCOUTING

Monitor insect populations to determine whether or not there is a need for application of **Liberty Indoxacarb EC** based on label recommendations and locally determined economic thresholds. More than one treatment of **Liberty Indoxacarb EC** may be required to control a population of pests.

PESTICIDE RESISTANCE MANAGEMENT

For resistance management, **Liberty Indoxacarb EC** contains the active ingredient indoxacarb which is a Mode of Action Group 22A insecticide. Insecticides with the same group number affect the same biological site of action on the target pest and when used repeatedly in the same treatment area, naturally-occurring resistant individuals may survive correctly applied insecticide treatments, reproduce, and become dominant.

To delay the development of insecticide resistance, a resistance management strategy should include incorporation of cultural and biological control practices, alternation to different mode of action insecticides

on succeeding generations, targeting the most susceptible life stage, and where possible controlling multiple life stages of the same pest.

Consult with your local or state agricultural authorities or product manufacturer, or visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irac-online.org> for more information about developing a resistance management strategy.

Unless directed otherwise in the specific crop/pest sections of this label, follow these guidelines to delay the development of insecticide resistance:

- Apply **Liberty Indoxacarb EC** and other Group 22A insecticides within a single “treatment window” to minimize exposing successive generations of a pest species to the same mode of action insecticides.
- A “treatment window” is defined as the period of residual insecticidal activity provided by one or more applications of products with the same mode of action not to exceed approximately 30 days.
- Within the Group 22A “treatment window”, make no more than 2 applications of **Liberty Indoxacarb EC** or other Group 22A insecticides.
- Following a Group 22A “treatment window”, rotate to a “treatment window” of effective insecticides with a different mode of Action Group Number. The period between Group 22A “treatment windows” should be at least 30 days.
- For short cycle crops (< 50 days), the duration of the crop cycle may be considered as the Group 22A “treatment window” if no Group 22A insecticides are used during the next crop cycle at the same farm location.
- If **Liberty Indoxacarb EC** is tank mixed with other insecticides, then apply rates that are individually registered for use against the target species. **DO NOT** rely on the same mixture repeatedly to control the same pest species and follow the same “treatment window” rotation recommendation described above for the tank-mixed products.
- Use labeled rates of **Liberty Indoxacarb EC** when applied alone or in tank mixtures.
- Monitor after application for unexpected target pest survival. If insect resistance is suspected consult with your manufacturer’s representative, local university specialist, or certified pest control advisor.

If resistance to **Liberty Indoxacarb EC** develops in your area other products with a similar mode of action (Mode of Action Group 22A) may not provide adequate control.

BENEFICIAL ARTHROPODS

Other than reducing the target pest species as a food source, **Liberty Indoxacarb EC** helps conserve certain beneficial arthropods, including parasitic wasps, predatory mites, big-eyed bugs, damsel bugs, minute pirate bugs, and spiders. While these beneficials arthropods cannot be relied upon to control pests, they are of potential value and can be monitored along with pests in pest management programs on these crops.

APPLICATION

Apply at the listed rates when insect populations reach locally determined economic thresholds. Consult the cooperative extension service, professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area.

Liberty Indoxacarb EC applications should target eggs and small instar larvae.

Follow-up treatments of **Liberty Indoxacarb EC** should be applied, as needed, to keep pest populations within threshold limits. Apply **Liberty Indoxacarb EC** at 5 to 7 day intervals or as needed to manage specific target insect pests, as specified in the specific crop sections, to maintain control.

Use sufficient water to obtain thorough, uniform coverage.

Because **Liberty Indoxacarb EC** is most effective through ingestion of treated plant material, thorough spray coverage is essential for optimum control of targeted pest insects. Using Increased water volumes

will typically result in better spray coverage, especially under adverse conditions such as dry, hot weather or dense plant foliage. **Liberty Indoxacarb EC** may be applied by ground, aerial or overhead sprinkler chemigation application equipment. For aerial application use the following directions unless otherwise specified in specific crop/pest sections of this label or EPA-approved supplemental labeling: use a minimum of 3 gal water per acre (gpa) [minimum of 5 gal water per acre in Arizona and California]. For ground application use the following directions unless otherwise specified in specific crop/pest sections of this label or EPA-approved supplemental labeling: use a minimum of 5 gal. water per acre. Higher gallonage will provide better coverage and performance. For overhead chemigation applications, see "Application by Chemigation" section of the label for guidance on water volumes to be used.

Use of Adjuvants: In some situations where coverage is difficult to achieve such as closed canopy, dense foliage, plants with waxy leaf surfaces, or less than optimum application equipment, an adjuvant may improve performance. Use only adjuvant products that are labeled for agricultural use and follow the directions on the manufacturer's label.

SPRAY PREPARATION

Spray equipment must be clean and free of previous pesticide deposits before applying **Liberty Indoxacarb EC**. Fill spray tank 1/4 to 1/2 full of water. Add **Liberty Indoxacarb EC** directly to spray tank. Mix thoroughly to fully disperse the insecticide; once dispersed continued agitation is required. Use mechanical or hydraulic means; **DO NOT** use air agitation.

Spray mix must not be stored overnight in spray tank. Observe the most restrictive of the labeling limitations and precautions of all products used in mixtures.

Tank Mixing and Compatibility - Since formulations may be changed and new ones introduced, it is a best practice that users premix a small quantity of a desired tank mix and observe for possible physical incompatibility (settling out, flocculation, crystallization, etc.). It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. **DO NOT** exceed label application rates. This product cannot be mixed with any product containing a label prohibition against such mixing.

Spray volumes of less than 3 gal of water and tank mixtures of more than two products can increase the chances of incompatible spray mixtures. A jar test (as described below) should be conducted when label guidance is not given or prior experience with a specific tank mixture is unknown. The jar test should follow the mixing sequence below in water volume proportional to the planned spray tank water volume to assure that the tank mixture is compatible. Constant agitation may be needed during mixing and spraying of mixtures. **Liberty Indoxacarb EC** is compatible with most commonly used plant protectants.

Steps to conduct a jar test to determine physical tank mix compatibility of **Liberty Indoxacarb EC** with other products:

- Add clean water to the jar in proportion to the planned water volume that will be used in the spray tank (a jar size of 16 oz is acceptable).
- While wearing the most restrictive PPE, mix proper proportional amounts of **Liberty Indoxacarb EC** and desired tank mix partner(s) as will be present in the spray tank. Add one product at a time following the sequence of addition according to formulation type provided in this label.
- Seal and shake mixture after each product is added.
- Allow to stand for 1 hour.
- View jar to determine if settling, flocculation, crystallization or any other undesirable changes have happened.
- If none of the above is observed or the solution can be easily remixed after shaking, the mixture is compatible with **Liberty Indoxacarb EC**.
- If the tank mixture is not compatible, a higher water volume, reduced rate of the tank mix partner(s), reduced number of tank mix partners or a compatibility agent may be needed.

Tank Mixtures and Crop Safety

Crop varieties can differ in their responsiveness to tank mixtures, and environmental conditions can influence product performance and crop response. It is not possible to test **Liberty Indoxacarb EC** alone or with all possible tank mix combinations on all varieties under all environmental conditions. When considering the use of a tank mixture on a labeled crop without prior experience, or which is not specifically described on **Liberty Indoxacarb EC** product labeling or in other Liberty Crop Protection, LLC product use instruction, it is important to check crop safety first. To test for crop safety, prepare a small volume of the intended tank mixture, apply it to an area of the target crop as directed by both this and the tank mix partner product labels, and observe the treated crop to ensure that a phytotoxic response does not occur.

Use of **Liberty Indoxacarb EC** in any tank mixture applications that is not specifically described on **Liberty Indoxacarb EC** product labeling or in other Liberty Crop Protection, LLC product use instructions, could potentially result in crop injury. Follow the precautions on this label and on the label for any other product to be used in tank mixtures before making such applications to your crops. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Tank Mixing Sequence - Add different formulation types in the sequence indicated below. * Allow time for complete mixing and dispersion after addition of each product.

1. Products in water soluble bags (WSB)
2. Water soluble granules (SG)
3. Water dispersible granules (WG, XP, DF)
4. Wettable powders (WP)
5. Water based suspension concentrates (SC)
6. Water soluble concentrates (SL)
7. Suspoemulsions (SE)
8. Oil Based suspension concentrates (OD)
9. **Liberty Indoxacarb EC** or other emulsifiable concentrates (EC)n
12. Drift retardants

*- Unless otherwise specified by manufacturer directions for use or by local expertise.

SPRAY TANK CLEANOUT

Prior to application, start with clean, well maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove. Drain spray equipment. Thoroughly rinse sprayer and flush hoses, boom and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. **DO NOT** clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

APPLICATION BY CHEMIGATION – ALFALFA, CORN (FIELD), CORN (GROWN FOR SEED), CORN (POP), COTTON, PEANUT, AND SOYBEAN

Instructions for the Use of Liberty Indoxacarb EC in Overhead Sprinkler Chemigation Systems.

Overhead chemigation applications offer the advantage of greater penetration and coverage of the target plant. However, typical chemigation applications are more dilute than ground or aerial applications. For best results, it is recommended to keep the concentration of **Liberty Indoxacarb EC** as high as possible in the application. Apply **Liberty Indoxacarb EC** in 0.1 to 0.2 inches of water per acre. **Liberty Indoxacarb EC** is most active as an ingestion insecticide, although it does have activity as a direct contact insecticide. For best results, applications of **Liberty Indoxacarb EC** should ensure thorough coverage of the target plant to maximize the opportunity for target insects to ingest **Liberty Indoxacarb EC**.

Types of Chemigation Systems:

Liberty Indoxacarb EC may be applied only through overhead sprinkler irrigation systems. Overhead irrigation systems include the following; center pivot, end tow, hand move, lateral move, side roll, solid set and wheel line. Center pivot and lateral move irrigation systems are preferred. Other overhead sprinkler systems may be used if they provide uniform water distribution. **DO NOT** apply **Liberty Indoxacarb EC** through any other type of irrigation system. **DO NOT** use filter screens smaller than 50 mesh throughout the system, due to possible buildup of material on 100 mesh or smaller screens.

Directions for Chemigation:

Preparation

A pesticide tank is recommended for the application of **Liberty Indoxacarb EC** in chemigation systems. Thoroughly clean the injection system and tank of any fertilizer or chemical residues using a standard clean-out procedure. Dispose of any residues in accordance with State and Federal laws. With the mix tank 1/4 to 1/2 full with water and the agitator running, measure the required amount of **Liberty Indoxacarb EC** and add it to the tank. Then add additional water to bring your total pesticide mixture up to the desired volume for your application. Note: Always add the **Liberty Indoxacarb EC** to water, never put **Liberty Indoxacarb EC** into a dry tank or other mixing equipment without first adding water. See "Tank Mixing Sequence" section of the container label for tank mixing sequence. Continue to agitate the mixture throughout the application process. Use mechanical or hydraulic agitation, **DO NOT** use air agitation. Highly alkaline water should be buffered so that the pH of the spray solution is in the range of neutral to slightly acidic.

Injection Into Chemigation Systems

Inject the proper amount of **Liberty Indoxacarb EC** into the irrigation water flow using a positive displacement injection pump. Injection should occur at a point in the main irrigation water flow to ensure thorough mixing with the irrigation water. For continuously moving systems, inject the solution containing **Liberty Indoxacarb EC** into the irrigation waterline continually and uniformly throughout the irrigation cycle. Apply in no more than 0.2 inches of water per acre. For overhead sprinkler systems that are stationary, add the solution containing **Liberty Indoxacarb EC** to the irrigation water line and apply no more than 0.2 inches of water per acre just before the end of the irrigation cycle.

Uniform Water Distribution

The irrigation system used for application of **Liberty Indoxacarb EC** must provide for uniform distribution of **Liberty Indoxacarb EC** treated water. Non-uniform distribution might result in crop injury, lack of effectiveness or illegal pesticide residues in or on the crop being treated. Ensure the irrigation system is calibrated to uniformly distribute the chemigation application to the crop. Contact the equipment manufacturer, the local University Extension agent or other experts if you have questions about achieving uniform distribution of the application.

Equipment Calibration

Calibrate the irrigation system and injector before applying **Liberty Indoxacarb EC**. Calibrate the injection pump while the system is running using the expected irrigation rate. If you have questions about calibration, you should contact your state extension service specialists, equipment manufacturer or other experts.

Monitoring of Chemigation Applications

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of a responsible person, shall shut the system down and make necessary adjustments should the need arise. Wear the personal protective equipment as defined in the PPE section of the label for applicators and other handlers when making adjustments or repairs on the chemigation system when **Liberty Indoxacarb EC** is in the irrigation water.

Required System Safety Devices

DO NOT connect any irrigation system used for pesticide applications to a public water system unless the pesticide label prescribed safety devices are in place. Public water system means a system for the provision

to the public of piped water for human consumption, if such a system has at least 15 service connections or regularly serves an average of at least 25 individuals at least 60 days out of the year.

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

Operation

Start the water pump and sprinkler, and let the system achieve the desired pressure and speed before starting the injector. Start the injector and calibrate the injection system according to the directions above. This procedure is necessary to deliver the desired rate per acre in a uniform manner. When the application is finished, allow the entire irrigation and injector system to be thoroughly flushed clean before stopping the system.

- End guns must be turned off during the application, if they irrigate nontarget areas or if they **DO NOT** provide uniform application and coverage.
- Plug nozzles in the immediate area of control panels, chemical supply tanks and system safety devices to prevent contamination of these areas.
- **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.
- **DO NOT** apply when system connections or fittings leak or when nozzles **DO NOT** provide uniform distribution.
- **DO NOT** allow irrigation water to collect or run-off during chemigation.

Cleaning the System Thoroughly clean the injection system and tank of any fertilizer or chemical residues using a standard clean-out procedure. Dispose of any residues in accordance with State and Federal laws. Consult your owner's manual or your local equipment dealer for cleanout procedures for your injection system.

MANDATORY SPRAY DRIFT MANAGEMENT

AERIAL APPLICATIONS

- **DO NOT** release spray at a height greater than 10 feet above the ground or vegetative canopy unless a greater application height is necessary for pilot safety.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the wind speed is greater than 10 mph, the boom length must be 65% or less for fixed-wing aircraft and 75% or less for helicopters. Otherwise, the boom length must be 75% or less for fixed-wing aircraft and 90% or less for helicopters.
- **DO NOT** apply during temperature inversions.

GROUND BOOM APPLICATIONS

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- **DO NOT** apply during temperature inversions.

AIRBLAST APPLICATIONS

- Sprays must be directed into the canopy.
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer row.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

The APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles – Follow nozzle manufacturer's recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT – Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, that it is configured properly, and that drift is not occurring.

Note: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the application equipment section of this label to determine if use of an air assisted sprayer is recommended.

CROP ROTATION

Crops on this label and the following crops or crop groups may be planted immediately following harvest: Bean, dried (crop subgroup 6C); Bean, succulent (crop subgroups 6A, 6B); Bushberries (crop subgroup 13-07B); Cucurbit vegetables (crop subgroups 9A, 9B); Fruiting vegetables (crop subgroups 8-10A, 8-10B, 8-10C); Garden beets; Grapes; Leafy green vegetables (crop subgroup 4A); Leafy petiole vegetables (crop subgroup 4B); Low growing berry (except strawberry) (crop subgroup 13-07H); Mint (peppermint & spearmint); Okra; Pome fruit (crop group 11); Small fruit vine climbing subgroup (except fuzzy kiwifruit) (crop subgroup 13-07F); Stone fruit (crop group 12); Sweet corn; Tuberous and corm vegetables (crop subgroup 1C: arracacha, arrowroot, Chinese artichoke, Jerusalem artichoke, edible canna (Queensland arrowroot), bitter and sweet cassava, chayote (root), chufa, dasheen (taro), ginger, leren, potato, sweet potato, taniel (cocoyam), tumeric, yam bean (jicama, manioc pea) and true yam).

DO NOT plant for food or feed any other crops not registered for use with indoxacarb for 30 days after last use.

USE SITES

Crops	Insects	Lb. AI	Fl Oz	Acres Treated per Gal of Liberty Indoxacarb EC	Last Application (Days to Harvest)	REI
Alfalfa: and alfalfa grown for seed	Cabbage looper [**]	0.045 - 0.11	4.6 - 11.3	11.5 - 27.8	7	12 hrs.
	Grasshoppers					
	Alfalfa caterpillar Alfalfa weevil larvae Beet armyworm	0.065 - 0.11	6.7 - 11.3	11.5 - 19	For alfalfa, harvest is defined as when the crop is cut.	
	Cabbage looper (CA Only)					
	Egyptian alfalfa weevil larvae					
	Granulated cutworm					
	Potato leafhopper* [**]	0.09 - 0.11	9.2 - 11.3	11.5 - 14		
Lygus bugs (Western U.S.)*						
Western yellowstriped armyworm						

ALFALFA USE RESTRICTIONS:

- **DO NOT** apply more than 11.3 fl. oz/A (0.11 lb. ai./A) of **Liberty Indoxacarb EC** or other indoxacarb-containing products per cutting.
- **DO NOT** apply more than 45 fl. oz/A (0.44 lb. ai./A) of **Liberty Indoxacarb EC** or other indoxacarb-containing products per calendar year.
- When **Liberty Indoxacarb EC** is used on alfalfa grown for seed, the seed may not be used for sprouts or livestock feed. All seed from treated crop must be tagged, "Not for Human or Animal Use" at the processing plant.
- Allow a minimum of 7 days between applications.

Note: Apply lower listed rates for light to moderate infestations. Use intermediate to high rates within the listed rate range on heavier infestations or when later instar larvae exist. Use the highest listed rate for controlling severe infestations or when longer residual control is desired.

*Suppression only.

[** Not Registered for Use in California]

Crops	Insects	Lb. AI	Fl Oz	Acres Treated per Gal of Liberty Indoxacarb EC	Last Application (Days to Harvest)	REI
Bean, (Dried and Succulent) (except Soybeans) Including: Dried cultivars of bean (<i>Lupinus</i>) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin); bean (<i>Phaseolus</i>) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); pea (<i>Pisum</i> spp.) (includes dwarf pea, edible pod pea, english pea, field pea, garden pea, green pea, snow pea, sugar snap pea); pigeon pea (<i>Cajanus cajan</i>); bean (<i>Vigna</i>) (includes adzuki bean, blackeyed pea, catjang, chinese longbean, cowpea, crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean); broad bean (dry); chickpea; guar; jackbean; lablab bean, lentil; sword bean; wax bean	Corn earworm European corn borer	0.065-0.11	6.7-11.3	11.5-19	7	12 hrs.
BEAN, (DRIED AND SUCCULENT) (EXCEPT SOYBEANS) USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 45 fl. oz/A (0.44 lb. ai/A) of Liberty Indoxacarb EC or other indoxacarb-containing products per crop. • DO NOT make more than 4 applications of Liberty Indoxacarb EC per acre per crop. • Allow a minimum of 7 days between applications. <p>Note: For ground applications, make a uniform application in approximately 20-100 gal/A of water.</p>						

Crops	Insects	Lb. AI	Fl Oz	Acres Treated per Gal of Liberty Indoxacarb EC	Last Application (Days to Harvest)	REI
Corn (field) Corn (grown for seed) Corn (sweet) Corn (pop)[***]	Beet armyworm Corn earworm* Corn rootworm(adult) European corn borer Fall armyworm Grasshoppers Japanese beetles (adult)** Western bean cutworm Yellowstriped armyworm	0.059 - 0.11	6.0 - 11.3	11.5 - 21.3	14 Days for Grain and Stover (field, pop, and corn grown for seed) 1 Day for Foliage, Fodder, Silage, (field and corn grown for seed only)	12 hrs.
	Brown stink bug** Green stink bug** Southern green stink bug**	0.09 - 0.11	9.2 - 11.3	11.5 - 14		
CORN USE RESTRICTIONS: <ul style="list-style-type: none">• DO NOT apply more than 22.6 fl. oz/A (0.22 lb. ai/A) of Liberty Indoxacarb EC or other indoxacarb-containing products per calendar year.• DO NOT make more than 2 applications of Liberty Indoxacarb EC per acre per crop.• Allow a minimum of 5 days between applications. <p>* Corn earworm control is only for treated foliage and silks. New foliage and new silks will not be protected with a single application. **Suppression only. [***Not Registered for Use in California]</p>						

Crops	Insects	Lb. AI	Fl Oz	Acres Treated per Gal of Liberty Indoxacarb EC	Last Application (Days to Harvest)	REI
Cotton	Cotton bollworm* Tobacco budworm*	0.11	11.3	11.5	14	12 hrs.
	Cotton bollworm in Transgenic Bt Cotton	0.09 - 0.11	9.2 - 11.3	11.5 - 14		
	Beet armyworm Fall armyworm Western yellowstriped	0.09 - 0.11	9.2 - 11.3	11.5 - 14		
	Cabbage looper Soybean looper	0.065 - 0.09	6.7 - 9.2	14 - 19		
	Cotton fleahopper** Lygus bugs (Western U.S.)*** Tarnished plant bug**	0.09 - 0.11	9.2 - 11.3	11.5 - 14		

COTTON USE RESTRICTIONS:

- **DO NOT** apply more than 45 fl oz/A (0.44 lb. ai/A) of **Liberty Indoxacarb EC** or other indoxacarb-containing products per calendar year.
- **DO NOT** make more than 4 applications of **Liberty Indoxacarb EC** per acre pre crop.
- Allow a minimum of 5 days between applications.

Note: Beet armyworm and Western yellowstriped armyworm (AZ & CA only) - **Liberty Indoxacarb EC** may be applied to seedling cotton (less than 18 inches high), at rates of 6.7 - 11.3 fl. oz/A in sufficient water to obtain thorough coverage (minimum of 5 gal/A).

*Cotton Bollworm and Tobacco Budworm - For the most effective control, applications of **Liberty Indoxacarb EC** should be made when the majority of the population is within the time of blackhead egg stage to egg hatch.

AL & GA only - **Liberty Indoxacarb EC** may be applied at 0.09 lb. ai/A (9.2 fl. oz/A of product) for control of low populations of tobacco budworm and cotton bollworm on conventional cotton varieties that **DO NOT** contain the transgenic Bt trait. Low populations are defined as less than 30 eggs per 100 terminals and/or less than 10 tobacco budworm/cotton bollworm larvae detected per 100 terminals, blooms, or squares. If tobacco budworm or cotton bollworm populations exceed the egg or larval threshold as described above, then increase the **Liberty Indoxacarb EC** use rate to 0.11 lb. ai/A (11.3 fl. oz/ A of product).

Tarnished Plant Bug and Cotton Fleahopper - A single application of **Liberty Indoxacarb EC will provide control of light to moderate populations of tarnished plant bug or cotton fleahopper. Heavy populations of tarnished plant bug or cotton fleahopper may require multiple applications. For the most effective control, fields should be scouted twice per week with application timing based on locally determined economic thresholds. Consult the cooperative extension service, professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area.

***Suppression only.

Crops	Insects	Lb. AI	Fl Oz	Acres Treated per Gal of Liberty Indoxacarb EC	Last Application (Days to Harvest)	REI
Peanut	Corn Earworm	0.065 – 0.11	6.7 – 11.3	11.5 - 19	14	12 hrs.
	Beet armyworm Fall armyworm Granulate cutworm Rednecked peanutworm (except California) Tobacco budworm	0.09 – 0.11	9.2 – 11.3	11.5 - 14		

PEANUT USE RESTRICTIONS:

- **DO NOT** apply more than 45 fl oz/A (0.44 lb. ai/A) of **Liberty Indoxacarb EC** or other indoxacarb-containing products per calendar year.
- **DO NOT** make more than 4 applications of **Liberty Indoxacarb EC** per acre per crop.
- Allow a minimum of 5 days between applications.

Crops	Insects	Lb. AI	Fl Oz	Acres Treated per Gal of Liberty Indoxacarb EC	Last Application (Days to Harvest)	REI
Soybean [***]	Beet armyworm Cabbage looper* Corn earworm Fall armyworm Grasshoppers Green cloverworm Soybean looper* Yellowstriped armyworm	0.045 - 0.11	4.6 - 11.3	11.5 - 27.8	21	12 hrs.
	Velvetbean Caterpillar**	0.055 - 0.11	5.6 - 11.3	11.5 - 22.8		
	Tobacco budworm Bean leaf beetle** Brown stink bug** Green stink bug** Southern green stink bug**	0.09 - 0.11	9.2 - 11.3	11.5 - 14		
SOYBEAN USE RESTRICTIONS: <ul style="list-style-type: none">• DO NOT apply more than 45 fl. oz/A (0.44 lb. ai/A) of Liberty Indoxacarb EC or other indoxacarb-containing products per calendar year.• DO NOT make more than 4 applications of Liberty Indoxacarb EC per acre per crop.• Allow a minimum of 5 days between applications.• DO NOT feed or graze livestock on treated fields. <p>* Use lower rate (4.6 fl. oz/A) for low to moderate populations of cabbage and soybean loopers. Use higher rates (5.6 fl. oz/A to 11.3 fl. oz/A) for higher populations or when crop canopy is dense.</p> <p>**Suppression only.</p> <p>[***Not Registered for Use in California]</p>						

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep this product in its tightly closed original container, when not in use. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals, and avoid excessive heat while in storage.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[NONREFILLABLE CONTAINER (EQUAL TO OR LESS THAN 5 GALLONS):DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

[NONREFILLABLE CONTAINER (GREATER THAN 5 GALLONS): DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by other procedures allowed by state and local authorities.

REFILLABLE CONTAINER: Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. After triple rinsing is complete, and the container is not suitable for refilling or reconditioning, offer the container for recycling if available, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of LIBERTY CROP PROTECTION LLC or Seller. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW all such risks shall be assumed by Buyer and User and Buyer and User agree to hold LIBERTY CROP PROTECTION LLC and Seller harmless for any claims relating to such factors.

LIBERTY CROP PROTECTION LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or LIBERTY CROP PROTECTION LLC, and TO THE EXTENT CONSISTENT WITH APPLICABLE LAW Buyer and User assume the risk of any such use. To the extent consistent with applicable law LIBERTY CROP PROTECTION LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF

FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE. To the extent consistent with applicable law, neither LIBERTY CROP PROTECTION LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF LIBERTY CROP PROTECTION LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LIBERTY CROP PROTECTION, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT. LIBERTY CROP PROTECTION LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of LIBERTY CROP PROTECTION LLC.

All trademarks are the property of the respective owners.

{LANGUAGE ON LABEL AFFIXED TO CONTAINER}

INDOXACARB

GROUP 22A

INSECTICIDE

Liberty Indoxacarb EC

[Emulsifiable Concentrate]

ACTIVE INGREDIENT:

Indoxacarb

(S)-methyl 7-chloro-2,5-dihydro-2-[[[(methoxycarbonyl)[4-

(trifluoromethoxy)phenyl]amino]carbonyl]indeno

[1,2-e][1,3,4]oxadiazine-4a-(3H)-

carboxylate.....15.84%

OTHER INGREDIENTS:.....84.16%

TOTAL.....100.00%

Contains 1.25 pounds of active ingredient per gallon.

% By WT.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand the label, find someone to explain it to you in detail.)

FIRST AID

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- **DO NOT** induce vomiting unless told to do so by the poison control center or doctor.
- **DO NOT** give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

HOTLINE NUMBERS

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergencies call the poison control center at **1-800-222-1222**. For non-emergency resource information concerning this product, call the National Pesticides Information Center (NPIC) at **1-800-858-7378** Monday – Friday 8 am – Noon Pacific Time, (NPIC Web site: www.npic.orst.edu). For Chemical Spill, Leak, Fire or Exposure, call CHEMTREC **800-424-9300**.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contact with skin or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to mammals, birds, fish, and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding application when rainfall is forecasted to occur within 24 hours.

Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems. **DO NOT** apply to any impervious surfaces which may contact or lead directly to surface water, storm drains, or urban runoff conveyance systems (gutters).

Pollinator Advisory: This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. **DO NOT** apply this product or allow it to drift to blooming crops or weeds while bees are foraging the treatment area.

PHYSICAL OR CHEMICAL HAZARDS:

DO NOT mix or allow to come in contact with any oxidizing agent. Hazardous chemical reaction may occur.

See inside label booklet for additional First Aid Statements, Precautionary Statements and Directions for Use.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep this product in its tightly closed original container, when not in use. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals, and avoid excessive heat while in storage.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[NONREFILLABLE CONTAINER (EQUAL TO OR LESS THAN 5 GALLONS):DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

[NONREFILLABLE CONTAINER (GREATER THAN 5 GALLONS): DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by other procedures allowed by state and local authorities.

REFILLABLE CONTAINER: Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. After triple rinsing is complete, and the container is not suitable for refilling or reconditioning, offer the container for recycling if available, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

Manufactured for:
Liberty Crop Protection, LLC
1880 Fall River Drive, Suite 100
Loveland, CO 80538

EPA Reg. No.: 89168-163

EPA Est. No.: _____

NET CONTENTS: _____

021324