



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460-0001

EPA Reg. Number:

89168-164

Date of Issuance:

5/17/24

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

LIBERTY INDOXACARB 30 DG

Name and Address of Registrant (include ZIP Code):

Daniel Sandoval
Regulatory Manager
Liberty Crop Protection, LLC
1880 Fall River Drive, Suite 100
Loveland, CO 80538

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 89168-164."

Continues page 2

Signature of Approving Official:

Tamica L. Cain, Product Manager 10
Invertebrate-Vertebrate Branch 2
Registration Division (7505T)

Date:

5/17/24

3. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF:

- Basic CSF, dated 10/18/23

If you have any questions, please contact Mr. Carmen J. Rodia, Jr. via e-mail at Rodia.Carmen@epa.gov.

*Enclosures: Stamped "Accepted" Master Label, dated 5/17/24
 Acute Toxicity Review, dated 1/9/24
 Product Chemistry Review, dated 10/27/23*

{Note to reviewer: [Text] in brackets denotes optional or explanatory language}
{Note to reviewer: {Text} in braces denotes where in the final label text will appear}
{BOOKLET FRONT PANEL LANGUAGE}

INDOXACARB	GROUP 22A	INSECTICIDE
------------	-----------	-------------

LIBERTY INDOXACARB 30 DG

[Contains indoxacarb, the active ingredient used in Avaunt® [eVo Insect Control].]

[Dispersible Granules]

ACTIVE INGREDIENT:

% BT. WT.

Indoxacarb*

(S)-methyl 7-chloro-2,5-dihydro-2-[[[(methoxycarbonyl)[4-(trifluoromethoxy)phenyl]amino]carbonyl]indeno

[1,2-e][1,3,4]oxadiazine-4a(3H)-carboxylate..... 30.00%

OTHER INGREDIENTS: 70.0%

TOTAL 100.0%

*Contains 0.3 lb. of Indoxacarb per pound of formulated product.

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you DO NOT understand the label, find someone to explain it to you in detail.)

[See inside booklet for First Aid, [additional] Precautionary Statements, and Directions for Use.]
[See other panels for additional precautionary statements.] [See below for additional Precautionary
Statements.]

[Liberty Indoxacarb 30 DG is not manufactured, or distributed by FMC Corporation, seller of Avaunt®
[eVo Insect Control].]

{Note to reviewer: if used by the registrant, the contains statement and corresponding disclaimer will both
appear on the front of the final product packaging}

ACCEPTED

05/17/2024

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under

EPA Reg. No. 89168-164

EPA Reg. No.: 89168-164

EPA Est. No.: _____

Net Contents: _____

Manufactured for:
LIBERTY CROP PROTECTION, LLC
1880 Fall River Drive, Suite 100
Loveland, CO 80538

051324

{LANGUAGE INSIDE BOOKLET}

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• DO NOT induce vomiting unless told to do so by the poison control center or doctor.• DO NOT give anything by mouth to an unconscious person.
IF INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For chemical emergencies (including spill, leak, fire, exposure, or accident), you may contact CHEMTREC at 1-800-424-9300 (within the USA and Canada) or at +1 703-527-3887 (collect calls accepted), twenty-four (24) hours per day seven (7) days per week within the USA and Canada. In addition, for additional information on this pesticide product (including health concerns or medical emergencies), you may contact the National Pesticides Information Center (NPIC) at 1-800-858-7378 , from 8:00am to 12:00 pm Pacific Standard Time, Monday through Friday, excluding holidays.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Harmful if swallowed or if inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants;
- Shoes plus socks;
- Protective eyewear; and
- Chemical-resistant gloves made of: Barrier Laminate ≥ 14 mils, Butyl Rubber ≥14 mils, Nitrile Rubber ≥14 mils, Neoprene Rubber ≥, Natural Rubber ≥14 mils, Polyethylene, Polyvinyl Chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils.

In addition, mixers and loaders supporting aerial applications to dried and/or succulent beans must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified

as specified in the WPS. **IMPORTANT:** when reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for “applicator and other handlers” and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside gloves before removing. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, birds, fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wash-water or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding application when rainfall is forecasted to occur within 24 hours. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems. **DO NOT** apply to any impervious surfaces which may contact or lead directly to surface water, storm drains, or urban runoff conveyance systems (gutters). Cover, incorporate, or clean up granules that are spilled.

Pollinator Advisory: This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. **DO NOT** apply this product or allow it to drift to blooming crops or weeds while bees are foraging in the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT mix or allow to come in contact with any oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

- **Liberty Indoxacarb 30 DG** must be used only in accordance with the directions on this label, or as otherwise permitted by FIFRA. Always read the entire label, including the Limitation of Warranty and Liability.
- **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- **DO NOT** use in home plantings.
- **DO NOT** formulate this product into any other end-use products without written permission of Liberty Crop Protection, LLC.
- **DO NOT** use an adjuvant on bushberries or garden beets.
- Use only in commercial and farm plantings.
- For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Liberty Indoxacarb 30 DG must be used only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours for all crops.

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Coveralls over long-sleeved shirt and long pants;
- Socks plus chemical resistant footwear; and
- Chemical-resistant gloves made of: Barrier Laminate \geq 14 mils, Butyl Rubber \geq 14 mils, Nitrile Rubber \geq 14 mils, Neoprene Rubber \geq , Natural Rubber \geq 14 mils, Polyethylene, Polyvinyl Chloride (PVC) \geq 14 mils, or Viton \geq 14 mils.

PRODUCT INFORMATION

Liberty Indoxacarb 30 DG is a water dispersible granule that can be applied as a foliar spray to control many important insects. **Liberty Indoxacarb 30 DG** is mixed with water for application.

For fields to which applications of **Liberty Indoxacarb 30 DG** will be made, construct a vegetative filter strip if one does not already exist. Existing and new filter strips must be, at a minimum, 10-foot-wide and composed of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds). Vegetative filter strips must be maintained to optimize their utility. Only apply products containing indoxacarb onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

CHEMIGATION: DO NOT apply this product through any type of irrigation system except for application to cranberries, mint, potatoes, spinach¹ and sweet corn (See **"APPLICATION BY CHEMIGATION"** section of the label.)

¹Use of spinach via overhead irrigation is allowed only by states of Arkansas, Georgia, Missouri, North Carolina, New Mexico, Oklahoma, and Texas unless otherwise permitted in supplemental labeling.

INTEGRATED PEST MANAGEMENT

Liberty Crop Protection, LLC supports the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an IPM program which can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other pest detection methods, correct target pest identification, population monitoring, rotation of insecticides with different modes-of-action, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants, product manufacturer or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop or site systems in your area.

SCOUTING

Monitor insect populations to determine whether or not there is a need for application of **Liberty Indoxacarb 30 DG** based on label recommendations and locally determined economic thresholds. More than one treatment of **Liberty Indoxacarb 30 DG** may be required to control a population of pests.

PESTICIDE RESISTANCE MANAGEMENT

For resistance management, **Liberty Indoxacarb 30 DG** contains the active ingredient indoxacarb which is a Mode of Action Group 22A insecticide. Insecticides with the same group number affect the same biological site of action on the target pest and when used repeatedly in the same treatment area, naturally-occurring resistant individuals may survive correctly applied insecticide treatments, reproduce, and become dominant.

To delay the development of insecticide resistance, a resistance management strategy should include incorporation of cultural and biological control practices, alternation to different mode of action insecticides on succeeding generations, targeting the most susceptible life stage, and where possible controlling multiple life stages of the same pest.

Consult with your local or state agricultural authorities or product manufacturer, or visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irac-online.org> for more information about developing a resistance management strategy.

Unless directed otherwise in the specific crop/pest sections of this label, follow these guidelines to delay the development of insecticide resistance:

- Apply **Liberty Indoxacarb 30 DG** and other Group 22A insecticides within a single “treatment window” to minimize exposing successive generations of a pest species to the same mode of action insecticides.
- A “treatment window” is defined as the period of residual insecticidal activity provided by one or more applications of products with the same mode of action not to exceed approximately 30 days.
- Within the Group 22A “treatment window”, make no more than 2 applications of **Liberty Indoxacarb 30 DG** or other Group 22A insecticides.
- Following a Group 22A “treatment window”, rotate to a “treatment window” of effective insecticides with a different mode of Action Group Number. The period between Group 22A “treatment windows” should be at least 30 days.
- For short cycle crops (< 50 days), the duration of the crop cycle may be considered as the Group 22A “treatment window” if no Group 22A insecticides are used during the next crop cycle at the same farm location.
- If **Liberty Indoxacarb 30 DG** is tank mixed with other insecticides, then apply rates that are individually registered for use against the target species. **DO NOT** rely on the same mixture repeatedly to control the same pest species and follow the same “treatment window” rotation recommendation described above for both tank-mixed products.
- Use labeled rates of **Liberty Indoxacarb 30 DG** when applied alone or in tank mixtures.
- Monitor after application for unexpected target pest survival. If insect resistance is suspected consult with your manufacturer's representative, local university specialist, or certified pest control advisor.

If resistance to **Liberty Indoxacarb 30 DG** develops in your area other products with a similar mode of action (Mode of Action Group 22A) may not provide adequate control.

APPLICATION

Apply at the listed rates when insect populations reach locally determined economic thresholds. Consult the cooperative extension service, professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area.

Follow-up treatments of **Liberty Indoxacarb 30 DG** should be applied, as needed, to keep pest populations within threshold limits. Apply **Liberty Indoxacarb 30 DG** on most crops every 3 to 5 days, as specified in the specific crop sections, to maintain control. For bushberry, cranberry, dry bean, pome and stone fruit, and tree nuts the minimum interval between treatments is 7 days. Use sufficient water to obtain thorough, uniform coverage.

Because **Liberty Indoxacarb 30 DG** is most effective through ingestion of treated plant material, thorough spray coverage is essential for optimum control of targeted pest insects. Using increased water volumes will typically result in better spray coverage, especially under adverse conditions such as dry, hot weather

or dense plant foliage. **Liberty Indoxacarb 30 DG** may be applied by ground, aerial or overhead sprinkler chemigation application equipment. For aerial application use the following directions unless otherwise specified in this label: use a minimum of 5 gallons per acre (gpa) of water, except in tree and vine crops use a minimum of 10 gpa. For ground applications, use the following directions unless otherwise specified in this label: use a minimum of 10 gallons per acre (gpa) of water, except in tree and vine crops use a minimum of 50 gpa and a maximum of 200 gpa of water.

Use of Adjuvants: In some situations where coverage is difficult to achieve such as closed canopy, dense foliage, plants with waxy leaf surfaces, or less than optimum application equipment, an adjuvant may improve performance. Use only adjuvant products that are labeled for agricultural use and follow the directions on the manufacturer's label. For uses in fruit crops, use a proven and recommended adjuvant that does not affect fruit finish.

DO NOT use an adjuvant on bushberries or garden beets.

SPRAY PREPARATION

Spray equipment must be well-maintained, clean and free of previous pesticide deposits before applying **Liberty Indoxacarb 30 DG**. **Liberty Indoxacarb 30 DG** is a water dispersible granule. Fill spray tank 1/4 to 1/2 full of water. Add **Liberty Indoxacarb 30 DG** directly to spray tank. Mix thoroughly to fully disperse the insecticide while adding the remaining water. Once **Liberty Indoxacarb 30 DG** is fully dispersed continued agitation is required. Use mechanical or hydraulic means; **DO NOT** use air agitation. Spray mix must not be stored overnight in spray tank. Observe the most restrictive of the labeling limitations and precautions of all products used in mixtures.

Tank Mixing and Compatibility - Since formulations may be changed and new ones introduced, it is a best practice that users premix a small quantity of a desired tank mix and observe for possible physical incompatibility (settling out, flocculation, crystallization, etc.). This product can be tank mixed with pesticide products labeled for use on crops on this label in accordance with the most restrictive of label limitations and precautions. **DO NOT** exceed label dosage rates for any products being mixed. This product cannot be mixed with any product containing a label prohibition against such mixing.

Spray volumes of less than 3 gal of water and tank mixtures of more than two products can increase the chances of incompatible spray mixtures. A jar test (as described below) should be conducted when label guidance is not given or prior experience with a specific tank mixture is unknown. The jar test should follow the proper sequence of addition at the spray water volume planned to assure that the tank mixture is compatible. Constant agitation may be needed during mixing and spraying of mixtures. **Liberty Indoxacarb 30 DG** is compatible with most commonly used plant protectants.

Steps to conduct a jar test to determine physical tank mix compatibility of **Liberty Indoxacarb 30 DG** with other products:

- Add clean water to the jar in proportion to the planned water volume that will be used in the spray tank (a jar size of 8-16 oz is acceptable).
- While wearing the most restrictive PPE, mix proper proportional amounts of **Liberty Indoxacarb 30 DG** and desired tank mix partner(s) as will be present in the spray tank. Add one product at a time following the sequence of addition according to formulation type provided in this label.
- Seal and shake mixture after each product is added.
- Allow to stand for 1 hour.
- View jar to determine if settling, flocculation, crystallization or any other undesirable changes have happened.
- If none of the above is observed or the solution can be easily remixed after shaking, the mixture is compatible with **Liberty Indoxacarb 30 DG**.
- If the tank mixture is not compatible, a higher water volume, reduced rate of the tank mix partner(s), reduced number of tank mix partners or a compatibility agent may be needed.

Tank Mixtures and Crop Safety

Liberty Indoxacarb 30 DG is a water dispersible granule. The crop safety of **Liberty Indoxacarb 30 DG** alone or in a tank mix with many common insecticides, fungicides, nutritionals and adjuvants has been found to be acceptable. Some materials including oils, surfactants, adjuvants, nutritionals and pesticide formulations when applied individually, sequentially or in tank mixtures may solubilize the plant cuticle, facilitate penetration into plant tissue, and increase the potential for crop injury.

Applying **Liberty Indoxacarb 30 DG** with any product that produces adverse crop response in a tank mixture may also cause adverse crop response when applied in a short time sequence (i.e. seven days apart or less between applications). Such uses should be tested as described below before broad application is made.

Crop varieties can differ in their responsiveness to tank mixtures, and environmental conditions can have an influence on product performance and crop response. It is not possible to test **Liberty Indoxacarb 30 DG** alone or with all possible tank mix combinations and sequences on all varieties under all environmental conditions. When considering the use of a tank mixture on a labeled crop without prior experience, or which is not specifically described on **Liberty Indoxacarb 30 DG** product labeling or in other Liberty Crop Protection, LLC product use instructions, or when applying any products in close sequence with **Liberty Indoxacarb 30 DG**, it is important to check crop safety first. To test for crop safety, prepare a small volume of the intended tank mixture or products to be applied in a sequence, apply to an area of the target crop as directed by both this and the other product labels, and observe the treated crop to ensure that a phytotoxic response does not occur.

Use of **Liberty Indoxacarb 30 DG** in any tank mixture or sequence of applications that is not specifically described on **Liberty Indoxacarb 30 DG** product labeling or other Liberty Crop Protection, LLC product use instructions could potentially result in crop injury. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Tank Mixing Sequence - Add different formulation types in the sequence indicated below. *Allow time for complete mixing and dispersion after addition of each product.

1. Products in water soluble bags (WSB)
2. Water soluble granules (SG)
3. **Liberty Indoxacarb 30 DG** and other water dispersible granules (WG, XP, DF)
4. Wettable powders (WP)
5. Water based suspension concentrates (SC)
6. Water soluble concentrates (SL)
7. Suspoemulsions (SE)
8. Oil Based suspension concentrates (OD)
9. Emulsifiable concentrates (EC)
10. Adjuvants, surfactants and oils
11. Soluble fertilizers
12. Drift retardants

*Unless otherwise specified by manufacturer directions for use or by local expertise.

SPRAY TANK CLEANOUT

Prior to application, start with clean, well maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove.

Drain spray equipment. Thoroughly rinse sprayer and flush hoses, boom and nozzles with clean water.

Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment.

DO NOT clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

APPLICATION BY CHEMIGATION – CRANBERRY, MINT, POTATOES, SPINACH* AND SWEET CORN

* Use on spinach via overhead sprinkler irrigation is allowed only in the states of Arkansas, Georgia, Missouri, North Carolina, New Mexico, Oklahoma, and Texas unless otherwise permitted in supplemental labeling.

Instructions for the Use of Liberty Indoxacarb 30 DG in Overhead Sprinkler Chemigation Systems.

Overhead chemigation applications offer the advantage of greater penetration and coverage of the target plant. However, typical chemigation applications are more dilute than ground or aerial applications. For best results, it is recommended to keep the concentration of **Liberty Indoxacarb 30 DG** as high as possible in the application. Apply **Liberty Indoxacarb 30 DG** in 0.1 to 0.2 inches of water per acre. **Liberty Indoxacarb 30 DG** is most active as an ingestion insecticide, although it does have activity as a direct contact insecticide. For best results, applications of **Liberty Indoxacarb 30 DG** should ensure thorough coverage of the target plant to maximize the opportunity for target insects to ingest **Liberty Indoxacarb 30 DG**.

Types of Chemigation Systems:

Liberty Indoxacarb 30 DG may be applied only through overhead sprinkler irrigation systems. Overhead irrigation systems include the following; center pivot, end tow, hand move, lateral move, side roll, solid set and wheel line. Center pivot and lateral move irrigation systems are preferred. Other overhead sprinkler systems may be used if they provide uniform water distribution. **DO NOT** apply **Liberty Indoxacarb 30 DG** through any other type of irrigation system. **DO NOT** use filter screens smaller than 50 mesh throughout the system, due to possible build up of material on 100 mesh or smaller screens.

Directions for Chemigation:

Preparation

Use a pesticide tank for the application of **Liberty Indoxacarb 30 DG** in chemigation systems. Thoroughly clean the injection system and tank of any fertilizer or chemical residues using a standard clean-out procedure. Dispose of any residues in accordance with State and Federal laws. With the mix tank 1/4 to 1/2 full with water and the agitator running, measure the required amount of **Liberty Indoxacarb 30 DG** and add it to the tank. Then add additional water to bring your total pesticide mixture up to the desired volume for your application. Note: Always add the **Liberty Indoxacarb 30 DG** to water, never put **Liberty Indoxacarb 30 DG** into a dry tank or other mixing equipment without first adding water. See container label for tank mixing sequence. Continue to agitate the mixture throughout the application process. Use mechanical or hydraulic agitation, **DO NOT** use air agitation. Highly alkaline water must be buffered so that the pH of the spray solution is in the range of neutral to slightly acidic.

Injection Into Chemigation Systems

Inject the proper amount of **Liberty Indoxacarb 30 DG** into the irrigation water flow using a positive displacement injection pump. Inject the mixture at a point in the main irrigation water flow to ensure thorough mixing with the irrigation water. For continuously moving systems, inject the solution containing **Liberty Indoxacarb 30 DG** into the irrigation water line continually and uniformly throughout the irrigation cycle. Apply in no more than 0.2 inches of water per acre. For overhead sprinkler systems that are stationary, add the solution containing **Liberty Indoxacarb 30 DG** to the irrigation water line and apply no more than 0.2 inches of water per acre just before the end of the irrigation cycle.

Uniform Water Distribution

The irrigation system used for application of **Liberty Indoxacarb 30 DG** must provide for uniform distribution of **Liberty Indoxacarb 30 DG** treated water. Non-uniform distribution might result in crop injury, lack of effectiveness or illegal pesticide residues in or on the crop being treated. Ensure the irrigation system is calibrated to uniformly distribute the chemigation application to the crop. Contact the equipment manufacturer, the local University Extension agent or other experts if you have questions about achieving uniform distribution of the application.

Equipment Calibration

Calibrate the irrigation system and injector before applying **Liberty Indoxacarb 30 DG**. Calibrate the injection pump while the system is running using the expected irrigation rate. If you have questions about calibration, you should contact your state extension service specialists, equipment manufacturer or other experts.

Monitoring of Chemigation Applications

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of a responsible person, shall shut the system down and make necessary adjustments should the need arise. Wear the personal protective equipment as defined in the PPE section of the label for applicators and other handlers when making adjustments or repairs on the chemigation system when **Liberty Indoxacarb 30 DG** is in the irrigation water.

Required System Safety Devices

DO NOT connect any irrigation system used for pesticide applications to a public water system unless the pesticide label- prescribed safety devices are in place. Public water system means a system for the provision to the public of piped water for human consumption, if such a system has at least 15 service connections or regularly serves an average of at least 25 individuals at least 60 days out of the year.

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

Operation

Start the water pump and sprinkler, and let the system achieve the desired pressure and speed before starting the injector. Start the injector and calibrate the injection system according to the directions above. This procedure is necessary to deliver the desired rate per acre in a uniform manner. When the application is finished, allow the entire irrigation and injector system to be thoroughly flushed clean before stopping the system.

- End guns must be turned off during the application, if they irrigate nontarget areas or if they **DO NOT** provide uniform application and coverage.
- Plug nozzles in the immediate area of control panels, chemical supply tanks and system safety devices to prevent contamination of these areas.
- **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.
- **DO NOT** apply when system connections or fittings leak or when nozzles **DO NOT** provide uniform distribution.
- **DO NOT** allow irrigation water to collect or run-off during chemigation.

Cleaning the System

Thoroughly clean the injection system and tank of any fertilizer or chemical residues using a standard clean-out procedure. Dispose of any residues in accordance with State and Federal laws. Consult your owner's manual or your local equipment dealer for cleanout procedures for your injection system.

MANDATORY SPRAY DRIFT MANAGEMENT

AERIAL APPLICATIONS

- **DO NOT** release spray at a height greater than 10 feet above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the wind speed is greater than 10 mph, the boom length must be 65% or less for fixed-wing aircraft and 75% or less for helicopters. Otherwise, the boom length must be 75% or less for fixed-wing aircraft and 90% or less for helicopters.
- **DO NOT** apply during temperature inversions.

GROUND BOOM APPLICATIONS

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- **DO NOT** apply during temperature inversions.

AIRBLAST APPLICATIONS

- Sprays must be directed into the canopy.
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer row.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- **Volume** – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- **Adjust Nozzles** – Follow nozzle manufacturer recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT – Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, that it is configured properly, and that drift potential has been minimized.

Note: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Read the specific crop use and application equipment instructions to determine if an air assisted field crop sprayer can be used.

AIR ASSISTED (AIR BLAST) - TREE AND VINE SPRAYERS

Air assisted tree and vine sprayers carry droplets into the canopy of trees and vines via a radially or laterally directed air stream. These sprayers are not suitable for applying herbicides. In addition to the general drift management principal already described, the following specific practices will further reduce the potential for drift:

- Adjust deflectors and aiming devices so that spray is only directed into the canopy.
- Block off upward pointed nozzles when there is no overhanging canopy.
- Use only enough air volume to penetrate the canopy and provide good coverage.
- Movement of spray that goes beyond the edge of the cultivated area may be minimized by practices such as spraying the outside row only from outside the planting.

SENSITIVE AREAS

Making applications when there is a sustained wind moving away from adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is an effective way to minimize the effects of spray drift.

DRIFT CONTROL ADDITIVES

Using product compatible drift control additives can reduce drift potential. When a drift control additive is used, read and carefully observe cautionary statements and all other information on the additive's label. If using an additive that increases viscosity, ensure that the nozzles and other application equipment will function properly with a viscous spray solution. Preferred drift control additives have been certified by the Council of Producers & Distributors of Agrotechnology.

CROP ROTATION

Crops that are on this label and alfalfa, cotton, peanuts and soybeans may be planted immediately following harvest. **DO NOT** plant for food or feed any other crops not registered for use with indoxacarb for 30 days after last use.

USE SITES

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Bean, Dry Seed Including: Dried cultivars of bean (Lupinus) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin); bean (Phaseolus) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); bean (Vigna) (includes adzuki bean, blackeyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean); broad bean (dry); chickpea; guar; lablab bean; lentil	Cabbage Looper [*] Corn Earworm European Corn Borer Soybean Looper [*]	0.065 - 0.11	3.5 - 6.0	7	12 hrs.
USE RESTRICTIONS: <ul style="list-style-type: none">• DO NOT apply more than 4 applications of Liberty Indoxacarb 30 DG per acre per crop.• DO NOT apply more than 24 oz/A (0.45 lb. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per crop.• DO NOT apply more than 72 oz/A (1.35 lbs. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per calendar year.• The minimum interval between sprays is 7 days. <p>Note: For ground applications, make a uniform application in approximately 20-100 gal of water per acre. [*Not Registered for Use in California]</p>					

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Bean, Succulent Including: bean (<i>Phaseolus</i>) (includes lima bean, green; broad bean, succulent; runner bean, snap bean, wax bean); bean (<i>Vigna</i>) (includes asparagus bean, blackeyed pea, Chinese longbean, cowpea, moth bean, southern pea, yardlong bean); jackbean; sword bean	Cabbage Looper [*] Corn Earworm European Corn Borer Soybean Looper [*] Looper Suppression Only: Leafminers [*]	0.065 - 0.11	3.5 - 6.0	3	12 hrs.
USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 4 applications of Liberty Indoxacarb 30 DG per acre per crop. • DO NOT apply more than 24 oz/A (0.45 lb. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per crop. • DO NOT apply more than 72 oz/A (1.35 lbs. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per calendar year. • The minimum interval between sprays is 7 days. <p>Note: For ground applications, make a uniform application in approximately 20-100 gal of water per acre. [*Not Registered for Use in California]</p>					

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Vegetable, Brassica, Leafy Group 5 Including: Broccoli, Chinese broccoli, broccoli raab, brussels sprouts, cabbage, Chinese cabbage (napa and bok choy), Chinese mustard cabbage, cauliflower, cavalo broccolo, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens and turnip tops ¹	Beet Armyworm Diamondback moth	0.065	3.5	3	12 hrs.
	Cabbage looper Cabbage webworm[*] Cross striped cabbageworm[*] Imported cabbageworm	0.046 - 0.065	2.5 - 3.5		
USE RESTRICTIONS: <ul style="list-style-type: none">• DO NOT apply more than 4 applications of Liberty Indoxacarb 30 DG per acre per crop.• DO NOT apply more than 14 oz/A (0.26 lb. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per crop.• DO NOT apply more than 56 oz/A (1.05 lbs. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per calendar year.• DO NOT apply to greenhouse or field grown brassica crops grown for transplant.• The minimum interval between sprays is 3 days. <p>Resistance Management for Diamondback Moth: DO NOT apply Liberty Indoxacarb 30 DG more than twice to any generation of diamondback moth larvae. After the second application, rotate to another insecticide with a different mode of action (ie. a product with a different IRAC group number). DO NOT apply less than 3.5 oz/A of Liberty Indoxacarb 30 DG. If applications of Liberty Indoxacarb 30 DG DO NOT result in reduction in diamondback moth larvae populations, immediately stop use of Liberty Indoxacarb 30 DG and apply a registered insecticide with a different mode of action. DO NOT make more than 6 total applications of Liberty Indoxacarb 30 DG per calendar year for control of diamondback moth per farm location. In the State of Georgia: DO NOT apply more than 4 applications of Liberty Indoxacarb 30 DG per calendar year for the control of diamondback moth per farm location.</p> <p>Note: Add a wetting agent to improve spray coverage. [*Not Registered for Use in California] ¹For use on turnips grown for tops or greens, not for the production of turnip roots.</p>					

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Bushberries, Subgroup 13-07B Including: Aronia berry, blueberries (highbush blueberry, and lowbush blueberry), Chilean guava, currants (black currant, buffalo currant, native currant and red currant), European barberry, elderberry, gooseberry, highbush cranberry, honeysuckle, huckleberry, jostaberry, junberry, salal, sea buckthorn; cultivars, varieties and/or hybrids of these.	Cranberry fruitworm Cherry fruitworm Winter moth	0.065 - 0.11	3.5 - 6.0	7	12 hrs.
	Bruce spanworm Cranberry weevil (adult) Plum curculio (adult)	0.11	6.0		
USE RESTRICTIONS: <ul style="list-style-type: none">• DO NOT apply more than 4 applications of Liberty Indoxacarb 30 DG per acre per crop.• DO NOT apply more than 24 oz/A (0.45 lb. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per calendar year.• DO NOT apply dilute applications of more than 200 gal/A of water.• DO NOT use adjuvants.• Minimum interval between treatments is 7 days. <p>Note: For best results apply in 20 - 50 gal/A of water by ground or 10 gal/A of water by air.</p>					

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Corn (sweet) For application through tassel push only	Beet armyworm [*] [¹] European corn borer[*] Fall armyworm Corn earworm Western bean cutworm	0.046 - 0.065	2.5 - 3.5	3 (35 for forage and stover)	12 hrs. (14 days for hand harvesting)
<p>USE RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 4 applications of Liberty Indoxacarb 30 DG per acre per crop. • Whorl stage through tassel push (prior to silking) application only. • DO NOT apply more than 14 oz/A (0.26 lb. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per crop. • DO NOT apply more than 42 oz/A (0.78 lb. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per calendar year. • The minimum interval between sprays is 3 days. <p>Note: Overhead Chemigation – Liberty Indoxacarb 30 DG may be applied to sweet corn by overhead chemigation. For specific guidance see label section titled APPLICATION BY CHEMIGATION – CRANBERRY, MINT, POTATOES, SPINACH² AND SWEET CORN.</p> <p>Begin application when sweet corn is in the V1 (1st collar) stage of growth up to tassel push (V15) when damage from larvae populations exceed recommended thresholds. For best results, a slurry of Liberty Indoxacarb 30 DG, vegetable oil and an emulsifier must be kept continuously agitated in the injection tank to keep the mixture in suspension and to ensure application of the proper rate per acre.</p> <p>For aerial applications, apply using a minimum of 3 gal/A of water.</p> <p>[*Not Registered for Use in California]</p> <p>[¹ For control of Beet Armyworm in California, use the higher rate of 3.5 oz (0.065 lb ai) per acre.]</p> <p>²Use on spinach via overhead sprinkler irrigation is allowed only by the states of Arkansas, Georgia, Missouri, North Carolina, New Mexico, Oklahoma, and Texas unless otherwise permitted in supplemental labeling.</p>					

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Low Growing Berry Subgroup 13-07H, (Except Lowbush Blueberry and Strawberry) Including: Bearberry; bilberry; cloudberry; cranberry; lingonberry; muntries; partridgeberry; cultivars, varieties and/or hybrids of these	Cranberry weevil ¹ Blackheaded fireworm Black vine weevil (adult) ² (OR and WA only) Spanworm	0.11	6.0	30	12 hrs.
USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply to flow through bogs or allow release of irrigation water from bogs for at least 1 day following application. • DO NOT apply more than 4 applications of Liberty Indoxacarb 30 DG per acre per crop. • DO NOT apply more than 24 oz/A (0.45 lb. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per season. • The minimum interval between sprays is 7 days. <p>Note: For CRANBERRY only – Liberty Indoxacarb 30 DG may be applied by overhead chemigation. For specific guidance see label section titled APPLICATION BY CHEMIGATION – CRANBERRY, MINT, POTATOES, SPINACH³ AND SWEET CORN.</p> <p>¹Apply up to two applications to the spring (overwintering) generation of adult cranberry weevil prior to bloom. DO NOT apply more than 12 oz/A of Liberty Indoxacarb 30 DG (0.22 lb. ai/A) per season for control of cranberry weevils.</p> <p>²Black vine weevil adults are nocturnal feeders - it is important to monitor adult emergence by regular sweeping or trapping in the evening hours. Make repeat applications on a 7 to 10 day schedule if monitoring indicates continued adult feeding activity. Broadcast applications may need supplemental spot treatments in localized areas of heavy insect pressure. Allow 5 to 7 days to achieve maximum results, then follow-up with night time monitoring.</p> <p>³Use on spinach via overhead sprinkler irrigation is allowed only by the states of Arkansas, Georgia, Missouri, North Carolina, New Mexico, Oklahoma, and Texas unless otherwise permitted in supplemental labeling.</p>					

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Vegetable, Cucurbit, Group 9 Including: Chayote (fruit), Chinese waxgourd (Chinese) preserving melon, citron melon, cucumber, gherkin, edible gourd (including hyotan, cucuzza, hechima and Chinese okra), momordica species (including balsam apple, balsam pear, bitter melon and Chinese cucumber), muskmelon (including true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon), pumpkin, summer squash (including crookneck squash, scallop squash, straightneck squash, vegetable marrow and zucchini), winter squash (including butternut squash, calabaza, hubbard squash, acorn squash and spaghetti squash) and watermelon	Cabbage Looper Melonworm Pickleworm	0.046 - 0.11	2.5 - 6.0	3	12 hrs.
	Beet Armyworm	0.065 - 0.11	3.5 - 6.0		
USE RESTRICTIONS: <ul style="list-style-type: none">• DO NOT apply more than 4 applications of Liberty Indoxacarb 30 DG per acre per crop season.• DO NOT apply more than 24 oz/A (0.45 lb. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per crop.• DO NOT apply more than 72 oz/A (1.35 lbs. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per calendar year. The minimum interval between sprays is 5 days. <p>Note: For ground applications, apply using a minimum of 10 gal/A of water.</p>					

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Fruiting Vegetables and Okra Including: eggplant, groundcherry, pepino, peppers (bell, chili, cooking, pimento and sweet), tomatillo and tomato	Beet armyworm European Corn Borer ¹ [*]- bell pepper only Fall armyworm [*] Leafminer (Use on Florida tomatoes only - suppression only) ² Southern armyworm Tomato fruitworm (corn earworm) Tomato pinworm Western yellowstriped armyworm	0.065 - 0.11	3.5 - 6.0	3	12 hrs.
	Hornworms Loopers	0.046 - 0.11	2.5 - 6.0		
	Pepper weevil[*]	0.11	6.0		
USE RESTRICTIONS: <ul style="list-style-type: none">• DO NOT apply more than 4 applications of Liberty Indoxacarb 30 DG per acre per crop.• DO NOT apply more than 24 oz/A (0.45 lb. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per crop.• DO NOT apply more than 72 oz/A (1.35 lbs. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per calendar year.• The minimum interval between sprays is 5 days.					
¹ European corn borer applications for use only on bell peppers - For best results, begin applications of Liberty Indoxacarb 30 DG following two applications of an organo-phosphate insecticide labeled for European corn borer control in bell pepper.					
² Suppression of leafminer on Florida tomatoes- Use of an adjuvant may improve performance.					
[*Not Registered for Use in California]					

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Garden Beet	Beet armyworm	0.065 - 0.11	3.5 - 6.0	7	12 hrs.
USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 4 applications of Liberty Indoxacarb 30 DG per acre per crop. • DO NOT apply more than 24 oz/A (0.45 lb. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per crop. • DO NOT apply more than 72 oz/A (1.35 lbs. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per calendar year. • The minimum interval between sprays is 3 days. • DO NOT use adjuvants. 					

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Grape	Grape leaffolder Japanese beetle[*] Omnivorous leafroller ¹ Western grapeleaf skeletonizer	0.065 - 0.11	3.5 - 6.0	7	12 hrs.
	European grapevine moth Grape berry moth[*] Leafhoppers (suppression only) Light brown apple moth	0.09 - 0.11	5.0 - 6.0		
	Katydid (nymphs) ²	0.11	6.0		

USE RESTRICTIONS:

- **DO NOT** apply more than 2 applications of **Liberty Indoxacarb 30 DG** per acre per crop.
- **DO NOT** apply more than 12 oz/A (0.22 lb. ai/A) of **Liberty Indoxacarb 30 DG** or other indoxacarb-containing products per calendar year.
- The minimum interval between sprays is 21 days.

Note: Make the first application at initiation of egg hatch or at the first signs of infestation. Use the higher application rate for moderate to heavy insect pressure. Make application before pests reach damaging levels. Monitor fields and make an additional application if populations rebuild to potentially damaging levels. Apply in sufficient water to obtain thorough coverage of foliage. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action threshold levels for these pests.

For best results, use an adjuvant to help increase coverage, penetration and thus performance.

[*Not Registered for Use in California]

¹Use the 5-6 oz/A rate for control of omnivorous leafroller and the 3.5-5 oz/A rate for SUPPRESSION of omnivorous leafroller.

²Forktailed bush katydid (*Scudderia furcata*) and Angularwinged katydid (*Microcentrum retinerve*). Correct timing of spray application is to the early nymphal stages; thorough spray coverage is critical to achieve best results. Make repeat applications if monitoring indicates continued feeding activity.

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Fruit, Small Fruit Vine Climbing, (Except Fuzzy Kiwifruit), Subgroup 13-07F Including: Amur river grape; gooseberry; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and/or hybrids of these	Grape leaffolder Japanese beetle[*] Western grapeleaf skeletonizer	0.065 - 0.11	3.5 - 6.0	7	12 hrs.
	Grape berry moth[*] Leafhoppers (suppression only) Omnivorous leafroller	0.09 - 0.11	5.0 - 6.0		
USE RESTRICTIONS: <ul style="list-style-type: none">• DO NOT apply more than 2 applications of Liberty Indoxacarb 30 DG per acre per crop.• DO NOT apply more than 12 oz/A (0.22 lb. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per calendar year.• The minimum interval between sprays is 21 days. <p>Note: Make the first application at initiation of egg hatch or at the first signs of infestation. Use the higher application rate for moderate to heavy insect pressure. Make application before pests reach damaging levels. Monitor fields and make an additional application if populations rebuild to potentially damaging levels. Apply in sufficient water to obtain thorough coverage of foliage. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action threshold levels for these pests.</p> <p>For best results, use an adjuvant to help increase coverage, penetration and thus performance.</p> <p>[*Not Registered for Use in California]</p>					

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Vegetable, Leafy Greens, (Except <i>Brassica</i> Vegetables, Spinach and Spinach Varieties), Group 4 Including: Arugula (roquette), chervil, edible-leaved chrysanthemum, garland chrysanthemum, corn salad, garden cress, upland cress (yellow rocket, winter cress), dandelion, dock (sorrel), endive (escarole), head and leaf lettuce, orach, parsley, garden purslane, winter purslane and radicchio (red chicory)	Beet armyworm Corn earworm Diamondback moth [*]	0.065 - 0.11	3.5 - 6.0	3	12 hrs.
	Cabbage looper	0.046 - 0.11	2.5 - 6.0		
USE RESTRICTIONS: <ul style="list-style-type: none">• DO NOT apply more than 3 applications of indoxacarb-containing products at the 0.11 lb. ai/A rate per acre per crop.• A total of 4 applications of indoxacarb-containing products per crop can be made but total amounts cannot exceed 0.33 lb. ai/A (1.35 lb. ai/A) of indoxacarb per crop year.• DO NOT apply more than 18 oz/A (0.33 lb. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per crop.• DO NOT apply more than 72 oz/A (1.35 lbs. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per calendar year.• The minimum interval between sprays is 3 days.					
[*Not Registered for Use in California]					

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Vegetable, Leaf Petioles (Except Brassica Vegetables), Group 4 Including: Cardoon, celery, Chinese celery, celtuce, florence fennel (finocchio), rhubarb and swiss chard	Beet Armyworm Cabbage looper	0.065 - 0.11	3.5 - 6.0	3	12 hrs.
USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 3 applications of Liberty Indoxacarb 30 DG per acre per crop season. • DO NOT apply more than 18 oz/A (0.33 lb. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per crop. • DO NOT apply more than 72 oz/A (1.35 lbs. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per calendar year. • The minimum interval between sprays is 3 days. 					

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Mint (Peppermint and Spearmint Tops)	Cabbage looper Spotted cutworm	0.065	3.5	7	12 hrs.
USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 4 applications of Liberty Indoxacarb 30 DG per acre per crop season. • DO NOT apply more than 14 oz/A (0.26 lb. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per calendar year. • The minimum interval between sprays is 3 days. <p>Note: For ground applications, apply using a minimum of 20 gal/A of water.</p> <p>Liberty Indoxacarb 30 DG may be applied to mint by overhead chemigation. For specific guidance see label section titled APPLICATION BY CHEMIGATION –CRANBERRY, MINT, POTATOES, SPINACH¹ AND SWEET CORN</p> <p>¹Use on spinach via overhead sprinkler irrigation is allowed only by the states of Arkansas, Georgia, Missouri, North Carolina, New Mexico, Oklahoma, and Texas unless otherwise permitted in supplemental labeling.</p>					

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Pear	Codling moth - East of the Rocky Mountains	0.09 - 0.11	5.0 - 6.0	28	12 hrs.
	Codling moth - West of the Rocky Mountains ¹	0.09 - 0.11	5.0 - 6.0		
	Light brown apple moth Oriental fruit moth Pandemis leafroller[*] Redbanded leafroller White apple leafhopper[*]	0.09 - 0.11	5.0 - 6.0		
	Earwigs (adults) (suppression only) ²	0.11	6.0		

USE RESTRICTIONS:

- **DO NOT** apply more than 3 applications of **Liberty Indoxacarb 30 DG** prior to hand-thinning. No hand-thinning after the 4th application.
- **DO NOT** apply more than 4 applications of **Liberty Indoxacarb 30 DG** per acre per crop.
- **DO NOT** apply more than 24 oz/A (0.45 lb. ai/A) of **Liberty Indoxacarb 30 DG** or other indoxacarb-containing products per calendar year.
- **DO NOT** apply dilute applications of more than 200 gal/A of water. For best results apply 50 - 150 gal/A of water.
- The minimum interval between treatments is 7 days.

Note: Codling Moth Resistance Management: To minimize resistance development by codling moth: Only apply **Liberty Indoxacarb 30 DG** (or other Group 22A insecticides) to one generation of codling moth per year. Application(s) to other generations of codling moth must be with an effective product with a different mode of action (i.e. a product with a different IRAC group number).

[*Not Registered for Use in California]

¹West of the Rockies. For use against low to moderate infestations in conjunction with alternate control measures such as established Mating Disruption blocks.

²Make foliar application of **Liberty Indoxacarb 30 DG** as a fruit protection spray to SUPPRESS adult earwigs feeding on fruit. **Liberty Indoxacarb 30 DG** will not be effective against adult earwigs that are not actively feeding on fruit. When adult earwig pressure is high, two sequential applications 10-14 days apart may be needed for best results.

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Pome Fruit (Except Pear), Group 11 Including: Apple, crabapple, loquat, mayhaw, and quince	Codling moth - East of the Rocky Mountains	0.09 - 0.11	5.0 - 6.0	14	12 hrs.
	Codling moth - West of the Rocky Mountains ¹	0.09 - 0.11	5.0 - 6.0		
	European apple sawfly[*] Green fruitworm[*] Lesser appleworm Light brown apple moth Oriental fruit moth Pandemis leafroller Plum curculio Potato leafhopper Redbanded leafroller Spotted tentiform leafminer-suppression only ² [*] Tarnished plant bug Tufted apple bud moth White apple leafhopper ³	0.09 - 0.11	5.0 - 6.0		
	Lacanobia fruitworm (except California)	0.056 - 0.11	3.0 - 6.0		
	Apple maggot ⁴ Earwigs (adults) (suppression only ⁵)	0.11	6.0		

USE RESTRICTIONS:

- **DO NOT** apply more than 3 applications of **Liberty Indoxacarb 30 DG** prior to hand-thinning. No hand-thinning after the 4th application.
- **DO NOT** apply more than 4 applications per crop.
- **DO NOT** apply more than 24 oz/A (0.45 lb. ai/A) of **Liberty Indoxacarb 30 DG** or other indoxacarb-containing products per calendar year.
- **DO NOT** apply dilute applications of more than 200 gal/A of water. For best results apply 50 - 150 gal/A of water.
- The minimum interval between treatments is 7 days.

Note: Codling Moth Resistance Management: To minimize resistance development by codling moth: Only apply **Liberty Indoxacarb 30 DG** (or other Group 22A insecticides) to one generation of codling moth per year. Application(s) to other generations of codling moth must be with an effective product with a different mode of action (i.e. a product with a different IRAC group number).

*West of the Rockies. For use against low to moderate infestations in conjunction with alternate control measures such as established Mating Disruption blocks.

[* Not Registered for Use in California]

²Use of an adjuvant may improve performance. For best results, especially when using the lower use rate, use an adjuvant.

³White apple leafhopper (OR and WA only)-application rates of 2.5 - 4.9 oz/A (0.046 - 0.089 lb. ai/A) may be used for suppression of light infestations.

⁴Apple maggot - apple maggot entering the orchard from border areas may not be controlled if they **DO NOT** feed on treated apples prior to oviposition.

⁵Make foliar applications of **Liberty Indoxacarb 30 DG** as a fruit protection spray to suppress adult earwigs feeding on fruit. **Liberty Indoxacarb 30 DG** will not be effective against adult earwigs that are not actively feeding on fruit. When adult earwig pressure is high, two sequential applications 10 to 14 days

apart may be needed for best results.

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Spinach, New Zealand spinach, Vine spinach and Amaranth (leafy amaranth, Chinese spinach amaranth)	Beet armyworm Cabbage looper	0.065	3.5	3	12 hrs.
USE RESTRICTIONS: <ul style="list-style-type: none">• DO NOT apply more than 4 applications of Liberty Indoxacarb 30 DG per acre per crop.• DO NOT apply more than 14 oz/A (0.26 lb. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per crop.• DO NOT apply more than 56 oz/A (1.05 lbs. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per calendar year.• The minimum interval between sprays is 3 days. <p>Note: Make sequential applications at 3 day intervals or until insect populations are brought below threshold. Use on spinach via overhead sprinkler irrigation is allowed only in the states of Arkansas, Georgia, Missouri, North Carolina, New Mexico, Oklahoma, and Texas unless otherwise permitted in supplemental labeling. For specific guidance see label section titled APPLICATION BY CHEMIGATION – CRANBERRY, MINT, POTATOES, SPINACH¹ AND SWEET CORN.</p> <p>¹Use on spinach via overhead sprinkler irrigation is allowed only by the states of Arkansas, Georgia, Missouri, North Carolina, New Mexico, Oklahoma, and Texas unless otherwise permitted in supplemental labeling.</p>					

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Stone Fruit, Group 12 Including: Apricot, sweet cherry, tart cherry, nectarine, peach, plum, Chickasaw plum, damson plum, Japanese plum, plumcot and prune	Light brown apple moth Plum curculio	0.09 - 0.11	5.0 - 6.0	14	12 hrs.
	Katydid (nymphs) ¹ Oriental fruit moth ² Peach twig borer ³ Earwigs (adults) (suppression only) ⁴	0.11	6.0		
USE RESTRICTIONS: <ul style="list-style-type: none">• DO NOT apply more than 3 applications of Liberty Indoxacarb 30 DG prior to hand-thinning. No hand-thinning after the 4th application.• DO NOT apply more than 4 applications of Liberty Indoxacarb 30 DG per acre per crop.• DO NOT apply more than 24 oz/A (0.45 lb. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per calendar year.• DO NOT apply dilute applications of more than 200 gal/A of water. For best results apply 50 - 150 gal/A of water.• The minimum interval between treatments is 7 days.					

¹Forktailed bush katydid (*Scudderia furcata*) and Angularwinged katydid (*Microcentrum retinerve*) - Correct timing of spray application is to the early nymphal stages; thorough spray coverage is critical to achieve best results. Make repeat applications on a 7 to 10 day schedule if monitoring indicates continued feeding activity.

²Oriental fruit moth (OFM) - For applications East of the Rockies: **Liberty Indoxacarb 30 DG** is effective for control of OFM when used as part of an effective IPM program. Rotate to a product with another mode of action after each **Liberty Indoxacarb 30 DG** application.

West of the Rockies: **Liberty Indoxacarb 30 DG** provides suppression only of OFM.

³Peach twig borer (dormant and delayed dormant, CA only) - **Liberty Indoxacarb 30 DG** may be used as a dormant or delayed dormant spray for the control of first generation peach twig borer. Make application with an EPA registered dormant oil; for specific recommendations on the use of oil consult the manufacturer's label. For best performance, ground application equipment is recommended.

⁴Make foliar applications of **Liberty Indoxacarb 30 DG** as a fruit protection spray to suppress adult earwigs feeding on fruit. **Liberty Indoxacarb 30 DG** will not be effective against adult earwigs that are not actively feeding on fruit. When adult earwig pressure is high, two sequential applications 10 to 14 days apart may be needed for best results.

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Tree Nuts, Group 14-12 [*] Including: African nut-tree; almond; beechnut; Brazil nut; Brazilian pine; bunya; bur oak; butternut; cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; dika nut; ginkgo; Guiana chestnut; hazelnut (filbert); heartnut; hickory nut; Japanese horse- chestnut; macadamia nut; mongongo nut; monkey- pot; monkey puzzle nut; okari nut; pachira nut; peach palm nut; pecan; pequi; Pili nut; pine nut; pistachio; sapucaia nut; tropical almond; walnut, black; walnut, English; yellowhorn; cultivars, varieties, and/or hybrids of these	Codling moth ¹ Navel orangeworm ² Peach twig borer ³	0.075 - 0.11	4.0 - 6.0	5	12 hrs.
USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 3 applications of Liberty Indoxacarb 30 DG per crop season. • DO NOT apply more than 18 oz/A (0.33 lb. ai/A) of Liberty Indoxacarb 30 DG or other indoxacarb-containing products per calendar year. • The minimum interval between treatments is 7 days. <p>Note: Spray Volume: Thorough coverage is essential to achieve best results. Select a spray volume appropriate for the size of trees or plants and density of foliage. Where higher spray volumes are used, apply a higher rate in the specified rate range.</p> <p>Ground Application: Apply in a minimum of 50 gal/A of water by conventional ground equipment. For best results, apply 100-150 gal/A of water. Aerial Application: Apply in a minimum of 10 gal/A of water. Applying by air can result in reduced canopy coverage, which may also reduce efficacy. For best results, use an adjuvant to help increase coverage, penetration and thus performance.</p> <p>[*Not Registered for Use in California]</p> <p>¹Codling moth – (Walnut) Make initial application at or before peak egg lay for targeted generation. Depending on level of infestation reapply 14 days later as needed. Use higher rates and ground application equipment to achieve thorough coverage.</p> <p>²Navel orangeworm – For NOW, use at May or hull split timing. If targeting a hull split application, apply at 1-5% hull split. Make a second application approximately 10 – 14 days later. Depending on level of pest infestation, use of higher rates in the labeled rate range and multiple applications may be needed.</p> <p>³Peach twig borer - For spring application to overwintering generation: Make application at late dormant (just prior to bud break) to early bloom. For "May spray" applications to the summer generation: Make applications at peak moth flight (timed at or before peak egg lay). Higher rates in the labeled rate range may be needed for high infestations levels and large, dense foliage trees.</p> <p>[¹ Not for use in California]</p>					

Crops	Insects	Liberty Indoxacarb 30 DG Rate per Acre		Last Application (Days to Harvest)	REI
		Lbs. AI	Ounces		
Tuberous and Corm Vegetables, Subgroup 1C Including: Arracacha, arrowroot, Chinese artichoke, Jerusalem artichoke, edible canna (Queensland arrowroot), bitter and sweet cassava, chayote (root), chufa, dasheen (taro), ginger, leren, potato, sweet potato, tanier (cocoyam), tumeric, yam bean (jicama, manioc pea), and true yam	Cabbage looper	0.046 - 0.11	2.5 - 6.0	7	12 hrs.
	Colorado potato beetle ¹ European corn borer[*]	0.065 - 0.11	3.5 - 6.0		
	Potato tuberworm ²	0.056 - 0.11	3.0 - 6.0		

USE RESTRICTIONS:

- **DO NOT** apply more than 4 applications of **Liberty Indoxacarb 30 DG** per acre per crop.
- **DO NOT** apply more than 24 oz/A (0.45 lb. ai/A) of **Liberty Indoxacarb 30 DG** or other indoxacarb-containing products per crop.
- **DO NOT** apply more than 72 oz/A (1.35 lbs. ai/A) of **Liberty Indoxacarb 30 DG** or other indoxacarb-containing products per calendar year.
- The minimum interval between sprays is 5 days.

Note: For POTATO only - **Liberty Indoxacarb 30 DG** may be applied by overhead chemigation. For specific guidance see label section titled **APPLICATION BY CHEMIGATION – CRANBERRY, MINT, POTATOES, SPINACH AND SWEET CORN**.

[*Not Registered for Use in California]

¹Colorado potato beetle - In situations where Colorado potato beetle populations are known or suspected to be difficult to control with current insect control products, the inclusion of piperonyl butoxide (PBO), a synergist, with **Liberty Indoxacarb 30 DG** may be necessary to achieve optimum control. In these situations, a combination of **Liberty Indoxacarb 30 DG** applied at a rate of 3.5 - 6.0 oz/A combined with 0.25 lb. ai/A of PBO may be necessary to achieve the most effective control of Colorado potato beetle larvae.

Apply the low rates on small plants, small insects and light infestations of insects. Use intermediate rates on large insects and heavier infestations of insects. Use the highest specified rate for controlling severe infestations. Apply **Liberty Indoxacarb 30 DG** insecticide as a thorough coverage spray using properly calibrated air or ground spray equipment. Use sufficient water to obtain thorough and uniform coverage. For aerial application, use a minimum of 5 gal/A of water.

²Potato tuberworm foliar feeding larvae - **Liberty Indoxacarb 30 DG** is most effective when applied by ground, air or overhead chemigation to vigorously growing plants through tuber bulking prior to the beginning of crop senescence. For control of potato tuberworm foliar feeding larvae, apply **Liberty Indoxacarb 30 DG** insecticide when tuberworm larvae and/or moth counts reach locally established treatment threshold populations. **Liberty Indoxacarb 30 DG** is absorbed into leaf tissue via translaminar movement and is most effective when applied to vigorously growing plants through tuber bulking (Growth Stage IV) prior to the beginning of crop senescence (Growth Stage V). Repeat applications of effective

insecticides may be needed to keep tuberworm larvae populations as low as possible prior to harvest in order to reduce the risk of tuber damage. Failure to adequately control tuberworm larvae prior to crop senescence or vinekill increases the risk of tuber damage. To improve control of adults (moths), apply **Liberty Indoxacarb 30 DG** in a tankmix with a pyrethroid insecticide.

Potato tuberworm is a difficult pest to control due to several factors; eggs can be laid deep in the canopy and on the underside of the leaf, and larvae feed inside the leaves prior to moving to the soil to feed on the tubers. An integrated spray approach is essential. Foliar sprays alone (ground or air) may not provide adequate control of larvae in the mid to lower crop canopy. For best results, apply via chemigation or integrate chemigation applications into the foliar spray program. Ensure thorough coverage by using sufficient spray volumes. For ground applications use at least 10 gal/A of water. For aerial applications, use at least 5 gal/A of water. For best results with foliar sprays, add Methylated Seed Oil (MSO) as a spray adjuvant at 1 gal per 100 gal of spray volume (1% v/v). For chemigation applications, apply in 0.1 to 0.2 acre inch of water and add MSO at 12 to 16 fl oz/A.

DO NOT make more than two sequential applications of **Liberty Indoxacarb 30 DG** for control of potato tuberworm before rotating to another registered insecticide having a different mode of action.

³Use on spinach via overhead irrigation is allowed only by the states of Arkansas, Georgia, Missouri, North Carolina, New Mexico, Oklahoma, and Texas unless otherwise permitted in supplemental labeling.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep this product in its tightly closed container, when not in use. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals, and avoid excessive heat while in storage.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[Bag: Nonrefillable outer bag. **DO NOT** reuse or refill the outer bag. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by other procedures allowed by state and local authorities.]

[Plastic Container: Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of LIBERTY CROP PROTECTION LLC or Seller. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW all such risks shall be assumed by Buyer and User and Buyer and User agree to hold LIBERTY CROP PROTECTION LLC and Seller harmless for any claims relating to such factors.

LIBERTY CROP PROTECTION LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or LIBERTY CROP PROTECTION LLC, and TO THE EXTENT CONSISTENT WITH APPLICABLE LAW Buyer and User assume the risk of any such use. To the extent consistent with applicable law LIBERTY CROP PROTECTION LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither LIBERTY CROP PROTECTION LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF LIBERTY CROP PROTECTION LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED

ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LIBERTY CROP PROTECTION, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.
LIBERTY CROP PROTECTION LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of LIBERTY CROP PROTECTION LLC.

All trademarks are the property of the respective owners.