



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Biopesticides and Pollution Prevention Division (7511P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:
 89361-2

Date of Issuance:
 12/12/2016

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:
 Unconditional

Name of Pesticide Product:
 RADIX

Name and Address of Registrant (include ZIP Code):

Intercontinental Import Export S.A. DE C.V.
 CERRADA II NO. 3006, COL.VILLAS DE IRAPUATO
 IRAPUATO, GUANAJUATO
 MEXICO

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA Registration Number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA or the Act).

Registration is in no way to be construed as an endorsement or recommendation of this product by the U.S. Environmental Protection Agency (EPA). In order to protect health and the environment, the Administrator, on his or her motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under the Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration or registration review of your product when the EPA requires all registrants of similar products to submit such data.
2. Submit storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) data as these data requirements are not satisfied. A one-year study is required to satisfy these data requirements. You have 18 months from the date of this registration to provide these data to the EPA.

Signature of Approving Official:

Andrew Bryceland, Team Leader
 Biochemical Pesticides Branch
 Biopesticides and Pollution Prevention Division (7511P)
 Office of Pesticide Programs

Date:

12/12/2016

3. Make the following labeling change before you release this product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 89361-2.”
4. Submit one (1) copy of the final printed labeling for the record before you release this product for shipment.

Should you wish to add/retain a reference to your company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6. A stamped copy of the labeling is enclosed for your records. Please also note that the record for this product currently contains the following acceptable Confidential Statement of Formula (CSF):

- Basic CSF dated 11/09/2016

Any CSF(s) other than those listed above are superseded.

If you have any questions, please contact Cheryl Greene of my team by phone at (703) 308-0352 or via email at greene.cheryl@epa.gov.

Sincerely,



Andrew Bryceland, Team Leader
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)
Office of Pesticide Programs

Enclosure

Master Label Containing

Sublabel 1: for use with 0.30 oz (8.6 grams) tablets

Sublabel 2: for use with 0.050 oz (1.43 grams) tablets

[Master label]

ACCEPTED

12/12/2016

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 89361-2



RADIX[®]
Plant Growth Regulator
Water soluble Tablets

For application in transplanted vegetables or directly sown grains, fodder, fruits, forestry and pasture, to promote the overall development of seedlings and accelerate the recovery of the plant after transplantation and / or seeding by inducing development of lateral roots. For indoor or outdoor use.

ACTIVE INGREDIENT:

Indole-3-Butyric Acid (4-(1 H-indol-3-yl) butyric acid)35.0%

OTHER INGREDIENTS:.....65.0%

TOTAL.....100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have a person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center for further treatment advice.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For 24-hour medical assistance call **1-800-222-1222**.

EPA REG. No. **Net Weight: 0.6 oz., 0.1oz (17.2 grams or 2.86 grams)**
2 Tablet Packets

EPA EST. No. **Batch No.**
Manufactured by
Intercontinental Import Export, Salamanca, Mexico

Master Label Containing
Sublabel 1: for use with 0.30 oz (8.6 grams) tablets
Sublabel 2: for use with 0.050 oz (1.43 grams) tablets

[Sublabel 1: for use with 0.30 oz. (8.6 grams) tablets]



**Plant Growth Regulator
 Water soluble Tablets**

For application in transplanted vegetables or directly sown grains, fodder, fruits, forestry and pasture, to promote the overall development of seedlings and accelerate the recovery of the plant after transplantation and / or seeding by inducing development of lateral roots. For indoor or outdoor use.

ACTIVE INGREDIENT:

Indole-3-Butyric Acid (4-(1 H-indol-3-yl) butyric acid)35.0%
 OTHER INGREDIENTS:.....65.0%
TOTAL.....100.0%

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center for further treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For 24-hour medical assistance call 1-800-222-1222 .	

EPA REG. No.

Net Weight: 0.6 oz. (17.2 grams)

2 Tablets Packets

EPA EST. No.

Batch No.

**Manufactured by
 Intercontinental Import Export, Salamanca, Mexico**

PRECAUTIONARY STATEMENTS HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed, absorbed through skin, or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Remove and wash contaminated clothing before reusing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Wear the appropriate PPE:

- Coveralls (over long-sleeved shirt and long pants)
- Chemical-resistant footwear
- Chemical-resistant gloves
- Goggles
- Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C), or a NIOSH approved respirator with any N,R,P or HE filter.

Remove and wash contaminated clothing before reusing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of washwaters or rinsate.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other water unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product may be applied alone, as a tank mix and through chemigation. Always dissolve tablets in water prior to any application.

Chemigation Directions:

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move; flood (basin); furrow; border or drip (trickle) irrigation & system(s). Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place."

SPRINKLER CHEMIGATION SAFETY REQUIREMENTS

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system(s) designated above must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.

"The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump."

"The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down."

"The system must contain functional inter-locking controls to automatically shut off the pesticide injection pump when the water pump motor stops."

"The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected."

"Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock."

Do not apply when wind speed favors drift beyond the area intended for treatment.

Unless otherwise stated below, prior to application, be sure to completely dissolve tablets in at least one gallon of water and then add it to application equipment, tank mix or chemigation system.

Crops	Doses	Timing	Method
AGAVE (Agave Teguilana)	1 tablet dissolved in 0.26 gal of water (3000ppm)	“Seed” treatment before sowing	Spray over the base of the cutting
	1 tablet dissolved in 26 gal of water (30ppm)	Apply every 4 months	Soil Drench 3.38 oz. per plant
Garlic (Allium sativum)	1 tablet dissolved in 52 gal of water (15ppm)	Treatment of the “seed”	Seed immersion for 30 minutes.
	1 Tablet per 2.5 acre	30 days after sowing	Foliar application, spray watering or trickle irrigation
Onion (allium cepa), Carrot (Daucus carota)	2 Tablets per 2.5 acre	30 days after sowing	Foliar application, spray watering or trickle irrigation.

Crops	Doses	Timing	Method
Banana (<i>Musa paradisiaca</i>)	Dissolve 1 Tablet in 8 gal of water (100ppm)	Apply every 4 months	Apply on the corm or by the soil drench with 2.71 to 3.38 oz. per plant
Broccoli (<i>Brassica oleracea</i> var. <i>italica</i>) Cabbage (var. <i>viridis</i>)	In seed nurseries: dissolve 1 tablet in 0.26 gal of water (24ppm)	Before sowing in the cultivating media	Moisten the cultivating media before sowing and before filling the trays
Cauliflower (var. <i>Botrytis</i>), Lettuce (<i>Lactuca sativa</i>)	1- 2 Tablets per 2.5 acre	7-13 days after transplanting	Foliar application, spray watering or trickle irrigation
Squash (<i>Cucurbita pepo</i>) Melon (<i>cucumis melo</i>) Watermelon (<i>citrullus lanatus</i>)	1-2 Tablets per 2.5 acre	7-13 days after transplanting	Foliar application, spray watering or trickle irrigation
Bell Pepper (<i>Capsicum annum</i>) Tomatoes (<i>Licopersicum esculentum</i>)	4 Tablets per 2.5 acre in four applications of 1 tablet each per 2.5 acre	Applications on the 10, 40, 130 and 190 days after transplanting	Foliar application, spray watering or trickle irrigation

Crops	Doses	Timing	Method
Barley (hordeum vulgare) Maize (Zea mays) Wheat (triticum aestivum) Sorghum (sorghum vulgare)	2 Tablets per 2.5 acre	30-40 days after emergence of plant	Foliar application, spray watering or trickle irrigation
Asparagus (Asparagus officinalis)	1 Tablet per 2.5 acre	Apply every 4 months	Foliar application, spray watering or trickle irrigation
Strawberry (Fragaria vesca) Raspberry (Rubus idaeus) Blackberry (Rubus fruticosus)	Dissolve 1 tablet in 52 gal of water (15 ppm)	Before sowing	Immersion of the stolons form 2 to 8 hours
	1 Tablet per 2.5 acre	30, 90, 150, and 210 days after sowing	Foliar application, spray watering or trickle irrigation
Potatoes (Solanum tuberosum)	Dissolve 1 tablet in 52 gal of water (15ppm)	Seed treatment	Spray over the "seed" or quick dip the "seed"
	1 Tablet per 2.5 acre	30 to 40 days after sowing	Foliar application, spray watering or trickle irrigation
Turf Pastures and Range Lands	1 Tablet per 2.5 acre	Apply every 30 to 60 days	Foliar application, spray watering or trickle irrigation.

Crops	Doses	Timing	Method
Avocado (Persea Americana) Apple (Malus domestica) Mango (Mangifera indica) Walnut (Juglans regia) Grapes (Vitis vinifera) Coffee (coffea arabica)	Dissolve 1 Tablet in 0.5 gal of water (1500ppm)	Cuttings and rootstock propagation	Quick dip on the cutting end
Cocoa (Theobroma cacao)	2 to 4 Tablets per 2.5 acre	Apply every 3 to 4 months	Foliar application, spray watering or trickle irrigation.
Citrus Fruits	Dissolve 4 tablets in 52 gal of water (60ppm)	Apply every 4 months	Soil drench 3.38 oz. per plant
Papaya (carica papaya)	Dissolve 2 Tablets in 52 gal of water (30ppm)	Apply every 30 to 60 days	Soil drench 6.09 oz. per plant
Pineapples (ananas comosus)	6 Tablets per 2.5 acre	30 to 45 days after sowing	Foliar application, spray watering or trickle irrigation.

Crops	Doses	Timing	Method
Ornamental Plants	1 Tablet dissolved in 26.42 gal of water (30 ppm)	10 to 15 days after transplant and every 60 days	Drench in between 1.69 to 3.38 oz. of dissolution per plant
	250 ppm (1 tablet in 3 gal) to 12000 ppm (8 tablets in 0.5 gal) depending on size of the cutting and species. Trials and literature are recommended for precise information on ideal concentrations	Cuttings and rootstock propagation	Quick dip on the cutting end

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Do not store with foodstuff.

PESTICIDE DISPOSAL: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or, by other procedures approved by state and local authorities.

WARRANTY STATEMENT

Intercontinental Import Export warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of Intercontinental Import Export. To the extent consistent with applicable law, Intercontinental Import Export shall not be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. To the extent consistent with applicable law, the exclusive remedy of any buyer or user of this product for any and all losses, injuries, or damages resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall not exceed the purchase price paid for this product or at Intercontinental Import Export election, the replacement of this product. Intercontinental Import Export **MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

Master Label containing

Sublabel 1: for use with 0.30 oz. (8.6 grams) tablets

Sublabel 2: for use with 0.050 oz. (1.43 grams) tablets

[Sublabel 2: for use with 0.050 oz. (1.43 gram) tablets]



**Plant Growth Regulator
Water soluble Tablets**

For application in transplanted vegetables or directly sown grains, fodder, fruits, forestry and pasture, to promote the overall development of seedlings and accelerate the recovery of the plant after transplantation and / or seeding by inducing development of lateral roots. For indoor or outdoor use.

ACTIVE INGREDIENT:

Indole-3-Butyric Acid (4-(1 H-indol-3-yl) butyric acid)35.0%

OTHER INGREDIENTS:.....65.0%

TOTAL.....100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center for further treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For 24-hour medical assistance call 1-800-222-1222 .	

EPA REG. No.

Net Weight: 0.1 oz. (2.86 grams)

2 Tablet Packets

EPA EST. No.

Batch No.

Manufactured by

Intercontinental Import Export, Salamanca, Mexico

PRECAUTIONARY STATEMENTS HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed, absorbed through skin, or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Remove and wash contaminated clothing before reusing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Wear the appropriate PPE:

- Coveralls (over long-sleeved shirt and long pants)
- Chemical-resistant footwear
- Chemical-resistant gloves
- Goggles
- Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C), or a NIOSH approved respirator with any N,R,P or HE filter.

Remove and wash contaminated clothing before reusing.

ENVIROMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of washwaters or rinsate.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other water unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product may be applied alone, as a tank mix and through chemigation. Always dissolve tablets in water prior to any application.

Chemigation Directions:

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move; flood (basin); furrow; border or drip (trickle) irrigation & system(s). Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place."

SPRINKLER CHEMIGATION SAFETY REQUIREMENTS

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system(s) designated above must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.

"The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump."

"The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down."

"The system must contain functional inter-locking controls to automatically shut off the pesticide injection pump when the water pump motor stops."

"The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected."

"Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock."

Do not apply when wind speed favors drift beyond the area intended for treatment.

Unless otherwise stated below, prior to application, be sure to completely dissolve tablets in at least one gallon of water and then add it to application equipment, tank mix or chemigation system.

Crops	Doses	Timing	Method
AGAVE (Agave Teguilana)	1 tablet dissolved in 0.5 gal of water (3000ppm)	“Seed” treatment before sowing	Spray over the base of the cutting
	1 tablet dissolved in 4.5 gal of water (30ppm)	Apply every 4 months	Soil Drench 3.38 oz. per plant
Garlic (Allium sativum)	1 tablet dissolved in 9 gal of water (15ppm)	Treatment of the “seed”	Seed immersion for 30 minutes.
	1-2 Tablets per acre	30 days after sowing	Foliar application, spray watering or trickle irrigation
Onion (allium cepa), Carrot (Daucus carota)	5 Tablets per acre	30 days after sowing	Foliar application, spray watering or trickle irrigation.

Crops	Doses	Timing	Method
Banana (Musa paradisiaca)	Dissolve 1 Tablet in 1 gal of water (100ppm)	Apply every 4 months	Apply on the corm or by the soil drench with 2.71 to 3.38 oz. per plant
Broccoli (Brassica oleracea var. italic) Cabbage (var. viridis)	In seed nurseries: dissolve 1 tablet in 0.26 gal of water (24ppm)	Before sowing in the cultivating media	Moisten the cultivating media before sowing and before filling the trays

Cauliflower (var. Botrytis), Lettuce (Lactuca sativa)	2-5 Tablets per acre	7-13 days after transplanting	Foliar application, spray watering or trickle irrigation
Squash (Cucurbita pepo) Melon (cucumis melo) Watermelon (citrullus lanatus)	2-5 Tablets per acre	7-13 days after transplanting	Foliar application, spray watering or trickle irrigation
Bell Pepper (Capsicum annum) Tomatoes (Licopersicum esculentum)	4 Tablets per acre in four applications of one tablet each application	Applications on the 10, 40, 130 and 190 days after transplanting	Foliar application, spray watering or trickle irrigation

Crops	Doses	Timing	Method
Barley (hordeum vulgare) Maize (Zea mays) Wheat (triticum aestivum) Sorghum (sorghum vulgare)	4-5 Tablets per acre	30-40 days after emergence of plant	Foliar application, spray watering or trickle irrigation
Asparagus (Asparagus officinalis)	2-3 Tablets per acre	Apply every 4 months	Foliar application, spray watering or trickle irrigation
Strawberry (Fragaria vesca) Raspberry (Rubus ideaus) Blackberry (Rubus fruticosus)	Dissolve 6 tablets in 52 gal of water (15 ppm)	Before sowing	Immersion of the stolons form 2 to 8 hours
	1-2 Tablets per acre	30, 90, 150, and 210 days after sowing	Foliar application, spray watering or trickle irrigation
Potatoes (Solanum tuberosum)	Dissolve 6 tablets in 52 gal of water (15ppm)	Seed treatment	Spray over the "seed" or quick dip the "seed"
	1-2 Tablets per acre	30 to 40 days after sowing	Foliar application, spray watering or trickle irrigation
Turf Pastures and Range Lands	2-3 Tablet per acre	Apply every 30 to 60 days	Foliar application, spray watering or trickle irrigation.

Crops	Doses	Timing	Method
Avocado (Persea Americana) Apple (Malus domestica) Mango (Mangifera indica) Walnut (Juglans regia) Grapes (Vitis vinifera) Coffee (coffea arabica)	Dissolve 6 Tablets in 0.5 gal of water (1500ppm)	Cuttings and rootstock propagation	Quick dip on the cutting end
Cocoa (Theobroma cacao)	4-10 Tablets per acre	Apply every 3 to 4 months	Foliar application, spray watering or trickle irrigation.
Citrus Fruits	Dissolve 24 tablets in 52 gal of water (60ppm)	Apply every 4 months	Soil drench 3.38 oz. per plant
Papaya (carica papaya)	Dissolve 12 Tablets in 52 gal of water (30ppm)	Apply every 30 to 60 days	Soil drench 6.09 oz. per plant
Pineapples (ananas comosus)	14 Tablets per acre	30 to 45 days after sowing	Foliar application, spray watering or trickle irrigation.

Crops	Doses	Timing	Method
Ornamental Plants	1 Tablet dissolved in 26.42 gal of water (30 ppm)	10 to 15 days after transplant and every 60 days	Drench in between 1.69 to 3.38 oz. of dissolution per plant
	250 ppm (1 tablet in 3 gal) to 12000 ppm (8 tablets in 0.5 gal) depending on size of the cutting and species. Trials and literature are recommended for precise information on ideal concentrations	Cuttings and rootstock propagation	Quick dip on the cutting end

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Do not store with foodstuff.

PESTICIDE DISPOSAL: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or, by other procedures approved by state and local authorities.

WARRANTY STATEMENT

Intercontinental Import Export warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of Intercontinental Import Export. To the extent consistent with applicable law, Intercontinental Import Export shall not be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. To the extent consistent with applicable law, the exclusive remedy of any buyer or user of this product for any and all losses, injuries, or damages resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall not exceed the purchase price paid for this product or at Intercontinental Import Export election, the replacement of this product. Intercontinental Import Export **MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**