


89442-2

11/20/2012

11/12



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number: **89442-2**
 Date of Issuance: **NOV 20 2012**

NOTICE OF PESTICIDE:
 Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance: **unconditional**
 Name of Pesticide Product:
Ethofumesate Select

Name and Address of Registrant:
Prime Source, LLC
4609 E. Boonville-New Harmony Road
Evansville, IN 47725

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.


The basic and alternate formulations #1 and #2 (all dated July 31, 2012) are acceptable and will be added to your file.

This product is registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration review/reregistration of your product when the Agency requires all registrants of similar products to submit data.
2. Make the following label revision(s):
 - a. Revise "EPA Reg. No." to "EPA Reg. No. 89442-2"
 - b. Assure that the EPA establishment number and net contents are also added to the final printed label.
3. Per 40 CFR 156.10(6), submit one copy of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec.6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. If you have any questions regarding this notice, please contact Beth Benbow of my staff at benbow.bethany@epa.gov.

Signature of Approving Official:
 Kathryn V. Montague
 Project Manager 23
 Herbicide Branch
 Registration Division (7505P)



Date:
NOV 20 2012

Ethofumesate Select

NOT FOR USE ON COMMERCIAL SOD FARMS IN ARIZONA
SUSPENSION CONCENTRATE HERBICIDE

Selective Herbicide Intended For Use By Professional Applicators On Ornamental Turf Only

ACTIVE INGREDIENT:	% by Weight
Ethofumesate (2-ethoxy-2,3-dihydro-3,3-dimethyl-5-benzofuranyl methanesulfonate).....	42.0%
OTHER INGREDIENTS:	58.0%
TOTAL:	100.0%

This product contains 4.0 lbs. active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUCION


Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail)

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time or your poison control center at 1-800-222-1222.	

EPA Reg. No.

Manufactured for:
Prime Source, LLC
4609 E. Boonville-New Harmony Road
Evansville, IN 47725

EPA Est. No.



Net Contents:

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, the pesticide registered under EPA Reg. No. 8942-a

3/12

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or inhaled. Avoid breathing vapor or spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this product are polyethylene or polyvinyl chloride. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All Mixers, Loaders, Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (such as natural rubber)
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Engineering Controls for: Enclosed Cabs for Aerial Applicators: Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

On-Site Closed Mixing and Loading System Engineering Controls for Liquid Formulations:

Mixers and loaders must use a closed system that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4)] for Dermal protection, and must:

- wear the personal protective equipment required in the PPE section of this labeling for mixers and loaders as well as a chemical-resistant apron
- wear protective eyewear, if the system operates under pressure, and
- be provided chemical-resistant footwear that is immediately available in an emergency, such as a broken package, spill or equipment breakdown.

USER SAFETY REQUIREMENTS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not graze livestock on treated turf. Do not feed treated grass clippings to livestock. Read entire Use Precautions and

Restrictions, Directions for Use and Condition of Sale and Limitation of Warranty and Liability before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency in your State responsible for pesticide regulation.

SHAKE CONTAINER WELL BEFORE USING

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours for all crops, except turf grown for sod. The REI for turf is 9 days. The REI for each crop is listed in the directions for use associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter until sprays have dried.

Professional pesticide applicators applying to residential turf, including home lawns, parks, and recreation areas must inform their customers that all persons and pets must be kept off the treated turf until sprays have dried.

PRODUCT INFORMATION

Ethofumesate Select is a herbicide intended for use on ornamental turf such as golf courses, sod farms, parks, cemeteries and homeowner or commercial lawns, and after overseeding specific grasses. It may be used on established perennial ryegrass, Kentucky bluegrass, creeping bentgrass, turf-type tall fescue, St. Augustinegrass, and dormant bermudagrass for the control and/or suppression of the annual grasses and broadleaf weeds listed in the tables below. Ethofumesate Select is intended for professional use only.

Ethofumesate Select has both preemergent and early (two-leaf stage) postemergence activity and works best in programs emphasizing both approaches. Ethofumesate Select application is most effective on healthy, vigorously growing turf.

SPRAY DRIFT MANAGEMENT

This chemical can contaminate surface water through spray drift. A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g.,

5/12

ground, aerial) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Wind Speed: Do not apply at wind speeds greater than 15 mph.

Temperature Inversions: Do not make applications into areas of temperature inversion of stable atmospheric conditions.

WEED SPECIES CONTROLLED

PREEMERGENCE

Annual Grasses

- Barnyardgrass (*Echinochloa crus-galli*)
- Bluegrass, annual (*Poa annua*)
- Canarygrass (*Phalaris canariensis*)
- Crabgrass, large (*Digitaria sanguinalis*)
- Crabgrass, smooth (*Digitaria ischaemum*)
- Foxtail, green (*Setaria viridis*)
- Foxtail, yellow (*Setaria glauca*)

Annual Broadleaves

- Burclover (*Medicago sp.*)
- Chickweed, common (*Stellaria media*)
- Purslane, common (*Portulaca oleracea*)
- Pigweed, redroot (*Amaranthus retroflexus*)

Ethofumesate Select will also reduce competition from:

- Nutsedge, purple (*Cyperus rotundus*)
- Nutsedge, yellow (*Cyperus esculentus*)

POSTEMERGENCE

Annual Grasses

- Bluegrass, annual (*Poa annua*)

Annual Broadleaves

- Chickweed, common (*Stellaria media*)
- Clover, white (*Trifolium repens*)

PREEMERGENCE/EARLY POSTEMERGENCE

Annual Grasses

- Crabgrass, large (*Digitaria sanguinalis*)
- Crabgrass, smooth (*Digitaria ischaemum*)

USE PRECAUTIONS and RESTRICTIONS

- Ethofumesate Select is not intended for use by homeowners and should be applied to homeowner lawns by licensed or certified applicators.
- DO NOT OVER APPLY Ethofumesate Select. Follow the instructions in this label or damage to non-target turf may result.
- Do not apply more than 1.5 lbs. ai (48 oz. product/A or 1.1 oz. product/1,000 sq. ft.) per year.
- Do not harvest treated sod for 16 days following application.
- Spray overlap can cause turf injury due to over application. Use of a spray colorant or indicator is recommended so that spray overlap can be avoided.
- Do not apply with hand-held or flood jet nozzles because treatments may not be uniform.
- Do not apply this product through any type of irrigation system.

SEE CHART BELOW FOR USE RATES AND TIMINGS AND READ TEXT BELOW FOR SPECIFIC DIRECTIONS FOR EACH TYPE OF GRASS.

Application Rates and Timings

Turf Type	Primary Targets	Application Timing	Use Rate ¹		# of Apps ⁵	Application Interval (Days)	Overseeding Safety Interval ³ (Weeks After Treatment)	Application Safety Interval ⁴ (Weeks After Emergence)
			Oz./ Acre	Oz/ 1,000 Sq. Ft.				
Creeping Bentgrass	Annual Bluegrass	Fall	24	9/16	0-2	21-28	4	4
		Spring ²	24	9/16	0-2	21-28		
Kentucky Bluegrass	Annual Bluegrass	Fall	24	9/16	2	21-28	6	8
Ryegrass	Annual Bluegrass	Fall Spring	32-48	3/4-1.1	0-1 0-1	N/A N/A	1-2	1-2
St. Augustinegrass (Est. Turf)	Bermudagrass Suppression	Spring	24	9/16	2	21-28	N/A	N/A
Turf-Type Tall Fescue	Annual Bluegrass	Fall	24-48	3/4-1.1	1-2	21-28	0	2-3
Overseeded (ryegrass) Bermudagrass: Dormant Bermudagrass	Annual Bluegrass	Fall	24-48	9/16 - 1.1	1-2	21-28	N/A	N/A
Nondormant Bermudagrass	Annual Bluegrass	Fall	32-36	3/4-7/8	1	N/A	N/A	N/A

N/A= Not applicable.

¹ Apply Ethofumesate Select in 20 to 60 gallons of water per acre or 1 to 3 gallons of water per 1000 sq. ft.

² Only make spring applications after previous fall treatments.

³ After the last treatment of Ethofumesate Select, the interval provided applies to overseeding of the specific grass on same type of grass to which Ethofumesate Select was applied.

⁴ Ethofumesate Select may be applied to specific grass following seed emergence after the recommended time interval.

⁵ Do not exceed the maximum single application rate or yearly maximum application rate of 48 oz./acre (i.e. a split application of 24 oz./A once in the spring and once in the fall OR a single application of 48 oz./A in either the spring or the fall).

INSTRUCTIONS FOR SPECIFIC GRASS TYPES

COOL SEASON TURFGRASSES

Creeping Bentgrass

Apply Ethofumesate Select to bentgrass as long as the turf is at a length typically found on fairways (or longer). When applying to bentgrass, use the following guidelines.

- Avoid using Ethofumesate Select in areas that are heavily shaded and/or poorly drained.
- Creeping bentgrass tolerance to Ethofumesate Select may be improved by tank mixing Ethofumesate Select with nitrogen fertilizer (controlled release or soluble) at a rate of 0.1 to 0.25 lb N per 10,000 ft². This may also improve creeping bentgrass conversion in the areas treated.
- Control of annual bluegrass is best achieved by making two applications of Ethofumesate Select: one in the fall followed by one in the spring.

- Ethofumesate Select works best in the spring if applied after creeping bentgrass has resumed active growth and is fully green.

When overseeding with creeping bentgrass, use the following guidelines:

- After applying Ethofumesate Select, wait a minimum of 3-4 weeks before overseeding with bentgrass.
- When an area has been renovated or overseeded with bentgrass, do not apply Ethofumesate Select until 3-4 weeks after seedling emergence.

Kentucky Bluegrass

- Delay application of Ethofumesate Select to bluegrass until at least 8 weeks after emergence.
- Bluegrass overseeded with ryegrass may be treated with Ethofumesate Select 1-2 weeks after ryegrass emergence.

When overseeding with Kentucky Bluegrass use the following guidelines:

- Wait at least six weeks after the last Ethofumesate Select application before reseeding with Kentucky bluegrass.
- Note that sod quality during the spring may be diminished following the fall applications of Ethofumesate Select. Alternatives to fall application of Ethofumesate Select include seeding with dormant bluegrass in the late fall or to delay reseeding until the spring.

Perennial Ryegrass

Control of annual bluegrass in perennial ryegrass is easiest when it is newly emerged, and application of Ethofumesate Select is made during the primary period of annual bluegrass germination (and up to 30 days after annual bluegrass emergence) in the fall and/or spring. Consult your local Extension Service or university weed specialist for the date(s) that annual bluegrass germinates in your area.

- For best results, apply Ethofumesate Select at 24 oz./A (0.5 oz./1,000 sq. ft.) once in the fall and once in the spring.
- Application in the fall or spring will also reduce competition from selected broadleaf weeds and crabgrass.

When overseeding with perennial ryegrass, use the following guidelines:

- Wait 1-2 weeks (or until the seedlings are approximately 1" tall) before applying Ethofumesate Select to the seeded area.
- To the extent possible, thatch should be removed from the area being seeded as it can diminish the effectiveness of the Ethofumesate Select treatment.
- Do not apply mulch or straw to the seeded areas until after Ethofumesate Select has been applied.

Turf-Type Tall Fescue

Make Ethofumesate Select applications to turf-type tall fescue in the fall. If overseeding with turf-type tall fescue, application of Ethofumesate Select may be done at the same time as seeding.

WARM SEASON TURFGRASSES

St. Augustinegrass (Established Turf)

Ethofumesate Select suppresses the development of actively growing Bermudagrass in established St. Augustine sod. Apply Ethofumesate Select to St. Augustinegrass in the spring and early summer to suppress Bermudagrass and Bermudagrass seedhead formation and/or for control of annual bluegrass.

Notes regarding applications to St. Augustinegrass:

- Do not treat St. Augustinegrass in the first six months after germination.
- Do not treat St. Augustinegrass that is under stress or injury to the turf may result.
- Temporary stunting and minor discoloration of St. Augustinegrass may occur after application. If the St. Augustinegrass shows signs of severe yellowing or stunting, discontinue applications.

Use the following guidelines when applying Ethofumesate Select to St. Augustinegrass:

- Begin Ethofumesate Select applications when Bermudagrass first breaks dormancy in the spring. Application timing is critical to achieve optimum results and will vary depending on location and temperature.
- For best results, make two applications of Ethofumesate Select at 24 oz./A (0.5 oz./1,000 sq. ft.) with the second application being made 21-28 days after the first application.
- To improve suppression, each application of Ethofumesate Select may be tank-mixed with (Atrazine at up to 2 lb ai/A for first application and 0.75 lb-1 lb ai/A triazine for second or third application).

Overseeded Dormant Bermudagrass

Notes regarding applications to Bermudagrass:

- Do not make more than two applications at 24 oz./A (0.5 oz./1,000 sq. ft.) OR one application at 48 oz./A (1.0 oz./1,000 sq. ft.) of Ethofumesate Select when treating nondormant bermudagrass. Treatment to turf that is not fully dormant may cause early injury and/or delayed spring green-up.
- Application of Ethofumesate Select to bermudagrass that is stressed due to shade, poorly drained soils, and high traffic may result in increased turf injury. Avoid use of Ethofumesate Selector use lower rates under these conditions.

Use the following guidelines when applying Ethofumesate Select to Bermudagrass:

- To control annual bluegrass, make applications of Ethofumesate Select in late fall (ideally 1-2 weeks after emergence of overseeded perennial ryegrass) on overseeded bermudagrass.
- When Ethofumesate Select is applied in late November or early December, applications of 32-36 oz. per acre have shown acceptable control of annual bluegrass with minimum injury to nondormant (or predormant) bermudagrass. If lower rates are used, reduced annual bluegrass control should be expected. If higher rates are used, increased injury to Bermudagrass may occur.
- Ethofumesate Select should not be applied to Bermudagrass in the 4 weeks prior to breaking winter dormancy. Applications made to Bermudagrass in the 4 weeks prior to breaking dormancy may temporarily delay the normal start of active growth.

When overseeding bermudagrass, use the following guidelines:

- Common bermudagrass in fairways or roughs may be more susceptible to herbicide injury than hybrid bermudagrass.
- When making applications to nondormant bermudagrass, Ethofumesate Select should be kept within areas that are overseeded so that the ryegrass will mask any early injury or late transition in the spring that may occur.
- Using a higher than normal seeding rate may be desirable in order to minimize the appearance of thin turf.

Ethofumesate Select Mixtures with Fertilizers

Ethofumesate Select Impregnation on Dry Bulk Fertilizers

Ethofumesate Select may be impregnated on many dry bulk fertilizers (See "1" below) and applied for the control of labeled grasses and broadleaf weeds. All Ethofumesate Select label and supplementary literature instructions and precautions regarding rates per acre, soil type, application, and other directions must be followed. All individual State regulations relating to dry bulk fertilizer blending, registration, labeling and application are the responsibility of the individual and/or company selling the Ethofumesate Select fertilizer mixtures. A minimum of 200 pounds and a maximum of 700 pounds of approved fertilizer ingredients impregnated with the appropriate amount of Ethofumesate Select must be applied per acre. For impregnating the pesticide on dry fertilizers, use a closed rotary drum type mixer equipped with suitable spraying equipment. The spray nozzles should be positioned inside of the mixer to provide uniform spray coverage of the tumbling fertilizer. The Ethofumesate Select should be sprayed uniformly onto the fertilizer using a fine spray pattern.

The physical properties of fertilizers vary, particularly in liquid absorptive capacity. When absorptivity is sufficient, simple spray impregnation of the fertilizer with Ethofumesate Select provides a satisfactory dry mixture. If the absorptivity is not adequate, use of a highly absorptive powder is required to provide a dry, free-

flowing mixture. Microcel E (Johns-Manville Products Corporation) is the recommended absorbent powder. It should be added separately and uniformly to the prepared pesticide/fertilizer mixture in a quantity that is sufficient to provide a suitably free-flowing mixture. Generally, less than 2% by weight of Microcel E is required.

The amount of Ethofumesate Select actually required in the formulation of specific fertilizer mixtures should be calibrated carefully for each production operation, This is necessary to ensure that the amount of Ethofumesate Select actually contained in the fertilizer mixture applied to the soil represents the correct dosage rate.

Bulk fertilizers impregnated with Ethofumesate Select should be applied immediately, NOT STORED.

1. Approved dry fertilizer ingredients for use with Ethofumesate Select.

Ethofumesate Select Impregnation on Dry Bulk Fertilizers

	N	P	K
Ammonium nitrate	34	0	0
Ammonium sulfate	21	0	0
Ammonium phosphate-sulfate	16	20	0
Diammonium phosphate	18	46	0
Monoammonium phosphate	11	56	0
Potassium chloride	0	0	60
Potassium sulfate	0	0	52
Single superphosphate	0	20	0
Triple superphosphate	0	46	0
Urea	45	0	0

Ethofumesate Select Physical Data	
Density	1.15 g/cm ³
Pounds/gallon	9.60
Flashpoint	Non-combustible

Ethofumesate Select with Liquid Fertilizer

The following procedure is suggested for evaluation of physical compatibility of Ethofumesate Select in mixtures with liquid fertilizers for spray tank applications.

Material Required

1. Ethofumesate Select- components of tank sizes if intended for use
2. Liquid fertilizer to be used.
3. Adjuvant for fertilizer tank mix: Compex* or E-Z Mix**
4. Two (or more) one quart wide mouth containers with lids or stoppers
5. Measuring spoons (25 ml pipette or graduated cylinder provides more accurate measurement)
6. Measuring cup, 8 fl. oz. (237 ml)

*Compex, Kalo Baloratories, Inc., Kansas City, MO

**E-Z Mix, United Agri-Products, Greeley, CO

Procedure

1. Pour one pint (473 ml) of the liquid fertilizer into each of the quart containers.
2. Add adjuvant(s) to one or more of the containers and mix; follow label directs of adjuvant.
3. Add the Ethofumesate Select and tank mix components to the containers (see rate table below).
4. Close the containers with lids or stoppers and mix contents by inverting the containers ten times.
5. Inspect the surface and body of mixture:
 - a. immediately after mixing

11/12

- b. after allowing mixtures to stand quietly for 30 minutes,
- c. immediately after mixing again (invert the containers ten more times).

If uniform mixture does not occur, the spray tank mixture should not be used. If any of the mixtures remain uniform for 30 minutes, that mixture may be used in spray tank applications. Should any of the mixtures separate after 30 minutes but remix readily into a uniform mixture with inversion of the container, the mixture may be used provided that the adequate agitation is maintained in the spray tank. If a Ethofumesate Select plus fertilizer mixture utilizing an adjuvant is satisfactory, but the one without adjuvant is not, be sure to use the adjuvant in the spray tank at the rate recommended on the label which was used in the test.

If non-dispersible oil, sludge, or clumps of solids form in the mixtures, those combinations should not be used for spray tank application.

Rate Table for Ethofumesate Select Mixtures with Liquid Fertilizers

Gal. of Liquid Fertilizer to be applied per acre	ml or tsp of Ethofumesate Select to be added to 1 pint of fertilizer	
	ml	tsp.
gal 20	17.6	3.6
30	12.0	2.4
40	9.0	1.9
50	7.1	1.5
60	6.0	1.2

* Based on field rate of 3.0 lb. ai/acre (3/4 gal/acre) in the fertilizer volumes indicated. Adjust amount of Ethofumesate Select added proportionately to correspond with intended field use rate listed on Ethofumesate Select label. Add the proportionate amount of tank mix component.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Protect product from freezing temperatures.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

(Nonrefillable container ≤ 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Nonrefillable > 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

12/
12

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Prime Source, LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Prime Source, LLC and Seller harmless for any claims relating to such factors.

Prime Source, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Prime Source, LLC, and Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, PRIME SOURCE, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent consistent with applicable law, neither Prime Source, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF PRIME SOURCE, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF PRIME SOURCE, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Prime Source, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Prime Source, LL