

89442-4

12/6/2012

1/50



U S ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (7504P)  
Ariel Rios Building  
1200 Pennsylvania Ave , NW  
Washington, D C 20460

EPA Registration Number  
89442-4

Date of Issuance  
DEC - 6 2012

NOTICE OF PESTICIDE  
 Registration  
 Reregistration

Term of Issuance Unconditional

Name of Pesticide Product  
PS Clethodim

(under FIFRA as amended)

Name and Address of Registrant (include ZIP Code)  
**Prime Source, LLC**  
**c/o Wagner Regulatory Associates, Inc**  
**P O Box 640**  
**Hockessin, DE 19707**

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant the above named pesticide is hereby registered/reregistered under the Federal Insecticide Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment the Administrator on his motion may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA sec 3(c)(5) provided that you

- 1 Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data
- 2 Make the following label revisions
  - a Revise the EPA Reg No to 89442-4
  - b Assure that the establishment number and net content are also added to the label
- 3 Submit one year storage stability (guideline 830 6317) and corrosion characteristics (guidelines 830 6320 studies) with observations made at 0 3 6 9 and 12 month intervals. The results from both study types must be submitted to the USEPA
- 4 Submit one copy of the revised final printed label before you release the product for shipment

If these conditions are not complied with the registration will be subject to cancellation in accordance with FIFRA sec 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records

If you have any questions regarding the Notice please contact Grant Rowland at (703) 347-0254 or [rowland.grant@epa.gov](mailto:rowland.grant@epa.gov)

Signature of Approving Official  
Kathryn Montague  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)

Date  
DEC - 6

2/50

### PS Clethodim

**ACTIVE INGREDIENT**

\*Clethodim

**By Wt**

26.4%

**OTHER INGREDIENTS**

73.6%

**TOTAL**

100.0%

Contains Petroleum Distillates

\*(E)-2-[1[[[(3-chloro-2-propenyl)-oxy]imino]propyl] 5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one

Contains 2.0 lbs clethodim per gallon

**KEEP OUT OF REACH OF CHILDREN**

**WARNING – AVISO**

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle  
(If you do not understand the label find someone to explain it to you in detail)

FIRST AID	
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes</li> <li>Remove contact lenses if present after the first 5 minutes then continue rinsing eye</li> <li>Call a poison control center or doctor for treatment advice</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes</li> <li>Call a poison control center or doctor for treatment advice</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>Immediately call a poison control center or doctor</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor</li> <li>Do not give any liquid to the person</li> <li>Do not give anything by mouth to an unconscious person</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>Move person to fresh air</li> <li>If person is not breathing call 911 or an ambulance then give artificial respiration preferably by mouth-to-mouth if possible</li> <li>Call a poison control center or doctor for further treatment advice</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment For emergency information concerning this product call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week 6:30 am to 4:30 pm Pacific Time or your poison control center at 1-800-222-1222	
NOTE TO PHYSICIAN	
Ingestion of this product or subsequent vomiting can result in aspiration of light hydrocarbon liquid which can cause pneumonitis If ingested probable mucosal damage may contraindicate the use of gastric lavage	

EPA Reg No 89442-

Manufactured for  
Prime Source LLC  
4609 E Boonville-New Harmony Road  
Evansville IN 47725

DEC - 6 2012  
89442-4

EPA Est No

Net Contents

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
WARNING**

Causes substantial but temporary eye irritation Avoid contact with skin Do not get in eyes on skin or on clothing Harmful if swallowed or inhaled Avoid breathing vapors or spray mist Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are listed below If you want more options follow the instructions for category G on an EPA chemical resistance category selection chart

**Applicators and other handlers must wear** long-sleeved shirt and long pants chemical-resistant gloves such as Barrier Laminate or Viton ≥ 14 mils shoes plus socks and protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate Do not reuse them Follow manufacturer's instructions for cleaning/maintaining PPE If no such instructions for washables exist use detergent and hot water Keep and wash PPE separately from other laundry

**USER SAFETY RECOMMENDATIONS**

**Users should**

- Wash thoroughly with soap and water after handling and before eating drinking chewing gum using tobacco or using the toilet
- Remove clothing/PPE immediately if pesticide gets inside Then wash thoroughly and put on clean clothing
- Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark Do not apply where runoff is likely to occur Do not apply where weather conditions favor drift from areas treated Do not contaminate water when disposing of equipment washwater or rinsate

The use of this product may pose a hazard to the federally designated endangered species of Solano Grass and Wild Rice Use of this product is prohibited in the following areas where the species are known to exist

Solano Grass Solano County California the vernal lakes are bounded by the Union Pacific Railroad and Hastings Road to the north Highway 113 to the east Highway 12 to the south and Travis Air Force Base to the west

Wild Rice Hays County Texas

**PHYSICAL OR CHEMICAL HAZARDS**

**Combustible** Do not use or store near heat or open flame

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

**READ THE ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS**

Do not apply this product in a way that will contact workers or other persons either directly or through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe consult the agency responsible for pesticide regulation

4/50

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170 This Standard contains requirements for the protection of agricultural workers on farms forests nurseries and greenhouses and handlers of agricultural pesticides It contains requirements for training decontamination notification and emergency assistance It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants soil or water is coveralls chemical-resistant gloves such as Barrier Laminate or Viton ≥ 14 mils shoes plus socks and protective eyewear

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170) The WPS applies when this product is used to produce agricultural plants on farms forest nurseries or greenhouses

Keep all unprotected persons out of operating areas or vicinity where there may be drift Do not enter treated areas without protective clothing until sprays have dried

### TANK MIXES

**NOTICE** Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor

Read and follow the entire label of each product to be used in the tank with this product

**[THE FOLLOWING STATEMENT ON CHEMIGATION WILL BE USED ONLY IF A SUPPLEMENTAL LABEL IS CREATED**

### CHEMIGATION

Do not apply this product through any irrigation system unless the supplemental labeling on chemigation is followed Refer to supplemental labeling entitled Application of PS Clethodim Onions (dry bulbs and green) and Garlic by Chemigation for use directions for chemigation

May be applied to onions and garlic by sprinkler irrigation systems Do not apply by chemigation to any other crop or to this crop using any other type of irrigation system ]

### PRODUCT INFORMATION

FOR USE ON Alfalfa Artichoke (Globe) Asparagus Bean and Pea (dry shelled)<sup>1</sup> Bean and Pea (succulent)<sup>2</sup> Broccoli Bushberry<sup>3</sup> Cabbage Caneberry<sup>4</sup> Canola\* Carrot Cauliflower (and other Head and Stem Brassica Vegetables)<sup>5</sup> Celery Clover (grown in Idaho Oregon and Washington only) Conifers Cotton Cranberry Cucurbit Eggplant (and other Fruiting Vegetables)<sup>6</sup> Fallow Land (and other non-producing agricultural areas) Flax\* Field Corn<sup>7</sup> Garden Beet Garlic Herbs<sup>8</sup> Hops Horseradish (and other Root Vegetables)<sup>9</sup> Legume Vegetables (edible podded)<sup>10</sup> Lettuce Head and Leaf (and other leafy greens)<sup>11</sup> Melons (including Cantaloupes and Watermelons)<sup>12</sup> Mint Mustard Greens (and other leafy brassica greens)<sup>13</sup> Mustard Seed\* Non-Bearing Food Crops Non-Crop or Non-Planted Areas Onions (dry bulb and green) Ornamentals

Peaches Peanut (including perennial) Peppers (bell and non-bell) Potato Radish Rhubarb (and other Leaf Petioles)<sup>14</sup> Safflower Sesame Shallots (dry bulbs and green) Soybeans Squash (including Pumpkins) Strawberry Sugar Beet Sunflower Sweet Potato Tomato Turnip Greens Yam (and other Tuberous and Corm Vegetables)<sup>15</sup>

**\*Not for use in California**

<sup>1</sup> Other Dry Shelled Bean and Pea crops approved for use with PS Clethodim include Bean (*Lupinus* spp ) grain sweet white and white sweet Bean (*Phaseolus* spp ) field kidney lima (dry) navy pinto and tepary Bean (*Vigna* spp ) adzuki bean black-eyed pea catjang cowpea crowder pea moth bean mung bean rice bean Southern pea urd bean broad (dry) chickpea (garbanzo) guar lablab bean and lentil Pea (*Pisum* spp ) field and pigeon

<sup>2</sup> Other Succulent Bean and Pea crops approved for use with PS Clethodim includes Bean (*Lupinus* spp ) grain sweet white and white sweet Bean (*Phaseolus* spp ) field kidney lima (dry) navy pinto and tepary Bean (*Vigna* spp ) adzuki bean black-eyed pea catjang cowpea crowder pea moth bean mung bean rice bean Southern pea urd bean broad (dry) chickpea (garbanzo) guar lablab bean and lentil Pea (*Pisum* spp ) field and pigeon

<sup>3</sup> Other bushberry crops approved for use with PS Clethodim include Aronia Berry Blueberry (highbush) Black Currant Buffalo Currant Chilean Guava Cranberry (highbush) Elderberry European Barberry Gooseberry Honeysuckle (edible) Huckleberry (edible) Jostaberry Juneberry (Saskatoon berry) Native Currant Salal Sea Buckthorn Red Currant cultivars varieties and/or hybrids of these

<sup>4</sup> Other caneberry crops approved for use with PS Clethodim include Blackberry Loganberry Raspberry (black and red) Wild Raspberry cultivars varieties and/or hybrids of these

<sup>5</sup> Other head and stem brassica vegetables approved for use with PS Clethodim include Chinese broccoli Brussels sprouts Chinese (napa) cabbage Chinese mustard cavalo broccoli and kohlrabi

<sup>6</sup> Other Fruiting Vegetables (except tomato) approved for use with PS Clethodim include eggplant groundcherry pepino peppers (all) and tomatillo

<sup>7</sup> For burndown of existing stand of Roundup Ready field corn or volunteer Roundup Ready field corn prior to replanting field corn

<sup>8</sup> Other Herb crops approved for use with PS Clethodim include angelica balm basil borage burnet chamomile catnip chervil (dried) chive Chinese chive clary coriander (leaf) costmary cilantro (leaf) curry (leaf) dill (dillweed) horehound hyssop lavender lovage (leaf) marigold marjoram (*origanum* spp ) nasturtium parsley (dried) pennyroyal rosemary rue sage and savory summer and winter

<sup>9</sup> Other root vegetables approved for use with PS Clethodim include burdock edible celeriac chervil turnip-rooted chicory ginseng parsley turnip-rooted parsnip radish oriental rutabaga salsify black salsify Spanish skirret and turnip

<sup>10</sup> Other Edible Podded Legume Vegetable crops approved for use with PS Clethodim include Bean (*Phaseoulus* spp ) runner snap and wax Bean (*Vigna* spp ) asparagus Chinese longbean moth yardlong jackbean Pea (*Pisum* spp ) dwarf edible-pod snow sugar snap pigeon and sword bean

<sup>11</sup> Other Leafy Greens crops approved for use with PS Clethodim include amaranth (Chinese spinach leafy amaranth and tampala) arugula (roquette), chervil chrysanthemum (edible-leaved and garland) corn salad cress (garden yellow rock and winter) dandelion dock (sorrel) endive (escarole) lettuce (head and leaf) orach parsley purslane (garden and winter) radicchio (red chicory) spinach (New Zealand and Vine (Indian and Malabar))

<sup>12</sup> Other cucurbit crops approved for use with PS Clethodim include Chayote (fruit) Chinese Wax Gourd Citron Melon Edible Gourd Gherkin and Muskmelons (all) including Honeydew Melon

<sup>13</sup> Other leafy brassica greens approved for use with PS Clethodim include broccoli raab Chinese (bok choy) cabbage collards kale mizuna mustard greens mustard spinach and rape greens

<sup>14</sup> Other leaf petiole crops approved for use with PS Clethodim include cardoon celtuce Chinese celery Florence fennel and Swiss chard

<sup>15</sup> Other tuber and corm vegetables approved for use with PS Clethodim include arracacha arrowroot Chinese artichoke Jerusalem artichoke edible burdock edible canna bitter and sweet cassava chayote (root) chufa dasheen (taro) ginger leren tanier turmeric and bean yam

6/50

PS Clethodim is not recommended for use on vegetable crops being grown for seed production unless specific use directions are provided

PS Clethodim is a selective postemergence herbicide for control of annual and perennial grasses PS Clethodim does not control sedges or broadleaf weeds

Repeated use of PS Clethodim (or similar postemergence grass herbicides with the same mode of action) may lead to the selection of naturally occurring biotypes that are resistant to these products in some grass species

If poor performance occurs and cannot be attributed to adverse weather or application conditions a resistant biotype may be present This is most likely to occur in fields where other control strategies such as crop rotation mechanical removal and other classes of herbicides are not used from year to year

Do not allow PS Clethodim to come in contact with desirable grass crops such as corn rice sorghum small grains or turf as these and other grass crops will be injured or killed Minor leaf spotting may occur on treated plants under certain environmental conditions New foliage is not affected

**Control Symptoms**

Treated grass weeds show a reduction in vigor and growth Early chlorosis/necrosis of younger plant tissue is followed by a progressive collapse of the remaining foliage Symptoms will generally be observed in 7 to 14 days after application depending on grass species treated and environmental conditions


**APPLICATION INFORMATION**

**Timing of Applications**

Apply PS Clethodim postemergence to actively growing grasses according to rate table instructions Applications made to grass plants stressed by insufficient moisture or hot or cold temperatures or to grass plants exceeding listed growth stages may result in unsatisfactory control Do not apply under these conditions

In arid regions where irrigation is used to supplement limited rainfall PS Clethodim should be applied as soon as possible after irrigation (within 7 days) In arid regions a second application of PS Clethodim will generally provide more effective control of perennial grass weeds than a single application Make second application to actively growing grass 2 to 3 weeks after emergence of new growth

Cultivation of treated grasses 7 days prior to or within 7 days after application of PS Clethodim may reduce weed control DO NOT APPLY PS Clethodim if rainfall is expected within one hour since control may be reduced



**ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE**

CROP	ADJUVANT USE INSTRUCTIONS
Alfalfa Cotton Dry Shelled Bean & Pea Edible Podded Legume Vegetable Peanuts (including perennial) Potato Soybean Succulent Bean & Pea Sugar Beet and Sunflower	Always use a crop oil concentrate* at 1.0 qt /A by ground or 1% v/v (but not less than 1 pt /A) in the finished spray volume by air  1 to 2 qts /A of liquid fertilizer (10-34-0 28%N or 32%N) or an equivalent amount (2.5 to 4.0 lbs /A) of spray grade ammonium sulfate (AMS) may be added to PS Clethodim applications in addition to the recommended rate of crop oil concentrate. The addition of AMS has shown improved grass control for difficult to control species including quackgrass rhizome Johnsongrass red rice wild oats volunteer cereals and volunteer corn
Artichoke (Globe) Asparagus Bushberry Caneberry Canola Carrot Clover Cranberry Cucurbits Flax Fruiting Vegetables (except Tomato) Garden Beet Garlic Head & Stem Brassica Vegetables Herbs Hops Leaf Petioles Leafy Brassica Greens Leafy Greens Mint Mustard Seed Onion (Dry Bulb & Green) Peaches Root Vegetables Safflower Sesame Shallots (Dry Bulb & Green) Strawberry Sweet Potato (Yam & other Tuberous and Corm Vegetables except Potato) and Tomato	Always use a crop oil concentrate at 1% v/v in the finished spray volume unless tank mix instructions indicate otherwise  Addition of liquid fertilizer is not recommended for these crops
Non-Bearing Food Crops Ornamental Plants	Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt per 50 gals (0.25% v/v). Use of crop oil concentrate is not recommended since it may injure flowers and foliage
Conifer Trees Fallow Land (and other non-producing agricultural areas) and Non-Crop or Non-Planted Areas	Always use a crop oil concentrate containing at least 15% emulsifier at 1% v/v (but not less than 1 pt /A) in the finished spray volume

\*Acceptable crop oil concentrates would be those that contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality, and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils.

**Ground Application**

Use of sufficient spray volumes and pressure is essential to ensure complete coverage. Use a minimum of 5 gals and a maximum of 40 gals of spray solution per acre. Under the following conditions a minimum of 10 gals per acre is required: ultra narrow row cotton, narrow row soybeans, broadleaf herbicide tank mixes, perennial grasses, volunteer corn, drought or stress conditions, heavy grass pressure or when grasses are at or near maximum height. Failure to use a minimum of 10 gals per acre under these conditions can result in poor coverage and reduced grass control requiring repeat applications. Spray pressures should reflect a minimum of 30 psi and a maximum of 60 psi at the nozzle. Do not use flood nozzles.

Applications to onions (dry bulbs and green), garlic, and shallots (dry bulbs and green) should be made in a minimum of 20 gals of spray solution per acre.

**Air Application**

Use a minimum of 3 gals of spray solution per acre unless otherwise directed in this label. Increase spray volumes up to 10 gals as grass or crop foliage becomes dense. For onions (dry bulbs and green) garlic or shallots (dry bulbs and green) When applying by air do not exceed 8 fl oz /A in a single application. In California air applications to onions garlic or shallots should be made in a minimum of 20 gals of spray solution per acre. In states other than California air application to onions garlic or shallots should be made in a minimum of 10 gals of spray solution per acre.

**NOTE Crop injury may occur when PS Clethodim is applied to onions, garlic or shallots with aerial equipment**

**Spot Treatment**

When using hand sprayers or high volume sprayers utilizing hand guns mix 1/4% to 1/2% (0.33 oz to 0.65 oz per gal) PS Clethodim and treat to wet vegetation while not allowing runoff of spray solution. For uses requiring crop oil concentrate include crop oil concentrate at 1% (1.3 oz per gal) by volume. For uses requiring non-ionic surfactant include non-ionic surfactant at 1.4% (0.33 oz per gal) by volume.

**NOTE If PS Clethodim is applied as a spot treatment, care should be taken to not exceed the maximum rate allowed on a "per acre" basis or crop injury may occur**

**CHEMIGATION – ONION (Dry Bulb and Green) AND GARLIC  
SPRINKLER IRRIGATION APPLICATION**

- **Do not apply PS Clethodim by chemigation in the states of Idaho, Montana, Oregon and Washington**

Apply PS Clethodim at the high rate recommended for annual grasses (16 fl oz per acre) when the grass height is at the low end of the range (application to larger grasses may not provide adequate control). Add a crop concentrate containing at least 15% emulsifier at 1 quart per acre.

Apply PS Clethodim in 0.1 to 0.2 acre-inch of water either at the end of a regular irrigation set or as a separate application not associated with a regular irrigation using the least amount of water that provides proper distribution and coverage. Application of more than label recommended quantities of irrigation water per acre may result in decreased product performance by removing the chemical from the zone of effectiveness. Use a metering device to inject the PS Clethodim into the irrigation water at a constant flow. Constant agitation must be maintained in the chemical supply tank during the entire period of herbicide application. Inject the product with a positive displacement pump into the main line ahead of a right angle turn to ensure adequate mixing. Allow time for all lines to flush the herbicide through all nozzles before turning off irrigation water. To ensure the lines are flushed and free of remaining herbicide a dye indicator may be injected into the lines to mark the end of the application period.

It is not recommended that PS Clethodim be applied through an irrigation system connected to a public water system. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

**Use Precautions**

- 1 Apply this product only through sprinkler irrigation systems including center pivot lateral move end tow side (wheel) roll travelers big gun solid set or hand move. Do not apply this product through any other type of irrigation system.
- 2 Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of treated water.
- 3 If you have any questions about calibration you should contact State Extension Service specialists, equipment manufacturers or other experts.
- 4 Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.



- 5 A person knowledgeable of the chemigation system and responsible for its operation or under supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise
- 6 The system must contain a functional check valve vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow
- 7 The pesticide injection pipeline must contain a functional automatic quick-closing check valve to prevent the flow of fluid back toward the injection pump
- 8 The pesticide injection pipeline must also contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down
- 9 The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops
- 10 The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected
- 11 Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock
- 12 Do not apply when wind speed favors drift beyond the area intended for treatment

**RESTRICTIONS AND LIMITATIONS**

Do not apply if rain is expected within 1 hour of application as control may be unsatisfactory

Do not apply a postemergence broadleaf herbicide within one day following application of PS Clethodim or reduced grass control may result

PS Clethodim is not recommended for use on vegetable crops being grown for seed production unless specific use directions are provided

For canola do not apply more than 6 fl oz of PS Clethodim per acre per season For clover flax mustard seed and radish crops do not apply more than 16 fl oz of PS Clethodim (0.25 lb ai) per acre per season For all other crops do not apply more than 32 fl oz of PS Clethodim (0.50 lb ai) per acre per season Application on Long Island New York is restricted to no more than 16 fl oz of PS Clethodim (0.25 lb ai) per acre per season

Do not apply more than 8 fl oz /A of PS Clethodim per application to the following crops asparagus brassica vegetables (head and stem) bean (succulent) carrot cranberry cucurbits flax fruiting vegetables (except tomato) garden beet green onion herbs hops leaf petioles leafy brassica greens leafy greens legume vegetables (edible podded) non-bearing food crops pea (dry shelled) pea (succulent) root vegetables safflower sesame and strawberry Do not apply more than 6 fl oz /A of PS Clethodim per application to canola or mustard seed For all other crops do not apply more than 16 fl oz of PS Clethodim (0.25 lb ai) per acre per application **Exceeding these rates may result in unacceptable crop injury**

Do not apply under conditions of stress Applying PS Clethodim under conditions that do not promote active grass growth will reduce herbicide effectiveness These conditions include drought excessive water extremes in temperature low humidity and grasses either partially controlled or stunted from prior pesticide applications Grasses under these kinds of stressful conditions will not absorb and translocate PS Clethodim effectively and will be less susceptible to herbicide activity

Optimal perennial grass control can be obtained if rhizomes or stolons are cut up by preplant tillage practices (discing plowing etc ) to stimulate maximum emergence of grass shoots Cultural practices such as continuous no-tillage in which the perennial grass rhizomes or stolons are not cut up result in a very

staggered non-uniform weed emergence Due to this non-uniform weed emergence no fewer than two PS Clethodim applications per season per year are recommended at the appropriate weed-growth stage rate under continuous no-till conditions

Grass crops such as corn rice sorghum small grains or turf etc are highly sensitive to PS Clethodim

While all the vegetable crops on this label have been tested and are tolerant to PS Clethodim not all specialty varieties of these crops have been tested It is advised that before applying PS Clethodim to specialty varieties of vegetable crops on this label crop tolerance be investigated first using a small section of the field It is possible that injury symptoms can occur Symptoms may appear as leaf speckling or stunting

Always read and follow the restrictions and limitations for all products whether used alone or in a tank mix The most restrictive labeling of any product used applies in tank mixtures including all crop rotational and other crop restrictions

Tank mixes of PS Clethodim and broadleaf herbicides may result in reduced grass control If grass regrowth occurs an additional application of PS Clethodim may be necessary **AVOID SPRAY DRIFT**

Do not allow spray from ground or aerial equipment to drift onto adjacent land or crops When drift may be a problem do everything possible to reduce spray drift including

- Do not spray if wind speeds are or become excessive
- Do not spray if wind speed is 10 mph or greater If sensitive crops or plants are downwind extreme caution must be used under all conditions
- Do not spray if winds are gusty
- Use extreme caution when conditions are favorable for drift (high temperatures drought low relative humidity) especially when sensitive plants are located nearby
- Do not apply when a temperature inversion exists If inversion conditions are suspected consult with local weather services before making an application
- Further reductions in drift can be obtained by
  - 1 Using large droplet size sprays Do not use nozzles that produce small droplets Orient nozzles downward and slightly backward as needed to reduce drift for ground applications
  - 2 Orienting nozzles straight back with windstream using straight stream orifices for aerial applications Use the lowest number of nozzles practical with the largest possible orifice size to obtain the minimum 3 GPA volume Application height and boom length should be set according to manufacturer s instructions to minimize drift
  - 3 Increasing the volume of spray mixture (for example a minimum of 10 GPA for ground applications) by using higher flow rate nozzles Using lower pressure with the appropriate nozzle to obtain higher volumes will also reduce drift
  - 4 Applying as close to target plants as practical while maintaining a good spray pattern for adequate coverage

**Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption**

**CROP SPECIFIC RESTRICTIONS AND LIMITATIONS FOR PS CLETHODIM**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Alfalfa</b> including Sainfoin Holy Clover Birdsfoot trefoil <sup>(3)</sup>	15 days before grazing feeding or harvesting (cutting) for forage or hay	6-16 fl oz <sup>(4)</sup>	1 qt by ground or 1% v/v (but not less than 1 pt /A) by air <sup>(5)</sup>	Refer to tank mix partners for feeding grazing and harvesting restrictions <sup>(5,6)</sup>  The addition of AMS has shown improved grass control for difficult to control species including quackgrass rhizome Johnsongrass red rice wild oats volunteer cereals and volunteer corn
<b>Artichoke (Globe)</b>	5 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application  For repeat applications make on a minimum of a 14 day interval  Do not apply more than 32 fl oz /A (0.5 lb ai/A) per season
<b>Asparagus</b>	1 day	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application  For repeat applications make on a minimum of a 14 day interval
<b>Bean, Dry Shelled</b> including Bean ( <i>Lupinus</i> spp ) Grain Sweet White White Sweet Bean ( <i>Phaseolus</i> spp ) Field Kidney Lima (dry) Navy Pinto Tepary Bean ( <i>Vigna</i> spp ) Adzuki Bean Black-eyed Pea Catjang Cowpea Crowder Pea Moth Bean Mung Bean Rice Bean Southern Pea Urd Bean Broad (dry) Chickpea (garbanzo) Guar Lablab Bean Lentil	30 days	6-16 fl oz	1 qt by ground or 1% v/v (but not less than 1 pt /A) by air <sup>(5)</sup>	Do not apply more than 16 fl oz /A per application  For repeat applications make on a minimum of 14 day interval  Refer to appropriate Table for reduced rate directions for the control of small annual grasses  The addition of AMS has shown improved grass control for difficult to control species including quackgrass rhizome Johnsongrass red rice wild oats volunteer cereals and volunteer corn

12/50

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Bean, Succulent</b> including Bean ( <i>Phaseolus</i> spp ) Broad Bean (succulent) Lima Bean (green) Bean ( <i>Vigna</i> spp ) Black-eyed Pea Cowpea Southern Pea	21 days	6-8 fl oz	1 qt by ground or 1% v/v (but not less than 1 pt /A) by air <sup>(5)</sup>	Refer to appropriate Table for reduced rate directions for the control of small annual grasses  Do not apply more than one (1) application per acre per season  The addition of AMS has shown improved grass control for difficult to control species including quackgrass rhizome Johnsongrass red rice wild oats volunteer cereals and volunteer corn
<b>Beet, Garden</b>	30 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application  For repeat applications make on a minimum of a 14 day interval
<b>Brassica Vegetables, Head &amp; Stem</b> including Broccoli Brussels sprouts Cabbage Cauliflower	30 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application  For repeat applications make on a minimum of a 14 day interval
<b>Bushberry,</b> including Aronia berry Blueberry highbush Chilean Guava Cranberry Highbush Currant Black Currant Buffalo Currant native Currant Red Elderberry European Barberry Gooseberry Honeysuckle edible Huckleberry Jostaberry Juneberry Saskatoon berry Native currant Salal Sea Buckthorn Cultivars varieties and/or hybrids of these	14 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application  Do not apply more than 32 fl oz (0.5 lb ai/A) per season  For repeat applications make on a minimum of a 14 day interval  Apply at the base of the plant where grassy weeds are growing near the ground  Do not apply to low growing berries  Do not apply to Bushberry grown for root stock

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Caneberry,</b> including Blackberry Loganberry Raspberry black Raspberry red Raspberry wild Cultivars varieties and/or hybrids of these	7 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application  Do not apply more than 32 fl oz /A (0.5 lb ai/A) per season  For repeat applications make on a minimum of a 14 day interval  Apply at the base of the plant where grassy weeds are growing near the ground  Do not apply to low growing berries  Do not apply to Caneberry grown for root stock
<b>Canola</b>	70 days	4-6 fl oz	1% v/v in the finished spray volume	Do not apply after crop has begun bolting. Crop injury may occur when PS Clethodim is applied during the bloom period  Do not apply more than 6 fl oz /A in a single application  Do not exceed 6 fl oz/A in a season
<b>Carrot</b>	30 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application  For repeat applications make on a minimum of a 14 day interval
<b>Clover</b>	15 days before grazing feeding or harvesting (cutting) for forage or hay	6-16 fl oz	1% v/v in the finished spray volume	For use on clover grown in the states of Idaho, Oregon and Washington only  Do not exceed 16 fl oz/A in a season
<b>Corn, Field</b> For burndown of existing stand of Roundup Ready field corn or volunteer Roundup Ready field corn prior to replanting field corn See Directions for Use in Roundup Ready Field Corn (Burndown) table	90 days	3 fl oz	Do not use COC or MSO with PS Clethodim in this use pattern	Do not make more than 1 application per season  Do not apply more than 3 fl oz /A per season  To control the existing stand replant no sooner than 6 days after application

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Cotton</b>	60 days	6-16 fl oz	1 qt by ground or 1% v/v (but not less than 1 pt /A) by air <sup>(5)</sup>	Do not graze treated fields or feed treated forage or hay to livestock  The addition of AMS has shown improved grass control for difficult to control species including quackgrass rhizome Johnsongrass red rice wild oats volunteer cereals and volunteer corn
<b>Cranberry</b>	30 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application  Do not apply between the hook stage and full fruit set  For repeat applications make on a minimum of a 14 day interval
<b>Cucurbits</b> including Chayote (fruit) Chinese Wax Gourd Citron Melon Cucumber Gherkin Gourd Edible Muskmelons (all) including Cantaloupes Honeydew Melon Pumpkin Squash (all) Watermelon	14 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application  For repeat applications make on a minimum of a 14 day interval
<b>Fallow Land</b> <b>Conifer Trees</b> (and other non-producing agricultural areas)  <b>Non-Crop or Non-Planted Areas</b>	N/A	6-16 fl oz	1% v/v (but not less than 1 pt /A) in the finished spray volume using a crop oil concentrate containing at least 15% emulsifier	Do not plant any crop for 30 days after application unless clethodim is registered for use in that crop
<b>Flax</b>	60 days	6-8 fl oz	1% v/v in the finished spray volume	Apply prior to bloom Crop injury may occur when PS Clethodim is applied during the bloom period  Do not apply more than 8 fl oz /A in a single application  Do not exceed 16 fl oz /A in a season

15/50

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Fruiting Vegetable</b> (except Tomato) including Eggplant Groundcherry Pepino Peppers (all) Tomatillo	20 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application  For repeat applications make on a minimum of a 14 day interval
<b>Herbs</b> including Angelica Balm Basil Borage Burnet Camomile Catnip Chervil (dried) Chive Chinese Clary Coriander (leaf) Costmary Cilantro (leaf) Curry (leaf) Dill (dillweed) Horehound Hyssop Lavender Lovage (leaf) Marigold Marjoram ( <i>Origanum</i> spp) Nasturtium Parsley (dried) Pennyroyal Rosemary Rue Sage Savory Summer and Winter	14 days	6-8 fl oz	1% v/v in the finished spray volume	PS Clethodim has not been tested on all herbs and herb varieties It is the responsibility of the user to test PS Clethodim on a small portion of the crop to be treated before treating the entire field  Crop tolerance should be verified to PS Clethodim on a small area of the herb crop at the desired PS Clethodim rate and with the same crop oil concentrate that will be used on the herb field If no crop response is evident seven (7) days after treatment PS Clethodim may be used on the entire field at the rate tested and with the same crop oil used in the tolerance test  Do not apply more than 8 fl oz /A in a single application  For repeat applications make on a minimum of a 14 day interval
<b>Hops</b>	21 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application  For repeat applications make on a minimum of a 14 day interval
<b>Leaf Petioles</b> including Cardoon Celery Celtuce Chinese Celery Fennel Florence (finochio) Rhubarb Swiss Chard	30 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application  For repeat applications make on a minimum of a 14 day interval

16/50

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Leafy Brassica Greens</b> including Broccoli Raab Cabbage Chinese (bok choy) Collards Kale Mizuna Mustard Greens Mustard Spinach Rape Greens	14 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in the single application  For repeat applications make on a minimum of a 14 day interval
<b>Leafy Greens</b> including Amaranth Chinese Spinach Leafy Amaranth Tampala Arugula (roquette) Chervil Chrysanthemum Edible-leaved Chrysanthemum Garland Corn Salad Cress Garden Upland (yellow rock and winter) Dandelion Dock (sorrel) Endive (escarole) Lettuce Head and Leaf Orach Parsley Purslane Garden Winter Radicchio (red chicory) Spinach New Zealand Vine (Indian and malabar)	14 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application  For repeat applications make on a minimum of a 14 day interval



171/50

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Legume</b> <b>Vegetables, Edible Podded</b> including Bean ( <i>Phaseolus</i> spp ) Runner Snap Wax Bean ( <i>Vigna</i> spp ) Asparagus Chinese Longbean Moth Yardlong Jackbean Pea ( <i>Pisum</i> spp ) Dwarf Edible-pod Snow Sugar Snap Pigeon Sword Bean	21 days	6-8 fl oz	1 qt by ground or 1% v/v (but not less than 1 pt /A) by air <sup>(5)</sup>	Do not apply more than 8 fl oz /A in a single application  Do not apply more than one (1) application per acre per season  <i>For peas apply before bloom but no later than 21 days before harvest</i>  Refer to appropriate Table for reduced rate directions for the control of small annual grasses  The addition of AMS has shown improved grass control for difficult to control species including quackgrass rhizome Johnsongrass red rice wild oats volunteer cereals and volunteer corn
<b>Mint</b>	21 days	6-16 fl oz <sup>(4)</sup>	1 qt by ground or 1% v/v (but not less than 1 pt /A by air )	Do not apply more than 16 fl oz /A in a single application  For repeat applications make on a minimum of a 14 day interval
<b>Mustard Seed</b>	75 days	4-6 fl oz	1% v/v in the finished spray volume	Do not apply after crop has begun bolting Crop injury may occur when PS Clethodim is applied during the bloom period  Do not apply more than 6 fl oz /A in a single application  Do not exceed 16 fl oz /A in a season
<b>Onion</b> (Dry Bulb Only) <b>Garlic</b> <b>Shallot</b> (Dry Bulb Only)	45 days	6-16 fl oz <sup>(7) (8)</sup>	1% v/v in the finished spray volume	Minimum of 20 gals /A spray volume by ground in entire U S  Minimum of 20 gals /A spray volume by air in California <sup>(9)</sup>  In states other than California air application to onions garlic or shallots should be made in a minimum of 10 gals /A

18/50

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Onion, Green</b> including Green Eschalot Japanese Bunching Onion Leeks Scallion or Spring Onion	14 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application  For repeat applications make on a minimum of a 14 day interval
<b>Ornamentals</b>	N/A	6-16 fl oz	Use of crop oil concentrate is not recommended since it may injure flowers and foliage See Special Use Instructions	Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt per 50 gals (0.25% v/v)  Sugar maples cannot be tapped for syrup within one year of PS Clethodim application  Do not apply more than 8 fl oz /A in a single application to non-bearing food crops
<b>Non-Bearing Food Crops</b>	N/A	6-8 fl oz <sup>(8)</sup>		
<b>Pea, Dry Shelled</b> including Pea ( <i>Pisum</i> spp ) Field Pigeon	30 days	6-8 fl oz	1 qt by ground or 1% v/v (but not less than 1 pt /A) by air <sup>(5)</sup>	Do not apply more than 8 fl oz /A per application  Do not apply more than one (1) application per acre per season  Apply before bloom but not later than 30 days prior to harvest <sup>(10)</sup>  Refer to appropriate Table for reduced rate directions for the control of small annual grasses  The addition of AMS has shown improved grass control for difficult to control species including quackgrass rhizome Johnsongrass red rice wild oats volunteer cereals and volunteer corn

19/50

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Pea, Succulent</b> including Pea ( <i>Pisum</i> spp ) English Pea Garden Pea Green Pea Pigeon Pea	21 days	6-8 fl oz	1 qt by ground or 1% v/v (but not less than 1 pt /A) by air <sup>(5)</sup>	Do not apply more than 8 fl oz /A per application  Do not apply more than one (1) application per acre per season  Apply before bloom but not later than 21 days prior to harvest <sup>(10)</sup>  Refer to appropriate Table for reduced rate directions for the control of small annual grasses  The addition of AMS has shown improved grass control for difficult to control species including quackgrass rhizome Johnsongrass red rice wild oats volunteer cereals and volunteer corn
<b>Peach</b>	14 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application  Do not apply more than 32 fl oz /A (0.5 lb ai/A) per season  For repeat applications make on a minimum of a 14 day interval  Do not apply to Peach grown for root stock
<b>Peanut (including Perennial)</b>	40 days	6-16 fl oz	1 qt by ground or 1% v/v (but not less than 1 pt /A) by air <sup>(5)</sup>	The addition of AMS has shown improved grass control for difficult to control species including quackgrass rhizome Johnsongrass red rice wild oats volunteer cereals and volunteer corn
<b>Potato</b>	30 days	8-16 fl oz	1 qt by ground or 1% v/v (but not less than 1 pt /A) by air <sup>(5)</sup>	The addition of AMS has shown improved grass control for difficult to control species including quackgrass rhizome Johnsongrass red rice wild oats volunteer cereals and volunteer corn
<b>Radish</b>	15 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application  Do not apply more than 16 fl oz /A (0.25 lb ai) per acre in a season  For repeat applications make on a minimum of a 14 day interval

20/50

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Root Vegetable</b> (except Radish) including Chicory Ginseng Horseradish Turnip	30 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application For repeat applications make on a minimum of a 14 day interval
<b>Safflower</b>	70 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application For repeat applications make on a minimum of a 14 day interval
<b>Sesame</b>	14 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply during flowering Do not apply more than 8 fl oz /A in a single application For repeat applications make on a minimum of a 14 day interval
<b>Soybean</b>	60 days	6-16 fl oz	1 qt by ground or 1% v/v (but not less than 1 pt /A) by air <sup>(5)</sup>	Do not graze treated fields or feed treated forage or hay to livestock Refer to appropriate Table for reduced rate directions for the control of small annual grasses The addition of AMS has shown improved grass control for difficult to control species including quackgrass rhizome Johnsongrass red rice wild oats volunteer cereals and volunteer corn
<b>Strawberry</b>	4 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application For repeat applications make on a minimum of a 14 day interval
<b>Sugar Beet</b>	40 days	6-16 fl oz	1 qt by ground or 1% v/v (but not less than 1 pt /A) by air <sup>(5)</sup>	Refer to appropriate Table for reduced rate directions for the control of small annual grasses The addition of AMS has shown improved grass control for difficult to control species including quackgrass rhizome Johnsongrass red rice wild oats volunteer cereals and volunteer corn
<b>Sunflower</b>	70 days	6-16 fl oz	1 qt by ground or 1% v/v (but not less than 1 pt /A) by air <sup>(5)</sup>	The addition of AMS has shown improved grass control for difficult to control species including quackgrass rhizome Johnsongrass red rice wild oats volunteer cereals and volunteer corn

21/50

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Sweet Potato, Yam and Other Tuberous and Corm Vegetables</b> (except Potato) including Artichoke Chinese Jerusalem Cassava Bitter Sweet Ginger	30 days	6-16 fl oz	1% v/v in the finished spray volume	The addition of AMS has shown improved grass control for difficult to control species including quackgrass rhizome Johnsongrass red rice wild oats volunteer cereals and volunteer corn
<b>Tomato</b>	20 days	6-16 fl oz	1% v/v in the finished spray volume	For repeat applications make on a minimum of a 14 day interval
<b>Turnip Greens</b>	14 days	6-8 fl oz	1% v/v in the finished spray volume	Do not apply more than 8 fl oz /A in a single application  For repeat applications make on a minimum of a 14-day interval

N/A = Not Applicable

- <sup>1</sup> PS Clethodim is not recommended for use on vegetable crops being grown for seed production unless specific use directions are provided
- <sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality, and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the Addition of Adjuvant and Crop Oil Concentrate section for further information.
- <sup>3</sup> PS Clethodim may be applied to seedling or established alfalfa grown for seed, hay, silage, green crop, or direct grazing.
- <sup>4</sup> For weed control in established alfalfa and mint, the minimum use rate is 10 fl oz /A.
- <sup>5</sup> 1 to 2 qts /A of liquid fertilizer (10-34-0, 28%N or 32%N) or an equivalent amount (2.5 to 4.0 lbs /A) of spray grade ammonium sulfate (AMS) may be added to PS Clethodim applications in addition to the recommended rate of crop oil concentrate.
- <sup>6</sup> Do not apply PS Clethodim plus 2,4-DB as a tank mix to alfalfa unless the 60 day feeding, grazing, and harvesting restriction on the 2,4-DB label can be observed.
- <sup>7</sup> For ground applications to garlic or shallot, do not exceed 8 fl oz /A in a single application. For air applications to onion, garlic, or shallot, do not exceed 8 fl oz /A in a single application. For garlic and shallot, do not exceed 2 applications per season. In CA, for air applications to onion, do not exceed 2 applications per season.
- <sup>8</sup> If PS Clethodim is applied as a spot treatment to onion, garlic, shallot, or non-bearing food crops, care should be taken to not exceed the maximum rate allowed on a per acre basis or crop injury may occur.
- <sup>9</sup> In California, do not apply PS Clethodim to onion, garlic, or shallot until the crop has at least 1/3 full leaves. In California, 14 day spray intervals are recommended between the application of PS Clethodim and liquid nitrogen or other herbicide applications. Injury to crop may occur when shorter intervals are observed.

<sup>10</sup> Applications of PS Clethodim to peas during the bloom period could result in severe crop injury including loss of yield and delayed maturity

**IMPORTANT**

Plant tolerance to PS Clethodim at labeled rates has been found to be acceptable for the indicated genera and species listed below. Due to variability within species, crop growth stage, environmental conditions, and application techniques, it is recommended that the user determine if the herbicide can be used safely on a few plants prior to widespread application. Neither the seller nor the manufacturer of PS Clethodim have investigated the safety factor to plants not listed on the label.

**NON-BEARING FOOD CROPS**

**PS CLETHODIM SHOULD NOT BE APPLIED TO NON-BEARING FRUIT OR NUT CROPS WHICH ARE GROWN FOR ROOT STOCK**

Crop injury to non-bearing fruit and nut crops can occur if PS Clethodim is improperly applied. PS Clethodim should not be applied directly over the top of these plant types. Instead, spray should be directed at the base of the plant where grassy weeds are growing near the ground.

Non-bearing fruit and nut crops are plants which will not bear fruit or nuts for at least one year following PS Clethodim application.

COMMON NAME	SCIENTIFIC NAME
Apples	<i>Malus</i> spp
Berries	<i>Vaccinium</i> spp <i>Rubus</i> spp
Cherry Sweet	<i>Prunus avium</i>
Citrus Fruits	<i>Citrus</i> spp
Grapes	<i>Vitis</i> spp
Olives	<i>Olea</i> spp
Peach	<i>Prunus persica</i>
Pears	<i>Pyrus communis</i>
Prunes	<i>Prunus</i> spp
Stone Fruits	<i>Prunus</i> spp
Strawberries	<i>Fragaria</i> spp
<b>Tree Nuts</b>	
Almond	<i>Prunus triloba</i>
Filbert	<i>Corylus maxima</i>
Pecan	<i>Carya illinoensis</i>
Pistachio	<i>Pistacia vera</i>
Walnut	<i>Juglans</i> spp

**CONIFER TREES**

PS Clethodim can be used to control labeled grasses in Christmas tree farms, conifer nurseries, and conifer plantations (but not in forests).

COMMON NAME	SCIENTIFIC NAME
Arborvitae American	<i>Thuja occidentalis</i>
Cedars	<i>Cedrus</i> spp
Cypress	<i>Taxodium</i> spp
Fir Douglas	<i>Pseudotsuga menziesii</i>
Firs	<i>Abies</i> spp
Hemlock Canadian/Eastern	<i>Tsuga Canadensis</i>

COMMON NAME	SCIENTIFIC NAME
Hemlock Western	<i>Tsuga heterophylla</i>
Pines	<i>Pinus spp</i>
Spruces	<i>Picea spp</i>
Yew	<i>Taxus spp</i>

**NON-CROP OR NON-PLANTED AREAS**

The following areas are considered non-crop or non-planted areas rights-of-way including railroads highways roads dividers medians pipelines public utility lines pumping stations transformer stations and substations Around airports electric utilities commercial buildings manufacturing plants storage yards rail yards fence lines parkways and post-harvest croplands Also beneath greenhouse benches and around golf courses

**ANNUAL GRASSES  
(EXCEPT FOR IN ESTABLISHED ALFALFA AND MINT)**

- Apply only to actively growing grasses at recommended weed heights
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height
- Do not apply more than 8 fl oz /A of PS Clethodim **per application** to the following crops asparagus carrot cranberry cucurbits flax fruiting vegetables (except tomato) garden beets green onion head and stem brassica vegetables herbs hops leafy brassica greens leafy greens non-bearing food crops root vegetables safflower sesame and strawberry Do not apply more than 6 fl oz /A of PS Clethodim per application to canola or mustard seed

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT* (inches)	RATE FL OZ / ACRE	HIGH RATE <sup>(1)</sup>
Barnyardgrass	<i>Echinochloa crus-galli</i>	2 to 8	6	8
Broadleaf Signalgrass	<i>Brachiaria platyphlla</i>	2 to 6	6	8
Brome				
California	<i>Bromus carinatus</i>	2 to 6	6	8
Cheat	<i>Bromus secalinus</i>	2 to 6	6	8
Downy	<i>Bromus tectorum</i>	2 to 6	6	8
Ripgut	<i>Bromus diandrus</i>	2 to 6	6	8
Canarygrass	<i>Phalaris canariensis</i>	1 to 4	6	8
Crabgrass				
Hairy	<i>Digitaria adscendens</i>	2 to 6**	6	8
Large	<i>Digitaria sanguinalis</i>	2 to 6**	6	8
Smooth	<i>Digitaria ischaemum</i>	2 to 6**	6	8
Southern	<i>Digitaria ciliaris</i>	2 to 6**	6	8
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	2 to 6**	6	8
Fall Panicum	<i>Panicum dichotomiflorum</i>	2 to 8	6	8
Field Sandbur	<i>Cenchrus incertus</i>	2 to 6	6	8
Foxtail				
Giant	<i>Setaria feberi</i>	2 to 12	6	8
Green	<i>Setaria viridis</i>	2 to 8	6	8
Yellow	<i>Setaria glauca</i>	2 to 8	6	8
Goosegrass	<i>Eleusine indica</i>	2 to 6**	6	8
Itchgrass	<i>Rottboellia cochinchinensis</i>	2 to 6	6	8
Junglerice	<i>Echinochloa colona</i>	2 to 6	6	8
Lovegrass (Stinkgrass)	<i>Eragrostis cilianensis</i>	2 to 6	6	8
Rabbitsfootgrass	<i>Polypogon monspeliensis</i>	1 to 4	6	8

24/50

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT* (inches)	RATE FL OZ / ACRE	HIGH RATE <sup>(1)</sup>
Red Rice	<i>Oryza sativa</i>	1 to 3	6	8
Ryegrass				
Hardy	<i>Lolium remotum</i>	2 to 6	6	8
Italian	<i>Lolium multiflorum</i>	2 to 6	6	8
Seedling Johnsongrass	<i>Sorghum halepense</i>	4 to 10	6	8
Shattercane	<i>Sorghum bicolor</i>	6 to 18	6	8
Southwestern Cupgrass	<i>Eriochloa gracilis</i>	2 to 6	6	8
Sprangle top				
Amazon	<i>Leptochloa panicoides</i>	2 to 6	6	8
Bearded	<i>Leptochloa fascicularis</i>	2 to 6	6	8
Mexican	<i>Leptochloa uninervia</i>	2 to 6	6	8
Red	<i>Leptochloa filiformis</i>	2 to 6	6	8
Texas Panicum	<i>Panicum texanum</i>	2 to 6	6	8
Volunteer Cereals <sup>(2)</sup>				
Barley	<i>Hordeum vulgare</i>	2 to 6	6	8
Oats	<i>Avena sativa</i>	2 to 6	6	8
Rye	<i>Secale cereal</i>	2 to 6	6	8
Wheat	<i>Triticum aestivum</i>	2 to 6	6	8
Volunteer Corn <sup>(3)</sup>	<i>Zea mays</i>	4 to 12	4	6
Volunteer Corn <sup>(3)</sup>	<i>Zea mays</i>	12 to 24	6	8
Volunteer Corn (S R) <sup>(4)</sup>	<i>Zea mays</i>	4 to 12	8 (suppression only)	
Volunteer Grain Sorghum	<i>Sorghum bicolor</i>	8 to 12	6	8
Wild Oats	<i>Avena fatua</i>	2 to 6	6	8
Wild Proso Millet	<i>Panicum miliaceum</i>	2 to 10	6	8
Witchgrass	<i>Panicum capillare</i>	2 to 8	6	8
Woolly Cupgrass	<i>Eriochloa villosa</i>	2 to 8	6	8

\* Generally occurs between 3-leaf stage and tillering


\*\* Length of lateral growth

(1) Rates higher than 8 fl oz /A may be applied in certain geographic areas cropping situations or environmental conditions where experience has shown that higher rates are needed for satisfactory control of annual grasses. In these situations rates from 8 to 16 fl oz /A may be applied. Do not apply more than 8 fl oz /A of PS Clethodim per application to the following crops: asparagus, carrot, cranberry, cucurbits, flax, fruiting vegetables (except tomato), garden beet, green onion, head and stem brassica vegetables, herbs, hops, leaf petioles, leafy brassica greens, leafy greens, non-bearing food crops, root vegetables, safflower, sesame and strawberry.

(2) When a cereal grain crop (such as wheat) is interseeded for crop establishment or is planted as wind breaks to aid crop establishment, the minimum PS Clethodim use rate for control is 8 fl oz /A.

(3) Includes Roundup Ready®, Liberty Link® and IMI-CORN® volunteer corn.

(4) Sethoxydim resistant volunteer corn.







26/50

ANNUAL BLUEGRASS CONTROL WITH PS CLETHODIM			
GRASS SPECIES	WEED STAGE	RATE FL OZ / ACRE	HIGH RATE
Annual Bluegrass ( <i>Poa annua</i> )	to 4-leaf	6*	16
<ul style="list-style-type: none"> <li>Apply under favorable soil moisture and humidity which exists within a few days after rainfall or within 7 days after irrigation Grass needs to be actively growing at time of application(s)</li> <li>Apply at weed stage indicated on the label as reduced control can be expected with more mature annual bluegrass</li> <li>Use the high rate under heavy grass pressure and/or when annual bluegrass is more mature</li> <li>Always add a crop oil concentrate at 1 qt /A by ground to the finished spray volume</li> </ul> <p>*Use a minimum of 10 fl oz /A to control annual bluegrass in seedling and established alfalfa and mint</p>			

FOR USE IN ROUNDUP READY FIELD CORN (BURNDOWN)		
GRASS SPECIES	WEED SIZE (inches)	Rate when applied alone or with glyphosate
Field Corn	Up to 12	3 fl oz /A
<ul style="list-style-type: none"> <li>For control of existing stand of Roundup Ready field corn or volunteer Roundup Ready field corn prior to replanting corn</li> <li>Care must be taken to avoid in-field boom (spray) overlaps or excessive crop injury may occur</li> <li>Replant no sooner than 6 days after application</li> <li>Adjuvant recommendations NIS at 0.25% v/v plus AMS at 2.5 to 4 lbs /A</li> <li>Do not use COC or MSO with PS Clethodim in this use pattern</li> </ul>		

**DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES IN CANOLA, DRY SHELLLED BEAN & PEA (INCLUDING SOYBEAN), EDIBLE PODDED LEGUME VEGETABLES, FLAX, MUSTARD SEED, SUCCULENT BEAN & PEA AND SUGAR BEET (REDUCED RATE DIRECTIONS NOT FOR USE IN CALIFORNIA)**

- Apply only to actively growing grasses at recommended weed heights
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment
- Regrowth by tillering may occur if application is made when plants are stressed by lack of moisture excessive moisture low or high temperatures and/or under very low humidity

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT (inches)	RATE FL OZ /ACRE <sup>(1)</sup>
Barnyardgrass	<i>Echinochloa crus-galli</i>	1 to 4	4
Broadleaf Signalgrass	<i>Brachiaria platyphlla</i>	1 to 4	5
Crabgrass			
Large	<i>Digitaria sanguinalis</i>	1 to 3*	4
Large	<i>Digitaria sanguinalis</i>	1 to 4*	5
Smooth	<i>Digitaria ischaemum</i>	1 to 3*	4
Smooth	<i>Digitaria ischaemum</i>	1 to 4*	5
Southern	<i>Digitaria ciliaris</i>	1 to 4*	5
Fall Panicum	<i>Panicum dichotomiflorum</i>	1 to 4	4
Foxtail			
Giant	<i>Setaria faberi</i>	1 to 4	4

27/50

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT (inches)	RATE FL OZ /ACRE <sup>(1)</sup>
Green	<i>Setaria viridis</i>	1 to 4	4
Millet	<i>Setaria italic</i>	1 to 4	5
Yellow	<i>Setaria glauca</i>	1 to 4	4
Seedling Johnsongrass	<i>Sorghum halepense</i>	1 to 6	5
Shattercane	<i>Sorghum bicolor</i>	4 to 10	4
Texas Panicum	<i>Panicum texanum</i>	1 to 4	5
Volunteer Cereals			
Barley	<i>Hordeum vulgare</i>	1 to 4	5
Oats	<i>Avena sativa</i>	1 to 4	5
Wheat	<i>Triticum aestivum</i>	1 to 4	5
Volunteer Corn**	<i>Zea mays</i>	4 to 12	4
Wild Proso Millet	<i>Panicum miliaceum</i>	1 to 6	4
Wild Oats	<i>Avena fatua</i>	1 to 4	5

\* Length of lateral growth

\*\* Not S R Corn

<sup>(1)</sup>Always add a crop oil concentrate at 1 qt /A by ground application to the finished spray volume

**PERENNIAL GRASSES**

- Apply only to actively growing grasses at recommended weed heights
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height
- Do not apply more than 8 fl oz /A of PS Clethodim per application to the following crops asparagus carrot cranberry cucurbits flax fruiting vegetables (except tomato) garden beet green onion head & stem brassica vegetables herbs hops leaf petioles leafy brassica greens leafy greens non-bearing food crops root vegetables safflower sesame and strawberry
- Do not apply more than 6 fl oz /A of PS Clethodim per application to canola or mustard seed

GRASS SPECIES	WEED HEIGHT (inches)	RATE FL OZ /ACRE	HIGH RATE
<b>Bermudagrass (<i>Cynodon dactylon</i>)</b>			
First Application	3 (or up to 6 runners)	8	16
Repeat Application(s) (if regrowth occurs)	3 (or up to 6 runners)	8	16
<b>Fescue Tall (<i>Festuca arundinacea</i>)</b>			
First Application	4 to 8	8	16
Repeat Application(s) (if regrowth occurs)	4 to 8	8	16
<b>Foxtail Barley (<i>Hordeum jubatum</i>)</b>			
First Application	2 to 6	8	16
Repeat Application(s) (if regrowth occurs)	2 to 6	8	16
<b>Orchardgrass (<i>Dactylis glomerata</i>)</b>			
First Application	4 to 8	8	16
Repeat Application(s) (if regrowth occurs)	4 to 8	8	16
<b>Quackgrass* (<i>Elytrigia repens</i>)</b>			
First Application	4 to 12	8	16
Repeat Application(s) (if regrowth occurs)	4 to 12	8	16
<b>Rhizome Johnsongrass (<i>Sorghum halepense</i>)</b>			

GRASS SPECIES	WEED HEIGHT (inches)	RATE FL OZ /ACRE	HIGH RATE
First Application	12 to 24	8	16
Repeat Application(s) (if regrowth occurs)	6 to 18	6	8
<b>Wirestem Muhly (<i>Muhlenbergia frondosa</i>)</b>			
First Application	4 to 8	8	16
Repeat Application(s) (if regrowth occurs)	4 to 8	8	16
<b>Perennial Bluegrass*</b>			
Roughstalk ( <i>Poa trivialis</i> )			
Kentucky ( <i>Poa prantensis</i> )			
First Application	2 to 4	8	16
Repeat Application(s) (if regrowth occurs)	2 to 4	8	16
<b>Bentgrass* (<i>Agrostis</i> spp )</b>			
First Application	2 to 4	-	16
Repeat Application(s) (if regrowth occurs)	2 to 4	-	16

\*Control of quackgrass perennial bluegrass and bentgrass with PS Clethodim may be enhanced by adding AMS at 2.5 to 4.0 lbs /A

### TANK MIXES

#### INFORMATION

The labels for each of the herbicides recommended for tank mixing with PS Clethodim are unique to the characteristics of those products and contain restrictions and limitations that may be more restrictive than the PS Clethodim label in certain considerations. Those concerns may include but are not limited to:

- 1 Geographic restrictions – all products are not registered for use in all areas and rates may vary from one region of labeled use to another
- 2 Crop rotation restrictions
- 3 Applicator certification requirements
- 4 Worker safety rules (e.g. protective clothing, reentry time, posting)
- 5 Soil type or soil characteristics (e.g. pH, OM)
- 6 Maximum dosage or number of applications per season
- 7 Rain free period required or
- 8 Application timing (e.g. pre-harvest interval)
- 9 Do not exceed the total season rates

#### THE MOST RESTRICTIVE LABELING OF ANY PRODUCT USED IN A TANK MIX MUST BE FOLLOWED

#### TANK MIX APPLICATION OF PS CLETHODIM AND BROADLEAF HERBICIDES FOR CONTROL OF GRASSES AND BROADLEAF WEEDS

- Apply only to actively growing grass and broadleaf weeds at recommended height or growth stage listed on each label
- Apply when the first grass or broadleaf weed species in a mixed population reaches the recommended height or growth stage for treatment
- Apply under favorable soil moisture and humidity that exist a few days after rainfall or within seven days after irrigation
- Always add the appropriate adjuvant to the spray mix at the rate recommended for each specific tank mix combination
- Tank mix applications may sometimes result in reduced grass control and possible increases in crop injury as compared to either product used alone. If regrowth occurs or an additional flush of new grass emerges, make a second application of PS Clethodim as specified in the respective size and rate tables.

29/58

- Do not tank mix PS Clethodim when broadleaf weeds are tall and/or dense enough to prevent proper grass coverage

**MIXING INSTRUCTIONS**

- 1 Fill clean spray tank 1/2 to 2/3 of desired level with clean water
- 2 While agitating add the correct amount of PS Clethodim Agitation should create a rippling or rolling action on the water surface
- 3 If tank mixing PS Clethodim with other labeled herbicides add water soluble bags first followed by dry formulations flowables emulsifiable concentrates and then solutions Prepare no more spray mixture than is required for the immediate spray operation
- 4 Add any required adjuvants (crop oil concentrate non-ionic surfactant and/or nitrogen solution)
- 5 Fill spray tank to desired level with water **Agitation should continue until all spray solution has been applied**

Failure to agitate the spray solution may result in improper mixing of the herbicides and unsatisfactory weed control Mixing and compatibility qualities should be verified by a jar test

**INFORMATION ON ANTAGONISM**

Tank mixes of PS Clethodim with postemergence broadleaf herbicides have shown some reduction or failure to control certain grass species which would have otherwise been controlled when PS Clethodim is applied alone Activity of the postemergence broadleaf herbicide in the tank mix is not affected

**ALFALFA**

**Table 1 PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR ALFALFA**  
(Refer to tables above for specific grasses and growth stages)

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND	AIR
	PS Clethodim + 2 4-DB <sup>(4)</sup>	10 to 16 fl oz + Refer to 2 4-DB label	10 to 16 fl oz + Refer to 2 4-DB label	1%
PS Clethodim + PURSUIT DG <sup>(5)</sup> or PURSUIT <sup>(5)</sup>	10 to 16 fl oz + 1 08 to 2 16 oz or 3 to 6 fl oz	-	1%	1%
PS Clethodim + BUCTRIL® 2L <sup>(6)</sup> or BUCTRIL GEL <sup>(6 7)</sup>	10 to 16 fl oz + 1 0 to 1 5 pts or 0 5 to 0 75 pt	-	0 5%	0 5%

<sup>(1)</sup> If grass regrowth occurs or an additional flush of new grass emerges make a second application of PS Clethodim alone (without a tank mix herbicide) according to the appropriate size and rate recommendations

<sup>(2)</sup> Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage Tank mixing is not recommended in these situations

<sup>(3)</sup> Always use a crop oil concentrate at the listed rate (but not less than 1 pt /A) in the finished spray volume

<sup>(4)</sup> PS Clethodim plus 2 4-DB may increase the severity of crop injury when tank mixed Alfalfa plants will generally outgrow this temporary crop injury within a few weeks

<sup>(5)</sup> Before using this tank mix read and understand the PURSUIT or PURSUIT DG labels for geographical restrictions and restrictions regarding alfalfa growth stage and type Failure to do so can result in crop injury to alfalfa Do not feed graze or harvest alfalfa for 30 days following an application of PURSUIT to alfalfa

<sup>(6)</sup> In the states of Washington Oregon Idaho Montana Wyoming Colorado Utah Nevada and the western

halves of North Dakota South Dakota Nebraska and Kansas The PS Clethodim plus BUCTRIL or BUCTRIL GEL tank mix must be applied in the fall or spring to seedling alfalfa when the majority of the field has a minimum of 2 trifoliates Unacceptable crop injury may occur to alfalfa seedlings less than the 2 trifoliolate leaf stage PS Clethodim plus BUCTRIL or BUCTRIL GEL applications made when temperatures are expected to exceed 80°F at (and 3 days following) application can result in unacceptable crop injury In the states not listed above apply in the fall or spring to seedling alfalfa when the majority of the field has a minimum of 4 trifoliolate leaves When the alfalfa stand is uneven and conditions favor leaf burn unacceptable crop injury may occur to alfalfa in the 2 trifoliolate or smaller stage of growth PS Clethodim plus BUCTRIL or BUCTRIL GEL applications made when temperatures are expected to exceed 70°F at (and 3 days following) application can result in unacceptable crop injury Crop leaf burn can occur following PS Clethodim plus BUCTRIL or BUCTRIL GEL application Warm humid conditions may enhance leaf burn New crop growth will not be affected

(7) Do not apply when alfalfa is under moisture temperature insect or disease stress or has been stressed by other pesticide carryover or application

**CANOLA**

**Table 2 REDUCED RATE PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR CANOLA (Refer to the tables above for specific grasses and growth stages )**

PRODUCT	APPLICATION RATES/ACRE		AMMONIUM SULFATE	
	ANNUAL GRASSES <sup>(1)</sup>	PERENNIAL GRASSES	GROUND	AIR
	PS Clethodim <sup>(2)</sup> + LIBERTY <sup>(3)</sup>	4 to 5 fl oz + 34 fl oz	-	3 0 lbs

(1) Annual grasses and sizes controlled with these tank mixtures are those that are identified in the DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES table

(2) Do not apply PS Clethodim tank mix during or after bolting or flowering or crop injury may occur

(3) For use only on LibertyLink® canola

**COTTON**

**Table 3 PS CLETHODIM TANK MIXED WITH COBRA<sup>®</sup> AND MSMA APPLIED POST DIRECTED TO COTTON**

PRODUCT <sup>(1)</sup>	APPLICATION RATES/ACRE <sup>(2)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> V/V	COMMENTS
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND	
PS Clethodim <sup>(4)</sup> + COBRA + MSMA (4 0 lbs /gal ) or MSMA (6 6 lbs /gal )	6 to 8 fl oz	8 to 16 fl oz	1%	Reduce broadcast rate in proportion to the band area actually treated
	See COBRA label for rates to control broadleaf weeds and height limitations for cotton Refer to the PS Clethodim label for weed height and species controlled			
	See MSMA label for rates to control broadleaf weeds and height limitations for cotton Refer to the PS Clethodim label for weed height and species controlled			

(1) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage Tank mixing is not recommended in these situations

(2) If grass regrowth occurs or an additional flush of new grass emerges make a second application of PS Clethodim alone (without a tank mix herbicide) according to the appropriate size and rate recommendations

(3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt /A) in the finished spray volume

(4) If at the time of application grass height is so tall that post-directed applications cannot get good coverage over the top of the grassy weeds then poor control may result and a second (non-post directed) application of



**DRY SHELLED AND SUCCULENT BEANS**

**Table 6 PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR DRY SHELLED AND SUCCULENT BEANS**

(Refer to the tables above for specific grasses and growth stages )

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup>	
	ANNUAL GRASSES	PERENNIAL GRASSES	(V/V)	
			GROUND	AIR
PS Clethodim + BASAGRAN®	8 to 10 fl oz + 1 0 to 2 0 pts	10 to 16 fl oz + 1 to 2 pts	1%	1%

<sup>(1)</sup> If grass regrowth occurs or an additional flush of new grass emerges make a second application of PS Clethodim alone (without a tank mix herbicide) according to the appropriate size and rate recommendations

<sup>(2)</sup> Broadleaf weed control may be reduced when grass populations are tall enough or dense enough to intercept the spray pattern and prevent them from receiving complete coverage Tank mixing is not recommended in these situations

<sup>(3)</sup> Always use a crop oil concentrate at the listed rate (but not less than 1 pt /A) in the finished spray volume

**FLAX**

**Table 7 REDUCED RATE PS CLETHODIM TANK MIXED WITH BROADLEAF HERBICIDES FOR FLAX**

(Refer to the tables above for specific grasses and growth stages)

PRODUCT	APPLICATION RATES/ACRE		CROP OIL CONCENTRATE	
	ANNUAL GRASSES <sup>(1)</sup>	PERENNIAL GRASSES	GROUND	AIR
PS Clethodim + BRONATE ADVANCED™ <sup>(2 3)</sup>	4 to 5 fl oz + 11 4 fl oz	-	1 pt	1 pt
PS Clethodim + BRONATE® <sup>(2 3)</sup>	4 to 5 fl oz + 0 9 pt	-	1 pt	1 pt
PS Clethodim + BUCTRIL <sup>(2 3)</sup>	4 to 5 fl oz + 1 0 pt	-	1 pt	1 pt
PS Clethodim + RHONOX® <sup>(2 3)</sup>	4 to 5 fl oz + 0 25 to 0 5 pt	-	1 pt	1 pt

<sup>(1)</sup> Annual grasses and sizes controlled with these tank mixtures are those that are identified in the DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES table

<sup>(2)</sup> Do not apply PS Clethodim tank mix during or after the bud stage or to ornamental flax or crop injury may occur

<sup>(3)</sup> Do not apply tank mixes if temperatures are expected to exceed 85°F at (or 3 days following) application or crop injury may occur



**SOYBEAN**

**Table 8 PS CLETHODIM TANK MIXES<sup>(3)</sup> TO CONTROL ANNUAL GRASSES WHEN USED AS A BURNDOWN IN NO-TILL SOYBEAN**

PRODUCT	PRODUCT RATE/ACRE <sup>(1)</sup>	GRASS HEIGHT (inches)	CROP OIL CONCENTRATE /ACRE <sup>(2)</sup>	28%N OR 32%N QTS /A OR 2 5 TO 4 0 LBS AMS
PS Clethodim + 2 4-D ester <sup>(3)</sup>	3 fl oz	Foxtail 1 to 3 Fall Panicum 1 to 3	1 qt	1 to 2 qts or 2 5 to 4 0 lbs AMS
	4 fl oz	Foxtail 1 to 4 Fall Panicum 1 to 4	1 qt	1 to 2 qts or 2 5 to 4 0 lbs AMS
	6 to 8 fl oz + 0 5 lb ai	(See Grass Chart for grasses claimed )	1 qt	1 to 2 qts or 2 5 to 4 0 lbs AMS

\* 2 4-D ester should not be used where drift sensitive crops may be grown

<sup>(1)</sup> If regrowth occurs or an additional flush of new grass emerges make a second application of PS Clethodim according to the appropriate size and rate recommendations

<sup>(2)</sup> Always use a crop oil concentrate at the listed rate in the finished spray volume

<sup>(3)</sup> The following products can be tank mixed with PS Clethodim plus 2 4-D ester VALOR<sup>®</sup> AUTHORITY<sup>®</sup> BROADLEAF CANOPY XL<sup>®</sup> DUAL<sup>®</sup> 8 E DUAL II<sup>®</sup> DUAL MAGNUM<sup>®</sup> PROWL<sup>®</sup> SENCOR<sup>®</sup> SENCOR plus the DUAL products and TURBO<sup>®</sup>

**Table 9 PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEAN (Refer to the tables above for specific grasses and growth stages )**

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND	AIR
PS Clethodim + COBRA	6 to 8 fl oz + 12 5 fl oz	8 to 16 fl oz + 12 5 fl oz	0 5 to 1%	1%
PS Clethodim + BASAGRAN 4 SL	8 to 10 fl oz + 1 to 2 pts	10 to 16 fl oz + 1 to 2 pts	1%	1%
PS Clethodim + Glyphosate (For use on Roundup Ready soybean only)	6 to 8 fl oz + 0 75 to 3 0 lb ai	8 to 16 fl oz + 0 75 to 3 0 lb ai	0 5 to 1% <sup>(4)</sup>	1% <sup>(4)</sup>
PS Clethodim + ULTRA BLAZER <sup>®</sup>	6 to 8 fl oz + 1 to 1 5 pts	6 to 8 fl oz + 1 to 1 5 pts	0 5% to 1%	1%
PS Clethodim + FLEXSTAR <sup>®</sup> HL <sup>(6)</sup>	6 to 8 fl oz + Refer to the FLEXSTAR HL label for specific application rates	8 to 16 fl oz + Refer to the FLEXSTAR HL label for specific application rates	1%	1%
PS Clethodim + CLASSIC <sup>®</sup> 25 DG	8 to 10 fl oz + 0 5 to 0 75 oz	10 to 16 fl oz + 1 44 oz	1%	1%

34/50

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND	AIR
PS Clethodim <sup>(4)</sup> + PURSUIT® 70 DG	6 to 8 fl oz + 1.44 oz	8 to 16 fl oz + 1.44 oz	1%	1%
PS Clethodim <sup>(5)</sup> + COBRA + CLASSIC 25 DG	8 to 10 fl oz + 6 to 8 fl oz + 0.5 to 0.75 oz	-	0.5%	1%
PS Clethodim <sup>(5)</sup> + COBRA + BASAGRAN 4 SL	8 to 10 fl oz + 6 to 10 fl oz + 1 to 1.5 pts	-	0.5%	1%
PS Clethodim <sup>(5)</sup> + COBRA + PURSUIT 70 DG	8 to 10 fl oz + 6 to 10 fl oz + 1.44 oz	-	0.5%	1%
PS Clethodim <sup>(5)</sup> + STORM®	8 to 10 fl oz + 1.5 pts	-	0.5%	1%
PS Clethodim <sup>(5)</sup> + RESOURCE® + PURSUIT 70 DG	8 to 10 fl oz + 4 fl oz + 1.44 oz	-	1%	1%
PS Clethodim <sup>(5)</sup> + RESOURCE + BASAGRAN	8 to 10 fl oz + 4 fl oz + 1 pt	-	1%	1%
PS Clethodim <sup>(5)</sup> + RESOURCE + CLASSIC	8 to 10 fl oz + 4 fl oz + 0.5 oz	-	1%	1%
PS Clethodim <sup>(5)</sup> + COBRA + RESOURCE	6 to 8 fl oz + 6 fl oz + 4 fl oz	-	0.5%	1%
PS Clethodim <sup>(5)</sup> + FIRSTRATE®	6 to 8 fl oz + 0.3 oz	8 to 16 fl oz + 0.3 oz	1%	-
PS Clethodim <sup>(5)</sup> + COBRA + FIRSTRATE	6 to 8 fl oz + 6 to 8 fl oz + 0.3 oz	8 to 16 fl oz + 6 to 8 fl oz + 0.3 oz	1%	-

35/50

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND	AIR
PS Clethodim <sup>(5)</sup> + RAPTOR® (1 AS)	6 to 8 fl oz + 4 to 5 fl oz	-	1%	-
PS Clethodim <sup>(5)</sup> + COBRA + RAPTOR (1 AS)	6 to 8 fl oz + 6 to 8 fl oz + 4 to 5 fl oz	-	1%	-
PS Clethodim <sup>(5)</sup> + SYNCHRONY® STS™	6 to 8 fl oz <sup>(7)</sup> + 0.5 oz /A	-	1 qt	-
PS Clethodim <sup>(5)</sup> + COBRA Herbicide + SYNCHRONY STS™	6 to 8 fl oz <sup>(7)</sup> + 4 to 8 fl oz + 0.5 oz	-	1 pt	-
PS Clethodim <sup>(5)</sup> + RESOURCE	6 to 8 fl oz + 4 to 12 fl oz	-	1 qt	-
PS Clethodim <sup>(5)</sup> + FRONTROW™	8 to 10 fl oz + Refer to FRONTROW label for use rates	-	1%	-
PS Clethodim + FIRSTRATE + FLEXSTAR HL <sup>(5)</sup>	6 to 8 fl oz + 0.3 oz + Refer to the FLEXSTAR HL label for specific application rates	8 to 16 fl oz + 0.3 oz + Refer to the FLEXSTAR HL label for specific application rates	1%	-

<sup>(1)</sup> If grass regrowth occurs or an additional flush of new grass emerges make a second application of PS Clethodim alone (without a tank mix herbicide) according to the appropriate size and rate recommendations

<sup>(2)</sup> Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage Tank mixing is not recommended in these situations

<sup>(3)</sup> Always use a crop oil concentrate at the listed rate (but not less than 1 pt /A) in the finished spray volume

<sup>(4)</sup> The addition of 2.5 lb of ammonium sulfate is required when PS Clethodim is tank mixed with glyphosate If the glyphosate formulation has a stand alone built in adjuvant add 0.125% v/v non-ionic surfactant in place of crop oil concentrate If the glyphosate formulation does not have a built in adjuvant add 0.5 to 1% crop oil concentrate for ground application and 1% v/v for aerial application

<sup>(5)</sup> The addition of 1 to 2 qts /A of liquid fertilizer (10-34-0 28%N or 32%N) is recommended when PS Clethodim is tank mixed with PURSUIT RESOURCE STORM FIRSTRATE SYNCHRONY RAPTOR FRONTROW COBRA plus CLASSIC COBRA plus BASAGRAN COBRA plus PURSUIT COBRA plus FIRSTRATE COBRA plus SYNCHRONY and COBRA plus RAPTOR An equivalent amount (2.5 to 4.0 lbs /A) of spray grade ammonium sulfate (AMS) may be added in place of liquid fertilizer Fertilizer adjuvants are to be added in addition to the crop oil concentrate

<sup>(6)</sup> Refer to FLEXSTAR HL label for geographic and rotational restrictions

<sup>(7)</sup> Annual grasses and sizes controlled with these tank mixes are those that are identified in the DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES table

**Table 10 REDUCED RATE PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEAN (Refer to table for reduced rate use in canola, dry shelled bean & pea, edible podded legume vegetables, flax, mustard seed, soybean, succulent bean & pea and sugar beet for small annual grasses for specific grasses and growth stages)**

PRODUCT	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3,4)</sup> (V/V)	
	ANNUAL GRASSES <sup>(2)</sup>	PERENNIAL GRASSES	GROUND	AIR
PS Clethodim + FIRSTRATE	4 to 8 fl oz + 0.3 oz	-	1%	1%
PS Clethodim + PURSUIT 70 DG	4 to 6 fl oz + 1.44 oz	-	1%	1%

<sup>(1)</sup> If grass regrowth occurs or an additional flush of new grass emerges make a second application of PS Clethodim alone (without a tank mix herbicide) according to the appropriate size and rate recommendations

<sup>(2)</sup> Annual grasses and sizes controlled with these tank mixtures are those that are identified in the DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES table

<sup>(3)</sup> Always use a crop oil concentrate at the listed rate (but not less than 1 pt /A) in the finished spray volume

<sup>(4)</sup> The addition of 1 to 2 qts /A of liquid fertilizer (10-34-0 28%N or 32%N) is required when PS Clethodim is tank mixed at reduced rates. An equivalent amount (2.5 to 4.0 lbs /A) of spray grade ammonium sulfate (AMS) may be added in place of liquid fertilizer. Fertilizer adjuvants are to be added in addition to the crop oil concentrate

**PEANUT (INCLUDING PERENNIAL)**

**Table 11 PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR PEANUT (INCLUDING PERENNIAL)**

(Refer to the tables above for specific grasses and growth stages )

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
	ANNUAL GRASSES <sup>(2)</sup>	PERENNIAL GRASSES	GROUND	AIR
PS Clethodim + BASAGRAN	8 to 10 fl oz + 1.0 to 2.0 pts	-	1%	1%
PS Clethodim + ULTRA BLAZER	8 to 10 fl oz + 0.5 to 1.5 pts	-	1%	1%
PS Clethodim + STORM	8 to 10 fl oz + 1.5 pts	-	1%	1%

<sup>(1)</sup> If grass regrowth occurs or an additional flush of new grass emerges make a second application of PS Clethodim alone (without a tank mix herbicide) according to the appropriate size and rate recommendations

<sup>(2)</sup> Broadleaf weed control may be reduced when grass populations are tall or dense enough to interrupt the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations

<sup>(3)</sup> Always use a crop oil concentrate at the listed rate (but not less than 1 pt /A) in the finished spray volume

37/50

GRASS SUPPRESSION FOR HARVEST EFFICIENCY IN PEANUT (INCLUDING PERENNIAL) WITH PS CLETHODIM			
GRASS SPECIES	WEED STAGE	RATE FL OZ /ACRE	HIGH RATE
Annual and perennial grasses that exceed height claimed for control on height charts " <b>ANNUAL GRASSES</b> " & " <b>PERENNIAL GRASSES</b> "	Up to and including grasses in the seed head stage	16	32
Do not apply as part of a tank mix when apply PS Clethodim for grass suppression			
Add a crop oil concentrate at 1 qt /A by ground to the finished spray volume			

**SUGAR BEET**

**Table 12 PS CLETHODIM TANK MIXED WITH STINGER® APPLIED TO SUGAR BEET (Refer to the tables above for specific grasses and growth stages)**

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND	AIR
PS Clethodim + STINGER	6 to 8 fl oz	8 to 16 fl oz	1%	1%
See STINGER label for rates				

- <sup>(1)</sup> If grass regrowth occurs or an additional flush of new grass emerges make a second application of PS Clethodim alone (without a tank mix herbicide) according to the appropriate size and rate recommendations
- <sup>(2)</sup> Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage Tank mixing is not recommended in these situations
- <sup>(3)</sup> Always use a crop oil concentrate at the listed rate (but not less than 1 pt /A) in the finished spray volume

**Table 13 PS CLETHODIM TANK MIXED WITH BETAMIX<sup>o</sup> or BETANEX<sup>o</sup> APPLIED TO SUGAR BEET**

PRODUCT <sup>(1)</sup>	WEEDS CONTROLLED		WEED HEIGHT (inches)	APPLICATION RATE/ACRE <sup>(2)</sup>
	COMMON NAME	SCIENTIFIC NAME		
PS Clethodim <sup>(3)</sup> + BETAMIX or BETANEX	Barnyardgrass	<i>Echinochloa crus-galli</i>	1 to 3	8 fl oz
	Foxtail	<i>Setaria spp</i>	1 to 3	
	Foxtail Millet	<i>Setaria italic</i>	1 to 3	
	Wild Oat	<i>Avena fatua</i>	1 to 3	
	Wild Proso Millet	<i>Panicum miliaceum</i>	1 to 3	
See BETAMIX label for rates to control broadleaf weeds <b>No additives are recommended in the tank mix</b>				
See BETANEX label for rates to control broadleaf weeds <b>No additives are recommended in the tank mix</b>				

- <sup>(1)</sup> Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage Tank mixing is not recommended in these situations
- <sup>(2)</sup> Do not use crop oil concentrate No additives are recommended in the tank mix If grass regrowth occurs or an additional flush of new grass emerges make a second application of PS Clethodim alone (without a tank mix herbicide) according to the appropriate size and rate recommendations
- <sup>(3)</sup> If grass regrowth occurs or an additional flush of new grass emerges make a second application of PS Clethodim at full label rate with appropriate rate of crop oil concentrate

38/50

**Table 14 PS CLETHODIM PLUS BETANEX OR BETAMIX TANK MIX FOR THREE SEQUENTIAL APPLICATIONS FOR ANNUAL GRASS CONTROL (MICRO-RATE APPLICATION)**

PRODUCT	APPLICATION RATES/ACRE <sup>(1)</sup>		METHYLATED SEED OIL <sup>(2)</sup> (V/V)	
	ANNUAL GRASSES	GRASSES CONTROLLED (inches)	GROUND	AIR
PS Clethodim + BETANEX or BETAMIX	2 to 3 fl oz + 8 to 12 fl oz <sup>(3)</sup> or 8 to 12 fl oz <sup>(3)</sup>	Green Foxtail (1-2) Yellow Foxtail (1-2) Barnyardgrass (1-2) Wild Oat (1-2) Volunteer Cereals (1-2)	1.5%	1.5%

<sup>(1)</sup> Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not recommended in these situations.

<sup>(2)</sup> Always use a methylated seed oil at the listed rate (but not less than 1 pt /A) in the finished spray volume.

<sup>(3)</sup> Use 8 fl oz /A rate when sugar beet are in the cotyledon to 4 leaf stage. Rate can be increased up to 12 fl oz /A when the smallest sugar beet plants in the field are in the 4 true leaf stage or larger.

**Directions for Use for Micro-Rate Applications to Sugar Beet Application Information**

Multiple micro-rate applications of PS Clethodim in tank mixtures with reduced rates of BETANEX or BETAMIX and methylated seed oils may be applied by air or ground equipment to sugar beet to control early germinating annual grasses listed above. The rate of BETANEX or BETAMIX must not exceed 0.12 lb ai /A (broadcast application) when in combination with these spray adjuvants. Note that maximum rate allowed varies depending on crop growth stage. The use of wetting agents or spray adjuvants with conventional rates (0.73 to 1.22 lb ai /A) or multiple low rate (0.24 to 0.73 lb ai /A) applications of BETANEX or BETAMIX is prohibited on the BETANEX and BETAMIX master label. Favorable climatic conditions (good conditions for plant growth and development) are essential for adequate weed control. All use precautions and restrictions on the BETANEX and BETAMIX master labels must be followed.

**Directions for Using Micro-Rate Multiple Applications of PS Clethodim Tank Mixes**

Apply PS Clethodim in broadcast applications only at a rate of 2 to 3 fl oz /A in tank mixture with either BETANEX or BETAMIX following the directions for use on the tank mix partner label. A minimum of three sequential applications of 2 fl oz /A or a minimum of 2 sequential applications of 3 fl oz /A should be utilized for PS Clethodim tank mixtures. A minimum of 3 sequential applications of BETAMIX or BETANEX should be used. Accurate timing is essential. Make initial application immediately after weeds emerge and make repeat applications on 5 to 7 day intervals. If weed control is not adequate due to climatic conditions, spray coverage or other factors, return to conventional application rates of PS Clethodim (6 to 8 fl oz /A) and add rates of BETANEX or BETAMIX as directed on their label. When using conventional rates of BETANEX or BETAMIX in tank mixtures with PS Clethodim, a spray adjuvant is not recommended.

**Use Precautions for Micro-Rate Applications (See PS Clethodim, BETANEX and BETAMIX master label for further use precautions)**

Not all weeds will be adequately controlled, even with favorable climatic conditions. Conventional rates of PS Clethodim, BETANEX or BETAMIX and/or hand labor may be required if multiple micro-rate applications do not adequately control weeds. Plugging of spray nozzles may be encountered due to the potential for formation of a precipitate in the spray solution that is often associated with micro-rate applications. Prime Source LLC will not be responsible for any nozzle plugging that may occur with the use of multiple micro-rate applications. Methylated seed oils must not be added if the BETANEX or BETAMIX rate exceeds 0.12 lb ai /A broadcast, as the addition of methylated seed oils could increase the possibility of crop injury at dosage rates greater than 0.12 lb ai /A.

**GROUND APPLICATION**

Use of sufficient spray volumes and pressure is essential to ensure complete coverage. Use a minimum of 10 gals and a maximum of 20 gals of spray solution per acre. Spray pressures should reflect a minimum of 30 psi and a maximum of 60 psi at the nozzle. Do not use flood nozzles.

39/50

**AERIAL APPLICATION**

Use of sufficient spray volumes is essential to ensure complete coverage Use a minimum of 5 gals and a maximum of 15 gals of spray solution per acre

**Table 15 TANK MIX APPLICATION OF PS CLETHODIM AND FUNGICIDES FOR CONTROL OF GRASS WEEDS AND DISEASES IN SUGAR BEET**

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)
	ANNUAL GRASSES	PERENNIAL GRASSES	
PS Clethodim + EMINENT <sup>®</sup>	6 to 8 fl oz + 13 fl oz	8 to 16 fl oz + 13 fl oz	1%

<sup>(1)</sup> If grass regrowth occurs or an additional flush of new grass emerges make a second application of PS Clethodim alone (without a tank mix fungicide) according to the appropriate size and rate recommendations

<sup>(2)</sup> Refer to PS Clethodim and fungicide label for rates and weeds and diseases controlled

<sup>(3)</sup> Always use a crop oil concentrate at the listed rate (but not less than 1 pt /A) in the finished spray volume

**Table 16 TANK MIX APPLICATION OF PS CLETHODIM AND INSECTICIDES FOR CONTROL OF GRASS WEEDS AND INSECTS IN ALFALFA, COTTON, MINT, PEANUT (INCLUDING PERENNIAL), SOYBEAN AND SUNFLOWER**

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE (V/V) <sup>(3)</sup>	CROP					
	ANNUAL GRASSES	PERENNIAL GRASSES		Alfalfa <sup>(4)</sup>	Cotton	Mint <sup>(4,5)</sup>	Peanut	Soybean	Sunflower
PS Clethodim + ORTHENE <sup>®</sup> 75 S or ORTHENE 97	6 to 8 fl oz + 0.33 to 1.33 lbs or 0.25 to 1.0 lb	6 to 8 fl oz + 0.33 to 1.33 lbs or 0.25 to 1.0 lb	1%		X	X	X		
PS Clethodim + ORTHENE 90 S <sup>(6)</sup>	6 to 8 fl oz + 0.25 to 1 lb	8 to 16 fl oz + 0.25 to 1 lb	1%		X	X	X	X	
PS Clethodim + DANITOL <sup>®</sup> 2.4 EC	6 to 8 fl oz + 10 2/3 to 16 fl oz	8 to 16 fl oz + 10 2/3 to 16 fl oz	1%		X		X		
PS Clethodim + ASANA XL <sup>®</sup>	6 to 8 fl oz + Refer to ASANA XL label	8 to 16 fl oz + Refer to ASANA XL label	1%						X
PS Clethodim + WARRIOR <sup>®</sup>	6 to 8 fl oz + Refer to WARRIOR label	8 to 16 fl oz + Refer to WARRIOR label	1%						X
PS Clethodim + WARRIOR	10 to 16 fl oz <sup>(7)</sup> + Refer to WARRIOR label	10 to 16 fl oz + Refer to WARRIOR label	1%	X					
PS Clethodim + BAYTHROID <sup>®</sup>	10 to 16 fl oz <sup>(7)</sup> + Refer to BAYTHROID label	10 to 16 fl oz + Refer to BAYTHROID label	1%	X					

40/50

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE (V/V) <sup>(3)</sup>	CROP					
	ANNUAL GRASSES	PERENNIAL GRASSES		Alfalfa <sup>(4)</sup>	Cotton	Mint <sup>(4,5)</sup>	Peanut	Soybean	Sunflower
PS Clethodim + DIMETHOATE <sup>®</sup>	10 to 16 fl oz <sup>(7)</sup> + Refer to DIMETHOATE label	10 to 16 fl oz <sup>(7)</sup> + Refer to DIMETHOATE label	1%	X					
PS Clethodim + LORSBAN <sup>®</sup>	10 to 16 fl oz <sup>(7)</sup> + Refer to LORSBAN label	10 to 16 fl oz + Refer to LORSBAN label	1 to 2 pt <sup>(8)</sup>	X					
PS Clethodim + POUNCE <sup>®</sup>	10 to 16 fl oz <sup>(7)</sup> + Refer to POUNCE label	10 to 16 fl oz + Refer to POUNCE label	1%	X					

<sup>(1)</sup> If grass regrowth occurs or an additional flush of new grass emerges make a second application of PS Clethodim alone (without a tank mix fungicide) according to the appropriate size and rate recommendations

<sup>(2)</sup> Refer to PS Clethodim and fungicide label for rates and weeds and diseases controlled

<sup>(3)</sup> Always use a crop oil concentrate at the listed rate (but not less than 1 pt /A) in the finished spray volume

<sup>(4)</sup> Certain insecticides may cause temporary phytotoxic symptoms on alfalfa and mint foliage Refer to the insecticide label for further information It is suggested that prior to using any of these insecticides/herbicide tank mixtures that a small area of the field be treated first and observations for crop injury be made prior to treating the whole field

<sup>(5)</sup> The PS Clethodim rate should be 6 to 8 fl oz /A for annual grass control in baby mint minimum of 8 fl oz /A for annual grass control in established mint and 8 to 16 fl oz /A for perennial grass control Crop oil concentrate should be added at the rate of 1.0 to 2.0 pts /A

<sup>(6)</sup> Insecticide tank mix use with ORTHENE 90 S in soybean is permitted only in a state having an approved Section 24(c) registration for ORTHENE 90 S use in soybean

<sup>(7)</sup> The PS Clethodim rate should be 6 to 8 fl oz /A for annual grass control in seedling alfalfa

<sup>(8)</sup> For the PS Clethodim plus LORSBAN tank mix reduce the adjuvant rate down to 1.0 pt /A when the LORSBAN rate is 1.0 pt /A or higher

**Table 17 ROUNDUP READY VOLUNTEER CORN CONTROL IN ROUNDUP READY SOYBEAN WITH PS CLETHODIM HERBICIDE TANK MIX**

Roundup Ready Volunteer Corn Height (Inches)	PS Clethodim Rate fl oz /A	Glyphosate <sup>(1)</sup> rate for formulations with built in adjuvant	Adjuvant
<12	4	1.0 to 2.0 lb ai /A (Approximately equivalent to 22 to 44 fl oz /A of ROUNDUP Weather MAX)	Non-ionic surfactant @ 0.125 to 0.25% v/v plus ammonium sulfate @ 3.5 to 17 lbs per 100 gals /of carrier
12 to 18	5		
18 to 24	6		



41/50

Roundup Ready Volunteer Corn Height (Inches)	PS Clethodim Rate fl oz /A	Glyphosate <sup>(1)</sup> rate for formulations <b>without</b> built in adjuvant	Adjuvant
<12	4	Up to 2.0 lb ai /A (Equivalent to 32 to 64 fl oz /A of ROUNDUP Original)	Crop oil concentrate @ 1 pt /A plus ammonium sulfate @ 8.5 to 17 lbs per 100 gals of carrier
12 to 18	5		
18 to 24	6		

<sup>(1)</sup> Glyphosate formulation must be labeled for use on Roundup Ready soybean

**THE MOST RESTRICTIVE LABELING OF ANY PRODUCT USED IN A TANK MIX MUST BE FOLLOWED**

- Apply only to actively growing grass and broadleaf weeds at recommended height or growth stage listed on each label
- Apply under favorable soil moisture and humidity which exist a few days after rainfall or within seven days after irrigation
- Tank mix applications may sometimes result in reduced grass control. If regrowth occurs or an additional flush of new grass emerges, make a second application of PS Clethodim as specified in the respective size and rate tables
- Do not tank mix PS Clethodim when broadleaf weeds are tall and/or dense enough to prevent proper grass coverage
- This tank mix may be applied postemergence to ROUNDUP READY soybean up through the full flowering stage. Do not apply less than 60 days before harvest
- Avoid contact with foliage, green stems, or fruit of crops or any desirable plants and trees other than soybean with the ROUNDUP READY gene as severe injury or destruction will result
- Do not allow the PS Clethodim plus ROUNDUP to mist, drip, drift, or splash onto desirable vegetation as minute quantities of the tank mix can cause severe damage or destruction to the crops, plants, or other areas on which treatment was not intended. The likelihood of injury occurring from drift of this product is greatest when winds are gusty or in excess of 5 miles per hour. Even under lesser wind velocities, avoid conditions that allow spray drift to occur such as combinations of spray pressure and nozzle type that will result in fine particles (mist) that are likely to drift.

**FALLOW LAND**

**DIRECTIONS FOR USE**

PS Clethodim may be used to control annual and perennial grasses in land that has been left fallow the previous year and other non-producing agricultural areas. Apply PS Clethodim at 6 to 8 fl oz /A for annual grasses and 8 to 16 fl oz /A for perennial grasses. When both grass and broadleaf weeds are the target pest, PS Clethodim may be tank mixed with 2,4-D ester or BANVEL® SGF for broad spectrum control. When both annual and perennial grasses occur in the same field, use a minimum of 8 fl oz /A PS Clethodim rate.

**PRODUCT INFORMATION**

- Use a minimum spray volume of 5 gals /A for aerial applications and 15 gals /A for ground applications
- Apply only to actively growing grasses when the first grass reaches the recommended weed height as specified by the Annual and Perennial Grasses section of this label
- Annual grasses that emerge after the PS Clethodim application will not be controlled and a second application may be necessary
- The control of perennial grasses may require more than 1 application in non-tilled areas
- Do not plant any crop for 30 days after application unless clethodim is registered for use in that crop
- Do not apply to grasses that have tillered, formed seedheads, or exceeded recommended growth stage
- Do not use flood jet nozzles
- Do not apply to drought stressed grasses
- Do not mow area for 2 weeks prior to or after the PS Clethodim application

42/50

**Table 18 PS CLETHODIM IN TANK MIXES TO CONTROL ANNUAL AND PERENNIAL GRASSES IN FALLOW LAND**

PRODUCT	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND	AIR
PS Clethodim + 2 4-D ester or BANVEL SGF	6 to 8 fl oz + 0.5 lb /A or See BANVEL SGF label for rates	8 to 16 fl oz	1%	1%

<sup>(1)</sup> Refer to PS Clethodim label for weed height and species control. Review BANVEL SGF and 2 4-D labels for crop restrictions, use rates and weeds controlled.

<sup>(2)</sup> Always use a crop oil concentrate or methylated seed oil containing at least 15% emulsifier at the listed rate (but not less than 1 pt /A) in the finished spray volume.

**GRASS SUPPRESSION IN NON-CROP AREAS WITH PS CLETHODIM**

GRASS SPECIES	WEED STAGE	RATE FL OZ /ACRE	HIGH RATE
Annual and perennial grasses that exceed height claimed for control on height charts above	Up to and including grasses in the seed head stage	12	16
Do not apply as part of a tank mix when applying PS Clethodim for grass suppression. Add a crop oil concentrate at 1 qt /A by ground to the finished spray volume.			

**Table 19 PS CLETHODIM FOR THE CONTROL AND/OR SUPPRESSION OF TALL FESCUE IN NATIVE PRAIRIE WARM-SEASON GRASS RESTORATION PROJECTS**

PRODUCT	PRODUCT RATES	GRASS WEEDS CONTROLLED/SUPPRESSED		WEED STAGES
		Common Name	Scientific Name	
PS Clethodim	10 to 12 fl oz /A	Tall Fescue	<i>Festuca arundinacea</i>	4 to 6 inches tall (40 to 60% green-up)

**Adjuvant** PS Clethodim must be applied with crop oil concentrate at 1 qt /A plus a spray grade ammonium sulfate at 2.5 to 4 lbs /A. Mixing Order: Thoroughly mix spray grade ammonium sulfate in water, add PS Clethodim, then add crop oil concentrate.

**SPECIAL APPLICATION INSTRUCTIONS/ RESTRICTIONS/ PRECAUTIONS**

Burn or mow fields a minimum of 3 weeks prior to application to remove excess crop residue. Apply in the spring at 40 to 60% tall fescue green-up prior to emergence of warm-season grasses. Do not mow area for 2 weeks after the PS Clethodim application.

Apply in a minimum of 15 to 20 gals of water per acre at a spray pressure of 40 to 60 PSI at the nozzle. Apply using flat fan or hollow cone nozzles. Do not use flood jet nozzles.

Apply only to fields that have warm-season grasses established for 2 years. Applications of PS Clethodim to emerged warm-season grasses may cause injury. Do not apply to warm-season grasses grown for seed.

Do not graze treated fields or feed treated forage and or hay to livestock. Do not plant any crop for 30 days after application unless clethodim is registered for use in that crop.

**NOTE** PS Clethodim applications are most effective if applied when average nighttime temperatures are consistently greater than or equal to 47°F.

43/50

**Table 20 PS CLETHODIM FOR THE SUPPRESSION OF TALL FESCUE SEED-HEADS IN NON-PRODUCING AGRICULTURAL AREAS**

PRODUCT	PRODUCT RATE	SUPPRESSION	APPLICATION TIMING
PS Clethodim	1 1/2 to 2 fl oz /A	Tall Fescue Seed-Heads ( <i>Festuca arundinacea</i> )	(50 to 90% Tall Fescue green-up)
ADJUVANT PS Clethodim must be applied with crop oil concentrate at 1 qt /A plus a spray grade ammonium sulfate at 2.5 to 4 lb /A <b>Mixing Order</b> Thoroughly mix spray grade ammonium sulfate in water add PS Clethodim Herbicide then add crop oil concentrate			

**SPECIAL APPLICATION INSTRUCTIONS/ RESTRICTIONS/ PRECAUTIONS**

Apply at 50 to 90% tall fescue green-up  
 Use the higher PS Clethodim rate if less tall fescue green matter is present  
 Do not mow area for two weeks after the PS Clethodim application  
 Apply in a minimum of 15 to 20 gals of water per acre at a spray pressure of 40 to 60 psi at the nozzle Apply using flat fan or hollow cone nozzles Do not use flood nozzles  
 2,4-D ester may be added to this tank mix for broadleaf control (see 2,4-D ester label for weeds controlled)  
 Do not graze treated fields or feed treated forage and/or hay to livestock Do not plant any crop for 30 days after application unless clethodim is registered for use in that crop

**DIRECTIONS FOR USE IN ORNAMENTALS**

For ornamental plant uses PS Clethodim can be used to control labeled grass weeds in greenhouses bathhouses shadehouses and around outdoor ornamentals including nurseries parks roadside plantings and structure landscapes

**IMPORTANT**

PS Clethodim successfully controls weeds in newly transplanted and established non-grassy ornamentals Plant tolerance to PS Clethodim at labeled rates has been found to be acceptable for the indicated genera and species listed below Due to variability within species crop growth stage environmental conditions and application techniques it is recommended that the user determine if herbicide can be used safely on a few plants prior to widespread application Neither the seller nor the manufacturer of PS Clethodim have investigated the safety factor to ornamental plants not listed on the label

The following plants have shown a tolerance for PS Clethodim applications

**ORNAMENTAL TREES**

COMMON NAME	SCIENTIFIC NAME
Alder Red	<i>Alnus rubra</i>
Ash	<i>Fraxinus spp</i>
Basswood	<i>Tilia spp</i>
Birch European White	<i>Betula pendula</i>
Birch River	<i>Betula nigra</i>
Birch White	<i>Betula papyrifera</i>
Crabapple Flowering	<i>Malus halliana</i>
Dogwood Flowering	<i>Cornus florida</i>
Golden Chain Tree	<i>Laburnum anagyroides</i>
Maples	<i>Acer spp</i>
Mulberry White	<i>Morus alba</i>
Oaks	<i>Quercus spp</i>
Olive Wild	<i>Elaeagnus angustifolia</i>

COMMON NAME	SCIENTIFIC NAME
Redbud Eastern	<i>Cercis Canadensis</i>
Sweet Gum American	<i>Liquidambar styraciflua</i>

**GROUND COVERS**

COMMON NAME	SCIENTIFIC NAME
Bugleweed Carpet	<i>Ajuga reptans</i>
Ivy English	<i>Hedera helix</i>
Japanese Spurge	<i>Pachysandra terminalis</i>
Lilyturf	<i>Liriope muscari</i>
Moneywort	<i>Lysimachia nummularia</i>
Mondo Grass White	<i>Ophiopogon jaburan</i>
Mondo Grass Dwarf	<i>Ophiopogon japonicas</i>
Periwinkle Lesser	<i>Vinca minor</i>

**GARDEN FLOWERS AND PLANTS**

COMMON NAME	SCIENTIFIC NAME
Ageratum	<i>Ageratum spp</i>
Alyssum* Sweet	<i>Lobularia maritima</i>
Asparagus Fern	<i>Asparagus setaceus</i>
Bleeding Heart	<i>Dicentra spectabilis</i>
Cast Iron Plant	<i>Aspidistra elatior</i>
Chrysanthemum	<i>Chrysanthemum spp</i>
Cinquefoil	<i>Potentilla spp</i>
Coleus	<i>Coleus spp</i>
Coralbells	<i>Heuchera sanguinea</i>
Cranesbill	<i>Geranium spp</i>
Dahlia	<i>Dahlia spp</i>
Daisy Trailing African	<i>Osteospermum fruticosum</i>
Daylily	<i>Hemerocallis spp</i>
Dusty Miller	<i>Senecio cineraria</i>
Euonymus	<i>Euonymus spp</i>
Gazania	<i>Gazania spp</i>
Geranium House	<i>Pelargonium hortorum</i>
Heather False	<i>Cuphea hyssopifolia</i>
Hosta	<i>Hosta fortunei</i>
Iris	<i>Iris spp</i>
Jasmine Tobacco	<i>Nicotiana alata</i>
Loosestrife	<i>Lythrum salicaria</i>
Marigold	<i>Tagetes spp</i>
Partridgeberry	<i>Mitchella repens</i>
Petunia*	<i>Petunia hybrida</i>
Phlox	<i>Phlox spp</i>
Pinks	<i>Dianthus spp</i>
Portulaca	<i>Portulaca grandiflora</i>
Salvia	<i>Salvia spp</i>
Saxifrage	<i>Saxifrage spp</i>
Sedum	<i>Sedum spp</i>
Selloum	<i>Philodendron selloum</i>
Snapdragon*	<i>Antirrhinum majus</i>
Sweet Flag	<i>Acorus gramineus</i>

COMMON NAME	SCIENTIFIC NAME
Tickseed	<i>Coreopsis grandiflora</i>
Touch-Me-Not	<i>Impatiens spp</i>
Verbena	<i>Verbena spp</i>
Violet	<i>Viola spp</i>
Yarrow Common	<i>Achillea millefolium</i>
Zinnia	<i>Zinnia elegans</i>

\* Slight foliage or flower speckling has been observed on these species

**SHRUBS**

COMMON NAME	SCIENTIFIC NAME
Abelia	<i>Abelia spp</i>
Anise Purple	<i>Illicium floridanum</i>
Aucuba	<i>Aucuba spp</i>
Azalea*	<i>Rhododendron spp</i>
Bamboo	<i>Bambusa spp</i>
Barberry Japanese	<i>Berberis thunbergii</i>
Barberry Magellan	<i>Berberis buxifolia</i>
Bayberry	<i>Myrica pensylvanica</i>
Bottlebrush	<i>Callistemon citrinus</i>
Boxwood Common	<i>Buxus sempervirens</i>
Camellia Common	<i>Camellia japonica</i>
Candytuft	<i>Iberis sempervirens</i>
Cleyera	<i>Cleyera japonica</i>
Coralberry	<i>Ardisia crenata</i>
Crape Myrtle	<i>Lagerstroemia indica</i>
Coyote Brush	<i>Baccharis pilularis</i>
Fig Creeping	<i>Ficus pumila</i>
Gardenia	<i>Gardenia spp</i>
Holly	<i>Ilex spp</i>
Honeysuckle	<i>Lonicera spp</i>
Indian Hawthorn	<i>Raphiolepis indica</i>
Jasmine	<i>Jasminum spp</i>
Jasmine Asiatic	<i>Trachelospermum asiaticum</i>
Jasmine Star	<i>Trachelospermum jasminoides</i>
Juniper	<i>Juniperus spp</i>
Lantana	<i>Lantana spp</i>
Nandina* Bamboo Heavenly	<i>Nandinia domestica</i>
Oleander Common	<i>Nerium oleander</i>
Oregon Grape	<i>Mahonia aquifolium</i>
Photinia	<i>Photinia spp</i>
Pittosporum	<i>Pittosporum spp</i>
Podocarpus	<i>Podocarpus spp</i>
Privet	<i>Ligustrum spp</i>
Pyracantha	<i>Pyracantha spp</i>
Rhododendron	<i>Rhododendron spp</i>
Rose	<i>Spiraea bumalda</i>
Sweet Olive	<i>Osmanthus fragrans</i>
Viburnum	<i>Viburnum tinus</i>
Wisteria	<i>Wisteria spp</i>
Yellow Sage/Shrub Verbena	<i>Lantana camara</i>

46/50

\* Slight foliage or flower speckling has been observed on these species

**ANNUAL GRASSES IN ORNAMENTALS**

- Apply only to actively growing grasses at recommended weed heights
- Apply when the first grass weed species in a mixed grass and weed population reaches the recommended growth stage for treatment
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height

GRASS SPECIES	SCIENTIFIC NAME	WEED* HEIGHT INCHES	RATE FL OZ /ACRE <sup>(1)</sup>	HIGH RATE <sup>(2)</sup>
Barnyardgrass	<i>Echinochloa crus-galli</i>	2 to 8	8	16
Broadleaf Signalgrass	<i>Brachiaria platyphylla</i>	2 to 6	8	16
Brome				
California	<i>Bromus carinatus</i>	2 to 6	8	16
Cheat	<i>Bromus secalinus</i>	2 to 6	8	16
Downy	<i>Bromus tectorum</i>	2 to 6	8	16
Ripgut	<i>Bromus diandrus</i>	2 to 6	8	16
Canarygrass	<i>Phalaris canariensis</i>	1 to 4	8	16
Crabgrass				
Hairy	<i>Digitaria adscendens</i>	2 to 6**	8	16
Large	<i>Digitaria sanguinalis</i>	2 to 6**	8	16
Smooth	<i>Digitaria ischaemum</i>	2 to 6**	8	16
Southern	<i>Digitaria ciliaris</i>	2 to 6**	8	16
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	2 to 6**	8	16
Fall Panicum	<i>Panicum dichotomiflorum</i>	2 to 8	8	16
Field Sandbur	<i>Cenchrus incertus</i>	2 to 6	8	16
Foxtail				
Giant	<i>Setaria faberi</i>	2 to 12	8	16
Green	<i>Setaria viridis</i>	2 to 8	8	16
Yellow	<i>Setaria glauca</i>	2 to 8	8	16
Goosegrass	<i>Eleusine indica</i>	2 to 6**	8	16
Itchgrass	<i>Rottboellia cochin</i>	2 to 6	8	16
Junglerice	<i>Echinochloa colona</i>	2 to 6	8	16
Lovegrass (Stinkgrass)	<i>Eragrostis cilianensis</i>	2 to 6	8	16
Rabbitsfootgrass	<i>Polypogon monspeliensis</i>	1 to 4	8	16
Red Rice	<i>Oryza sativa</i>	1 to 3	8	16
Ryegrass				
Hardy	<i>Lolium remotum</i>	2 to 6	8	16
Italian	<i>Lolium multiflorum</i>	2 to 6	8	16
Seedling Johnsongrass	<i>Sorghum halepense</i>	4 to 10	8	16
Shattercane	<i>Sorghum bicolor</i>	6 to 18	8	16
Southwestern Cupgrass	<i>Eriochloa gracilis</i>	2 to 6	8	16
Sprangletop				
Amazon	<i>Leptochloa panicoides</i>	2 to 6	8	16
Bearded	<i>Leptochloa fascicularis</i>	2 to 6	8	16
Mexican	<i>Leptochloa uninervia</i>	2 to 6	8	16
Red	<i>Leptochloa filiformis</i>	2 to 6	8	16
Texas Panicum	<i>Panicum texanum</i>	2 to 6	8	16
Volunteer Cereals				
Barley	<i>Hordeum vulgare</i>	2 to 6	8	16
Oats	<i>Avena sativa</i>	2 to 6	8	16

47/50

GRASS SPECIES	SCIENTIFIC NAME	WEED* HEIGHT INCHES	RATE FL OZ /ACRE <sup>(1)</sup>	HIGH RATE <sup>(2)</sup>
Rye	<i>Secale cereale</i>	2 to 6	8	16
Wheat	<i>Triticum aestivum</i>	2 to 6	8	16
Volunteer Corn	<i>Zea mays</i>	4 to 12	6	8
Volunteer Corn	<i>Zea mays</i>	12 to 24	8	16
Volunteer Grain Sorghum	<i>Sorghum bicolor</i>	8 to 12	8	16
Wild Oats	<i>Avena fatua</i>	2 to 6	8	16
Wild Proso Millet	<i>Panicum miliaceum</i>	2 to 10	8	16
Witchgrass	<i>Panicum capillare</i>	2 to 8	8	16
Woolly Cupgrass	<i>Eriochloa villosa</i>	2 to 8	8	16

\* Generally occurs between 3-leaf stage and tillering

\*\* Length of lateral growth

<sup>(1)</sup> 8 fl oz /A = approximately 0.2 fl oz /1000 sq ft

<sup>(2)</sup> 16 fl oz /A = approximately 0.4 fl oz /1000 sq ft

Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt per 50 gals (0.25% v/v)

ANNUAL BLUEGRASS CONTROL WITH PS CLETHODIM IN ORNAMENTALS			
GRASS SPECIES	WEED STAGE	RATE FL OZ /ACRE	HIGH RATE
Annual Bluegrass ( <i>Poa annua</i> )	to 4-leaf	6	16

Apply under favorable soil moisture and humidity that exists within a few days after rainfall or within 7 days after irrigation. Grass needs to be actively growing at time of application(s)

Apply at weed stage indicated on the label as reduced control can be expected with more mature annual bluegrass

Use the high rate under heavy grass pressure and/or when annual bluegrass is more mature

Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt per 50 gals (0.25% v/v)

**PERENNIAL GRASSES**

- Apply only to actively growing grasses at recommended weed heights
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height

GRASS SPECIES	WEED HEIGHT (inches)	RATE FL OZ /ACRE <sup>(1)</sup>	HIGH RATE <sup>(2)</sup>
<b>Bermudagrass (<i>Cynodon dactylon</i>)</b>			
First Application	3 (or up to 6 runners)	8	16
Repeat Application(s) (if regrowth occurs)	3 (or up to 6 runners)	8	16
<b>Quackgrass (<i>Elytrigia repens</i>)</b>			
First Application	4 to 8	8	16
Repeat Application(s) (if regrowth occurs)	4 to 8	8	16

48/50

GRASS SPECIES	WEED HEIGHT (inches)	RATE FL OZ /ACRE <sup>(1)</sup>	HIGH RATE <sup>(2)</sup>
Rhizome Johnsongrass ( <i>Sorghum halepense</i> )			
First Application	12 to 24	8	16
Repeat Application(s) (if regrowth occurs)	6 to 18	6	8
Wirestem Muhly ( <i>Muhlenbergia frondosa</i> )			
First Application	4 to 8	8	16
Repeat Application(s) (if regrowth occurs)	4 to 8	8	16

<sup>(1)</sup> 8 fl oz /A = approximately 0.2 fl oz /1000 sq ft

<sup>(2)</sup> 16 fl oz /A = approximately 0.4 fl oz /1000 sq ft

Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt per 50 gals (0.25% v/v)

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

#### PESTICIDE STORAGE

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in cool, dry place. Do not store diluted spray.

#### PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### CONTAINER HANDLING

##### [NONREFILLABLE CONTAINERS]

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

(Nonrefillable plastic container ≤ 5 gallons) Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Nonrefillable plastic container > 5 gallons) Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Nonrefillable metal container > 5 gallons) Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures approved.



49/50

by state and local authorities

**[REFILLABLE CONTAINERS]**

Refillable container Refill this container with pesticide only Do not reuse this container for any other purpose  
Cleaning the container before final disposal is the responsibility of the person disposing of the container  
Cleaning before refilling is the responsibility of the refiller

To clean the container before final disposal empty the remaining contents from this container into application  
equipment or mix tank Fill the container about 10 percent full with water Agitate vigorously or recirculate  
water with the pump for 2 minutes Pour or pump rinsate into application equipment or rinsate collection  
system Repeat this rinsing procedure two more times Then offer for recycling if available or puncture and  
dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities by burning If  
burned stay out of smoke

**CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability  
before buying or using this product If the terms are not acceptable return the product at once unopened and  
the purchase price will be refunded

The Directions for Use of this product must be followed carefully It is impossible to eliminate all risks  
inherently associated with the use of this product Ineffectiveness or other unintended consequences may  
result because of such factors as manner of use or application weather presence of other materials or other  
influencing factors in the use of the product which are beyond the control of Prime Source LLC or Seller To  
the extent consistent with applicable law all such risks shall be assumed by Buyer and User and Buyer and  
User agree to hold Prime Source LLC and Seller harmless for any claims relating to such factors

Prime Source LLC warrants that this product conforms to the chemical description on the label and is  
reasonably fit for the purposes stated in the Directions for Use subject to the inherent risks referred to above  
when used in accordance with directions under normal use conditions This warranty does not extend to the  
use of this product contrary to label instructions or under abnormal conditions or under conditions not  
reasonably foreseeable to or beyond the control of Seller or Prime Source LLC and Buyer and User assume  
the risk of any such use **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW PRIME SOURCE LLC  
MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE  
NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE**

To the extent consistent with applicable law neither Prime Source LLC nor Seller shall be liable for any  
incidental consequential or special damages resulting from the use or handling of this product **TO THE  
EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER,  
AND THE EXCLUSIVE LIABILITY OF PRIME SOURCE, LLC AND SELLER FOR ANY AND ALL CLAIMS,  
LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY,  
CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE  
OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE  
PRODUCT OR, AT THE ELECTION OF PRIME SOURCE, LLC OR SELLER, THE REPLACEMENT OF THE  
PRODUCT**

Prime Source LLC and Seller offer this product and Buyer and User accept it subject to the foregoing  
Conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written  
agreement signed by a duly authorized representative of Prime Source LLC

Asana® is a registered trademark of DuPont  
Authority Broadleaf® Canopy® Classic® and Synchrony® are registered trademarks of E I DuPont de Nemours  
& Co Inc  
STS™ is a trademark of E I DuPont de Nemours & Co Inc  
Banvel® IMI-CORN® and Storm® are registered trademarks of BASF Corporation  
Basagran® is a registered trademark of BASF Ag  
Baythroid® and Sencor® are registered trademarks of Bayer Corporation

