



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

March 6, 2017

Kt Woodall
For Prime Source LLC
P.O. Box 250
10025 Hwy. 264 Alternate
Middlesex, NC 27557

Subject: Label Amendment – To incorporate re-pack restrictions
Product Name: Imidacloprid 75WSP Select
EPA Registration Number: 89442-11
Application Date: September 8, 2016
Decision Number: 521385

Dear Ms. Woodall,

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 89442-11
Decision No. 521385

with FIFRA section 6. If you have any questions, please contact Jessica Rogala by phone at 703-347-0263, or via email at rogala.jessica@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Elyssa Arnold". The signature is written in a cursive style with a loop at the end of the last name.

Elyssa Arnold, Acting Product Manager 01
Invertebrate & Vertebrate Branch 3
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

IMIDACLOPRID 75WSP Select

Insecticide for foliar and systemic insect control in turfgrass (including sod farms),
landscape ornamentals, listed fruit and nut trees and interior plantscapes.

ACTIVE INGREDIENT:	% BY WT.
Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimin	75.0%
OTHER INGREDIENTS:	<u>25.0%</u>
TOTAL:	100.0%

Keep water soluble packets in this container and store in a cool dry place but not below freezing (32°F).

Do not remove packets from container except for immediate use.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

PRECAUCION AL USUARIO: Si usted no puede leer o entender ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)


FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minute. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 Mon.-Fri., 8:00 a.m. to 12:00 p.m. Pacific Time (NPIC Website: www.npic.orst.edu). Outside of these times call your poison control center at 1-800-222-1222.</p>	
<p>Note to Physician: No specific antidote is available. Treat the patient symptomatically.</p>	

EPA Reg. No. 89442-11

EPA Est. No. xxxxxx-xx-xxx

NET CONTENTS: _____

Manufactured For:
Prime Source, LLC
P.O. Box 250
10025 Hwy. 264 Alternate
Middlesex, NC 27557



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or vapor.

Applicators and Other Handlers Must Wear:

1. Long-sleeved shirt and long pants
2. Chemical-resistant gloves made of barrier laminate, butyl rubber > 14 mils, nitrile rubber > 14 mils, neoprene rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinylchloride (PVC) \geq 14 mils, or viton \geq 14 mils
3. Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment, PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

User should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops/plants or weeds. Do not apply this product or allow it to drift to blooming crops/plants or weeds if bees are foraging the treatment area.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications.
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at:

<http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: www.aapco.org/officials.html. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

See individual crops for specific pollinator protection application restrictions. If none exist under the specific site, for outdoor foliar applications, follow these application directions.



- **Do not apply while bees are foraging.**
- **Do not apply this product to plants that are flowering.**
- **Only apply after all flower petals have fallen off.**

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If the product is applied by soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

1. Coveralls
2. Chemical-resistant gloves made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinylchloride (PVC) \geq 14 mils, or viton \geq 14 mils
3. Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and pets off treated area until dry.

RESTRICTION: Do not formulate this product into other end-use products.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE STORAGE: Keep pesticide in original container. Keep container tightly closed when not in use. Protect from excessive heat. Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or dispose of empty container in a sanitary landfill or by incineration.

APPLICATION TO TURFGRASS

IMIDACLOPRID 75WSP SELECT will control the following soil inhabiting pests found in turfgrass

PEST	SCIENTIFIC NAME
Northern & Southern masked chafers	<i>Cyclocephala borealis</i> , <i>C. immaculata</i> , and/or <i>C. lurida</i>
Asiatic garden beetle	<i>Maladera castanea</i>
European chafer	<i>Rhizotroqus majalis</i>
Green June beetle	<i>Cotinis nitida</i>
May or June beetle	<i>Phyllophaga</i> spp.
Japanese beetle	<i>Popillia japonica</i>
Oriental beetle	<i>Anomala orientalis</i>
Billbugs	<i>Spherophorus</i> spp.
Annual bluegrass weevil	<i>Hyperodes</i> spp.
Black turfgrass ataenius	<i>Ataenius spretulus</i> and <i>Aphodius</i> spp.
European Crane Fly	<i>Tipula paludosa</i>
Mole crickets	<i>scapteriscus</i> spp.

IMIDACLOPRID 75WSP SELECT will suppress cutworms and chinch bugs.

IMIDACLOPRID 75WSP SELECT can be applied on turfgrass in the following sites:

Home lawns	Cemeteries
Business and office complexes	Parks
Shopping complexes	Playgrounds
Multi-family residential complexes	Athletic fields
Golf courses	Sod farms
Airports	

IMIDACLOPRID 75WSP SELECT has adequate residual activity that applications can be made preceding the egg laying activity of the target pests.

Control is achieved when applications are made prior to egg hatch of the pests and when rainfall or irrigation after application will penetrate vertically in the soil column carrying the active ingredient into the zone where insects are normally located. In order to move the active ingredient through the thatch, irrigate if rainfall does not occur within 24 hours after application.

Apply 1.6 ounces (1 packet per 8,250 to 11,000 sq. ft.). Make application prior to egg hatch of grubs, billbugs, annual bluegrass weevil, and European Crane Fry to maximize control.

For chinch bugs (suppression) and mole crickets apply 1.6 ounces (1 packet) per 8,250 sq. ft. For suppression of chinchbugs, make application before the hatching of the first instar nymphs.

For control of mole crickets make application before or during the peak egg hatch period. If adults or large nymphs are actively tunneling, combine applications of IMIDACLOPRID 75WSP SELECT with a remedial insecticide. Follow label instructions for other insecticides when tank-mixing.

Consult your local turf, state Agricultural Experiment Station, or State Extension Service Specialists for more specific Information regarding timing of application.

Application Equipment for Use on Turfgrass

Dilute IMIDACLOPRID 75WSP SELECT with enough water to provide adequate volume to promote thorough distribution into the pest zone. Use only accurately calibrated equipment for application to turfgrass. Apply a uniform, coarse droplet spray, using a low pressure setting to eliminate off target drift. Perform calibration on a regular basis to ensure that equipment is distributing product properly.

RESTRICTIONS FOR TURF GRASS:

- Do not exceed a total of 8.6 ounces application (0.4 lb. of active ingredient) per acre per year.
- Do not allow this product to contact plants in bloom if bees are foraging in the treatment area.
- Do not mow turf or lawn area, until after adequate irrigation or rainfall has occurred.
- Keep people and pets off treated area until dry.

- Not for use on grasses grown for seed, or on commercial fruit and nut trees.
- Do not graze treated areas or use clippings from treated areas for feed or forage.
- Do not apply through any irrigation system.
- Do not apply to areas which are water logged or saturated, which will not allow penetration into the root zone of the plant.
- Use not permitted in Arizona unless otherwise directed by state 24(c) labeling.

APPLICATION TO ORNAMENTALS

IMIDACLOPRID 75WSP SELECT can be applied to ornamental plants in commercial and residential landscapes and interior plantscapes. IMIDACLOPRID 75WSP SELECT is a systemic insecticide that is transported within the plant system from the roots to upper foliage. IMIDACLOPRID 75WSP SELECT must be applied into a growing area of the plant that allows absorption of the active ingredient. Adding soluble nitrogen type fertilizers to the spray solution when appropriate can promote the uptake of the active ingredient.

Application can be made by foliar application or soil applications; including soil injection, drenches, and broadcast sprays. Foliar applications offer locally systemic activity against listed insect pests.

The systemic translocation of active ingredient will be slower when applied to woody plants with soil applications. This delay can take 60 days or longer depending on species and size of plant. To offset this, make applications before anticipated pest infestation.

ANT MANAGEMENT PROGRAMS

Using IMIDACLOPRID 75WSP SELECT to control aphids, scale insects, mealybugs and other sucking pests on ornamentals will reduce ant populations because it removes honeydew as a food source. To enhance control of ants, supplement with residual sprays, bait placements or other ant control tactics.

APPLICATION EQUIPMENT FOR FOLIAR APPLICATIONS

IMIDACLOPRID 75WSP SELECT mixes readily with water and may be used in many types of application equipment. Add a commercial spreader/sticker promote coverage on hard to wet foliage such as holly, pine, or ivy.

IMIDACLOPRID 75WSP SELECT is compatible with many commonly used fungicides, miticides, liquid fertilizers, and other insecticides. If applicator has no prior experience with a particular tank mix, check physical compatibility by making a small clear jar test using correct proportions of products to be tank mixed.

RESTRICTIONS FOR ALL ORNAMENTALS:

- Keep people and pets off treated area until dry.
- Do not apply more than 8.6 ounces (0.4 lb. of active ingredient) per acre per year.
- Not for use in commercial greenhouses, nurseries, or on grasses grown for seed, or on commercial fruit and nut trees.
- Do not apply through any irrigation system.
- Do not apply to areas which are water logged or saturated, which will not allow penetration into the root zone of the plant.
- Do not apply this product, by any application method, to linden, basswood or other *Tilia* species.
- Do not allow puddling of irrigation water following application.

FOR FOLIAR APPLICATION TO ORNAMENTAL AND NON-BEARING FRUIT & NUT TREES, SHRUBS, EVERGREENS, FLOWERS, FOLIAGE PLANTS, GROUNDCOVERS AND INTERIOR PLANTSCAPES

(Only for Industrial, Commercial Buildings, Residential and Landscaped Areas)

TO CONTROL:

Adelgids	(including Rhododendron gall midge,
Aphids	Honeylocust pod gall midge)
Honeylocust Plant Bug	Froghoppers
Diptera	Galls (Including Hickory stem gall)

Japanese beetles	Scales
Lace bugs	(including Lecanium, Azalea bark,
Leaf beetles	Calico, Cottony Camellia, Cottony maple,
(including elm and viburnum leaf beetles)	Cottony taxus)
Leafhoppers	Spittlebugs
(including glassy-winged sharpshooter)	Thrips (suppression)
Leafminers	(including Flower, Pear and Pine thrips)
(including Boxwood leafminer)	Treehoppers
Mealybugs	Weevils
Planthoppers	(including white Pine and Black vine)
Psyllids	Whiteflies
Sawfly larvae	

Apply 1.6 ounces (1 packet) of Imidacloprid 75WSP SELECT per 300 gallons of water. Make foliar applications before high pest populations become established. Reapply on an as needed basis.

To control White grub larvae, (such as Japanese beetle larvae, Chafers, *Phyllophaga* spp. Asiatic garden beetle, Oriental beetle) apply 1.6 ounces (1 packet) per 8,250 to 11,000 sq. ft.

Make broadcast applications by mixing specified rate in enough water to uniformly cover the treated area. Do not use less than 2 gallons of water per 1,000 sq. ft. If necessary, irrigate thoroughly to incorporate IMIDACLOPRID 75WSP SELECT into the upper soil profile.

Refer to use directions specific for FLOWERS and GROUND COVERS concerning additional use directions.

RESTRICTIONS FOR PLANTSCAPES:

- Do not apply more than 8.6 ounces (0.4 lb. of active ingredient) per acre per year.
- Do not apply this product, by any application method, to linden, basswood, or other *Tilia* species.

FOR SOIL APPLICATION TO ORNAMENTAL AND NON-BEARING FRUIT & NUT TREES, SHRUBS, FLOWERS AND GROUNDCOVERS

(Only For Industrial, Commercial Buildings, Residential, and Landscaped Planting Areas)
(Including State, National, and Private Wooded and Forested Areas)

TO CONTROL:

Adelgids	Leafminers
Aphids	Mealybugs
Armored scales (suppression)	Pine tip moth larvae
Black vine weevil larvae	Plant bug
Eucalyptus longhorned borer	Psyllids
Flatheaded borers	Roundheaded borers
(including bronze, alder and emerald ash borers)	(including Asian longhorned beetles)
Japanese beetles	Royal Palm Bugs
Lace bugs	Sawfly larvae
Leaf beetles	Soft scales
(including elm and viburnum leaf beetles)	Thrips (suppression)
Leafhoppers	White grub larvae
(including glassywinged sharpshooter)	Whiteflies

For TREES –

Use the following rates as a function of tree diameter at breast height (DBH):

Apply 1.6 ounces (1 packet) of Imidacloprid 75WSP SELECT per 12 to 48 inches of cumulative trunk diameter for tree(s) intended to be treated.

Use the higher rate only for trees greater than 15 DBH to control the following pests:

Asian longhorned beetle
Emerald Ash Borer
Eucalyptus longhorned borer
Bronze birch borer
Alder borer

To calculate the higher rates divide the cumulative trunk diameter by 12-23 inches.

EXAMPLES OF HOW TO CALCULATE:

To calculate the lower rate:

- If you have three trees having DBH of 8, 10 & 14 inches, the total cumulative inches of trunk diameter will be 34 inches ($8 + 10 + 16 = 34$).

$$34/48 = .708 \times 1.6 \text{ fl. oz. (1 packet)} = 1.13 \text{ oz. per tree}$$

OR

$$34/24 = 1.417 \times 1.6 \text{ fl. oz. (1 packet)} = 2.27 \text{ oz. per tree}$$

- If you have a single tree with a DBH of 12 inches, the lower rate range will be:

$$12/48 = 0.25 \times 1.6 \text{ oz. (1 packet)} = 0.4 \text{ oz. per tree}$$

OR

$$12/24 = 0.5 \times 1.6 \text{ oz. (1 packet)} = 0.8 \text{ oz. per tree}$$

To calculate the higher rate:

- If you have three trees having DBH of 15, 20 & 25 inches, the total cumulative inches of trunk diameter will be 60 inches ($15 + 20 + 25 = 60$).

$$60/23 = 2.6 \times 1.6 \text{ fl. oz. (1 packet)} = 4.17 \text{ oz. per tree}$$

OR

$$60/12 = 5 \times 1.6 \text{ fl. oz. (1 packet)} = 8 \text{ oz. per tree}$$

- If you have a single tree with a DBH of 30 inches, the higher rate range will be:

$$30/23 = 1.3 \times 1.6 \text{ oz. (1 packet)} = 2.08 \text{ oz. per tree}$$

OR

$$30/12 = 2.5 \times 1.6 \text{ oz. (1 packet)} = 4.0 \text{ oz. per tree}$$

RESTRICTION: Do not apply more than 8.53 oz. (0.4 lb. of active ingredient) per acre per year.

Diameter at Breast Height (D.B.H.) = is measured at 4.5 feet from the ground.

APPLICATION TECHNIQUE

SOIL INJECTION:

Mix required dosage in sufficient water to inject an equal amount of solution in each hole. For concentrate injectors mix required dose with up to 1 gallon of water per DBH inch. Maintain a low pressure and use sufficient solution for distribution of the liquid into treatment zone. Dilution rate may vary depending on equipment used, tree size and application rate. Refer to the instructions for injector equipment being used for guidance.

- GRID SYSTEM:** Make applications in a grid pattern on 2.5 foot centers within the drip line of the tree.
- CIRCLE SYSTEM:** Make applications in holes evenly spaced approximately 2 – 3 feet apart in a circle within the drip line of the tree. Larger trees may require additional application circles.
- BASAL SYSTEM:** Make applications into holes evenly spaced around the base of the tree trunk no more than 6 to 12 Inches out from the base.

SOIL DRENCH:

Apply in no less than 10 gallons of water per 1,000 square feet as a drench around the base of the tree, directed to the root zone. Any plastic or other barrier that can prevent drench solution from reaching the root zone must be removed.

FOR ALL APPLICATION TECHNIQUES:

- Inject an equal amount of water and solution in each hole.
- Use low pressure and sufficient solution for thorough distribution into the treatment zone.
- Maintain soil moisture for 7 to 10 days.
- Application to trees already heavily infested with borers listed may not prevent the eventual loss of the trees.

RESTRICTIONS FOR TREES:

- Do not use less than 4 holes per tree.
- No Soil Injection Applications Allowed In Nassau or Suffolk Counties of New York.
- Do not apply more than 8.53 oz. (0.4 lb. of active ingredient) per acre per year.
- Do not apply this product, by any application method, to linden, basswood or other *Tilia* species.

For SHRUBS –

Apply 1.6 ounces (1 packet) of Imidacloprid 75WSP Select per 24 to 48 foot of cumulative shrub height.

APPLICATION TECHNIQUES**SOIL INJECTION:**

Apply to individual plants using dosage Indicated.

SOIL DRENCH:

Uniformly apply the dosage in no less than 10 gallons of water per 1,000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.

- Mix required dosage in sufficient water to inject an equal amount of solution in each hole.
- Use low pressure and sufficient solution for distribution of the liquid into the treatment zone.
- Maintain soil moisture for 7 to 10 days.

RESTRICTIONS FOR SHRUBS:

- Do not use less than 4 holes per shrub.
- No Soil Injection Applications Allowed In Nassau or Suffolk Counties of New York.
- Do not apply more than 8.53 oz. (0.4 lb. of active ingredient) per acre per year.

For FLOWERS and GROUNDCOVERS -

Apply 1.6 ounces (1 packet) of Imidacloprid 75WSP Select per 8,250 to 11,000 sq. ft.

Apply as a broadcast treatment and incorporate into the soil before piling or apply after plants are established. If application is made to established plants, best control is achieved by irrigating after application.



POME FRUITS
(For Residential Areas Only)

Includes: Apple, Crabapple, Loquat, Mayhaw, Pear, Pear (oriental), Quince

TO CONTROL:

Aphids (except Woolly apple aphid)
Leafhoppers (including glassy-winged
sharpshooter)

Leafminer
Mealybugs
San Jose Scale

Apply 1.6 ounces (1 packet) per 300 gallons of water OR 2.1 ounces per acre¹ as foliar spray after petal-fall is complete.

¹The amount of Imidacloprid 75WSP Select required per acre depends on tree size and/or volume of foliage. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees with full foliage.

- For control of rosy apple aphid, apply prior to leafrolling caused by the pest.

- For first generation leafminer control, make first application as soon as petal fall is complete. Greatest leafminer control will result from the earliest possible application. For second and succeeding generations of leafminer, optimal control is obtained from applications made early in the adult flight against egg and early instar larvae. A second application may be required 10 days later if severe pressure continues or if generations are overlapping a single application may result in suppression only. IMIDACLOPRID 75WSP SELECT will not control late stage larvae.
- For San Jose Scale, time applications to the crawler stage. Treat each generation.
- For late season (preharvest) control of leafhopper species, apply IMIDACLOPRID 75WSP SELECT while most leafhoppers are in the nymphal stage.
- For optimal control of mealybug, insure good spray coverage of the trunk and scaffolding limbs or other resting sites of the mealybug.

RESTRICTIONS FOR RESIDENTIAL POME FRUITS:

- Do not apply more than 2.1 ounces per acre in a single application.
- Do not make more than 4 applications per year.
- Allow 10 or more days between applications.
- Allow at least 7 days between last application and harvest.
- Not for use in California for control on pears.
- Do not apply pre-bloom or during bloom or when bees are foraging.

**PECAN
(For Residential Areas Only)****TO CONTROL:**

Yellow pecan aphid
Pecan spittlebug
Black margined aphid

Pecan stem phylloxera
Pecan leaf phylloxera

Apply 1.6 ounces (1 packet) per 300 gals. OR 2.1 ounces per acre as a foliar application as pest pressure begins to increase.¹

Make a second application 10 to 14 days after first if field scouting reveals continued pest pressure. Use of an organosilicone based spray adjuvant at specified rate can insure thorough coverage of foliage.

RESTRICTIONS FOR RESIDENTIAL PECAN TREES:

- Do not apply more than a total of 6.3 ounces of IMIDACLOPRID 75WSP SELECT per acre per year.
- Use on pecans not permitted in California unless directed by state specific 24(c) labeling.
- Do not make more than 3 applications per year.
- Allow 10 or more days between applications.
- Do not apply pre-bloom or during bloom or when bees are foraging.

¹The amount of Imidacloprid 75WSP Select required per acre vary depending on tree size or volume of foliage. The rates given are based on a standard of 400 gallons of dilute spray solutions per acre for large trees with full foliage. DO NOT apply more than 6.3 fluid ounces of Imidacloprid 75 Select per acre per year.

**GRAPES
(For Use Only For Industrial, Commercial Buildings and Residential Planting Areas)****TO CONTROL:**

Leafhoppers (including glassy-winged sharpshooter)
Mealybugs

Apply 1.6 ounces (1 packet) per 300 gallons of water OR 1.0 ounces per acre as a foliar spray using 200 gallons of water per acre.

RESTRICTIONS FOR RESIDENTIAL GRAPES:

- Do not apply more than a total of 2.0 ounces of IMIDACLOPRID 75WSP SELECT per acre per year.
- Allow at least 14 days between applications.
- Applications may be applied up to and including day of harvest.

CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of PRIME SOURCE, LLC, its Supplemental Distributors, or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

PRIME SOURCE, LLC, its Supplemental Distributors and the Seller warrant that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NEITHER PRIME SOURCE, LLC, NOR ITS SUPPLEMENTAL DISTRIBUTORS MAKE ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS FOR ANY PARTICULAR PURPOSE OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND THE EXCLUSIVE LIABILITY OF PRIME SOURCE, LLC, ITS SUPPLEMENTAL DISTRIBUTORS AND THE SELLER FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. To the extent consistent with applicable law, When Buyer suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), Buyer must promptly notify Seller in writing of any claims to be eligible to receive either remedy stated above. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL PRIME SOURCE, LLC, ITS SUPPLEMENTAL DISTRIBUTORS, OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. PRIME SOURCE, LLC, its Supplemental Distributors, and the Seller offer this product, and the Buyer accepts it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of PRIME SOURCE, LLC.

No employee or agent of PRIME SOURCE, LLC, its Supplemental Distributor, or the Seller is authorized to vary or exceed the terms of this Warranty in any other manner.