



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

89442-54

Date of Issuance:

8/14/20

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Azoxy D Select

Name and Address of Registrant (include ZIP Code):

Prime Source, LLC
c/o Wagner Regulatory Associates, Inc.
P.O. Box 640
Hockessin, DE 19707

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Shaja B. Joyner, Product Manager 20
Fungicide-Herbicide Branch
Registration Division 7505P

Date:

8/14/20

2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 89942-54.”
3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 04/01/2020

If you have any questions, please contact Manjula Unnikrishnan by phone at 703-347-8520, or via email at unnikrishnan.manjula@epa.gov

Enclosure

[MASTER LABEL]

AZOXYSTROBIN	GROUP	11	FUNGICIDE
DIFENOCONAZOLE	GROUP	3	FUNGICIDE

Azoxy D Select

Contains difenoconazole and azoxystrobin, the active ingredient used in [Briskway™].

A broad-spectrum fungicide for prevention and control of listed diseases in golf course turfgrasses only.

ACTIVE INGREDIENTS:	(% by weight)
Azoxystrobin*	18.2%
Difenoconazole**	11.4%
OTHER INGREDIENTS:	70.4%
TOTAL:	100.0%

*CAS No. 131860-33-8
 **CAS No. 119446-68-3

Azoxy D Select is a suspension concentrate (SC) formulation that contains 1.67 lbs. of azoxystrobin and 1.05 lbs. of difenoconazole per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 .	

[Optional referral statements when booklets and container labels are used:]


[See label booklet for [complete] [additional] [First Aid,] [Precautionary Statements,] [Directions For Use,] and [Storage and Disposal].]

EPA Reg. No.: 89442-XX

EPA Est. No.: _____

Net Contents: _____ [Gal./L.]

Manufactured For:
 Prime Source, LLC
 P.O. Box 250
 10025 Hwy. 264 Alternate
 Middlesex, NC 27557



Azoxy D Select is not manufactured or distributed by Syngenta Crop Protection, LLC, seller of [Briskway™].

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminate clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber, or butyl rubber

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Difenoconazole is toxic to fish, mammals and aquatic invertebrates. Drift and runoff may be hazardous to estuarine/marine organisms in water adjacent to treated area. Azoxystrobin is toxic to freshwater and estuarine/marine fish and aquatic invertebrates.

Azoxystrobin can be persistent for several months or longer. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash water or rinsate.

Groundwater Advisory

Azoxystrobin and a degradate of azoxystrobin are known to leach through soil to groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of azoxystrobin and a degradate of azoxystrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Notify State and/or Federal authorities and Prime Source, LLC immediately if you observe any adverse environmental effects due to use of this product.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow coming into contact with oxidizing agents, hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN TURF INJURY AND/OR POOR DISEASE CONTROL.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

PRODUCT INFORMATION

Azoxy D Select is a broad-spectrum systemic product containing two fungicides. It has preventative and curative properties. **Azoxy D Select** provides excellent disease control of many major turfgrass diseases. **Azoxy D Select** may be applied as a foliar spray and integrated into a resistance management program. All applications need to be made according to the use directions that follow.

POLLINATOR ADVISORY STATEMENT: This product may adversely impact the forage and habitat of local pollinators, including the monarch butterfly (and its larvae), birds, or bats if reaches non-target areas. Protect pollinators by following label directions to minimize spray drift.

USE PRECAUTIONS AND RESTRICTIONS

- **DO NOT** apply by air.
- **DO NOT** apply this product through any type of irrigation system (CHEMIGATION).
- **DO NOT** spray **Azoxy D Select** where spray drift may reach apple trees. **Azoxy D Select** is extremely phytotoxic to certain apple varieties.

USE INFORMATION

Application: Thorough coverage is necessary to provide good disease control. Make no more spray solution than is needed for application.

For control of foliar diseases, allow **Azoxy D Select** applications to completely dry prior to irrigating. For control of soil-borne diseases, **Azoxy D Select** can be watered in after application.

Adjuvants: When an adjuvant is to be used with this product, Prime Source, LLC advises the use of a Chemical Producers and Distributors Association certified adjuvant.

Efficacy: Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if the maximum amount of **Azoxy D Select** has been used. If resistant isolates to Group 3 or Group 11 fungicides are present, efficacy can be reduced for certain diseases. The higher rates in the rate range and/or shorter spray intervals may be required under conditions of heavy infection pressure with highly susceptible varieties, or when environmental conditions are conducive to disease.

Azoxy D Select controls certain pathogens causing foliar, stem and root diseases including leaf and stem blights, leaf spots, patch diseases, mildews, anthracnose, fairy rings, molds, and rusts of turfgrass plants.

Integrated Pest Management (IPM): **Azoxy D Select** needs to be integrated into an overall disease and pest management strategy whenever the use of a fungicide is required.

Sound turf management resulting in healthy, vigorous turf is the foundation of a good IPM program. Cultural practices such as proper choice of turf variety, nutrient management, proper cutting height, thatch management, and proper watering, drainage, and moisture stress management need to be integrated with the use of fungicides to increase turf vigor and reduce susceptibility to disease. Immunoassay detection kits and Extension service diagnostic services can assist in the early and accurate identification of causal organisms and corresponding selection of the proper fungicide when required.

RESISTANCE MANAGEMENT

AZOXYSTROBIN	GROUP	11	FUNGICIDE
DIFENOCONAZOLE	GROUP	3	FUNGICIDE

For resistance management, please note that **Azoxy D Select** contains both a Group 11/azoxystrobin and Group 3/difenoconazole fungicide. Any fungal population may contain individuals naturally resistant to **Azoxy D Select** and other Group 11 or Group 3 fungicides. A gradual of total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies need to be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of **Azoxy D Select** or other Group 11 and Group 3 fungicides. Do not apply more than two sequential applications of **Azoxy D Select** for control of gray leaf spot. For all other diseases when gray leaf spot is not present, do not apply more than three sequential applications of **Azoxy D Select**.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting; uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM directions for specific crops and pathogens.
- For further information or to report suspected resistance, contact Prime Source, LLC. You can also contact your pesticide distributor or university extension specialist to report resistance.

Some turf disease pathogens are known to have developed resistance to products used repeatedly for their control. **Azoxy D Select** should be alternated with other registered fungicides that have a different mode of action and to which pathogen resistance has not developed. Since **Azoxy D Select** is a QoI fungicide (Group 11 fungicide) and a DMI fungicide (Group 3 fungicide), avoid alternations with other QoIs such as pyraclostrobin, fluoxastrobin and trifloxystrobin or other DMI's such as propiconazole, tebuconazole and metaconazole. Addition of, or alternation to, a fungicide with Pythium activity such as Subdue® MAXX fungicide may be required if Pythium diseases are a major concern.

Turfgrass Resistance: Azoxy D Select plant resistance has been found to be acceptable for all turfgrass species; however, not all possible tank-mix combinations have been tested under all conditions. When possible, it is advised to test the combinations on a small portion of the turf to ensure that a phytotoxic response will not occur as a result of application.

MANDATORY SPRAY DRIFT

Ground Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy unless making a turf, pasture, or rangeland application, in which case applicators may apply with a nozzle height no more than 4 feet above the ground.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Boom-less Ground Applications:

- Applicators are required to use a medium or coarser droplet size (ASABE S572.1) for all applications.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.
- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.

Spray Drift Advisories

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT - Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

MIXING AND APPLICATION METHODS

Spray Equipment

Nozzles

- Equip sprayers with nozzles that provide accurate and uniform application.
- Nozzles need to be the same size and uniformly spaced across the boom.
- Calibrate sprayer before use.
- It is suggested that screens be used to protect the pump and to prevent nozzles from clogging.
- Screens placed on suction side of pump need to be 16-mesh or coarser.
- **DO NOT** place a screen in the recirculation line.
- Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles.
- Check nozzle manufacturer's directions.

Pump

- Use a pump with capacity to:
 - Maintain 35-40 psi at nozzles.
 - Provide sufficient agitation in tank to keep mixture in suspension – this requires recirculation of 10% of tank volume per minute.
- Use a jet agitator or liquid sparge tube for agitation.
- **DO NOT** air sparge.

For more information on spray equipment and calibration, consult sprayer manufacturers and State directions. For specific local directions and spray schedules, consult the current State agricultural directions.

Mixing Instructions

Azoxy D Select is a suspension concentrate (SC) formulation.

Thoroughly clean spray equipment before using this product.

Make up only the amount of spray required for the immediate use.

To prepare a spray solution, partially fill the spray tank with clean water and begin agitation. Add the specified amount of **Azoxy D Select** to the tank, allowing time for good dispersion, then add an adjuvant, if specified.

DO NOT use silicone-based products with **Azoxy D Select** due to possible phytotoxicity.

If tank mixes are required, products need to be added to the spray tank in the following order: **Azoxy D Select**, WG or dry flowable formulations, wettable powders and flowable (aqueous suspensions) products. Finish filling the spray tank to the desired volume to obtain the proper spray concentration.

Maintain agitation throughout the spraying operation. Do not allow spray mixture to stand overnight or for prolonged periods. If spray tank mixture is unsprayed for more than 18 hours (overnight), re-suspend product with agitation for 20 minutes.

Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

Azoxy D Select Alone (No Tank Mix)

- Add 1/2 - 2/3 of the required amount of water to the spray or mixing tank.
- With the agitator running, add **Azoxy D Select** to the tank.
- Continue agitation while adding the remainder of the water.
- Begin application of the spray solution after **Azoxy D Select** has completely dispersed into the mix water.
- Maintain agitation until all of the mixture has been sprayed.

Azoxy D Select + Tank Mixtures

Azoxy D Select is usually compatible with many tank-mix partners registered for use on turf. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions

for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

If tank mixes are desired, observe all directions, precautions and limitations on labeling of all products used. Consult compatibility charts or your local or State turf authority for compatibility information. **DO NOT** combine **Azoxy D Select** in the spray tank with pesticides, surfactants, or fertilizers, unless compatibility charts or your own prior use has shown that the combination is physically compatible, effective, and non-injurious under your conditions of use.

To determine the physical compatibility of **Azoxy D Select** with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Mixing in the Spray Tank

- Add 1/2 - 2/3 of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank-mix partner(s) into the tank in the same order as described above.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and **Azoxy D Select** to the spray tank.
- Allow **Azoxy D Select** to completely disperse.
- Spray the mixture with the agitator running.

Application Instructions

- **DO NOT** apply by air.
- **DO NOT** apply this product through any type of irrigation system. (CHEMIGATION).

Azoxy D Select may be applied with all types of spray equipment commonly used for making ground applications. Proper adjustments and calibration of spraying equipment to give good coverage is essential for good disease control.

Ground Application

- Apply in sufficient water to provide good coverage. Typical application volumes range from 30 to 450 gallons of spray per acre.
- Thorough coverage is necessary to provide good disease control.

SPECIFIC DIRECTIONS FOR USE

Target Diseases	Use Rate (fl. oz. product per 1,000 sq. ft.)	Use Rate (fl. oz. product per acre)	Application Interval (days)	Application Instructions*
Anthracnose (<i>Colletotrichum cereale</i>) (formerly known as <i>C. graminicola</i>)	0.3 to 0.725 (0.0039 lb. azoxystrobin/ 0.0024 lb. difenoconazole to 0.0094 lb. azoxystrobin/ 0.0058 lb. difenoconazole)	13.1 to 31.6 (0.17 lb. azoxystrobin/ 0.1 lb. difenoconazole to 0.41 lb. azoxystrobin/ 0.25 lb. difenoconazole)	14	Use preventatively. Begin applications when conditions are favorable for disease infection, prior to disease symptom development. For optimum disease control, alternate Azoxy D Select with contact fungicides such as Daconil® brand products.
Brown Patch and Large Patch/Zoysia Patch (<i>Rhizoctonia solani</i>)	0.3 to 0.725 (0.0039 lb. azoxystrobin/ 0.0024 lb. difenoconazole to 0.0094 lb. azoxystrobin/ 0.0058 lb. difenoconazole)	13.1 to 31.6 (0.17 lb. azoxystrobin/ 0.1 lb. difenoconazole to 0.41 lb. azoxystrobin/ 0.25 lb. difenoconazole)	14-28	For cool-season turfgrasses, apply when conditions are favorable for disease development. For large patch of all warm-season turfgrasses, make 1 or 2 applications in fall prior to infection or when conditions are favorable for infection. A spring application prior to green-up may be required based on length of the infection period. Initiate first application for large patch/zoysia patch when soil temperature at a 2-4" depth averages 65° F.
Leaf and Sheath Spot (<i>Waitea circinata</i> var. <i>zeae</i> / <i>Chrysorhiza zeae</i> , formerly <i>R. zeae</i>)	0.5 to 0.725 (0.0065 lb. azoxystrobin/ 0.004 lb. difenoconazole to 0.0094 lb. azoxystrobin/	21.8 to 31.6 (0.28 lb. azoxystrobin/ 0.17 lb. difenoconazole to 0.41 lb. azoxystrobin/	14-28	For leaf and sheath spot, apply when conditions are favorable for infection such as sequential days, periods of temperatures at or above 90°F. Curative control may necessitate several applications.

	0.0058 lb. difenoconazole)	0.25 lb. difenoconazole)		Use higher rates at shorter intervals for curative control. Direct spray applications at crown of turfgrass.
Brown Ring Patch (<i>Waitea circinata</i> var. <i>circinata</i>)	0.5 to 0.725 (0.0065 lb. azoxystrobin/ 0.004 lb. difenoconazole to 0.0094 lb. azoxystrobin/ 0.0058 lb. difenoconazole)	21.8 to 31.6 (0.28 lb. azoxystrobin/ 0.17 lb. difenoconazole to 0.41 lb. azoxystrobin/ 0.25 lb. difenoconazole)	14-28	Apply when conditions are favorable for disease development.
Cool Weather Brown Patch, Yellow Patch (<i>Ceratorhiza cerealis</i> , formerly <i>Rhizoctonia cerealis</i>)	0.5 to 0.725 (0.0065 lb. azoxystrobin/ 0.004 lb. difenoconazole to 0.0094 lb. azoxystrobin/ 0.0058 lb. difenoconazole)	21.8 to 31.6 (0.28 lb. azoxystrobin/ 0.17 lb. difenoconazole to 0.41 lb. azoxystrobin/ 0.25 lb. difenoconazole)	14-28	Make 1 or 2 applications in fall or when conditions are favorable for disease development.
Dollar Spot (<i>Sclerotinia homoeocarpa</i>)	0.3 to 0.725 (0.0039 lb. azoxystrobin/ 0.0024 lb. difenoconazole to 0.0094 lb. azoxystrobin/ 0.0058 lb. difenoconazole)	13.1 to 31.6 (0.17 lb. azoxystrobin/ 0.1 lb. difenoconazole to 0.41 lb. azoxystrobin/ 0.25 lb. difenoconazole)	14-21	Apply preventatively when conditions are favorable for disease development. For optimum disease control, alternate Azoxy D Select with contact fungicides such as a Daconil brand product. If dollar spot is active, use higher rates and combine with a Daconil brand product.
Fairy Ring (<i>Lycoperdon</i> spp., <i>Agrocybe pediades</i> , and <i>Bovista plumbea</i>)	0.5 to 0.725 (0.0065 lb. azoxystrobin/ 0.004 lb. difenoconazole to 0.0094 lb. azoxystrobin/ 0.0058 lb. difenoconazole)	21.8 to 31.6 (0.17 lb. azoxystrobin/ 0.1 lb. difenoconazole to 0.41 lb. azoxystrobin/ 0.25 lb. difenoconazole)	14-28	For preventative control of fairy ring, apply early in the spring prior to the development of symptoms. Apply in 2-4 gallons of water per 1,000 sq. ft. Irrigate into the thatch prior to the spray drying. Repeat the application within 14 to 28 days after the first application. For curative control, apply as soon as possible after fairy ring symptoms develop. Apply in 2-4 gallons of water per 1,000 sq. ft. and irrigate lightly after application. Add the specified rate of a wetting agent to the final spray. Severely damaged or thin turf may require reseeding. Fairy ring symptoms may take 2 to 3 weeks to disappear following application. If area is hydrophobic, use wetting agents and irrigate prior to application(s) of Azoxy D Select . Reapplication after 28 days may be required in some cases.
Microdochium Patch (Formerly known as Fusarium Patch) (<i>Microdochium nivale</i>)	0.5 to 0.725 (0.0065 lb. azoxystrobin/ 0.004 lb. difenoconazole to 0.0094 lb. azoxystrobin/ 0.0058 lb. difenoconazole)	21.8 to 31.6 (0.17 lb. azoxystrobin/ 0.1 lb. difenoconazole to 0.41 lb. azoxystrobin/ 0.25 lb. difenoconazole)	14-28	Use preventatively. Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
*Gray Leaf Spot (<i>Pyricularia grisea</i>)	0.5 to 0.725 (0.0065 lb.	21.8 to 31.6 (0.17 lb.	14-21	Use Azoxy D Select in a preventative disease control program. Begin

	azoxystrobin/ 0.004 lb. difenoconazole to 0.0094 lb. azoxystrobin/ 0.0058 lb. difenoconazole)	azoxystrobin/ 0.1 lb. difenoconazole to 0.41 lb. azoxystrobin/ 0.25 lb. difenoconazole)		applications before disease is present and alternate with other fungicide chemistries that control gray leaf spot.
Leaf Rust Stem Rust Stripe Rust (<i>Puccinia</i> spp.)	0.5 to 0.725 (0.0065 lb. azoxystrobin/ 0.004 lb. difenoconazole to 0.0094 lb. azoxystrobin/ 0.0058 lb. difenoconazole)	21.8 to 31.6 (0.17 lb. azoxystrobin/ 0.1 lb. difenoconazole to 0.41 lb. azoxystrobin/ 0.25 lb. difenoconazole)	14-28	Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
Leaf Spot (<i>Bipolaris sorokiniana</i> , <i>Bipolaris cynodontis</i> , <i>bipolaris</i> spp.)	0.5 to 0.725 (0.0065 lb. azoxystrobin/ 0.004 lb. difenoconazole to 0.0094 lb. azoxystrobin/ 0.0058 lb. difenoconazole)	21.8 to 31.6 (0.17 lb. azoxystrobin/ 0.1 lb. difenoconazole to 0.41 lb. azoxystrobin/ 0.25 lb. difenoconazole)	14-21	Apply when conditions are favorable for disease development.
Melting Out (<i>Drechslera poae</i>)	0.5 to 0.725 (0.0065 lb. azoxystrobin/ 0.004 lb. difenoconazole to 0.0094 lb. azoxystrobin/ 0.0058 lb. difenoconazole)	21.8 to 31.6 (0.17 lb. azoxystrobin/ 0.1 lb. difenoconazole to 0.41 lb. azoxystrobin/ 0.25 lb. difenoconazole)	14-21	Apply when conditions are favorable for disease development
Necrotic Ring Spot (<i>Ophiosphaerella korrae</i>)	0.725 (0.0094 lb. azoxystrobin/ 0.0058 lb. difenoconazole)	31.6 (0.41 lb. azoxystrobin/ 0.25 lb. difenoconazole)	14-28	Apply when conditions are favorable for disease development
Pink Patch (<i>Limonomyces roseipellis</i>)	0.5 to 0.725 (0.0065 lb. azoxystrobin/ 0.004 lb. difenoconazole to 0.0094 lb. azoxystrobin/ 0.0058 lb. difenoconazole)	21.8 to 31.6 (0.17 lb. azoxystrobin/ 0.1 lb. difenoconazole to 0.41 lb. azoxystrobin/ 0.25 lb. difenoconazole)	14-28	Apply when conditions are favorable for disease development
Powdery Mildew (<i>Erysiphe graminis</i>)	0.5 to 0.725 (0.0065 lb. azoxystrobin/ 0.004 lb. difenoconazole to 0.0094 lb. azoxystrobin/ 0.0058 lb. difenoconazole)	21.8 to 31.6 (0.17 lb. azoxystrobin/ 0.1 lb. difenoconazole to 0.41 lb. azoxystrobin/ 0.25 lb. difenoconazole)	14-28	Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
Red Thread (<i>Laetisaria fuciformis</i>)	0.5 to 0.725 (0.0065 lb. azoxystrobin/ 0.004 lb. difenoconazole to 0.0094 lb. azoxystrobin/ 0.0058 lb. difenoconazole)	21.8 to 31.6 (0.17 lb. azoxystrobin/ 0.1 lb. difenoconazole to 0.41 lb. azoxystrobin/ 0.25 lb. difenoconazole)	14-28	Apply when conditions are favorable for disease development

Southern Blight (<i>Sclerotium rolfsii</i>)	0.5 to 0.725 (0.0065 lb. azoxystrobin/ 0.004 lb. difenoconazole to 0.0094 lb. azoxystrobin/ 0.0058 lb. difenoconazole)	21.8 to 31.6 (0.17 lb. azoxystrobin/ 0.1 lb. difenoconazole to 0.41 lb. azoxystrobin/ 0.25 lb. difenoconazole)	14-28	Apply when conditions are favorable for disease development
Summer Patch (<i>Magnaporthe poae</i>)	0.5 to 0.725 (0.0065 lb. azoxystrobin/ 0.004 lb. difenoconazole to 0.0094 lb. azoxystrobin/ 0.0058 lb. difenoconazole)	21.8 to 31.6 (0.17 lb. azoxystrobin/ 0.1 lb. difenoconazole to 0.41 lb. azoxystrobin/ 0.25 lb. difenoconazole)	14-28	Initiate applications when soil temperatures reach 65°F at a 2inch soil depth.
Take-All Patch (<i>Gaeumannomyces graminis var. avenae</i>)	0.5 to 0.725 (0.0065 lb. azoxystrobin/ 0.004 lb. difenoconazole to 0.0094 lb. azoxystrobin/ 0.0058 lb. difenoconazole)	21.8 to 31.6 (0.17 lb. azoxystrobin/ 0.1 lb. difenoconazole to 0.41 lb. azoxystrobin/ 0.25 lb. difenoconazole)	28	Begin applications prior to disease symptom development. Make two applications 28 days apart in the spring and two applications 28 days apart in the fall.

Restrictions

- ***DO NOT** apply more than two sequential applications of **Azoxystrobin D Select** for control of gray leaf spot at the 31.6 fl. oz. per acre rate. For all other diseases when gray leaf spot is not present, **DO NOT** apply more than three sequential applications of **Azoxystrobin D Select** at the 21.8 fl. oz. per acre rate.
- The annual limit for the active ingredient azoxystrobin in turf is 5.0 lbs. ai per acre per year (0.115 lb. a.i. per 1,000 square feet).
- The annual limit for the active ingredient difenoconazole in turf is 0.52 lb. ai per acre per year (0.012 lb. a.i. per 1,000 square feet).
- DO NOT** apply any product or products containing difenoconazole labeled for turf use in a manner that would exceed the annual limit of 0.52 lb. difenoconazole per acre per year.
- DO NOT** apply any product or products containing azoxystrobin labeled for turf use in a manner that would exceed the annual limit of 5.0 lbs. azoxystrobin per acre per year.
- One gallon of **Azoxystrobin D Select** (128 fl. oz.) contains 1.67 lbs. azoxystrobin and 1.05 lbs. difenoconazole.
- DO NOT** apply more than one-half gallon (64 fl. oz.) **Azoxystrobin D Select** per acre per year (1.47 fl. oz. **Azoxystrobin D Select** per 1,000 square feet per year). This annual rate delivers 0.84 lb. azoxystrobin/acre/year and the annual maximum of 0.52 lb. difenoconazole/acre/year.
- Additional applications of registered turf products containing azoxystrobin are allowed until the annual limit of 5.0 lbs. azoxystrobin per acre per year is achieved.
- Retreatment Interval (RTI): 14 days for all diseases except Take-All Patch (RTI: 28 days)
- DO NOT** apply more than 31.6 fl. oz. of **Azoxystrobin D Select** per acre (0.725 fl. oz. of **Azoxystrobin D Select** per 1,000 sq. ft.) in a single application.
- DO NOT** make more than 4 applications per year at reduced rate of 13.1 fl. oz. of **Azoxystrobin D Select** per acre (0.3 fl. oz. of **Azoxystrobin D Select** per 1,000 sq. ft.).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[For plastic containers ≤ 5 gallons: **Nonrefillable Container.** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix

tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

[For plastic containers > 5 gallons: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Prime Source, LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Prime Source, LLC and Seller harmless for any claims relating to such factors. Prime Source, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Prime Source, LLC, and Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, PRIME SOURCE, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Prime Source, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF PRIME SOURCE, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF PRIME SOURCE, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Prime Source, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Prime Source, LLC.

[Azoxy D Select is a trademark of Prime Source, LLC.]

[Briskway™, Daconil® (Chlorothalonil), Subdue® MAXX, the ALLIANCE FRAME the SYNGENTA Logo, and the PURPOSE ICON are Trademarks of a Syngenta Group Company.]

[EPA approval date]