

# UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

October 16, 2015

Kristin Hood Regulatory Affairs Specialist Central Garden & Pet Company 1501 East Woodfield Road, Suite 200 West Schaumburg, IL 60173

# Subject: Notification per PRN 98-10 – ABN and other label changes Product Name: PRENTOX PYRONYL OIL CONCENTRATE # OR-3610-A EPA Registration Number: 89459-27 Application Date: 10/6/2015 Decision Number: 509884

Dear Ms. Hood:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped "Notification" and will be placed in our records.

The alternate brand name "Pyronyl UL-300 Oil Concentrate" has been added to the product record.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you have any questions, you may contact Angela Hollis at 703-347-0216 or via email at <u>hollis.angela@epa.gov</u>.

Page 2 of 2 EPA Reg. No. 89459-27 Decision No. 509884

Sincerely,

V BD For

Jennifer Urbanski, Ph.D., Product Manager 4 Invertebrate and Vertebrate Branch 1 Registration Division (7505P) Office of Pesticide Programs FRONT PANEL



**KEEP THIS BOOKLET WITH CONTAINER AT ALL TIMES** 

# PYRONYL<sup>™</sup> OIL CONCENTRATE # OR-3610-A

Alternate brand names:

Prentox® Pyronyl UL-300 Pyronyl UL-300 Oil Concentrate

Controls: Ants, Confused Flour Beetles, Cockroaches, Flies, Gnats, Granary Weevils, Hornets, Mosquitoes, Rice Weevils, Flying Moths, Wasps, and Other Listed Insect Pests For Use As: Diluted and Undiluted Indoor Spray, Undiluted Outdoor Ground Applications For Use In: Public Gathering Places, Tall Grass, Shrubbery and Around Lawns, Indoor Cracks and Crevices, Behind Shelves and Drawers and Around Containers of Stored Foods

# RECOMMENDED FOR COMMERCIAL/INDUSTRIAL/INSTITUTIONAL USE

Authorized by U.S.D.A for use in Federally inspected meat and poultry plants.

# **ACTIVE INGREDIENTS:**

Pyrethrins	<del>)0</del> %
Piperonyl Butoxide, Technical*	<del>)0</del> %
N-octyl bicycloheptene dicarboximide	<del>)0</del> %
OTHER INGREDIENTS**:	<del>)0</del> %
TOTAL	<del>)0</del> %

\*Consists of 4.80% (butylcarbityl) (6-proplylpiperonyl) ether and 1.20% related compounds.

\*\*Contains Petroleum Distillates

Effective in the control of insects listed on this label if the applicator follows the Directions for Use set forth in this label.

# KEEP OUT OF REACH OF CHILDREN

# CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS, FIRST AID STATEMENTS AND DIRECTIONS FOR USE INSIDE THIS BOOKLET

# NOTIFICATION

#### 89459-27

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

10/16/2016

EPA Reg. No.-655-50189459-27 EPA Est. No. Net Contents:

# **PRECAUTIONARY STATEMENTS**

# HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed. Keep out of reach of children. Avoid breathing spray mist. Avoid skin and eye contact. Wash thoroughly after handling and before eating or smoking. Do not allow children or pets to contact treated surfaces until spray has dried. Do not use on plants. Do not allow spray to contact food, feedstuffs or water supplies. Thoroughly wash dishes and food handling utensils contaminated with this product. Remove pets and cover fish bowls before spraying.

FIRST AID		
lf	Immediately call a poison control center or doctor.	
swallowed	<ul> <li>Do not induce vomiting unless told to do so by a Poison Control Center or physician.</li> </ul>	
	• Do not give <b>any</b> liquid to the person.	
	• Do not give anything by mouth to an unconscious or convulsing person.	
If on skin or	Take off contaminated clothing.	
clothing	• Rinse skin immediately with plenty of water for 15-20 minutes.	
	Call a poison control center or doctor for treatment advice.	
If in eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes.	
	Remove contact lenses, if present, after the first 5 minutes, then continue	
	rinsing eye.	
	Call a poison control center or doctor for treatment advice.	
If inhaled	Move person to fresh <b>air.</b>	
	<ul> <li>If person is not breathing, call 911 or an ambulance, then give artificial</li> </ul>	
	respiration, preferably mouth-to-mouth if possible.	
	Call a poison control center or doctor for treatment advice.	
NOTE TO PHYSI	<b>CIAN:</b> Contains petroleum distillate – vomiting may cause aspiration pneumonia.	
Have the produ	Have the product container or label with you when calling a poison control center or physician, or going	
for treatment. You may contact 1-800-248-7763 257-4118 for emergency medical treatment		
information.		

For Chemical Spill, Leak, Fire, or Exposure, Call: CHEMTREC 1-800-424-9300

For Medical Emergencies Only, Call: 866-257-4118

## **ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds, and other wildlife. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater. Apply this product only as specified on this label.

## PHYSICAL HAZARDS

Do not use, or store near heat or open flame. Exposure to temperatures above 130° F may cause bursting.

# DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**INDOOR USE AS A SPRAY UNDILUTED:** The use of this product in food processing or food handling establishments should be confined to time periods when the plants or facilities are not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Close windows and doors and shut off ventilating systems. Use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at a rate of ½ fluid ounce per 1,000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary. Sweep up and destroy fallen insects.

**FOR COCKROACHES ONLY:** Apply at a rate of 1 fluid ounce per 1,000 cubic feet of space. Open all closets, cabinets, drawers, and other potential harborages. Direct spray toward ceiling and upper corners of the area and into, under and behind obstructions, cabinets, appliances and furnishings if the equipment and constructions of the treatment so allow. Keep area closed for at least 1 hour before ventilation. Repeat treatment as necessary. Sweep up and destroy fallen insects.

This method of application will control ROACHES and exposed, accessible stages of FLIES (including FRUIT FLIES), MOSQUITOES, SMALL FLYING MOTHS, GNATS, WASPS, HORNETS, CLOVER MITES, CHEESE MITES, CHEESE SKIPPERS, BOXELDER BUGS, EARWIGS, ANTS and the following common stored product pests: GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, SPIDER BEETLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, ANGOUMOIS GRAIN MOTHS, MEDITERRANEAN FLOUR MOTHS, INDIAN MEAL MOTHS, TOBACCO MOTHS, YELLOW MEALWORMS, DARKMEALWORMS, GRAIN MITES, CADELLES, and RED FLOUR BEETLES.

**CONTACT AND SURFACE SPRAY:** For the control of roaches, dilute 4 ¼ fluid ounces to make one gallon of finished solution in oil. Use a good sprayer adjusted to deliver a coarse, wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around bases of machinery, behind shelves and drawers. Spray roaches directly whenever possible. Repeat application as necessary.

**DRIED FRUIT PRODUCTS** in storage or being processed, for control of CONFUSED FLOUR BEETLES, DRIED FRUIT BEETLES, INDIAN MEAL MOTHS, and SAW-TOOTHED GRAIN BEETLES: Remove and destroy infested products. As noted above disperse this product as a fine mist in the air above the trays and shelves. Do not apply to fruit directly. Use <u>one-half-½ fluid</u> ounce per 1,000 cubic feet of space. Repeat application at this dosage in any given month. Leave the room closed for one hour. Do not remain in treated areas and ventilate the area before re-entry. Sweep up and destroy fallen insects.

**STORED FOOD:** (Stored in multi-wall bags or cloth bags) in warehouses, storage rooms, flour mills, grain elevators, granaries for control of ANGOUMOIS GRAIN MOTHS, CADELLES, CONFUSED FLOUR BEETLES, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS, RICE WEEVILS, SAW-TOOTHED GRAIN BEETLES, YELLOW MEALWORMS, CHEESE MITES, CHEESE SKIPPERS, CIGARETTE BEETLES, GRAIN MITES, GRANARY WEEVILS, DARK MEALWORMS, RED FLOUR BEETLES: Remove and destroy infested products. As noted above, disperse this product first by directing spray into corners, under pallets, and around stacks of packaged foods hitting as many insects as possible and then treat as a space spray. Infested store products should be fumigated or treated by other effective methods using an approved product intended for this purpose. Use a dosage of ½one-half fluid ounce per 1,000 cubic feet of space. Do not remain in treated areas and ventilate area before re-entry. This application will kill accessible, expose stages of those stored product pests listed above. Sweep up and destroy fallen insects.

**INDOOR USE AS A SPACE SPRAY, DILUTED:** For use in conventional mechanical fogging equipment to kill FLIES, FRUIT FLIES, MOSQUITOES, SMALL FLYING MOTHS, AND GNATS: Cover or remove exposed food and food handling surfaces. Close room and shut off all air conditioning or ventilating system equipment. Dilute 1 part concentrate plus 29 parts oil (refined kerosene) and mix well. Apply at the rate of 1 to 2 fluid ounces per 1,000 cubic feet filling the room with mist. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when re-infestation occurs.

**SURFACE SPRAY:** To insure maximum control of crawling insects (ANTS, BOXELDER BUGS, CLOVER MITES, DRUGSTORE BEETLES, EARWIGS, LICE, SPIDER BEETLES), dilute 1 part concentrate plus 29 parts oil (refined kerosene) and mix well and apply as a coarse wetting spray using approximately 1 gallon of spray mixture to 750 square feet of surface. Spray walls, floor and other surfaces of bins and storage and handling areas thoroughly, being especially careful to treat cracks, crevices and similar hiding places. Treat unloading, handling and processing areas and inside conveying processing and handling equipment. Spray thoroughly around the base of heavy machinery and equipment. Spray insects directly when possible.

For in transit protection, clean and spray truck beds, boxcars, and ships' holds before loading. Thoroughly treat walls and floors and spray each layer of cartons or containers lightly as stacked.

To control INDIAN MEAL MOTHS and ANGOUMOIS GRAIN MOTHS in stored grain, apply to the surface of binned grain at a rate of 1 gallon per 1,000 square feet. Apply at monthly intervals with the first appearance of moths and continuing until early fall. In case of severe infestation, break up webbing with rake before spraying and make a second application in two weeks.

As an aid in control of MOSQUITOES, GNATS, WASPS, AND SKIPPERS: treat walls, ceilings, moldings, screens, door and window frames, beams, light cords and similar resting places.

**ANIMAL QUARTER USE:** (Cattle barns, horse barns, poultry houses, swine houses, zoos): As a space spray diluted for use in conventional mechanical fogging equipment to kill FLIES, MOSQUITOES, SMALL FLYING MOTHS, AND GNATS, dilute 1 part concentrate plus 29 parts oils (refined kerosene) and mix well. Apply at the rate of 2 fluid ounces per 1,000 cubic feet of space above the animals. Direct spray towards the upper portions of the enclosure filling the room with mist. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when re-infestation occurs.

**FOR USE IN FEDERALLY INSPECTED MEAT AND POULTRY PLANTS:** Dilute one part concentrate with 3 parts base oil (refined kerosene) and use with a ULV spraying device at the rate of one fluid ounce per 1,000 cubic feet of room space. Apply the spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects shelter. Contact as many insects as possible. Repeat as necessary for control of exposed or accessible stages of ROACHES, SILVERFISH, WATERBUGS, EARWIGS, ANTS, SPIDERS, FLIES, FRUIT FLIES, MOSQUITOES, SMALL MOTHS AND GNATS. Before using exposed food products and packaging materials must be covered or removed. After use, equipment and utensils must be washed with and effective cleaning compound and rinsed with potable water after use of the product. Use as a space spray only when plant is not in operation.

**OUTDOOR GROUND APPLICATION, UNDILUTED:** To control ADULT MOSQUITOES, GNATS AND SMALL FLYING MOTHS, for best results, apply when there is a light breeze of approximately 5 mph and temperature is cool (75° or less). With a ULV spraying device, spray at a dosage rate of 4-5 fluid ounces per minute up-wind of the area to be controlled. If the equipment is truck mounted, keep the speed to approximately 5 mph with a swath width of no more than 300 feet or apply at a vehicle speed of 10 mph and flow rate of 8-10 fl. ounces per minute. To control MOSQUITO LARVAE, dilute at the rate of 15 fluid ounces per 50 gallons of oil and apply at the rate of 20-25 gallons per acre. Apply as a coarse wetting spray to the border or on shallow water of wet lands, swamps and marshes where mosquito larvae are known to breed.

**TEMPORARY REDUCTION OF ANNOYANCE** from FLIES, MOSQUITOES AND FLYING MOTHS OUTDOORS. The aforementioned application for outdoor ground application will afford temporary reduction of annoyance from these pests in public gathering places (drive-in restaurants, drive-in theaters, golf courses, municipalities, parks, playgrounds and recreational areas). Direct application into tall grass, shrubbery and around lawns where these pests may hover or rest, apply while air is still. Avoid wetting foliage. Application should be made prior to attendance. Repeat as necessary.

In additional outdoor areas (corrals, feed lots, swine lots and zoos), cover water, drinking fountains and animal feed before use. Fill area with mist, directing application into tall grass, shrubbery and around lawns where these pests may hover or rest, apply while air is still. Avoid wetting foliage. In zoos, avoid exposure of reptiles to the product. Repeat as necessary.

**FOR USE ON ANIMALS:** To protect beef and dairy cattle, and horses from HORN FLIES, HOUSE FLIES, MOSQUITOES, AND GNATS, dilute 1 part of concentrate plus 29 parts oil (refined kerosene) and mix well. Apply a light mist, sufficient to wet the tips of the hair. To control STABLE FLIES, HORSE FLIES, and DEER FLIES on beef and dairy cattle and horses, apply 2 fluid ounces per adult animal, sufficient to wet the hair but not to soak the hide. Repeat treatment once or twice daily or at intervals to give continued protection.

To control the motile stages of BLOOD SUCKING LICE on beef and dairy cattle, apply the spray to thoroughly wet the hair. Use 2 fluid ounces or less per animal and repeat 2 to 3 weeks as required.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. **STORAGE:** If container is damaged: Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material, such as clay, diatomaceous earth, sand, or sawdust. Sweep the contaminated absorbent on a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Place any leaking container into a similar drum or glass container. Do not store, use, pour, or spill near heat or open flame.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available or reconditioning, if appropriate. Clean container promptly after emptying.

[Containers 5 gallons or less] Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with oil used to dilute this product and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**[Container greater than 5 gallons]** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with oil used to dilute the product. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store for later use or disposal. Repeat this procedure two more times.

#### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY NOTICE:

Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Prentiss LLC Central Garden & Pet Company or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Central Garden & Pet CompanyPrentiss LLC and Seller harmless for any claims relating to such factors. Central Garden & Pet Company Prentiss LLC-warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Central Garden & Pet CompanyPrentiss LLC, and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, CENTRAL GARDEN & PET COMPANYPRENTISS LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIES WARRANTY EXCEPT AS STATED ABOVE. To the extent consistent with applicable law, in no event shall Central Garden & Pet CompanyPrentiss LLC or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. To the extent consistent with applicable law, **THE EXCLUSIVE** REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CENTRAL GARDEN & PET COMPANYPRENTISS LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACBH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CENTRAL GARDEN & PET COMPANYPRENTISS LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Central Garden & Pet CompanyPrentiss LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Central Garden & Pet CompanyPrentiss LLC.

PRENTOX<sup>®</sup> - Registered Trademark of Prentiss LLCWellmark International
PYRONYL<sup>™</sup> - Trademark of Wellmark InternationalPrentiss LLC
Manufactured byfor:
PRENTISS LLC
Office: 501 Cascade Pointe Lane, Suite 103, Cary, NC 27513
Central Garden & Pet Company
1501 East Woodfield Road 200W
Schaumburg, IL 60173