



## OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

November 6, 2025

Noelle Sanfilippo  
Regulatory Affairs Specialist  
Central Garden & Pet Company  
1501 East Woodfield Road, 200W  
Schaumburg, Illinois 60173

Subject: Label Amendment - Registration Review Mitigation for Permethrin  
Product Name: PRENTOX PERM-X AQUA 30-30  
EPA Registration Number: 89459-50  
Case Number: 478139  
Application Date: 3/16/2021

Dear Noelle Sanfilippo,

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Permethrin Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for

shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Alex McKee by phone at 202-566-1939, or via email at [mckee.alex@epa.gov](mailto:mckee.alex@epa.gov).

Sincerely,

A handwritten signature in blue ink that reads "Dana L. Friedman". The signature is written in a cursive style with a large, stylized 'D' and 'F'.

Dana L. Friedman, Chief  
Risk Management and Implementation Branch 1  
Pesticide Re-Evaluation Division  
Office of Pesticide Programs  
U.S. Environmental Protection Agency

ENCLOSURE: Stamped label

**ACCEPTED**

Nov 06, 2025

Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under

EPA Reg. No. 89459-50

**PERMETHRIN**

**GROUP**

**3A**

**INSECTICIDE**

## **PRENTOX PERM-X AQUA 30-30**

For use only by federal, state, tribal or local government officials responsible for public health or vector control, or by persons certified in the appropriate category or otherwise authorized by the state or tribal lead pesticide regulatory agency to perform adult mosquito control applications or by persons under their direct supervision

### **A Synergized Permethrin Formulation**

- For effective control of adult mosquitoes (including organophosphate resistant mosquitoes), midges (biting and non-biting) and black flies
- For ULV, thermal and barrier applications using ground and aerial equipment for control of adult mosquitoes and other biting flies
- Formulation allows dilution with water or oil
- May be applied undiluted
- For surface and barrier applications using handheld equipment
- Quick knockdown and effective control of target insects
- For ULV applications at rates up to 0.007 lb ai per acre
- For barrier application at rates up to 0.1 lb a.i. per acre
- Compatible with injector mixing systems or other insecticide mixing systems

### **ACTIVE INGREDIENTS:**

Permethrin\*: ..... 30%

Piperonyl Butoxide\*\* ..... 30%

**OTHER INGREDIENTS:** ..... 40%

**TOTAL** ..... 100%

\* Cis/trans ratio: Min. 35% ± cis and max. 65% ± trans

\*\* (butylcarbityl)(6-propylpiperonyl) ether and related compounds

Contains 2.52 lbs. a.i. Permethrin per gallon and 2.52 lbs. a.i. Piperonyl Butoxide per gallon

## **KEEP OUT OF REACH OF CHILDREN CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label find someone to explain it to you in detail.)

See inside label booklet for First Aid, Precautionary Statements, and Directions for Use including Storage and Disposal instructions

For chemical spill, leak, fire, or exposure, call: CHEMTREC 1-800-424-9300

For Medical Emergencies only, call: 800-xxx-xxxx

EPA Reg. No. 89459-50

EPA Est. No.

Manufactured for:

Central Garden & Pet Company

1501 East Woodfield Road 200W

Schaumburg, IL 60173

Batch/Lot No.

Net Contents:

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove contaminated clothing and wash before reuse.

FIRST AID	
If swallowed:	<ul style="list-style-type: none"><li>• Immediately call a poison control center or doctor.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
If on skin or clothing:	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15- 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
HOT LINE NUMBER	
Have this product container or label with you when calling a poison control center or doctor or going for treatment. You may contact 1-800-248-7763 for emergency medical treatment information. .	

### Personal Protective Equipment (PPE):

Mixers, loaders, applicators and other handlers including must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves (barrier laminate, viton ≥ 14 mils, neoprene rubber ≥ 14 mils, or nitrile rubber ≥ 14 mils for all handlers except for applicators using motorized ground equipment, pilots, and flaggers
- Chemical-resistant apron for mixers/loaders, persons cleaning equipment and persons exposed to the concentrate

See Engineering Controls for additional requirements.

### User Safety Requirements:

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

### Engineering Controls:

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)]. Human flagging is prohibited. Flagging to support aerial application is limited to use of the Global Positioning System (GPS) or mechanical flaggers.

### User Safety Recommendations:

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Runoff from treated areas or deposition of spray droplets into a body of water may be hazardous to fish and aquatic invertebrates. Do not apply over bodies of water (lakes, rivers, permanent streams, natural ponds, commercial fish ponds, swamps, marshes or estuaries), except when necessary to target areas where adult mosquitoes are present, and weather conditions will facilitate movement of applied material away from the water in order to minimize incidental deposition into the water body. Do not contaminate bodies of water when disposing of equipment rinsate or washwaters.

**Bee Warning:** This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment areas, except when applications are made to prevent or control a threat to public and/or animal health determined by a state, tribal or local health or vector control agency on the basis of documented evidence of disease causing agents in vector mosquitoes or the occurrence of mosquito-borne disease in animal or human populations, or if specifically approved by the state or tribe during a natural disaster recovery effort..

## PHYSICAL AND CHEMICAL HAZARDS

Combustible. Do not use, pour, spill, or store near heat or open flame.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### RESISTANCE-MANAGEMENT

For resistance management, Prentox Perm-X Aqua 30-30 contains a Group 3A insecticide. Any insect population may contain individuals naturally resistant to Prentox Perm-X Aqua 30-30 and other Group 3A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly. Appropriate resistance-management strategies should be followed. To delay insecticide resistance, take the following steps:

- Rotate the use of Prentox Perm-X Aqua 30-30 or other Group 3A insecticides with different groups that control the same pests.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, record keeping and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance, contact Central Garden & Pet Company at 1-800-248-7763.

## HOW TO APPLY AND WHERE TO USE

For outdoor use only. This product is designed for application as an Ultra Low Volume (ULV) or Thermal aerosol to control adult mosquitoes, and flies of nuisance, public health or disease vector importance in areas including residential areas, industrial areas, commercial areas, urban areas, parks, campgrounds, woodlands, athletic fields, golf courses, playgrounds, recreational areas, overgrown waste areas, roadsides, and other areas where adult mosquitoes and flies may be found.

Apply this product with ULV cold aerosol generators, thermal aerosol generators, mist-blowers, backpack sprayers, or handheld sprayers. Mist-blowers, backpack sprayers, handheld sprayers and low-pressure hand sprayers may be used for barrier or surface applications.

FOR BEST RESULTS, treat when mosquitoes or insects are most active and weather conditions are conducive to keeping the fog near the ground, e.g., cool temperatures, and consistent wind speeds are not greater than 10 mph. Applications during the cool hours of night or early morning are usually preferable. Apply this product only when wind speed is greater than 1 mph.

Do not retreat the same site more than once in 3 days. Do not exceed 25 applications at 0.007 lbs permethrin per acre per application or 0.025 lbs PBO per acre per application, or 0.18 lb of permethrin per acre per season or 2.0 lb PBO per acre per year. When targeting *Aedes taeniorhynchus* and other difficult species, applications of PBO may be made up to 0.08 lbs PBO per acre. More frequent treatments may be made to prevent or control a threat to public and/or animal health determined by a state, tribal or local health or vector control agency on the basis of documented evidence of disease causing agents in vector mosquitoes or the occurrence of mosquito-borne disease in animal or human populations, or if specifically approved by the state or tribe during a natural disaster recovery effort.

Apply this product over specific growing crops and range grasses prior to harvest for the control of adult mosquitoes and biting flies within or adjacent to these areas. Application can only be made where the following crops are present:

Alfalfa, Almonds, Apples, Artichokes - Globe, Asparagus, Avocados, Broccoli, Brussels sprouts, Cabbages, Cauliflower, Celery, Cherries, Corn, Fodder, Corn, Grain (field and pop), Corn - Stover, Corn - sweet kernel + cob with husks removed, , Eggplants, Filberts, Garlic, Kiwifruit, Horseradish, Leafy Vegetables (except Brassica), Lettuce - Head, Mushrooms, Onions -dry bulb, Peaches, Pears, Peppers - Bell, Pistachios, Potatoes, Range Grasses, Soybeans, Spinach, Tomatoes, Vegetables – Cucurbit, Walnuts, and Watercress.

Do not allow spray treatment to drift on cropland (other than crops listed) or water supplies. In treatment of corrals, feedlots, animal confinements/houses, swine lots and zoos, cover any exposed drinking water, drinking fountains and animal feed before application.

This product provides effective control of adult mosquitoes, black flies, gnats, biting and non-biting midges, stable flies, horse flies, deer flies, sheep flies, horn and heel flies, and nuisance flying insects such as house flies.

Air temperatures should be greater than 50° F when conducting all types of applications.

Do not apply directly to sewers or drains, or to any areas like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur, except as directed by this label. Not for use in outdoor residential misting systems. Not for use in metered release systems.

## **GROUND APPLICATION**

### **Vehicle-Mounted ULV Cold Aerosol Generators (Cold Fog)**

Apply through non-thermal ULV application equipment and base all acreage calculations on the equipment manufacturer's recommended swath width. Spray equipment must be adjusted so that the volume median diameter or mass medium diameter is less than 30 microns ( $D_{v0.5} < 30 \mu m$ ) and that 90% of the spray is contained in droplets smaller than 50 microns ( $D_{v0.9} < 50 \mu m$ ). Directions from the equipment manufacturer or vendor, pesticide registrant or a test facility using laser based measurement instrument must be used to adjust equipment to produce acceptable droplet size spectra. Application equipment must be tested at least annually to confirm that pressure at the nozzle and nozzle flow rate(s) are properly calibrated. Apply at a rate not to exceed 0.007 lb of permethrin per acre. An optimum swath is created when this product is applied from a truck being driven perpendicular to the wind direction. Direct the spray head of the equipment to ensure an optimum distribution of the spray cloud throughout the area. For best results, apply when insects are most active and meteorological conditions are conducive to keeping the spray cloud near the ground. An inversion of air temperatures and a light breeze of approximately 1 to 3 mph are preferable. During applications, consistent ground wind speeds should not exceed 10 mph. Application of this product during cooler hours of the night or early morning is recommended.

This product may be applied through all ULV Cold Aerosol Generators, or other similar equipment designed for non-thermal ULV aerosol applications. The desired application rate may be obtained from the chart below or by altering the dilution rate, the flow rate of the insecticide from the application equipment and the vehicle speed. Some examples are provided in the following chart. When an alternate rate is required, adjust the flow rate accordingly, provided the droplet size and spectrum remain within the label specifications.

FLOW RATE (Fluid Ounces Per Minute)						
Dilutions Based on a 15-Gallon Tank***						
Application Rate (Pounds of Permethrin A.I./Acre)	Vehicle Speed (MPH)	Undiluted	Diluted 1:4	Diluted 1:5	Diluted 1:6.5	Diluted 1:9
0.0015	10	0.46	2.32	2.77	3.50	4.7
	15	0.70	3.47	4.15	5.20	7.0
	20	0.93	4.64	5.60	7.00	9.3
0.00175	10	0.55	2.70	3.24	4.05	5.4
	15	0.82	4.05	4.84	6.07	8.1
	20	1.09	5.40	6.48	8.10	10.8
0.0035	10	1.09	5.40	6.48	8.10	10.8
	15	1.65	8.10	9.68	12.1	16.1
	20	2.19	10.8	12.9	16.1	21.5
0.007	10	2.19	10.7	12.9	16.1	21.5
	15	3.28	16.1	19.4	24.2	
	20	4.38	21.4	25.8		

\*\*\*Dilution Instructions: Fill tank with the appropriate amount of water for the desired dilution, then add this product and mix well. The use of metering pumps, injector systems or other mixing systems is optional. Note: When using oil as a diluent, agitate well before use.

When dense vegetation is present, use higher rates of application and/or a slower speed for best results.

**BACKPACK OR HANDHELD SPRAYER APPLICATION:** Apply through non-thermal ULV application equipment and base acreage calculations on a 300-foot swath. To apply undiluted at 0.007 lb of permethrin per acre, adjust equipment to an output rate of 0.86 ounces per minute and walk at 4 mph. To apply 0.0035 lb. of permethrin per acre, dilute 1 part of this product with 6 parts of diluent and apply at a rate of 3 ounces per minute and walk at 4 mph.

**THERMAL AEROSOL APPLICATION:** For use in truck-mounted, handheld or other thermal fogging equipment. Spray equipment must be adjusted so that the volume median diameter or mass median diameter is less than 30 microns ( $D_{v0.5} < 30 \mu\text{m}$ ) and that 90% of the spray is contained in droplets smaller than 48 microns ( $D_{v0.9} < 48 \mu\text{m}$ ). Directions from the equipment manufacturer or vendor, pesticide registrant or a test facility using a laser-based measurement instrument must be used to adjust equipment to produce acceptable droplet size spectra. Application equipment must be tested at least annually to confirm that pressure at the nozzle and nozzle flow rate(s) are properly calibrated. Apply this product mixed with water or oil at a rate not to exceed 0.007 lb per acre. Base acreage calculations on 300 feet swath. For best results, direct the thermal fog into and allow to drift through areas where adult mosquitoes are found.

**BARRIER APPLICATIONS:** To control resting or flying adult mosquitoes in areas including animal quarters, roadsides, urban areas, residential areas, commercial areas, industrial areas, golf courses, parks, recreational areas, tire piles and all public areas where adult mosquitoes occur or find harborage. May be used to treat aggregation and harborage areas such as the perimeter of residential buildings, commercial buildings, storage buildings, crawlspaces, transport trailers, abandoned or uninhabited buildings and other areas where adult mosquitoes find harborage.

To apply this product using low pressure hand sprayers or other similar equipment, for the control of mosquitoes on resting or aggregation surfaces, mix 0.0875 oz (2.59 ml) in one gallon of water and apply sufficient spray to dampen the surface – approximately 700-750 square feet of treated area. Use equipment that is capable of delivering an even distribution of the labeled rate over the treatment area. For application using backpack or truck-mounted mist-blower equipment, apply this product undiluted at a rate not to exceed 5.08 oz per acre. If diluted, apply at a rate not to exceed 0.1 lb permethrin a.i. per acre. Diluted or undiluted applications may be made onto foliage at the perimeter of areas such as recreational areas, parks, or golf courses from which mosquitoes or flies are to be excluded. Applications may be directed into harborage areas or other resting areas where adult mosquitoes or flies may be found.

**URBAN ULV MOSQUITO CONTROL APPLICATIONS:** For control of resting or flying adult mosquitoes, biting flies and biting midges in harborage areas such as but not limited to utility tunnels, sewers, storm drains and catch basins, pipe tunnels, basements, underground passages, parking decks, crawlspaces under buildings, abandoned buildings or uninhabited buildings. Apply this product diluted or undiluted using mechanical foggers, handheld or truck-mounted ULV equipment, thermal foggers or other spray equipment suitable for this type of application. This product may be applied at rates up to but not exceeding 0.007 lb permethrin per acre.

#### **MANDATORY SPRAY DRIFT MANAGEMENT**

##### **Ultra Low Volume Applications**

Apply when ground wind speeds are equal or to greater than 1 mph.

All types of applications should be conducted when temperatures at ground level are at or above 50°F.

##### **For Ground Applications:**

- Create an optimum swath when possible. An optimum swath width can be achieved when Prentox Perm-X Aqua 30-30 is applied from a truck that is being driven perpendicular to the wind directions. Direct the spray head of equipment to ensure even distribution of the spray cloud throughout the area.
- FOR BEST RESULTS treat when mosquitoes or insects are most active and weather conditions are conducive to keeping the spray cloud in the air column close to the ground.
- An inversion of air temperatures and a light breeze is preferable. Application during the cooler hours of the night or early morning is recommended.

##### **For Aerial Applications:**

- Do not apply by fixed wing aircraft at a nozzle height less than 100 feet (30.5 m) above ground or canopy, or by helicopter at a height less than 75 feet (22.9 m) above the ground or canopy, unless specifically approved by the state or tribe based on public health needs.

Adult mosquito control applications should be limited to trained personnel.

For use only by federal, state, tribal or local government officials responsible for public health or vector control, or by persons certified in the appropriate category or otherwise authorized by the state or tribal lead pesticide regulatory agency to perform adult mosquito control application or by persons under their supervision, or as allowed by state regulations for persons treating private property.

This pesticide is extremely toxic to aquatic organisms. Runoff from treated areas or deposition of spray droplets into a body of water may be hazardous to aquatic organisms.

Do not apply over bodies of water (lakes, rivers, permanent streams, natural ponds, commercial fish ponds, swamps, marshes, or estuaries), except when necessary to target areas where adult mosquitoes are present, and weather conditions will facilitate movement of applied material beyond the body of water to minimize incidental deposition into the water body. Do not contaminate bodies of water when disposing of equipment rinsate or washwaters.

Before making the first application in a season, it is advisable to consult with the state or tribal Agency with primary responsibility for pesticide regulations to determine if other regulatory requirements exist.

Do not treat a site with more than 0.007 lb permethrin and 0.08 lb PBO per acre in a single application or in any 24-hour period. Do not exceed 0.18 lb. permethrin or 2 lb PBO in any site in one year. More frequent applications may be made to prevent or control a threat to public and/or animal health determined by a state, tribal, or local health or vector control Agency on the basis of documented evidence of disease-causing agents in vector mosquitoes or the occurrence of mosquito-borne disease in animal or human populations, or if specifically approved by the state or tribe during a natural disaster recovery effort.

#### **AERIAL APPLICATION**

Apply this product either diluted or undiluted at rates up to 0.007 lbs. permethrin/PBO per acre by fixed wing or rotary aircraft equipment with a spraying system capable of making a ULV application.

For aerial applications of this product, use ULV equipment and nozzles such as rotary atomizers or similar spray equipment. Spray equipment must be adjusted so that the volume median diameter produced is less than 60 microns ( $D_{v0.5} < 60 \mu\text{m}$ ) and that 90% of the spray is contained in droplets smaller than 115 microns ( $D_{v0.9} < 115 \mu\text{m}$ ). Directions from the equipment manufacturer or vendor, pesticide registrant or a test facility using a wind tunnel and laser-based measurement instrument must be used to adjust equipment to produce acceptable droplet size spectra. Application equipment must be tested at least annually to confirm that pressure at the nozzle and nozzle flow rate(s) are properly calibrated.



Aerial application of this product shall be made at altitudes ranging from 75 feet to 300 feet. Do not apply by fixed wing aircraft at a height less than 100 feet, or by helicopter at a height less than 75 feet unless specifically approved by the state or tribe based on public health needs. Do not apply when ground wind speed exceeds 10 mph. Aerial applications of this product shall only be made when recommended by public health officials and trained personnel of mosquito abatement districts and other mosquito control programs.

For best results, apply this product when insects are most active and meteorological conditions are conducive to keeping the spray cloud near the ground. Aerial applications using aircraft equipped with Global Positioning Systems (GPS) may be needed to compensate for windy conditions and to ensure drift into target areas.

#### **IN FLORIDA:**

Aerial applications of this product require trained personnel to perform industry accepted assays to monitor resistance formation in targeted mosquitoes.

### **STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak-up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

**PESTICIDE DISPOSAL:** Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container promptly after emptying. Container  $\geq$  5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Container  $>$  5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure-rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after flow begins to drip. Offer for recycling, if available.

### **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this Product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Central Garden & Pet Company or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Central Garden & Pet Company and Seller harmless for any claims relating to such factors.

Central Garden & Pet Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Central Garden & Pet Company, and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, CENTRAL GARDEN & PET COMPANY MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, in no event shall Central Garden & Pet Company or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. To the extent consistent with applicable law, **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CENTRAL GARDEN & PET COMPANY AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CENTRAL GARDEN & PET COMPANY OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Central Garden & Pet Company and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Central Garden & Pet Company.