



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

May 5, 2016

Kristin Hood  
Regulatory Affairs Specialist  
Central Garden & Pet Company  
1501 East Woodfield Road, Suite 200W  
Schaumburg, IL 60173

Subject: Notification per PRN 98-10 – Minor changes requested by California  
Product Name: EQUIL LAMBDA 9.7 CS INSECTICIDE  
EPA Registration Number: 89459-51  
Application Date: 3/22/2016  
Decision Number: 515318

Dear Ms. Hood:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “Notification” and will be placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, you may contact Angela Hollis at 703-347-0216 or via email at [hollis.angela@epa.gov](mailto:hollis.angela@epa.gov).

Sincerely,

A handwritten signature in black ink, appearing to read "Kable Bo Davis", enclosed within a hand-drawn oval.

Kable Bo Davis, Product Manager 3  
Invertebrate and Vertebrate Branch 1  
Registration Division (7505P)  
Office of Pesticide Programs

# Equil Lambda 9.7 CS Insecticide

Alternate brand name: Lambda 9.7 CS

(Contains Signa Technology) [OR (With CapVantage™ Technology)] [Optional Marketing Language]

Controls: Ants, Cockroaches, Crickets, Flies, Litter Beetles (adults/immature stages ~~such as~~ including Darkling, Hide, and Carrion), Mosquitoes, Spiders, Ticks and other listed insects

For Use in: Food Handling Establishments, Food Serving Areas, Livestock/Poultry Housing Structures, Lawns/Grass and Ornamentals

For use in, on and around buildings and structures for the control of listed pests, including on lawns, ornamental trees and shrubs around residential, institutional, public, commercial, agricultural and industrial buildings; and parks, recreational areas and athletic fields.

Refer to the label for additional use sites and pests controlled

GROUP	3	INSECTICIDE
-------	---	-------------

## ACTIVE INGREDIENT:

Lambda-cyhalothrin<sup>1</sup>:

[1  $\alpha$ (S\*),3  $\alpha$ (Z)]-(±)-cyano-(3-phenoxyphenyl)methyl-3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate.....9.7%

**OTHER INGREDIENTS:**.....90.3%

**TOTAL:** .....100.0%

<sup>1</sup>Synthetic pyrethroid

## KEEP OUT OF REACH OF CHILDREN

### CAUTION

See additional PRECAUTIONARY STATEMENTS and Directions for Use in booklet.  
Si usted no entiende la etiqueta, busque a alguien para que la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

EPA Reg. No. 89459-51

EPA Est. No.

Country of origin Austria

Net Contents:

## NOTIFICATION

89459-51

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

05/05/2016

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Causes moderate eye irritation. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum or using tobacco, or using the toilet.

<b>FIRST AID</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continuing rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Do not give any liquid to the person.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-347-8272 for emergency medical treatment information.	

**ENVIRONMENTAL HAZARDS**

This product is extremely toxic to fish. To protect the environment, do not allow pesticide to enter or runoff into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Apply this product only as specified on this label. When making applications, care should be used to avoid household pets, particularly fish and reptile pets.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

**PHYSICAL AND CHEMICAL HAZARDS**

Do not use this product in or on electrical equipment due to the possibility of shock hazard.

**General Information**

Equil Lambda 9.7 CS Insecticide is a unique formulation utilizing Sigma Technology [CapVantage™] which is a proprietary blend of ingredients for use with the active ingredient, Lambda-cyhalothrin.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**IMPORTANT:** Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior landscapes, ornamental gardens or parks, or on lawns or grounds.

This product is RESTRICTED for use in New York State.

### Application Restrictions for Residential Outdoor Surface and Space Sprays:

- All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:
  - (1) Treatment to soil or vegetation around structure;
  - (2) Applications to lawns, turf, and other vegetation;
  - (3) Application to building foundations, up to a maximum height of 3 feet.
- Other than applications to building foundations, all outdoor applications to impervious surfaces ~~such as including~~ sidewalks, driveways, patios, porches and structural surfaces (~~such as including~~ windows, doors, and eaves) are limited to spot and crack-and-crevice applications only.
- Do not water the treated area to the point of runoff.
- Do not make applications during rain.
- Do not apply directly to sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur, except as directed by this label.

### PERIMETER PEST CONTROL

This product is for use as a general surface (nonfood/nonfeed areas), crack and crevice, or spot treatment in, on, and around buildings and structures and their immediate surroundings, and on modes of transport. Areas of use include, industrial buildings, houses, patios, porches, closets, furniture, apartment buildings, laboratories, buses, greenhouses (non-commercial), stores, factories, warehouses, wineries, vessels, railcars, trucks, trailers, aircraft (cargo and other non-cabin areas only), schools, nursing homes, hospitals, mausoleums, restaurants, hotels, correctional facilities, livestock/poultry housing, pet kennels, food granaries, food grain mills and food manufacturing, processing, and servicing establishments.

Nonfood/nonfeed areas of food/feed handling establishments include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage (after canning or bottling).

For indoor applications, this product can be reapplied at 21-day intervals if necessary.

### Mixing Instructions

Dilute this product with water for application using hand-held or power-operated application equipment as a coarse spray for crack and crevice or spot and general surface (non-food/nonfeed areas only) treatments. Application equipment that delivers low volume treatments, such as the Micro-Injector<sup>®</sup> or Actisol<sup>®</sup> applicator, may also be used to make crack and crevice or spot and general surface treatments. Fill applicator tank with the desired volume of water and add this product. Close and shake before use in order to ensure proper mixing. Shake or reagitator applicator tank before use if application is interrupted. Mix only amount of treatment volume as required. This product may be applied by using a paint brush or other porous applicator attached to a handle as a general surface treatment.

### Tank Mixing

This product may be tank mixed with other currently registered pesticides unless expressly prohibited by the product label. A small volume mixing test with the other products is recommended to ensure compatibility. If other chemicals are added to the applicator tank, this product should be added last. If mixed with EC formulations, use within 24 hours. Fill tank to desired volume and continue to agitate while making applications.

This product may be tank mixed with an Insect Growth Regulator (IGR). Observe all restrictions and precautions which appear on the labels of these products.

### Foam Applications

This product may be converted to a foam and the foam used to treat structural voids to control or prevent pests including ants, bees, termites (above ground only), wasps, or other arthropods harboring in walls, under slabs, or in other void areas.

### RATES FOR PESTS (HAND APPLICATION EQUIPMENT)

Pests	Dilution Rate (Concentration of AI)	Specific Use Directions
Ants Bed Bugs (adult) Bees Boxelder Bugs Carpenter Bees Carpet Beetles Centipedes Cigarette Beetles Clover Mites Cockroaches <sup>1</sup> Confused Flour Beetles Crickets Earwigs Firebrats Fleas <sup>2</sup> Flies Lesser Grain Borers Millipedes Mosquitoes Red Flour Beetles Rice Weevils Saw-toothed Grain Beetles Silverfish Sowbugs Spiders Stink Bugs Ticks Wasps	0.2 fl oz (6 mL) to 0.4 fl oz (12 mL) per gallon of water  (0.015-0.030% AI)	<p><b>Ants:</b> Apply to any trails around doors and windows and other places where ants may be found. Locate and treat nests. Where ants are trailing inside, apply as a residual surface treatment to active areas <del>such as</del>including baseboards, corners, around pipes, in and behind cabinets, behind and under refrigerators, sinks, furnaces and stoves, cracks and crevices. When combining baits and residual surface insecticides, apply surface insecticides in cracks and crevices, along baseboards, and infested surfaces and outside barrier treatments. Use baits in other areas that are untreated by residual insecticides; also see <b>Outdoor Surfaces Use</b>.</p> <p><b>Bed Bugs:</b> Clean floors and surfaces by vacuuming. Apply as a coarse, low-pressure spray to harborage areas including crevices, baseboards, loose plaster, behind bed frames and headboards, beneath beds and furniture, and to bed springs and bed frames. Do not apply to furniture surfaces or mattresses where people will be laying or sitting. Infested bedding should not be treated, but should be removed, placed in sealed plastic bags, and taken for laundering and drying at high temperatures.</p> <p><b>Cockroaches, Crickets, Earwigs, Firebrats, Silverfish, Spiders and Stink Bugs:</b> Apply as a coarse, low pressure spray to areas where these pests hide, <del>such as</del>including baseboards, corners, storage areas, closets, around water pipes, doors and windows, attics and eaves, cabinets, behind and under refrigerators, sinks, furnaces and stoves, the underside of shelves, drawers and similar areas. Pay particular attention to cracks and crevices; also see <b>Outdoor Surfaces Use</b>.</p>
Cockroaches <sup>1</sup> Crickets <sup>6</sup> Flies <sup>5</sup> Litter Beetles <sup>3</sup> (adults/ immature stages <del>such</del> <del>as</del> including Darkling, Hide, and Carrion) Mosquitoes <sup>4</sup> Pillbugs Scorpions Spiders <sup>6</sup> Spider Mites (Two-spotted, Spruce) Ticks <sup>6</sup>	0.80 fl. oz. (24 mL) per gallon of water  (0.060% AI)	<p><b>Bees, Flies, Mosquitoes, and Wasps:</b> Apply directly to walls, ceilings, window screens, and other resting areas as a residual surface treatment. May be used inside residential buildings as well as in and around carports, garages, and storage sheds; also see <b>Outdoor Surfaces Use</b>. Use caution when treating nests of stinging insects as this product does not provide instant knockdown. Protective equipment for the applicator may be required. Treat bee, wasp and hornet nests late in the day when most insects will be present. Allow 2-3 days for the colony to die and repeat after 21 days, if necessary.</p> <p>Yards or other frequented areas enclosed by landscaping can benefit from the creation of a mosquito barrier to reduce invading mosquitoes by the treatment of <b>perimeter</b> vegetation. Apply this product at specified rates in 2-5 gallons of water per 1,000 sq ft.</p> <p><b>Carpenter Bees:</b> Apply coarse spray to thoroughly wet wood</p>

Pests	Dilution Rate (Concentration of AI)	Specific Use Directions
		<p>surfaces where bees have been previously active or to provide protection against further damage. Apply early in the spring to prevent bees from invading wood. When bees have infested wood, surface applications can help control embedded larvae and bees that emerge from the wood.</p> <p><b>Pantry Pests</b> (i.e., Carpet Beetles, Cigarette Beetles, Confused Flour Beetles, Lesser Grain Borers, Red Flour Beetles, Rice Weevils, and Saw-toothed Grain Beetles): Apply to cupboards, shelving, and storage areas. Remove all utensils, uncovered foodstuffs (or any having original package opened), and shelf paper before making application. Allow treated surfaces to dry and cover shelves with clean paper before replacing any utensils, foodstuffs, or other items. Any foodstuffs accidentally contaminated with treatment solution should be destroyed.</p> <p><b>Boxelder Bugs, Centipedes, Millipedes, Pillbugs, and Sowbugs:</b> Apply around doors and windows and other places where these pests may be found or where they may enter premises. Treat baseboards, storage areas, and other locations. Apply treatments to prevent infestation as described below; also see <b>Outdoor Surfaces Use</b>.</p> <p><b>Fleas and Ticks:</b> To control nuisance fleas and ticks (e.g., dog ticks) apply to kennels, yards, runs, and other areas where pets may frequent. To control ticks, apply using a coarse fan spray to vegetation brush, branches, rock walls, and other areas near habitation where ticks may harbor or frequent. Treat entire area rather than making spot treatments, and retreat as necessary to maintain control. Do not apply to pasture or cropland, and do not allow animals and people access to treated areas until the deposit has dried. Applications can begin in the spring and can continue until frost to control both larvae and adult ticks; also see <b>Outdoor Surfaces Use</b>.</p> <p><b>Cluster Flies:</b> Apply in late summer or early fall before flies are observed alighting on surfaces. Apply thoroughly on siding, under eaves, and around windows and doors, paying particular attention to south-facing surfaces, but do not allow runoff to occur. Heavy precipitation prior to frost may require retreatments to maintain protection. In winter and spring when flies become active and are emerging, interior crack and crevice and void treatments can help reduce the infestation, along with ULV or general surface application in infested attics or unoccupied lofts.</p> <p><b>Litter Beetles (Darkling, Hide, and Carrion Beetles) and Flies In Animal Housing (Such As Including Poultry Houses):</b> To control adult litter beetles, apply this product to walls and floors at cleanout, before reintroduction of animals. This will suppress beetles that escaped earlier treatment and will help delay onset of future infestations. Pay attention to areas where beetles frequently occur, such as including walls, supports, cages, stalls, and around feeders. To help control flies, apply a directed application to horizontal surfaces and overhead areas and allow to dry before reintroduction of animals; also see <b>Livestock/Poultry Housing Structures and Pet Kennels</b>.</p>
Termites (above ground only)	0.20 fl oz (6 mL) to 0.4 fl oz (12 mL) per	This product may be used on termites to kill workers or winged reproductive forms which may be present in treated channels at the time of treatment. These applications are

Pests	Dilution Rate (Concentration of AI)	Specific Use Directions
	gallon of water  (0.015 - 0.030% AI)	not a substitute for mechanical alteration to control subterranean termites, soil treatment labeled termiticides, or fumigation for extensive infestation of drywood termites or other wood-infesting insects, but are considered supplemental. This product is not recommended as sole protection against termites. For active termite infestations, get a professional inspection.
<sup>1</sup> For cockroaches, the recommended rate for maintenance treatments is 0.015% and for clean-out treatments is 0.030%. For control of severe infestations, use 0.060% rate. <sup>2</sup> For outdoor use only and use 0.030% rate. <sup>3</sup> For control of LIGHT beetle infestations, use 0.030% rate. <sup>4</sup> For residual control, use 0.060% rate. <sup>5</sup> In all states except California, rates for flies may be increased to 0.060% when environmental conditions are severe and/or populations are high. <sup>6</sup> For clean-out/severe infestations, use 0.060% rate.		

### Application Within Food Handling Establishments

For places other than private residences in which exposed food is held, processed, prepared, or served including, ~~but not limited to,~~ areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing foods, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups), and serving areas: Use as a crack and crevice or spot treatment in and around both food and nonfood areas. Apply in small amounts directly into cracks and crevices, using equipment capable of delivering a pin stream of insecticide, in points between different elements of construction, between equipment and floor, openings leading to voids and hollow spaces in walls, equipment and bases. Food contact surfaces and equipment should be cleaned with an effective cleaning compound and rinsed with potable water before using.

Limit individual spot treatments to an area no larger than 20% of the treated surface. Individual spot treatments may not exceed two sq ft. Take extreme care that the product is not introduced into the air. Avoid contamination of food and food processing surfaces.

### Application Within Food Serving Areas

For facilities where foods are served, ~~such as including~~ dining rooms: Apply as a crack and crevice or spot treatment to selective surfaces ~~including such as~~ baseboards, under elements of construction, and into cracks and crevices. Avoid treating surfaces likely to be contacted by food. (Do not apply when facility is in operation or foods are exposed.) Food must be covered or removed in area being treated. Do not apply directly to food or allow applications to contaminate food.

**NOTE:** Application of this product in the Food Areas and/or Food Serving Areas of Food Handling Establishments other than as a spot and/or crack and crevice treatment is not permitted. Use of application equipment such as the Micro-Injector® or Actisol® applicator in food areas should be limited to crack and crevice treatment only.

### Livestock/Poultry Housing Structures and Pet Kennels

Apply as a general surface (including directed sprays) and/or crack and crevice treatment. Make interior and exterior perimeter applications in and around the livestock, poultry, and pet housing structures. Normal cleaning of the structure must also be done along with applications of this product to effectively control the crawling and flying insect pests listed in the table.



With the exception of cattle and calves that may be present, only apply this product to livestock barns or housing structures when unoccupied by animals. Apply to floors, vertical, and overhead surfaces where crawling or flying insect pests are or may be present. Feeders, waterers, and feed carts should be covered before application to prevent contamination. Do not apply to milk rooms or feed rooms. Pay attention to animal areas including stanchions, pipes, windows and doors, and areas where insect pests hide or congregate. Exterior applications to south facing walls and foundation perimeters can help prevent interior infestations of flying and crawling insect pests.

For poultry houses, apply to floor area (birds grown on litter) or to walls, posts, and cage framing (birds grown in cages). Application should also be made into cracks and crevices around insulation. Reapply after each growout or sanitization procedure. Indoor control can be enhanced by making perimeter treatments around the outside of building foundations to prevent immigrating adult beetles. Apply in a uniform band 1 - 3 ft. up and 2 - 6 ft. out from foundation. Maintaining a year-round treatment program will prevent background populations from reaching problem levels.

#### **RESTRICTIONS:**

- **DO NOT** make interior applications of this product in areas where animals, except cattle and calves are present in the facility. Allow treated surfaces to completely dry before restocking the facility.
- **DO NOT** make applications to any animal feedstuffs, water, or watering equipment.
- **DO NOT** contaminate any animal food, feed, or water in and around livestock, poultry, or pet housing when making applications.

#### **Outdoor Surfaces Use**

For control of ants, bees, centipedes, cockroaches, crickets, fleas, flies, millipedes, mosquitoes, scorpions, sowbugs, stink bugs, pillbugs, spiders, termites (above ground only), ticks, wasps, and other similar perimeter arthropod pests.

Apply with either hand or power application equipment as a residual treatment to ornamental plants next to foundations of buildings and to surfaces of buildings, porches, screens, window frames, eaves, patios, garages, refuse dumps, and other similar areas where these insect pests are active.

For termites, this type of application is not intended as a substitute for soil treatment labeled termiticides or mechanical alteration to control subterranean termites, or fumigation for extensive infestation of drywood termites or other wood-infesting insects. The purpose of such application of this product for termites is to kill workers or winged reproductive forms which may be present in treated channels at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but are merely a supplement. This product is not recommended as sole protection against termites. For active termite infestations, get a professional inspection.

#### **Perimeter Treatments**

Apply a continual band of insecticide around building foundations and around windows, doors, service line entrances, eaves, vents, and other areas to reduce the potential for entry by crawling pests. Before application, remove debris and leaf litter from next to the foundation, cut back vegetation and branches that touch the foundation, and move or rake back rocks, deep mulch, or other potential pest harborage next to the foundation. Apply the band up to 10 ft. wide around the structure (or according to state regulations governing commercial pest control) and upwards along the foundation to 3 ft. and around windows, doors, and roof overhangs. Apply as a coarse spray to thoroughly and uniformly wet the foundation and/or band area so that the insecticide will reach the soil or thatch level where pests may be active.

Amount of concentrate is dependent upon pest species (see pest table and comments), infestation levels, and service interval desired.



**Rate Table For Perimeter Barrier Applications**

<b>Application Rate: Fl oz (mL) of Equil Lambda 9.7 CS Insecticide</b>	<b>Gallons of Water<sup>1</sup></b>	<b>Area of Coverage (sq. ft.)</b>
0.2 fl oz (6 mL)	1-5 gallons	800-1600 sq ft
0.4 fl oz (12 mL)	1-5 gallons	800-1600 sq ft
0.8 fl oz (24 mL)	1-5 gallons	800-1600 sq ft

<sup>1</sup>Application volume may be greater than 5 gallons/800-1600 sq ft if required under heavy vegetative or landscaping materials in order to obtain desired coverage.

**Examples Of Dilutions For Perimeter Barrier Applications**

<b>Application Volume Gallons of Solution per 1,000 sq ft</b>	<b>Application Rate Fl oz (mL) of Equil Lambda 9.7 CS Insecticide per 1,000 sq ft</b>	<b>Fl oz (mL) of Equil Lambda 9.7 CS Insecticide to Dilute in Water According to Spray Tank Volumes</b>		
		<b>5 gallons</b>	<b>10 gallons</b>	<b>50 gallons</b>
1 gallon per 1,000 sq ft	0.2 fl oz (6 mL) 0.4 fl oz (12 mL) 0.8 fl oz (24 mL)	1 fl oz (30 mL) 2 fl oz (60 mL) 4 fl oz (120 mL)	2 fl oz (60 mL) 4 fl oz (120 mL) 8 fl oz (240 mL)	10 fl oz (300 mL) 20 fl oz (600 mL) 40 fl oz (1200 mL)
2 gallons per 1,000 sq ft	0.2 fl oz (6 mL) 0.4 fl oz (12 mL) 0.8 fl oz (24 mL)	0.5 fl oz (15 mL) 1 fl oz (30 mL) 2 fl oz (60 mL)	1 fl oz (30 mL) 2 fl oz (60 mL) 4 fl oz (120 mL)	5 fl oz (150 mL) 10 fl oz (300 mL) 20 fl oz (600 mL)
5 gallons per 1,000 sq ft	0.2 fl oz (6 mL) 0.4 fl oz (12 mL) 0.8 fl oz (24 mL)	0.2 fl oz (6 mL) 0.4 fl oz (12 mL) 0.8 fl oz (24 mL)	0.4 fl oz (12 mL) 0.8 fl oz (24 mL) 1.6 fl oz (48 mL)	2 fl oz (60 mL) 4 fl oz (120 mL) 8 fl oz (240 mL)

**Example calculation:** To apply the mid-rate of this product at a volume of 5 gallons/1,000 sq ft, mix 4 fl oz of concentrate in 50 gallons of water. The percent active ingredient in the finished product dilution can be calculated with the following formula:

mL needed to add times 9.7% active in concentrate, divided by gallons finished dilution times 3785 mL/gallon = % active in dilution. (Example: 4 fl oz in 50 gallons is 120 mL times 9.7 equals 1164, and 50 gallons times 3785 is 189,250. Dividing 1164 by 189,250 equals 0.006% active in the tank dilution.)

**NOTE:** Do not use water based sprays of this product in conduits, motor housings, junction boxes, switch boxes, or other electrical equipment because of possible shock hazard. Thoroughly wash out sprayer and screen with water and detergent before using this product. This product has not stained or caused damage to painted or varnished surfaces, plastics, fabrics, or other surfaces where water applied alone causes no damage. However, treat a small area and allow to dry, to determine whether staining will occur.

**LET TREATED SURFACES DRY BEFORE ALLOWING HUMANS AND PETS TO CONTACT SURFACES.**

### **RESTRICTIONS:**

- DO NOT use this product with oil.
- DO NOT apply this product in any room being used as living, eating, sleeping, or recovery area by patients, the elderly, or infirm when they are in the room.
- DO NOT apply to institutions (including libraries, sports facilities, etc.,) in the immediate area when occupants are present.
- DO NOT apply to classrooms when in use.
- DO NOT apply this product to edible growing crops or stored raw agricultural commodities used for food or feed.
- DO NOT allow applications to contact water inhabited by fish, such as in aquariums and ornamental fish ponds that are located in/around structures being treated.

### **LAWNS/TURFGRASS AND ORNAMENTALS**

This product may be used for applications to maintain indoor or outdoor areas where turf and ornamentals are grown ~~such as~~ including residential landscaped areas and non-residential landscapes around institutional, public, commercial and industrial buildings, parks, recreational areas, and athletic fields (Including trees, shrubs, flowers, evergreens, foliage plants and groundcovers). Application rates for turf and ornamentals are lower than structural pest control rates, reflecting that treatment intervals are more frequent.

Applicators must ensure that they are certified in the necessary pesticide certification categories to allow application of this product away from structures, ~~including such as~~ to turf and ornamental plantings. Structural pest control certification categories may limit the distance away from structures for pesticide application. Consult your state extension office or pesticide regulatory officials for further information.

**IMPORTANT:** Time application to flowering plants during periods when pollinating insects are not present, such as early morning or late evening.

### **RESTRICTIONS:**

- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply this product to edible crops.
- **DO NOT** apply this product by aerial application.
- Use of this pesticide adjacent to water may affect aquatic organisms. To protect these organisms, **DO NOT** apply this pesticide within 25 ft of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.
- **DO NOT** make outdoor broadcast applications to turf and ornamentals when wind speed is 15 mph or greater.
- In the state of New York, **DO NOT** apply within 100 ft of coastal marshes or streams that drain into coastal marshes.

### **Mixing Instructions**

This product is to be mixed with water and may be used in all types of standard application equipment. Fill applicator tank with the desired volume of water and add this product. It is suggested that the water be 5 - 7 pH. Adjust water pH with a buffering agent if necessary. Slowly add this product to applicator tank water with maximum agitation. Close and shake or reagitator applicator tank before use if application is interrupted. Make up only amount of treatment volume as required.

### **Tank Mixing**

This product may be tank mixed with other currently registered pesticides unless expressly prohibited by the product label. A small volume mixing test with the other products is recommended to ensure compatibility. If other chemicals are added to the applicator tank, this product should be added last. Fill tank to desired volume and continue to agitate while making applications. If mixed with EC formulations, use within 24 hours.

Observe all restrictions and precautions which appear on the labels of these products.

**INSECT CONTROL on ORNAMENTAL PLANTS**

This product is for use to control the pests listed in the table below ~~including such areas as~~ residential landscaped areas and landscaped areas around institutional, public, commercial and industrial buildings, parks, recreational areas, and athletic fields (including trees, shrubs, flowers, evergreens, foliage plants and groundcovers).

**NOTE:** While phytotoxicity testing has been carried out on a wide range of ornamental plants under various environmental conditions, and no phytotoxicity has been observed, certain cultivars may be sensitive to the final spray solution.

It is advised to prespray a selection of ornamental plants and observe them for 7-10 days prior to treating large areas if local use experience is unavailable.

**RESTRICTIONS:**

- **DO NOT** apply more than 0.36 lbs of the a.i. (52.4 fl oz of concentrate) per acre per year.

### TANK DILUTION RATES FOR ORNAMENTAL PESTS

Pest	Amount of Equil Lambda 9.7 CS Insecticide	Application Instructions
<p>Ants (Including Imported Fire Ants)  Armyworms  Azalea Caterpillars  Aphids  Bagworms  Black Vine Weevils (adults)  Boxelder Bugs  Budworms  California Oakworms  Cankerworms  Cockroaches  Crickets  Cutworms  Eastern Tent Caterpillars  Elm Leaf Beetles  European Sawflies  Fall Webworms  Flea Beetles  Forest Tent Caterpillars  Gypsy Moth Larvae  Japanese Beetles (adults)  June Beetles (adults)  Lace Bugs  Leaf-feeding Caterpillars  Leafhoppers  Leafminers (adults)  Leafrollers  Leaf Skeletonizers  Midges  Mosquitoes  Oleander Moth Larvae  Pillbugs  Pine Sawflies  Pine Shoot Beetles  Pinetip Moths  Plant Bugs  Root Weevils  Sawflies  Scale insects (Crawlers)  Spiders  Spittlebugs  Stink Bugs  Striped beetles  Striped oakworms  Thrips  Tip moths  Tussock Moth Larvae  Wasps</p>	<p>1.5 – 5.0 fl oz per 100 gallons</p> <p>or</p> <p>44 - 148 mL per 100 gallons</p>	<p>Application to ornamentals should be started prior to the establishment of high insect pest populations.</p> <p>Apply at 7-day intervals if retreatment is necessary. Limit more frequent treatment to spot treatments.</p> <p>Good spray coverage is necessary to provide the most effective level of control. Addition of a spreader-sticker at specified rates may enhance the control of insects on certain species of ornamentals having waxy, hard-to-wet foliage.</p> <p>For spot treatments use 0.50 fl oz of this product per 1 - 2.5 gallons of water.</p> <p>Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area.</p> <p><b>Scale:</b> Thoroughly cover the plant with this product's spray, including trunks, stems, twigs, and foliage for control of scale insects (crawler stage).</p> <p><b>Bagworm:</b> Apply this product when bagworm larvae begin to hatch. Spray directly on the larvae. Application is the most effective when the larvae are young.</p>
<p>Broadmites  Brown Soft Scales  California Red Scales (Crawlers)</p>	<p>3 - 5 fl oz per 100 gallons</p>	

Pest	Amount of Equil Lambda 9.7 CS Insecticide	Application Instructions
Clover Mites Mealybugs Pine Needle Scales (Crawlers) Spider Mites Whiteflies	or 88 -148 mL per 100 gallons	

**Equil Lambda 9.7 CS Insecticide Mixing Chart for Ornamental Insect Pest Control**

(Equil Lambda 9.7 CS Insecticide to add per spray tank)

Rate of Equil Lambda 9.7 CS Insecticide	25 Gallons	50 Gallons	100 Gallons	200 Gallons	300 Gallons
1.5 fl oz <sup>1</sup>	0.4 fl oz	0.8 fl oz	1.5 fl oz	3.0 fl oz	4.5 fl oz
3 fl oz <sup>2</sup>	0.8 fl oz	1.5 fl oz	3.0 fl oz	6.0 fl oz	9.0 fl oz
5 fl oz <sup>3</sup>	1.3 fl oz	2.5 fl oz	5.0 fl oz	10.0 fl oz	15.0 fl oz
<sup>1</sup> Equivalent to 3.5 mL per 1,000 sq ft (5 fl oz per acre) when applied at 8 gallons per 1,000 sq ft. <sup>2</sup> Equivalent to 7.0 mL per 1,000 sq ft (10 fl oz per acre) when applied at 8 gallons per 1,000 sq ft. <sup>3</sup> Equivalent to 9.5 mL per 1,000 sq ft (14 fl oz per acre) when applied at 8 gallons per 1,000 sq ft.					

**LAWN/TURFGRASS INSECT PEST CONTROL**

This product is for use to control the pests listed in the table below including areas around such areas as residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

**RESTRICTIONS:**

- **KEEP CHILDREN AND PETS OFF TREATED AREAS UNTIL SPRAY HAS DRIED FOLLOWING APPLICATION.**
- **DO NOT** apply when turfgrass is waterlogged or when soils are saturated with water (i.e., will not accept irrigation).
- **DO NOT** apply more than 0.36 lbs of a.i. (52.4 fl oz of concentrate) per acre per year.

### RATES FOR LAWN AND TURFGRASS PESTS

Pest	Amount Of Equil Lambda 9.7 CS Insecticide*	Application Instructions
Ants (Including Imported Fire Ants) Armyworms Centipedes Chinch Bugs <sup>1</sup> Crickets Cutworms Earwigs Fleas (adult) Grasshoppers Japanese Beetles (adult) Millipedes Mites Mosquitoes (adult) Pillbugs Sod Webworms Sowbugs Stink Bugs Ticks (including species which transmit Lyme disease)	3.4 – 7.0 mL per 1,000 sq ft  or  5 -10 fl oz per acre	<p>Application to turf should be started prior to the establishment of high insect pest populations and significant turf damage. Make reapplications as necessary to keep pest populations under control, using higher rates as pest pressure increases.</p> <p>Apply at 7-day intervals if retreatment is necessary. More frequent treatments should be limited to spot treatments.</p> <p>For <b>spot treatments</b>, use 0.50 fl oz of this product per 1 - 2.5 gallons of water.</p> <p><b>Surface Insect Control (Armyworm, Cutworms, Fleas, etc.):</b> Apply this product at specified rates in 2 - 5 gallons of water per 1,000 sq ft. The use of a spreader-sticker may be useful if high rainfall amounts are forecast, otherwise the addition of adjuvants is not necessary under normal conditions for surface insect control in turf. Delay watering or mowing for 12 - 24 hours to control surface-feeding insect pests.</p>
Bluegrass Billbugs (adult) Black Turfgrass Ataenius (adult) Chiggers Fleas (adult) Grubs (suppression) Hyperodes Weevils (adult) Mole Crickets (nymphs and young adults)	7 mL per 1,000 sq. ft  or  10 fl oz per acre	<p><b>Thatch Inhabiting Insect Control (Chinch Bugs, Billbugs, etc.):</b> Apply this product at specified rate in 2-10 gallons of water per 1,000 sq ft. The use of a nonionic wetting agent, penetrant or similar adjuvant is recommended at label rates. Lightly irrigate after application with up to ½ inch of water to move this product into the thatch layer. If irrigation is not available, then use high water application rates.</p> <p><b>Subsurface Insect Control (Mole Crickets, Grubs, etc.):</b> Apply this product at specified rates in 4-10 gallons of water per 1,000 sq ft. The use of a nonionic wetting agent, penetrant or similar adjuvant is strongly recommended following label rates. Use the highest water application rates possible with your sprayer. Apply this product to turf wet with dew, rain or irrigation. Water-in immediately after application with ¼ - ½ inch of water.</p> <p><b>Fire Ant Control:</b> Treat individual mounds with a drench application using a watering can. Use 0.50 fl oz of this product per 2.5 gallons of water. Thoroughly soak each mound and a 3 ft diameter circle around each mound. Gently apply the mixture to avoid disturbing the mound. Disturbing the mound may cause the ants to migrate and reduce the effectiveness of the treatment. Apply in early morning or late evening hours. You may</p>



Pest	Amount Of Equil Lambda 9.7 CS Insecticide*	Application Instructions
		<p>make applications on a monthly basis to maintain protection of treated areas.</p> <p><b>Mosquito Control:</b> Apply as a general spray around landscape plantings, turf, and building foundations to control mosquitoes. Apply this product at specified rates in 2 - 5 gallons of water per 1,000 sq ft.</p>

- Not for use on chinch bugs in New York state.

\* Example: To treat the listed turf pests at the 7 mL per 1,000 sq ft rate, determine the gallons of dilution per 1,000 sq ft needed to adequately cover the turf. At a 5 gallon per 1,000 sq ft dilution, take the 7 mL and divide it by the 5 (gallon) to get 1.4 mL to be added per gallon. For a 50 gallon tank, therefore, this would be 1.4 mL X 50 = 70 mL (or 2.5 fl oz) in 50 gallons of water.

### Equil Lambda 9.7 CS Insecticide Mixing Chart for Turf Insect Pest Control

(Equil Lambda 9.7 CS Insecticide to add per 100 gallons spray tank)

Rate of Equil Lambda 9.7 CS Insecticide	Application Rate Per 1,000 Square Feet of Turf				
	2 gallons	4 gallons	6 gallons	8 gallons	10 gallons
5 fl oz per acre	5.7 fl oz	2.9 fl oz	1.9 fl oz	1.4 fl oz	1.2 fl oz
10 fl oz per acre	11.5 fl oz	5.7 fl oz	3.8 fl oz	2.9 fl oz	2.3 fl oz

Conversion Rate: 1 fluid ounce (fl oz) equals 29 milliliters (mL).

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE AND SPILL PROCEDURES:** Store upright at room temperature. Keep container closed when not in use. Do not store near food or feed. Avoid exposure to extreme temperatures. In case of spillage or leakages, soak up with an absorbent material such as sand, sawdust, earth, Fuller's earth, etc. Dispose of with chemical waste.

**PESTICIDE DISPOSAL:** Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

**CONTAINER DISPOSAL: For Containers equal to or less than 5 gallons:** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available. If recycling is not available, then dispose of container in a sanitary landfill, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**For Containers greater than 5 gallons:** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling if available. If recycling is not available, then dispose of container in a sanitary landfill, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**For bulk containers: (Refillable Container):** Refill this container with pesticides only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

## **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this Product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Central Garden & Pet Company or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Central Garden & Pet Company and Seller harmless for any claims relating to such factors.

Central Garden & Pet Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Central Garden & Pet Company, and Buyer and User assume the risk of any such use. **CENTRAL GARDEN & PET COMPANY MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the fullest extent permitted by law Central Garden & Pet Company or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CENTRAL GARDEN & PET COMPANY AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CENTRAL GARDEN & PET COMPANY OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Central Garden & Pet Company and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Central Garden & Pet Company.

Actisol® is a registered trademark of Roussel-Uclaf.

Micro-Injector® is a registered trademark of Whitmire Micro-gen Research Laboratories, Inc.

### **Manufactured for:**

Central Garden & Pet Company  
1501 East Woodfield Road 200W  
Schaumburg, IL 60173