



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

90354-2

Date of Issuance:

6/29/17

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Yellow Jacket Killer!

Name and Address of Registrant (include ZIP Code):

David H. Dawe
President, D.H. Dawe & Associates, LLC (Agent)
2 Star Solutions Inc.
1954 East Bay Street
Hoschton, GA 30548

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Kable Bo Davis, Product Manager 03
Invertebrate & Vertebrate Branch 1, Registration Division (7505P)

Date:

6/29/17

2. You are required to comply with the data requirements described in the DCIs identified below:

- a. Fipronil GDCI-129121-1305
 GDCI-129121-1512

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCIs listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division: <http://www.epa.gov/pesticide-contacts/contacts-office-pesticide-programs-pesticide-re-evaluation-division>

3. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. A one year study is required to satisfy these data requirements. You have 18 months from the date of registration to provide these data.
4. Make the following label changes before you release the product for shipment:
- Revise the EPA Registration Number to read, “EPA Reg. No. 90354-2.”
5. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 04/17/2017

If you have any questions, please contact Tim Ciarlo by phone at 703-347-8082, or via email at Ciarlo.Timothy@epa.gov.

ACCEPTED

06/29/2017

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 90354-2

FRONT

Yellow Jacket Killer!™
Kills Yellow Jacket Scavengers

4 Bait Stations Included

KEEP OUT OF REACH OF CHILDREN

CAUTION (PRECAUCION)

See back panel and enclosed insert for additional First Aid.

Directions for use and Precautionary Statements on Top and Back Panel

Active Ingredient:

Fipronil*9.1%
Other Ingredients.....90.9%
Total.....100.00%

*(5-amino-1-(2,6-dichloro-4-(trifluoromethyl) phenyl)-4-(1,R,S)
(trifluoromethyl) sulfinyl)-1-H-pyrazole-3-carbonitrile

Net contents: 0.3ml (0.01 oz.)

EPA Reg. No. 90354-X

EPA Est. No. 90354-GA-001

TOP and BACK

FIRST AID

If Swallowed: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a Poison Control Center or doctor for treatment advice, If Inhaled: Move person to fresh air. If person is not breathing call 911 or ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a Poison Control Center or doctor for treatment advice.

If in Eye(s): Hold eye open and rinse gently with water for 15 to 20 minutes. If wearing contacts, rinse eye for 5 minutes, remove contact and continue rinsing. Call a Poison Control Center or doctor immediately for treatment advice

Note to Physician: There is no specific antidote. All treatment should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred. In severe cases of overexposure by oral ingestion, lethargy, muscle tremors, and in extreme cases, possibly convulsions may occur.

Have the product container or label with you when calling a Poison Control Center or doctor, or going for medical treatment. You may also call SafetyCall at 866-897-6050 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Caution: Harmful if swallowed, absorbed through skin or inhaled. Do not get in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash any contaminated clothing.

NOTE: If you are allergic to yellow jacket stings, do not personally use this product.

Have someone other than yourself handle and maintain the Yellow Jacket Killer!

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store unused product in original container only, out of reach of children and animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Non-refillable Container: Do not reuse or refill this container. Triple rinse promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

ENVIRONMENTAL HAZARDS

The pesticide contained in the bottle is extremely toxic to fish and other aquatic invertebrates. Cover or collect spilled bait during loading. Do not discharge this product into lakes, streams, ponds, estuaries, oceans or other waters or sewage systems. Do not use within 25 feet of any body of water.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product. See enclosed instructions for Bait Station and complete Directions For Use.

LEFT SIDE

FINALLY! Now you have an easy-to-use product that kills yellow jacket scavengers.

Simply hang each Yellow Jacket Killer!™ bait station with treated bait inside and replace the bait daily for 5 consecutive days. **It's that easy!**

Bait Stations are made of transparent plastic that hang easily from a tree branch, shrub or other hanging location with the included twist ties.

Place Bait Stations near Yellow Jacket activity.

To kill yellow jackets, use multiple Bait Stations.

- Bait Stations should be placed no more than 60-75 feet apart.
- 1 bait station kit contains 4 bait stations.
- Use 4 bait stations for every 10,000 sq. ft. (100'x100') of area.

@2017 2 Star Solutions Inc. All rights reserved

RIGHT SIDE

Everything's Included!

The YELLOW JACKET KILLER!™ is the easiest way to solve your yellow jacket problem...because it's a product that kills Yellow Jacket scavengers.

CAUTION: If you are allergic to yellow jacket stings, do not personally use this product. Have someone other than yourself hang and maintain the Bait Stations.

U.S. Patent Pending

Made in the U.S.A.

EPA Reg. No. 90354-Pending EPA Est. No. 90354-GA-001

SOLD BY:

2 Star Solutions Inc.

970 Peachtree Ind. Blvd.

Suite 100

Suwanee, GA 30024

Questions or Comments? Call 1-888-448-1669 Batch No. _____



© 2017, 2 Star Solutions, Inc. All rights reserved.

YELLOW JACKET KILLER!™

Instruction Insert

YELLOW JACKET KILLER!™ allows the user to kill the yellow jackets without the dangers of applying an aerosol or powder pesticide to the nest. YELLOW JACKET KILLER!™ is not a trapping device. This unique "Bait Station" is designed to attract yellow jackets and allow them to enter and feed on the pesticide-treated bait. The yellow jackets will leave the bait stations and die. Best of all, you do not need to know where the nest is located.

| | | | |
|---------------------------|------------------------|--------|--|
| Active Ingredient: | Fipronil*..... | 9.1% | |
| | Other Ingredients..... | 90.9% | * (5-amino-1-(2,6-dichloro-4-(trifluoromethyl) phenyl)-4-(1,R,S) |
| | Total..... | 100.0% | (trifluoromethyl) sulfinyl)-1-H-pyrazole-3-carbonitrile |

STOP! READ ENTIRE DIRECTIONS BEFORE USE

Keep Out of Reach of Children

CAUTION

FIRST AID

If Swallowed: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

If on Skin of Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a Poison Control Center or doctor for treatment advice.

If in Eyes: Hold eye open and rinse gently with water for 15 to 20 minutes. If wearing contacts, rinse eye for 5 minutes, remove contact and continue rinsing. Call a Poison Control Center or doctor immediately for treatment advice.

Note to Physician: There is no specific antidote. All treatment should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred. In severe cases of overexposure by oral ingestion, lethargy, muscle tremors and, in extreme cases, possibly convulsions may occur.

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. You may also contact the Poison Help Line at 1-800-222-1222 for treatment advice.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed, absorbed through skin or inhaled. Do not get in eyes, on skin or on clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PRECAUTION. Yellow jacket stings can be life threatening to persons who are allergic to the venom. People who develop hives, difficulty breathing or swallowing, wheezing or similar symptoms of allergic reaction should seek medical attention immediately if stung by a yellow jacket.

ENVIRONMENTAL HAZARDS

The insecticide contained in the dropper bottle is extremely toxic to fish and other aquatic invertebrates. Cover or collect spilled bait during loading. Do not discharge this product into lakes, streams, ponds, estuaries, oceans, or other waters or sewage systems. Do not contaminate water by disposal of waste. Do not use the product within 25 feet of any body of water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling. For residential and commercial use. For non-food use only. **Do not hang bait station or replace bait if you are allergic to yellow jacket stings.**

INITIAL SET-UP: DAY 1

1. Open Bait Station and remove all contents.
2. Remove empty pesticide bottle and fill with clean water to fill line (88.7ml) marked on the bottle. Open pesticide blister package by bending back and forth at the scored line at the top of the bottle imprint. Carefully hold the blister packaging on the flat areas next to the top of the bottle imprint and tear off the top of the blister package. Squeeze entire contents into the water bottle. Insert dropper tip onto bottle top and replace bottle cap.
3. Remove four (4) bait trays, bait food packet and spoon. Add bait food to each of the bait trays (4 cubes each tray). Add one (1) level spoonful of water to each bait tray. Wait 1 to 2 minutes for bait to absorb the water. **Shake insecticide bottle well (minimum 30 seconds).** Add 2 drops of diluted insecticide to each cube of bait for a total of 8 drops per bait tray.

IMPORTANT: Yellow jackets will not take spoiled meat.


Replace bait trays and bait every morning for 5 consecutive days.

4. Insert loaded Bait Tray into Bait Tray receptacle - as shown in right diagram **A**.
5. Close the Bait Station and press your fingers together on the button snap closures **B**.
6. Take the loaded Bait Stations and Twist Ties outdoors to hanging locations.
7. Attach the Twist Tie to the hanger opening at the top of the Bait Station. Wrap the Twist Tie at least two (2) times around itself for secure attachment.
8. Hang the Bait Stations a minimum of 10 feet away from any known nest. Keep the Bait Stations away from your house or any outdoor living area.
9. Hang the Bait Stations in a **SHADED AREAS** for best results. NOTE: Using the Twist Tie, hang the Bait Station from a tree branch, shrub (or other hanging location) high enough from the ground that it cannot be reached by children, family pets or scavenging wildlife.
10. Wash hands thoroughly with soap and water immediately after handling Bait Station(s).

DAILY OPERATION: DAY 2 AND BEYOND

Perform the following steps every morning after Day 1:

1. Check Bait Station for yellow jackets. **If yellow jackets are present, do not attempt to change the bait.**
2. If yellow jackets are not present, remove the Bait Station from the Twist Tie.
3. Open Bait Station and remove used Bait Tray. Use a NEW Bait Tray and add 4 cubes of bait food to the tray. Add one (1) level spoonful of water to each bait tray. Wait 1 to 2 minutes for bait to absorb the water. **Shake Insecticide bottle for a minimum of 30 seconds.** Add 2 drops of diluted Insecticide to each bait cube for a total of eight (8) drops. **IMPORTANT: Yellow jackets will not eat spoiled meat.** Replace bait tray and bait every morning for 5 consecutive days.
4. Re-hang the Bait Station with the Twist Tie from the bait station hanger opening.
5. Dispose of the used Bait Trays and bait into your trash receptacle.
6. Wash hands thoroughly with soap and water immediately after re-hanging the Bait Stations and after handling the used Bait Trays.



Bait Station - OPENED **A**
CLOSED **B**

Bait Tray -
20 TOTAL



Bait Food
Packet



Twist Tie



Oxygen Absorber

Insecticide
DROPPER
BOTTLE



Spoon

BAIT STATION PLACEMENT

1. Hang Bait Stations in shaded areas, if possible, away from house and outdoor living areas.
2. Bait Stations should be placed no more than 60 to 75 feet apart. You will need 4 Bait Stations for every 10,000 sq.ft. of area.
3. **Move Bait Stations to another location if yellow jackets are not finding and taking the bait.**

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store unused product in its original container only, out of the reach of children and animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide is in violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

CONTAINER HANDLING: Non-refillable Container: Do not reuse or refill this container. Triple rinse promptly after emptying; then offer for recycling, if available; or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by the state and local authorities.

WARRANTY STATEMENT


2 STAR SOLUTIONS, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. To the extent consistent with applicable law, the exclusive remedy of any buyer or user of this product for any and all losses, injuries, or damage resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall be damages not exceeding the purchase price paid for this product or, at 2 Star Solutions, Inc.'s option, the replacement of this product.

NET CONTENTS: One (1) blister packet of Insecticide 0.3ml (0.01 oz.)
One (1) blister packet of freeze-dried chicken (1.12 oz.)

EPA Reg. No. 90354-Pending EPA EST. No. 90354-GA-001 U.S. Patent Pending

SOLD BY: 2 Star Solutions, Inc.

970 Peachtree Industrial Blvd., Suite 100, Suwanee, GA 30024 1-888-448-1669



**Instructions
Instrucciones**

MADE IN THE U.S.A.



© 2017, 2 Star Solutions, Inc. All rights reserved.

Dropper Bottle Label

Yellow Jacket Killer!™

Active Ingredient:

Fipronil.....9.1%

Other Ingredients.....90.9%

Total.....100.00%

Keep out of reach of children

CAUTION

Read directions and precautionary statements before using.

EPA Reg No. 90354-Pending

Net Contents: 88.75 ml