



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

October 7, 2020

Premjit Halamkar, Ph.D., MPA
Global Head of Regulatory Affairs
CH Biotech R&D Co. Ltd,
c/o CH Biotech LLC
601 Kettering Dr.
Ontario, CA 91761

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment – Revision to the Labeling to update First Aid, precautionary statement, use directions sections; in addition to adding hemp and crop groups (Field crops, berry and small fruits, vegetables, vines, pome fruits, stone fruits, tree crops and tree nuts, brassica vegetables group, citrus fruits group, cucurbit vegetables group, fruiting vegetables group, herbs group, leafy vegetable group, legume vegetable group, pome/stone fruits group, root vegetable group, small fruits group, small grains group, tree nut group, and tuber vegetable group). The amendment also added “Roundup Ready® is a registered trademark of the Monsanto Company” and “LibertyLink® is a registered trademark of Bayer CropScience” to the label.

Product Name: HAPPYGRO

EPA Registration Number: 90866-12

Application Date: 07/20/2020

Submission Number: 1054867

Case Number: 00016900

Dear Dr. Halamkar:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling

or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the U.S. Environmental Protection Agency (EPA). If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.


If you have any questions, please contact James Parker by phone at (703) 306-0469 or via email at parker.james@epa.gov.

Sincerely,



Gina Burnett, Senior Regulatory Advisory
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)
Office of Pesticide Programs

Enclosure



HappyGro™

A Plant Growth Regulator for Listed Commodities

ACTIVE INGREDIENT:.....		By WT
Cytokinin, as Kinetin		0.5%
OTHER INGREDIENTS:.....		99.5%
	TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

First Aid	
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15-20 minutes • Call a poison control center or doctor for treatment advice
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye • Call a poison control center or doctor for treatment advice
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For non-emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 (NPIC Web site: www.npic.orst.edu).</p> <p>For general product information, call CH Biotech LLC at (909) 472-3033 between the hours of 9a.m. – 4 p.m. Pacific Time.</p>	

EPA Reg. No. 90866-12

EPA Est. No. 90866-CA-1

Product for CH Biotech R&D Co., Ltd.

By CH Biotech, LLC
 601 Kettering Drive
 Ontario, CA 91761
 Tel: 909.472.3033
 Email: info@chbio.com

Net Content: 1 Gallon (3.785L)
 Batch No.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

For Terrestrial Uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean highwater mark. Do not contaminate water when disposing of equipment wash water or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

This product is not compatible with strong oxidizers.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during applications. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and

restricted-entry interval. The requirements in this box only apply to uses that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves Category A, such as butyl rubber \geq 14 mils, or natural rubber \geq 14 mils, or neoprene rubber \geq 14 mils or nitrile rubber \geq 14 mils
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter without appropriate protective clothing until sprays have dried.

GENERAL INFORMATION

Kinetin (a cytokinin), the active ingredient in HappyGro™, is a plant growth hormone that can occur naturally. Kinetin has shown to increase cell division, cell differentiation and cell growth. HappyGro can enhance plant growth and development when applied as directed.

HappyGro is not a fertilizer; therefore, incorporate good fertilization program practices. Under certain circumstances, kinetin may delay senescence of the leaves on some crops. Make applications at the proper timing and when the crop is actively growing.

CHEMIGATION

Apply this product only through the following types of irrigation systems:

1. Sprinkler including side (wheel) roll, solid set or hand move irrigation systems.
2. Calibrated overhead watering booms

Do not apply this product through any other types of irrigation systems. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems), used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of

pesticide introduction. As an option to the RPZ, discharge the water from the public water system into a reservoir tank prior to pesticide introduction. A complete physical break (air gap) must occur between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must contain normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment. Agitate the pesticide supply tank throughout the application of HappyGro. Except for turfgrass, apply HappyGro at the end of the irrigation period in a sufficient amount of water to allow proper coverage of plant or crop. Fill the supply tank one-half full with water, add the appropriate amount of HappyGro to the tank and finish filling the tank with water.

SPRINKLER CHEMIGATION

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Agitate the pesticide supply tank throughout the application of HappyGro. Except for turfgrass, apply HappyGro at the end of the irrigation period in a sufficient amount of water to allow proper coverage of plant or crop but not to exceed 8 fluid ounces of HappyGro per acre per application. Fill the supply tank one-half full with water, add the appropriate amount of HappyGro to the tank and finish filling the tank with water.

APPLICATION DIRECTIONS

For all crops, unless otherwise specified, tank mix HappyGro by adding 13 fluid ounces per 100 gallons of water and spray crop canopy with sufficient volume to ensure uniform coverage. For more specific instructions and rates, see the following table of crops; the rate is expressed as fluid ounces of HappyGro per acre in a corresponding volume of water per acre. Water volume is usually determined by the grower and/or the particular circumstances that affect uniform crop coverage; however, sufficient volume to wet leaf is required as opposed to just misting it. Use lower rates with the corresponding lower volumes and higher rates with higher volumes.

If sufficient rain to wash leaf occurs within 2 hours of application, re-apply. Compatibility has not been fully determined for all agrochemicals.

Due to reduced coverage with gallonage of 5 gallons per acre or less, aerial application requires a minimum dose of 2 oz./acre (see footnote ⁽³⁾ below).

GENERALLY RECOMMENDED SPRAY VOLUME (GAL/ACRE)

CROP/CROP GROUP	GROUND		AIR	
	Adequate Coverage	Best Coverage	Adequate Coverage	Best Coverage
Field Crops, Miscellaneous	10.0	20.0	2.0	5.0
Berry and Small Fruits, Vegetables, Vines	25.0	100.0	5.0	10.0
Pome Fruits, Stone Fruits, Tree Crops and Tree Nuts	50.0	200.0	10.0	20.0

Crop/Crop Group	Application Timing	HappyGro Rate/Acre	Water Volume/Acre**
Alfalfa, clover and vetch (for seed)	Make one application, 5 to 8 days after full bloom	2.0 to 3.0 fl oz	15 to 25 gal
Brassica Vegetables Group. [Crops such as Broccoli, Cauliflower, Cabbage and Mustard greens]	1st application: At flowering stage	2.0 fl oz	15 gal
	2nd application: 10 to 14 days after first application	2.0 to 3.0 fl oz	15 to 25 gal
	3rd application: 7 to 10 days before harvest		
Citrus Fruits Group. [Crops such as Grapefruit, Lemon, Sweet orange, and Tangelos]	1st application: At small fruit stage when fruit size is approx. 6 to 8 mm	6.0 to 12.0 fl oz	50 to 100 gal
	2nd application: 40 days prior to harvest	6.0 to 12.0 fl oz	50 to 100 gal
Corn (Sweet, Field ⁽¹⁾⁽²⁾ , and Popcorn)	Make one application, V5 to VT (Full tassel)	2.0 to 3.0 fl oz	10 to 25 gal
Cotton including short staple, acala & pima varieties	1st application: Apply at the pin head square stage	2.0 to 3.0 fl oz	10 to 25 gal
	2nd application: Apply at first bloom	3.0 to 4.0 fl oz	10 to 30 gal
Cucurbit Vegetables Group. [Crops such as Cucumber, Cantaloupe, Honeydew, Muskmelon, Summer squash, and Watermelon]	Make 1st application at early fruiting stage	2.0 to 3.0 fl oz	15 to 25 gal
	Make 2 or more applications at 10 to 14-day intervals	3.0 to 5.0 fl oz	20 to 40 gal
Fruiting Vegetables Group. [Crops such as Eggplant, Pepper, and Tomato]	Make 1st application at early fruiting stage	2.0 to 3.0 fl oz	15 to 25 gal
	Make 2 or more applications at 10 to 14-day intervals	3.0 to 12.8 fl oz	25 to 100 gal
Grapes including table, wine, and raisin varieties	1st application: 10-14 days before bud break	3.0 to 5.0 fl oz	25 to 40 gal
	2nd application: Small berry stage (3 to 5 mm in size)	5.0 to 9.0 fl oz	40 to 60 gal
	3rd application: 40 days prior to harvest		
Herbs Group. [Crops such as Basil, Dill, Mustard and Sage]	1st application: During mid-season growth	3.0 to 5.0 fl oz	25 to 40 gal
	2nd application: As flowers or buds begin to emerge		

Hemp	1st application: At early bloom	2.0 to 6.0 fl oz	15 to 50 gal
	2nd application: As needed, 21 days after first application		
Hops	1st application: At inflorescence emergence	3.0 to 9.0 fl oz	25 to 75 gal
Leafy Vegetable Group. [Crops such as Celery, Head lettuce, Leaf lettuce and Spinach]	1st application: During mid-season growth	3.0 to 5.0 fl oz	25 to 40 gal
	2nd application: 10 to 14 days prior to harvest		
Legume Vegetables Group. [Crops such as Dry beans, Green beans, Lentils, Peas]	1st application: Apply at the 3 to 5 trifoliolate leaf stage	2.0 to 3.0 fl oz	15 to 25 gal
	2nd application: Apply at 5 to 10% bloom		
Oil Seed Crops including Canola ⁽¹⁾⁽²⁾ , Flax and Sunflower	1st application: First bloom	2.0 fl oz	10 to 15 gal
	2nd application: 14 to 21 days after 1st application		
Peanuts including all commercial varieties	1st application: Apply at the 3 to 5 leaf stage	2.0 to 3.0 fl oz	15 to 25 gal
	2nd application: Apply at initial pegging		
	3rd application: Apply during pod fill	3.0 to 5.0 fl oz	25 to 40 gal
Pome/Stone Fruits Group. [Crops such as Apple, Apricot, Cherry, Plum, Plumcot and Peach]	1st application: At small fruit stage when fruit size is approximately 3 to 5 mm	6.0 to 12.0 fl oz	50 to 100 gal
	2nd application: 40 days prior to harvest		
Root Vegetables Group. [Crops such as Carrot, Ginseng, Horseradish, Parsley (turnip-rooted) Radish, Sugar beet, and Turnip]	1st application: At plant thinning stage	2.0 fl oz	15 gal
	2nd application: 21 days after first application	2.0 to 3.0 fl oz	15 to 25 gal
	3rd application: Apply 10 to 14 days prior to anticipated harvest		
Small Fruits Group. [Crops such as Blackberry, Blueberry, Raspberry, and Strawberry]	1st application: At early flowering stage	2.0 to 5.0 fl oz	15 to 40 gal
	2nd application: 10 to 14 days after first application		
	3rd application: Repeat applications every 10 to 14 days		
Small Grains Group. [Crops such as Barley, Rice Sorghum, Rye, and Wheat]	Make one application: Boot stage to anthesis. Wheat Feekes stage 10 to 10.5	2.0 to 3.0 fl oz	10 to 25 gal
Soybeans ⁽¹⁾⁽²⁾	1st application: First bloom	2.0 fl oz	15 gal
	2nd application: 14 to 21 days after 1st application	2.0 to 3.0 fl oz	10 to 25 gal
Tree Nuts Group. [Crops such as Almond, Pecan, and Pistachio]	1st application: 2 weeks prior to bloom	6.0 to 12.0 fl oz	50 to 100 gal
	2nd application: 2 weeks following petal fall		

	3rd application: 30 days after 2nd application		
Tuber Vegetables Group. [Crops such as Potato Sweet Potato, and Yam]	1st application: At tuber initiation stage	2.0 to 3.0 fl oz	15 to 25 gal
	2nd application: 14 to 21 days after first application		
	3rd application: At early bloom stage	3.0 to 5.0 fl oz	25 to 40 gal

**Note: If using any other volume of water, use the tank mix dilution rate of 13 fluid ounces of HappyGro per 100 gallons of water and use sufficient water volume to obtain uniform coverage.

(1) This product can be tank mixed with glyphosate products registered for use on Roundup Ready® crops.

(2) This product can be tank mixed with products registered for use on LibertyLink® crops.

(3) Carrier volumes with 10 gallons per acre or less requires the use of 2 oz/acre minimum due to reduced coverage, which includes aerial applications.

TURFGRASS

For Sod Grass: Apply HappyGro by ground using 20 to 40 gallons of water per acre. Apply 2.5 to 5.0 fluid ounces product in 20 to 40 gallons of water, respectively, at a 1:1000 dilution rate.

For Turfgrass: Apply HappyGro by ground according to the table below using 1 to 10 gallons of water per 1000 square feet.

Turf	Application & Timing	HappyGro Rate/1000 ft. ²	Water Volume/1000 ft. ²
Cool Climate grasses such as Bluegrass, Fescue, Rye, and similar grasses	1st application: When seeded grass becomes established or at the beginning of the season for perennials	0.13 to 0.65 fl oz	1 to 5 gal
	Repeat as necessary		
Dichondra	1st application: When turf greens up in the spring	0.65 to 1.30 fl oz	5 to 10 gal
	Repeat at 14 to 21 days intervals during the growing season		
Warm Climate grasses such as St. Augustine, Bermuda, Bermuda hybrids, Centipede & similar grasses	1st application: When turf greens up in the spring	0.13 to 0.65 fl oz	1 to 5 gal
	Repeat at 14 to 21 days intervals during the growing season		

ORNAMENTALS

Greenhouse and nursery grown ornamentals

Differences in responsiveness may vary from one cultivar to another or from one set of growing conditions to another. Unless previous experience dictates otherwise, prior to widespread use, test a small number of plants from each cultivar to verify desired efficacy.

APPLICATION INSTRUCTIONS			
HappyGro is applied to foliage of ornamentals and leaves of bedding plants. Mixing instructions: dilute 1.3 fluid ounces of HappyGro in 10 gallons (37.8 mL/10 gal) of water (1:1000 dilution); for smaller quantities, mix 0.13 fluid ounces HappyGro (¾ tsp or 3.7 mL) per 1 gallon of water. Use sufficient volume of water to obtain uniform coverage. Thorough coverage is necessary for best results.			
Plant	Timing	HappyGro Rate	Volume of Water
Foliage Plants: Aglaonema, Ajuga, Anthurium, Aphelandra, Caladium, Cissus, Dieffenbachia, Dracaena, Ficus, Fittonia, Gynura, Hoya, Maranta, Palms, Peperomia, Philodendron, Pilea, Pothos, Schefflera, Schlumbergera, Spathiphyllum, Syngonium, Tradescantia and Similar foliage plants	1st application: At the beginning of the season	¾ tsp	1 gal
	Subsequent applications: Spray foliage 2 to 3 times throughout the year at even intervals	1.3 fl oz	10 gal
Bedding and Flowering Plants: Abutilon, Aglais, Alyssum, Calceolaria, Canna, Carnation, Champaca, Chrysanthemum Cineraria, Columbine, Coral Bells, Cyclamen, Dahlia, Delphinium, Dianthus, Foxglove, Fuchsia, Gardenia, Gazania, Geranium, Gladiolus, Gloxinia, Impatiens, Iris, Jasminum, Lily, Lupine, Marigold, Michelia, Monarda, Osmachus, Petunia, Poinsettia, Portulaca, Roses, Salvia, Scabiosa, Sedum, Sempervivum, Tulips, Vinca, Zinnia, and Similar plants	Bedding Plants: 1st application: At 3 to 5 leaf stage.	¾ tsp	1 gal
	2nd application: 10 to 14 days prior to sale	3.0 tsp	4 gal
	Established Plants: 1st application: 21 to 30 days after transplanting or in the Spring as re-growth begins 2nd application: At flowering	1.3 fl oz	10 gal
Woody Ornamentals: Arborvitae, Aucuba, Azalea, Boxwood, Carissa, Chinese magnolia, English Ivy, Holly, Juniper, Maple, Pine, Podocarpus, Rhododendron, Viburnum, and Similar plants	1st application: Early in the season as new growth begins Subsequent applications: every 30 days as necessary	1.3 fl oz	10 gal

Garden Grown Tree Fruits Apple, Asian pear, Apricot, Cherry, Fig, Guava, Grape, Jujube, Kumquat, Lemon, Litchi, Longara, Mango, Orange, Peach, Persimmon, Plum, Prunus, Starfruit and Similar plants	1st application: When fruit size is approximately 3 to 5 mm 2nd application: 40 days prior to harvest	3.0 tsp	4 gal
--	---	---------	-------

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Protect from freezing. Store out of direct sunlight. Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at www.acrecycle.org. If not recycled, then puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

For packages up to 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container% full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For packages greater than 56 gallons: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For refillable containers: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump

for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For help with any spill, leak, fire or exposure involving this material, call day or night
CHEMTREC- 1-800-424-9300.

LIMITED WARRANTY AND DISCLAIMER

NOTICE: CH Biotech R&D Co., Ltd. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use. Buyer assumes all risks of use and handling which is a variance in any way with the directions herein. To the extent consistent with applicable law, CH Biotech R&D Co., Ltd. makes no other express or implied warranty of fitness or merchantability. To the extent consistent with applicable law, in no case shall CH Biotech R&D Co., Ltd. or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. CH Biotech R&D Co., Ltd. and Seller offer this product and the Buyer and user accept it, subject to the foregoing Limited Warranty and Disclaimer which may be varied only by agreement in writing signed by a duly authorized representative of CH Biotech R&D Co., Ltd.

HappyGro™ is a trademark of CH Biotech R&D Co., Ltd.

Roundup Ready® is a registered trademark of the Monsanto Company

LibertyLink® is a registered trademark of Bayer CropScience

Produced for CH Biotech R&D Co., Ltd.

By CH Biotech, LLC.

601 Kettering Drive

Ontario, CA 91761

Tel: 909.472.3033

Email: info@chbio.com