



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

June 28, 2018

Premjit Halarnkar, Ph.D., MPA  
Global Head of Regulatory Affairs  
CH Biotech R&D Co. Ltd  
c/o CH Biotech, LLC  
601 Kettering Dr.  
Ontario, CA 91761

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment - Revision to product label to update Emergency Contact Information, company name (including contact information), EPA Registration Number, EPA Establishment Number, Personal Protection Equipment (PPE) section, Storage and Disposal section, and Warranty section.  
Product Name: GA<sub>3</sub> 20%  
EPA Registration Number: 90866-14  
Application Date: 06/06/2018  
OPP Decision Number: 541809

Dear Dr. Halarnkar:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the U.S. Environmental Protection Agency (EPA). If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or

misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact James Parker by phone at (703) 306-0469 or via email at [parker.james@epa.gov](mailto:parker.james@epa.gov).

Sincerely,

A handwritten signature in blue ink that reads "Andrew C. Bryceland". The signature is fluid and cursive, with the first name being the most prominent.

Andrew Bryceland, Team Leader  
Biochemical Pesticides Branch  
Biopesticides and Pollution  
Prevention Division (7511P)  
Office of Pesticide Programs

Enclosure

**ACCEPTED**

06/28/2018

Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No. 90866-14

# GA<sub>3</sub> 20%

## Plant Growth Regulator Soluble Powder

ACTIVE INGREDIENT:	BY WT
Gibberellic Acid .....	20.0%
OTHER INGREDIENTS: .....	80.0%
<b>TOTAL</b>	<b>100.0%</b>

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

<b>FIRST AID</b>	
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For non-emergency information concerning this product, call the National Pesticide Information Center at 1-800-858-7378 (NPIC Web site:www.npic.orst.edu). <b>For general product information, call CH Biotech LLC at (909)472-3033 between the hours of 9 a.m. to 4 p.m. Pacific Time.</b>	

EPA REG. NO. 90866-14  
EPA EST. NO. 90866-CA-001  
Net Contents 22.5 oz (640 grams)  
Batch No.

Produced for  
CH Biotech R&D., Ltd.  
By CH Biotech LLC  
601 Kettering Drive  
Ontario, CA 91761  
Tel: 909-472-3033  
Email: info@chbio.com

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wear goggles. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, or using tobacco.

### ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters or rinsate. Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long sleeved shirt and long pants,
- Chemical-resistant gloves in Category A, such as butyl rubber > 4 mils, or natural rubber > 14 mils, or neoprene rubber > 14 mils or nitrile rubber > 14 mils,
- Shoes plus socks, and
- Goggles

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

#### Users should:

- Wash hands before eating drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during applications. For any requirements specific to your State or Tribe, consult the State or Tribal Agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours unless wearing appropriate PPE.**

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:

- Coveralls,
- Chemical-resistant gloves,
- Shoes plus socks, and
- Goggles

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

### **GENERAL INFORMATION**

Use only as directed. Read the label thoroughly and understood before making applications. Keep out of reach of children.

Do not apply this product through any type of irrigation system.

#### **Application Instructions:**

GA<sub>3</sub> 20% contains gibberellic acid which is an extremely potent plant growth regulator; when applying plant growth regulators, follow the label directions for rates, timings, and water volumes. Do not apply untested spray mixes.

- Do not apply to plants under pest, nutritional, or water stress.
- Effectiveness requires that all parts of plant or crop receive thorough spray coverage or desired result will not occur. Prepare solution concentrations by mixing the required amount of product with water in a clean, empty spray tank. Dispose of any unused spray material at the end of each day following local, state or federal law.
- For best results, use water with a neutral pH and always below 8.5.
- GA<sub>3</sub> 20% applications made under slow drying conditions (cool to warm temperatures, medium to high relative humidity, and no wind) will increase absorption by the plant, thus optimizing effectiveness. Night-time applications are encouraged when day-time conditions are not conducive to slow drying conditions.
- Product persistence: Reapply GA<sub>3</sub> 20% if significant rain occurs within 2 hours of application.
- No preharvest interval is required for this product.
- DO NOT apply using ULV application methods. For aerial applications use spray volumes of 2 gallons per acre or greater (10 gallons per acre for tree crops).

#### **Compatibility Test for Tank Mix Components**

The use of surfactants and other additives has been reported to be beneficial. CH Biotech R&D Co., Ltd. does not assume responsibility for unexpected results due to the tank mixing of GA<sub>3</sub> 20% with other products. Conduct a jar test before tank mixing to ensure compatibility of GA<sub>3</sub> 20% with other pesticides or products if the tank mix combination has not been used previously.

To test for compatibility, use a small container (jar) and mix a small amount (0.5 to 1 quart) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5 to 15 minutes after mixing. To ensure maximum crop safety and product performance, follow all precautions and limitations on this label and labels of products used in the tank mixture with GA<sub>3</sub> 20 %.

## SPRAY INSTRUCTIONS FOR CROP CATEGORIES

### GRAPE

For all grapes, apply by ground sprayer. Apply in sufficient water volume to ensure thorough wetting. It is important to wet all flower clusters or berries thoroughly. For specific spray rates and timings, by variety, see accompanying tables. Do not exceed maximum rates.

#### SEEDLESS GRAPE

*For cluster elongation and looser cluster forms ("Stretch"). To reduce costs of thinning, allow better air circulation to aid in the control of bunch rot, and increase light penetration to aid in sugar development:*

Guide: Apply one to three applications before bloom when flower clusters are 2-to-7 inches long.

*For decreased berry set ("Thinning"), reduced hand-thinning costs, and hastened maturity:*

Guide: Apply one to four applications during bloom. Make only 1-to-2 applications for "Other Seedless Grapes". When the bloom period is extended, make subsequent sprays 1-to-7 days after the first application.

NOTE: Higher amounts or multiple applications cause an excess of shot berries or overthinning, especially in young vines or vines with high vigor. For "Other Seedless Grapes", new cultivars are very responsive and are over-thinned easily. Consult a local specialist before thinning unfamiliar cultivars.

*To help initiate the beginning of the berry growth period in Thompson Seedless variety "bump spray":*

Guide: Apply 16-to-24 grams a.i./acre as a single application during the period between the last thinning spray and the first sizing spray.

*For larger berries ("Sizing") and larger clusters when used in conjunction with established girdling and thinning practices:*

Guide: Apply one to four applications beginning when the average berry size reaches "target" diameter (See Table 1). Timing of the subsequent sprays is dictated by experience in the vineyard and temperatures occurring between sprays. Potential effect is reduced if the final spray occurs more than two weeks after the first application. Consult a local specialist before sizing unfamiliar cultivars.

**TABLE 1 Application Rates (Grams A.I./Acre) for Seedless Grape, Including Target Berry Diameters**

Seedless Grape	Stretch	Thinning	Sizing	
	grams a.i./acre	grams a.i./acre	"Target" Diameter	grams a.i./acre
Perlette	8-to-24	Not applicable	4-to-5 mm	32-to-128
Flame	8-to-24	3-to-16	6-to-9 mm	20-to-128
Thompson	8-to-24	8-to-20	3-to-5 mm	32-to-128
Raisin	8-to-24	3-to-12	3-to-5 mm	4-to-20
All Other Seedless Grape	Not applicable	0.5-12	3-to-14 mm	8-to-60

NOTE: High amounts of gibberellic acid reduces fruitfulness (cluster counts) the following year and delays berry skin color development, sugars accumulation and overall maturation.

#### SEEDED GRAPE

##### Emperor Grape

*For reducing berry shrivel. This can also increase berry size:*

Guide: Make applications as a whole vine spray, or as a spray or dip directly to the cluster.

Whole vine spray - Apply 20 grams a.i./acre as one application when the predominant berry diameter is 12-16 mm.

Directed spray to grape clusters or cluster dip - Prepare a spray solution of 40 to 50 ppm (16 to 20 grams a.i. per 100 gallons water) and apply as a direct spray to clusters or dip the clusters.

NOTE: Whole vine application reduces fruitfulness (cluster counts) the following year. High amounts of gibberellic acid also delay berry skin color development, sugars accumulation and overall maturation. Consult a local specialist before sizing unfamiliar cultivars.

##### Red Globe, Calmeria, Christmas Rose, Rogue and Queens

*To increase berry size:*

Guide: Make application as a whole vine spray, or as a spray or dip directly to the cluster.

Whole vine spray – Apply 20 grams a.i./acre as one application when the average berry size reaches the “target” diameter (See Table 2).

Directed spray to grape clusters or cluster dip – Prepare a spray solution of 40 to 50 ppm (16 to 20 grams a.i. per 100 gallons water) and apply as a direct spray to the cluster only or dip the clusters.

NOTE: Whole vine application reduces fruitfulness (cluster counts) the following year. High amounts of gibberellic acid delays berry skin color development, sugars accumulation and overall maturation. Consult a local specialist before sizing unfamiliar cultivars.

**TABLE 2 Application Rates for Seeded Grapes, Including Target Berry Diameters**

Seeded Grape	“Target” Diameter	Whole Vine Spray (grams a.i./acre)	Direct Spray to Cluster or Cluster Dip (rate in ppm of a.i.)
Emperor	12-16	20	40-50
Red Globe	12-18	20	40-50
Calmeria	12-16	20	40-50
Christmas Rose	12-16	20	40-50
Rogue	12-16	20	40-50
Queens	12-15	20	40-50

### **Black Corinth (Zante Currant) Grape**

*For improving berry size:*

Guide: Apply 1-to-12 grams a.i./acre as one application 3-to-5 days after full bloom, but before shatter begins.

### **Wine Varieties**

(Not for use in California)

*To increase cluster length, improve air circulation and light penetration within the cluster and help to reduce the incidence of bunch and sour rot:*

Guide: Apply one spray when clusters found in the dominant shoots arising from buds on count spurs are starting to elongate, average 3-to-4 inches in length, and show separation of the uppermost flower groups. Use 100 gallons of water per acre. Concentrations for registered varieties are shown below (See Table 3).

NOTE: Do not make applications less than three weeks before bloom. IT IS IMPORTANT that the proper rate be used on each variety; if late applications are made or if indicated rates are exceeded, reduction in yield may occur during the year of application and subsequent years. This reduction in yield may result from an increase in shot berries in the year of application and reduction in fruitfulness (cluster counts) in the first and second year following application. If growers have no experience with this application, consult local agricultural specialists before making this application.

**TABLE 3 Application Rates (Grams A.I./Acre) and Water Volume on Seeded Wine Varieties.**

Variety	Grams a.i./acre	Gallons/acre
Palomino Sauvignon Blanc Tinta Madeira	0.4-to-1	100
Aleatico Carignane Chardonnay Chenin Blanc French Colombard Pinot Noir Valdepenas	1-to-2	100
Barbera Petite Sirah Zinfandel	2-to-4	100

Variety	Grams a.i./acre	Gallons/acre
Green Hungarian	4-to-8	100
Grenache Alicante	8	100
Salvadore	8-to-16	100

## CITRUS

For all citrus, apply in sprays of sufficient water volumes to ensure thorough fruit wetting. Do not apply to trees of low vigor or under stress (pest, nutritional, or water, etc.) to avoid severe leaf and/or fruit drop. In most cases, some drop of older mature leaves will occur after application. Do not apply in white wash sprays in which lime or other caustic material has produced a high pH in the spray tank. Do not make applications of copper fungicides and/or oils within three weeks (before or after) application of GA<sub>3</sub> 20% to avoid significant leaf and fruit drop.

### NAVEL ORANGE

*To delay rind aging, reduce physiological disorders (e.g., rind staining, water spotting, sticky or tacky surface, puffy rind and rupture under pressure), and to produce a more orderly harvesting pattern. The delay in rind aging is greatest when an early spray is applied. This spray timing produces the firmest rind possible:*

Guide: Apply 16-to-48 grams a.i./acre in sufficient water volume to ensure thorough wetting.

**EARLY SPRAY:** Apply one spray approximately two weeks prior to color break, which normally occurs August through November.

**AND/OR**

**LATE SPRAY:** Apply one spray after marketable color has developed, normally from October through December.

**This late application may cause fruit re-greening.**

**NOTE:** Do not apply the early spray to groves that are harvested early, as fruit coloring will be delayed. Do not apply from January through July, as production will be reduced the following year. Expect slower color development in the target crop. After marketable color is achieved, treatment effects will be reduced the longer treated fruit remains on the tree.

**(For Florida Use Only)**

*To enhance fruit set and yield:*

Guide: Make a single application of 15-to-25 grams a.i./acre during December or January in 125-to-175 gallons of water per acre. Use a pure organo-silicone type surfactant at 0.05% (6 fl. oz./100 gallons).

### VALENCIA ORANGE

**(For California and Arizona use only)**

*To reduce rind creasing and to delay rind aging and softening:*

Guide: Apply a single spray in August to October to trees with a target crop of young fruit. Apply 40-to-80 grams a.i./acre in sufficient water volume to ensure thorough wetting.

**NOTE:** Do not apply the early spray to groves that are harvested early, as fruit coloring will be delayed. Do not apply from January through July, as production will be reduced the following year. Expect slower color development in the target crop. Increased re-greening of mature fruit may occur. After marketable color is achieved, treatment effects will be reduced the longer treated fruit remains on the tree.

**(For Florida Use Only)**

*To enhance fruit set and yield:*

Guide: Make a single application of 15-to-25 grams a.i./acre during December or January in 125 to 175 gallons of water per acre. Use a pure organo-silicone type surfactant at 0.05% (6 fl. oz./100 gallons).

### OTHER ROUND ORANGES

**(For Florida Use Only)**

*To reduce rind creasing and puffiness, and to delay aging and softening of the rind:*

Guide: Apply a single spray in August to October to trees with a target crop of young fruit. Apply 20-to-60 grams a.i./acre in sufficient water volume to ensure thorough wetting. Use a pure organo-silicone type surfactant at 0.05% (6 fl. oz. in 100 gallons).



## **AMBERSWEET ORANGE** **(For Florida Use Only)**

*To enhance fruit set and yield:*

Guide: Make a single application of 15-to-25 grams a.i./acre during January in 125-to-175 gallons of water per acre with a pure organo-silicone type surfactant at 0.05% (6 fl. oz./100 gallons).

### **LEMON/LIME**

*To decrease the amount of small ripe fruit and to produce a more desirable production pattern relative to market demand:*

Guide: Apply one spray when target crop is ½-to-¾ full size, but still green. Use 10-to-32 grams a.i./acre in sufficient water volume to ensure thorough wetting.

When applied two years in a row, an even larger difference in harvest pattern and maturity occurs.

### **TANGERINE HYBRIDS**

*To delay disorders associated with rind aging, puffiness, and softening, and to increase peel strength of tangerine hybrids such as Orlando, Robinson, Minneola and Sunburst:*

Guide: Apply 20-to-40 grams a.i./acre approximately two weeks prior to color break. Apply in sufficient water volume to ensure thorough wetting.

NOTE: Do not apply if early harvest is planned. Do not apply after coloring as preharvest rind staining may occur. Application during coloring may cause variation in rind color of development.

**(All States except California)**

*To increase fruit set and yields on tangerine hybrids with pollination problems such as the Orlando, Robinson, Minneola and Sunburst:*

Guide: Apply 8-to-30 grams a.i./acre during full bloom. Make one to two applications. Apply in sufficient water volume to ensure thorough wetting.

NOTE: Expect reduced fruit sizes and slightly retarded color development. A slight increase in mature leaf drop will occur in trees under stress.

### **GRAPEFRUIT**

**(All States except California)**

*To delay disorders associated with rind aging (e.g., puffiness, softening, and orange coloration), to prevent preharvest drop of mature fruit, to increase peel strength and reduce water loss during storage, and to produce a more orderly harvesting pattern. The delay in rind aging is greatest when an early spray is applied before color change. This spray timing produces the firmest rind possible:*

Guide: Apply 16-to-48 grams a.i./acre in a minimum of 250 gallons per acre.

EARLY SPRAY: Apply one spray approximately two weeks prior to color break, which normally occurs August through September.

**AND/OR**

LATE SPRAY: Apply one spray after marketable color has developed which is normally from October through December.

**This late application may cause fruit re-greening.**

NOTE: Do not apply the early spray to groves that are harvested early as fruit coloring will be delayed. Spot pick heavy crops to aid early marketing and to avoid reduction of yields, which generally follow late held crops. Fully colored fruit to which applications have been made will begin to re-green if allowed to remain on the tree for extended periods.

Application made after December, or when trees begin to break dormancy, will adversely affect new crop. Do not use concentrate sprays. Results vary from season to season depending on environmental conditions.

*To enhance fruit set, size and yield:*

Guide: Make a single application of 15-to-25 grams a.i./acre during December or January in 125-to-175 gallons of water per acre with a pure organo-silicone type surfactant at 0.05% (6 fl. oz./100 gallons).

### **GRAPEFRUIT, STAR RUBY VARIETY**

**(All States except California)**

*To reduce early-season small fruit drop of Star Ruby Variety thereby increasing yields:*

Guide: Apply a single spray during the bloom period. Use 25-to-35 grams a.i./acre in a minimum of 250 gallons of water per acre.

NOTE: Results vary from season to season depending on environmental conditions. Maintain a well-balanced fertilization and watering program.

### **CLEMENTINE MANDARIN**

*To increase fruit set and yield:*

Guide: Apply 1 to 8 grams per 100 gallons of water (2.5-20ppm) in 1 to 2 applications starting at 10% bloom up to 3 weeks after petal fall. Make applications as a dilute spray in sufficient gallonage to ensure thorough wetting. The number and timing of applications is dependent upon the amount of fruit set desired. In general, more fruit will set with more than 1 application and higher rates tend to set more fruit. Differences in the strain of Clementine Mandarin will also interact with the above factors to affect the degree of fruit set achieved.

NOTE: Applications may cause severe leaf drop.

### **POSTHARVEST APPLICATIONS**

(All States except California)

#### **LEMON**

*To delay fruit senescence and prolong storage life:*

Guide: Mix 2 to 4 grams of a.i. in 10 gallons of storage wax which has been diluted as per the wax label instructions. The incidence of sour rot is reduced by delaying senescence.

#### **YELLOW LEMONS AND OTHER MATURE CITRUS FRUIT**

*To delay rind senescence and color changes:*

Guide: Mix 2 to 4 grams of a.i. in 10 gallons of storage wax which has been diluted as per the wax label instructions. The incidence of sour rot is reduced by delaying senescence.

---

## **FRUIT CROPS**

---

### **BANANA**

(All States except California)

*To stimulate plant growth and overcome the effects of stress caused by insects, disease or adverse weather:*

Guide: Apply 1-to-6 grams a.i./acre by ground or aerial application in sufficient water volume to adequately cover foliage. Make applications once every 30 to 90 days throughout the year. Make more frequent applications (monthly) for 6 months prior to anticipated weather stress periods.

### **BLUEBERRY**

(All States except California)

*To improve fruit set. For natural fruit set problems due to insufficient natural honeybee pollination, adverse weather conditions, or physiological factors.*

*Highbush blueberry – (for varieties such as Coville, Jersey, Stanley, Earliblue, Weymouth, Walcott, Berkeley, Blueray, Bluecrop, 1316A, Concord, and others).*

Guide: Make one or two applications in 40 to 100 gallons of water. If a single application is made, apply 80 grams a.i./acre at full bloom (when 75% of the flowers are fully open). When 2 applications are made, apply 40 grams a.i./acre per application. Make the first application at full bloom, and the second one within 10-14 days of the first one. For Weymouth, delay application up to two weeks after bloom to increase size of “shot” berries.

*Rabbiteye blueberry – (for varieties such as Aliceblue, Beckyblue, Bonita, Brightwell, Climax, Delite, Tiftblue, Woodward, and others).*

Guide: Make a single application of 40 grams a.i./acre in 100-to-300 gallons of water per acre when most of the flowers are elongated but not yet open (bloom Stage 5).

**OR**

**MULTIPLE APPLICATIONS:** Make two to four applications 10-to-14 days apart starting at bloom Stage 5. Spray 20-to-40 grams a.i./acre in 40-to-100 gallons of water per application.

### **SWEET CHERRY**

*To produce larger, brighter colored, firmer fruit:*

Guide: Apply a single spray when the fruit is light green to straw colored. Use 16-to-48 grams a.i./acre in sufficient water volume to ensure thorough wetting.

NOTE: Color development and harvest will be slightly delayed.

### **RED TART CHERRY**

**(All States except California)**

*To maintain and extend high fruiting capacity of bearing tart cherry trees and reduce the occurrence of "blind" nodes. Treatment will cause bud differentiation, which is apparent the year after application. Therefore, changes in shoot, spur, and flower production will not be evident until two or three years after program initiation. Make annual applications to ensure vegetative development and subsequent yield improvement year after year.*

Guide: Apply one spray 14-to-28 days after bloom. Optimum timing is defined as that stage when 3-to-5 terminal leaves have fully expanded, or, at least 1-to-3 inches of terminal shoot extension has occurred. Use 4-to-18 grams a.i./acre, depending on tree age and vigor (See Table 4). Apply in sufficient water volume to ensure thorough wetting.

**TABLE 4 Application Rates (Grams A.I./Acre) for Tart Cherry Trees by Age**

<b>Tree Age (years)</b>	<b>Rate (grams a.i./acre)</b>
6-to-10	4-to-6
11-to-15	8-to-10
16-to-20	11-to-14
20 + years	14-to-18

NOTE: Rates are based on expected normal tree vigor at various ages. Adjust rate according to tree vigor. If trees are vigorous, use the lowest rates. Use lowest rates on trees that have been heavily pruned or hedged. Use higher rates for trees low in vigor and weak in shoot and spur production. Excessive application rates will increase vegetative growth at the expense of fruit production the following year. Applications will not improve growth of trees under stress conditions, such as nutritional, moisture, or pest. Best results are obtained when combined with good cultural practices.

### **STONE FRUIT GROUP**

*To improve fruit quality and increase firmness in the season of application:*

Guide: Apply a single spray of 16-to-32 grams a.i./acre in sufficient water volume to achieve complete coverage of fruits and foliage. Make application one to four weeks prior to the beginning of harvest.

NOTE: This application causes reduction in flower counts the year following application, particularly if application is made during the months of May through July.

### **ITALIAN PRUNE**

**(All States except California)**

*To reduce internal browning, improve quality, and increase size:*

Guide: Apply 4-to-5 weeks before expected harvest. Apply a single spray at 16-to-48 grams a.i./acre in sufficient water volume to ensure thorough wetting.

NOTE: Color development and harvest will be slightly delayed. Application can reduce bloom the following season.

### **NON-BEARING FRUIT TREES**

**(All States except California)**

*To reduce flowering and fruiting in young stone fruit trees in order to minimize the competitive effect of early fruiting on tree development:*

NOTE: DO NOT SPRAY TREES IN THE FIRST YEAR. Treat in the second season for reduction of flowering in the third season, and again in the third season if a reduction of flowering and fruiting is desired in the fourth season. Treat only trees that are in good physiological condition. Discontinue treatment the year before desired harvest.

Guide: Apply a single application of 20-to-80 grams a.i./acre during the period of flower bud initiation for the following year. Use sufficient water to achieve good coverage of the canopy. A local horticulturist can provide timings and rates for specific cultivars in your area.

---

## OTHER FRUIT

---

### STRAWBERRY

(All States except California)

*To increase runner production of mother plants:*

Guide: Apply a single spray of 15-to-25 grams a.i./acre to mother plants 10-to-30 days after planting when plants have 1-to-6 leaves. Apply 100 gallons spray/acre to thoroughly wet new foliage to the point of run-off.

NOTE: Not for use on fruiting plants. Treatments will not be effective on plantings set out after mid-May. Response varies with cultivar and location. Consult a local horticulturist for specific information in your area.

### CRANBERRY

(All States except California)

*To reduce or eliminate crop in the year of application:*

Guide: Make a single application of 10-to-50 grams a.i./acre at early bloom (2-5% scatter bloom) in sufficient water volume to ensure thorough coverage.

NOTE: Applications made later than the early bloom stage will have no effect or result in increased fruit set. Responses vary with cultivar, age of the bog and location. Consult a local specialist for specific information in your area.

---

## VEGETABLE CROPS

---

### ARTICHOKE

*To accelerate maturity and shift harvest to an earlier date:*

Guide: For perennials, apply one to three applications at bud initiation stage. For annuals, apply one to four applications at 2-week intervals, beginning at the fourth true leaf. Use 10-to-20 grams a.i./acre per application in sufficient water volume to ensure thorough wetting of the entire plant (leaves, stems and buds).

### CARROT

#### Fresh and Processing

*To delay leaf senescence:* Maintaining vigorous foliage may reduce the incidence of infection by *Alternaria dauci*.

Guide: Make the first application 4-to-6 weeks after emergence using ground or aerial spray equipment with spray concentrations of 20-to-30 ppm. A second spray 14 days later may be required to achieve the desired amount of foliar recovery in severe disease situations or cool weather. Do not apply more than twice per crop.

NOTE: Dilutions of greater concentration increase the risk of excessive top growth, particularly with a second application.

### CELERY

*To increase plant height and yield and overcome stress due to cold weather conditions or saline soils, and to obtain earlier maturity:*

Guide: Apply a single spray one to four weeks prior to harvest at a rate of 2.5-to-10 grams a.i./acre. Use 25-to-50 gallons of water per acre by ground application or 5-to-10 gallons of water per acre for aerial application\*. Use lower concentrations applying 3-to-4 weeks before harvest and higher concentrations within 1-to-2 weeks before harvest.

\*Do not apply by air in California.

NOTE: Do not apply earlier than 4 weeks before harvest as bolting (seed stalk formation) may occur.

### LETTUCE FOR SEED

*To obtain uniform bolting and increase seed production:*

Guide: Apply one to four applications at 2-week intervals, beginning at the fourth true leaf. Use 1-to-4 grams a.i./acre per application in sufficient water volume to ensure thorough wetting.

### PEPPER

(All States except California)

*To Promote Plant Growth:*

Guide: Apply one to two sprays of 1-to-3 grams a.i./acre in 25-to-50 gallons of water per acre at two-week intervals. Begin sprays 2 weeks after transplanting.

NOTE: This use is for areas with short growing seasons, or when low temperatures slow plant growth.

*To Increase Fruit Set and Promote Fruit Growth:*

Guide: Apply one to two sprays of 1-to-3 grams a.i./acre in 25-to-50 gallons of water per acre at weekly intervals during the flowering period. The high rate is suggested for areas and/or varieties with pollination and/or fruit set problems.

*To Increase Fruit Size:*

Guide: Apply 1-to-3 grams a.i./acre in 25-to-50 gallons of water per acre at the beginning of the picking period. The high rate is for plants with heavy fruit loads.

**MELON AND CUCUMBER**  
**(All States except California)**

*To stimulate fruit set during periods of cool temperatures:*

Guide: Use 1-to-4 grams a.i./acre in sufficient water volume for thorough coverage of exposed foliage. Make one application prior to bloom followed by two additional applications at intervals of 10-to-14 days on cantaloupes and watermelons. On cucumbers, up to four applications are required.

For maximum benefits, vines in good condition are required, except for reduced rate of growth due to cool temperatures.

**RHUBARB**

*To break dormancy on plants receiving insufficient chilling and to increase marketable yield of forced rhubarb:*

Guide: Make a single application of 2 fluid ounces (60 ml) of a solution containing 20 grams a.i. in 10 gallons of water to each cleaned crown, when the rest period is not completely broken. When the rest period is broken by cold weather, apply 2 fluid ounces (60 ml) of a solution containing 10 grams a.i. in 10 gallons of water to each cleaned crown.

NOTE: Keep forcing house temperatures at 40° F-to-50° F for 24 hours after application. If house is warmer than 50° F, cover the crowns with plastic. Temperatures in the forcing house above 50° F may lower yields and cause poor stalk color.

**SEED POTATO**

*To stimulate uniform sprouting to aid in maximum production, more uniform development, fewer late maturing plants, and to break dormancy of newly harvested potatoes that have not had a full rest period.*

Guide: Dip whole or cut seed pieces in a solution containing 0.2-to-0.4 grams a.i. in 100 gallons of water prior to planting.

NOTE: Under high soil temperatures use the minimum concentration for dormant seed. Do not treat rested seed pieces.

**SPINACH**

**(All States except California)**

*To facilitate harvest, increase yield and improve quality of fall and over-winter spinach.*

Guide: Apply a single spray 10-to-18 days before each anticipated harvest on fall or over-winter spinach, ideally when daytime temperatures are 40° F-to-70° F and during early morning hours when dew is present on crop. Apply 6-to-10 grams a.i./acre in 10-to-50 gallons of water per acre by ground sprayer or in a minimum of 5-to-10 gallons of water per acre by air. When applied to promote growth of second cutting, wait until some regrowth has started before spraying. Maximum benefit is obtained when below normal temperatures predominate following application and growth is otherwise slowed in untreated spinach.

NOTE: Since the promotion of bolting may occur, do not apply after the mid-winter period or if temperatures are expected to exceed 75° F within several days of application. Do not apply on spring plantings.

---

**ORNAMENTAL CROPS, CUT FLOWERS AND TURFGRASS**

---

The following suggestions are based on results with common cultivars. Differences in responsiveness vary between cultivars, growing conditions, and cultural management systems. Therefore, prior to widespread usage, test a small number of plants from each cultivar under a specific set of growing and cultural management conditions to verify desired efficacy.

When applying foliar applications of GA<sub>3</sub> 20%, spray plants to run-off. The actual spray application rate will vary, depending on plant size and spacing density. Thorough spray coverage is essential for uniform flowering.

NOTE: A representative spray application rate which has been proven effective for 6 inch potted plants spaced at a density of 1 per square foot is 1-gallon spray solution/200 square feet.

## **ORNAMENTALS** **(All States except California)**

### **AZALEA**

*As a partial replacement of cold treatment to break flower dormancy:*

Guide: Apply three sprays of 250-to-500 ppm a.i. at weekly intervals after 3-to-4 weeks of chilling. At the earliest, initiate treatment at Stage 5 of floral development (i.e., style elongated and open). A representative spray schedule would consist of applications made at 3, 10, and 17 days after four weeks of chilling. Flowers will not develop properly if applied prior to Stage 5.

NOTE: Thorough spray coverage is essential for uniform flowering. Do not apply after flower buds show color. On some cultivars (e.g., 'Gloria', 'Prize', and 'Redwing'), a single spray of 1000-ppm a.i. after 4 weeks of chilling has proven effective in breaking dormancy.

*As a complete substitution of cold treatment to break flower dormancy:*

Guide: Apply four to six sprays of 1000-ppm a.i. at weekly intervals. At the earliest, apply first spray when plants are at Stage 5 of floral development (style elongated and open). Flowers will not develop properly if applied prior to Stage 5 of floral development.

NOTE: Thorough spray coverage is essential for uniform flowering. Do not apply after flower buds show color.

*To inhibit flower bud initiation during vegetative growth:*

Guide: After each pinch, apply two to three sprays of 100-to-750 ppm a.i. at intervals of 2-to-3 weeks. Continue applications on a weekly basis for 1-to-2 weeks after the first application.

NOTE: Apply a maximum of three applications.

### **CALLA LILY**

*For increased flowering:*

Guide: Soak rhizome or tuber in a 500 ppm solution of GA<sub>3</sub> 20% for 10 minutes prior to planting.

NOTE: Flower or leaf stretching occurs in some cultivars. Reduce rates if this occurs.

### **CAMELIA**

*For substitution of chilling requirements and to increase bloom size:*

Guide: Prepare a 2.0% (a.i.) solution of GA<sub>3</sub> 20% by mixing 10 g GA<sub>3</sub> 20% in 100 ml water. Remove the vegetative bud immediately adjacent to or below the floral bud. Place a single drop of the prepared solution to the vegetative bud scar.

NOTE: Adding a deposition aid (such as carboxymethylcellulose) to thicken the solution will decrease run-off.

### **CYCLAMEN**

*For uniform flowering:*

**Bud Application** – Apply 8 ml (0.25 fl. oz.) of a 10-to-15 ppm solution directly to the crown when buds are pinhead size in the leaf axils.

**Foliar Application** – Apply a single foliar application of 25 ppm directly toward the crown and adjacent leaves when buds are pinhead size in the leaf axils. Thoroughly wet the crown.

NOTE: Both bud and foliar applications have been shown to promote uniform flowering. Late or excessive applications result in weakened floral stems or poorly formed flowers.

### **FUCHSIA**

*To produce tree forms of common fuchsia cultivars by stem elongation:*

Guide: Apply a single foliar application of 250 ppm for four consecutive weeks. Begin applications after the plant has reached the desired size. Spray the entire plant to the point of run-off.

NOTE: Staking is required after treating plants. Concentrations higher than 250 ppm may cause plants to become stretched and spindly, with weakened stems.

## **GERANIUM**

**Cuttings** - *To increase flower number and size of geranium cuttings:*

Guide: Apply a single foliar application of 1-to-5 ppm when inflorescence first begins to show color. Direct spray at the developing inflorescence.

NOTE: Concentrations above 5 ppm or treatments prior to inflorescence showing color cause peduncle stretching.

**Seedlings** – *To advance flowering:*

Guide: Make a single foliar application of 5-to-15 ppm when first flower bud set is noted. Spray the entire plant to the point of run-off.

NOTE: Incorrect timing or concentrations above 15 ppm cause plant stretching.

**Tree Forms** – *To produce tree forms of common geranium cultivars by stem elongation:*

Guide: Make a single foliar application of 250 ppm for four consecutive weeks. Spray the entire plant to the point of run-off.

NOTE: Plants require staking after treatment.

## **HYDRANGEA**

*To substitute for chilling requirements to break flower bud dormancy:*

Guide: Apply a single foliar application of 2-to-5 ppm for one to four consecutive weeks. Begin applications at the start of forcing. Thoroughly cover all growing points containing flower buds.

NOTE: Overuse or concentrations above 5 ppm may result in stretched, spindly, and weak stems.

## **POMPOM CHRYSANTHEMUM**

*For elongating peduncles on Pompom chrysanthemum.*

Guide: Apply a single foliar application of 25-to-60 ppm 4-to-5 weeks after initiation of short day conditions. Apply directing the spray solution towards the flower buds.

NOTE: Overuse of incorrect timing may cause long, spindly, and weak stems.

## **SPATHIPHYLLUM**

*To accelerate bloom and increase the number of flowers per plant:*

Guide: Make a single foliar application of 150-to-250 ppm approximately 9-to-12 weeks prior to the expected sale date. Spray to the point of run-off and thoroughly wet all growing points.

NOTE: Flower distortion or leaf stretching may occur on some cultivars such as 'Petite', 'Starlight', 'Tasson', and 'Mauna Loa'. Reduce rates if this occurs. On other cultivars, prior to application on a commercial basis, evaluate the effects of GA<sub>3</sub> 20% on a small number of plants.

## **AGLAONEMA, ANTHURIUM, DIFFENBACHIA (Dumb Cane) AND SYNGONIUM**

*To accelerate bloom and increase the number of flowers per plant:*

Guide: Apply a single foliar application of 250-to-500 ppm a.i. for one to four consecutive weeks beginning at the start of forcing for Aglaonema, Anthurium and Diffenbachia. Apply a single foliar application of 500-to-2000 ppm a.i. for one to four consecutive weeks beginning at the start of forcing for Syngonium. Thoroughly apply solution to all growing points containing flower buds.

NOTE: Application of GA<sub>3</sub> 20% can increase flower yield and decrease time to flowering. Make 1 or 2 applications during the vegetative phase of plant development to induce bloom. On other cultivars, prior to application on a commercial basis, evaluate the effects of GA<sub>3</sub> 20% on a small number of plants.

## **CUT FLOWERS**

**(All States except California)**

Apply GA<sub>3</sub> 20% to ornamental plants grown for cut flowers to promote stem elongation and flowering. GA<sub>3</sub> 20% is very active and application at excessive rates results in undesirable effects. Evaluate the effects of GA<sub>3</sub> 20% on a small number of plants prior to application on a widespread basis.

## **ASTER**

Monte Carlo-type, Novi-type and Belgi-type – *To promote stem elongation and break dormancy:*

Guide: Make 1-to-3 applications of 50-to-100 ppm during the early vegetative period when plants are 2” to 6” tall. Make applications at 2-to-3 week intervals.

#### **BABY’S BREATH (Gypsophila)**

*To accelerate plant growth, increase flower yield and uniformity:*

Guide: Make 3-to-4 applications of 150-to-500 ppm a.i. solution at 4 weeks of growth (after pinching). Make applications at 2-week intervals.

#### **BELLS OF IRELAND (Moluccella)**

*To promote plant growth and stem elongation:*

Guide: Apply a 50-to-100 ppm a.i. solution when plants are 4” to 8” tall. Make applications at 2-to-3 week intervals.

#### **BUPLUREUM**

*To promote plant growth and stem elongation:*

Guide: Apply a 50-to-100 ppm a.i. solution as a foliar spray when plants are 4” to 8” tall. Make applications at 2-to-3 week intervals.

#### **CAMPANULA**

*To promote plant growth and stem elongation:*

Guide: Apply a 50-to-100 ppm a.i. solution as a foliar spray when plants are 4” to 8” tall. Make applications at 2-to-3 week intervals.

#### **CANDY TUFT (Iberis)**

*To promote plant growth and stem elongation:*

Guide: Apply a 50-to-100 ppm a.i. solution as a foliar spray when plants are 4” to 8” tall. Make applications at 2-to-3 week intervals.

#### **COLUMN STOCK (Matthiola)**

*To promote plant growth and stem elongation:*

Guide: Apply a 50-to-100 ppm a.i. solution as a foliar spray when plants are 4” to 8” tall. Make applications at 2-to-3 week intervals.

#### **DELPHINIUM**

Including: *D. elatum*, *D. grandiflorum*, *D. belladonna*, *D. cardinale*, *D. nudicale*, and Delphinium hybrids

- *To promote plant growth and stem elongation:*

Guide: Apply a 50-to-100 ppm a.i. solution as a foliar spray when plants are 4” to 8” tall. Make applications at 2-to-3 week intervals.

#### **DIDISCUS (Trachyme)**

*To promote plant growth and stem elongation:*

Guide: Apply a 50-to-100 ppm a.i. solution as a foliar spray when plants are 4” to 8” tall. Make applications at 2-to-3 week intervals.

#### **HYDRANGEA**

*To promote plant growth and stem elongation:*

Guide: Apply a 50-to-100 ppm a.i. solution as a foliar spray when plants are 4” to 8” tall. Make applications at 2-to-3 week intervals.

#### **LARKSPUR**

*Consolida ambigua*, *C. orientalis*, *Delphinium ajacis* - *To promote plant growth and stem elongation:*

Guide: Apply a 50-to-100 ppm a.i. solution as a foliar spray when plants are 4” to 8” tall. Make applications at 2-to-3 week intervals.

#### **LISIANTHUS (Eustoma)**

*To promote plant growth and stem elongation:*

Guide: Apply a 50-to-100 ppm a.i. solution as a foliar spray when plants are 4” to 8” tall. Make applications at 2-to-3 week intervals.

#### **PHLOX**

*Phlox paniculata* and *Drummondii hybrida* - *To promote plant growth and stem elongation:*

Guide: Apply a 50-to-100 ppm a.i. solution as a foliar spray when plants are 4” to 8” tall. Make applications at 2-to-3 week intervals.



### **QUEEN ANNE'S LACE (Ammi)**

*To promote plant growth and stem elongation:*

Guide: Apply a 50-to-100 ppm a.i. solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2-to-3 week intervals.

### **SAFFLOWER**

*To promote plant growth and stem elongation:*

Guide: Apply a 50-to-100 ppm a.i. solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2-to-3 week intervals.

### **SOLIDASTER**

*To promote plant growth and stem elongation:*

Guide: Apply a 50-to-100 ppm a.i. solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2-to-3 week intervals.

### **STATICE**

*To promote earlier flowering and to increase flower yield:*

Guide: Apply as a foliar spray consisting of 10 ml (0.33 fl. oz.) of a 400-to-500 ppm a.i. solution to each plant when plants are 10 inches or more in diameter (approximately 90-to-110 days after normal seeding time).

NOTE: Do not exceed specified rates. Do not apply repeated sprays. Accelerated flowering is influenced by extended photoperiod, adequate nutrition, and reduced night temperatures. This treatment reduces the cold requirement and/or the long photoperiod.

*To promote plant growth and stem elongation:*

Guide: Apply a 50-to-100 ppm a.i. solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2-to-3 week intervals.

### **SUNFLOWER**

*To promote plant growth and stem elongation:*

Guide: Apply a 50-to-100 ppm a.i. solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2-to-3 week intervals.

### **SWEET WILLIAM (Dianthus)**

*To promote plant growth and stem elongation:*

Guide: Apply a 50-to-100 ppm a.i. solution as a foliar spray when plants are 4" to 8" tall. Make applications at 2-to-3 week intervals.

## **TURF**

(All states except California)

### **(Golf Courses, Parks and Turf Farms)**

Application of GA<sub>3</sub> 20% to Bermudagrass grown in golf courses, parks and turf farms has been shown to initiate or maintain growth and prevent color change during periods of cold stress.

NOTE: Do not exceed specified rates. Maintain adequate moisture and proper fertilization programs for your local area. Keep high rate applications at least 2-weeks apart. Do not use on dormant turf. Stop treatments if thinning is observed. More frequent mowing may be necessary.

#### **Cool Weather Application**

### **BERMUDAGRASS (Tidwarf, Tifgreen, and other cultivars)**

*To initiate or maintain growth and prevent color change during periods of cold stress and light frosts:*

Guide: Apply 10 grams a.i./acre weekly or 25 grams a.i./acre biweekly in 25-to-100 gallons of water/acre.

#### **Warm Weather Application**

### **BERMUDAGRASS (Tidwarf, Tifgreen, and other cultivars)**

*To maintain or enhance regrowth of golf course Bermudagrass during summer months:*

Guide: Apply 1-to-3 grams a.i./acre weekly in 25-to-100 gallons of water/acre.

## **BEDDING PLANTS, ANNUAL AND PERENNIAL POTTED CROPS, FIELD GROWN ORNAMENTALS AND BULB CROPS**

(All States except California)

*To promote plant growth and/or overcome the effects of overuse of a gibberellin-inhibiting plant growth regulator:*

Guide: Apply a single application of 1-to-25 ppm a.i. solution directly to plant foliage. When applying GA<sub>3</sub> 20% to promote plant growth, start with a 1 ppm a.i. solution unless previous experience dictates a higher rate is warranted. If desired results are not achieved, a reapplication or higher rate is necessary.

NOTE: GA<sub>3</sub> 20% is very active and excessive application rates result in undesirable stem elongation. Do not use more than 25 ppm a.i. Evaluate the effects of GA<sub>3</sub> 20% on a small number of plants prior to application on a widespread basis.

---

## OTHER CROPS

---

### HOPS

(Northwestern U.S. only)

For seeded and seedless Fuggle hop and similar varieties adapted to the Northwestern states.

*To increase yield and fruit set:*

Guide: Apply a single spray when vine growth is 5-to-8 feet in length. Use 4-to-6 grams a.i./acre in 100-to-150 gallons of water/acre.

### COTTON

(All states except California)

*To promote early plant growth and increase early seedling vigor on young cotton plants:*

Guide: Apply 1-to-6 grams a.i./acre via in-furrow application to seed, or as a foliar application from the cotyledon stage through the 7-leaf/node stage. Repeat applications to a maximum of 3 applications.

NOTE: Use higher rates when temperatures will likely average 75° F or less during the 14 days following the applications. Do not tank mix with herbicides. Do not apply to cotton plants under drought stress. If cotton plants are under continuous stress, delay application until the stress is alleviated and plants are beginning to recover. Do not apply more often than necessary to achieve the desired height, as overdosage results in excessive growth.

#### Mixing Instructions

Fill the treatment tank with half of the final tank mix volume. Add the required amount of GA<sub>3</sub> 20% and mix thoroughly while adding water to the desired final volume. Dispose of any unused spray material at the end of the day.

#### Application Equipment

Apply GA<sub>3</sub> 20% by aerial or ground spray equipment. As an aerial spray, use a spray system capable of producing a uniform spray pattern of medium to fine spray droplets at 10 gallons per acre (GPA). Apply no less than 3 GPA of total spray volume. Low pressure ground sprayers equipped with boom and flat fan nozzles using 10 to 15 GPA spray volume may be used.

#### Compatibility with Other Chemicals

Compatibility information regarding tank mixtures of GA<sub>3</sub> 20% with herbicides used in cotton is not available.

### GRAIN SORGHUM

(All States except California)

*For use as a seed treatment to break dormancy and allow germination under cold soil conditions:*

Guide: Apply 0.25 to 1.00 grams a.i. per 100 pounds of seed. GA<sub>3</sub> 20% can be applied to dry seed with standard mist-treating equipment. Make certain the seed is completely and uniformly covered with GA<sub>3</sub> 20%. Fill the seed treatment tank with water to one-half the final tank mix volume. Add the required amount of GA<sub>3</sub> 20%, mixing thoroughly while adding water and other seed treatment products to the desired final volume.

DO NOT USE TREATED SEED FOR FOOD, FEED OR OIL PURPOSES. Add an approved dye to distinguish GA<sub>3</sub> 20% treated seed and prevent inadvertent use for food, feed or oil purposes. Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the federal and state seed laws. GA<sub>3</sub> 20% is compatible with most commonly used fungicide seed treatments such as Vitavax® and Dithane®, standard dyes and sticker-binding agents. When preparing tank mixes, ensure adequate physical compatibility and mixing characteristics. Refer to the Compatibility Test for Tank Mix Components under the GENERAL INFORMATION section of this label for additional information.

## **RICE**

**(All States except California)**

### **FOLIAR APPLICATION**

Early season foliar application of GA<sub>3</sub> 20% promotes vigorous and more uniform seedling growth of rice prior to permanent flood establishment. This may permit earlier flooding (5 to 10 days earlier) of drill or broadcast-seeded rice and is particularly effective on semi-dwarf varieties. Early flooding may reduce the additional flushing costs associated with a delay in establishing the permanent flood, reduce weed infestations and the number of herbicide applications, and/or promote earlier and more uniform grain maturity.

Late season foliar applications of GA<sub>3</sub> 20% between split-boot and 100% heading increases panicle height of rice. This may facilitate harvest efficiency in the field by allowing the rice grain to be cut above the leaf canopy at faster combine speeds and at reduced vegetative load. Grain quality and maturity are advanced with the promotion of tiller panicle development. Heading applications to the first crop accelerates regrowth of second crop rice. This results in earlier second crop maturity and maximized grain yield.

#### **Timing and Rate Instructions**

##### **Seedling Applications (Early Season)**

Apply GA<sub>3</sub> 20% at a rate of 1 to 3 grams a.i. per acre to rice between the 1-2 and 4-5 leaf stages of growth. Timing and dosage is based upon environmental conditions, tank mix combinations with herbicides, and preferred permanent flood practice in relation to rice leaf stage.

For best results, apply GA<sub>3</sub> 20% at a rate of 1 to 2 grams a.i. per acre using either a non-ionic surfactant known to be non-phytotoxic to rice or in tank mix combination with rice herbicides (See Compatibility with Other Chemicals section). Use higher rates of 1.5 to 3 grams a.i. per acre with some dry and water-based herbicide formulations, or when temperatures will likely average 75°F or less during 14 days after application.

NOTE: Do not apply when rice is subjected to drought stress conditions. Accelerated growth rates following GA<sub>3</sub> 20% application can temporarily cause foliage to appear lighter green in color.

##### **Panicle Extension Applications (Late Season)**

Apply GA<sub>3</sub> 20% at a rate of 3 to 8 grams a.i. per acre between split-boot and 100% panicle heading to promote main culm and tiller panicle extension. Tank mix with a non-ionic surfactant known to be non-phytotoxic to rice. Timing and dosage is based upon environmental conditions, tank mix combinations with herbicides, and preferred permanent flood practice in relation to rice leaf stage.

NOTE: Do not apply when rice is subjected to drought stress conditions. Accelerated growth rates following GA<sub>3</sub> 20% application can temporarily cause foliage to appear lighter green in color.

#### **Mixing Instructions**

Fill the treatment tank with half of the final tank mix volume. Add the required amount of GA<sub>3</sub> 20% and mix thoroughly while adding water to the desired final volume. Dispose of any unused spray material at the end of the day.

#### **Application Equipment**

Apply GA<sub>3</sub> 20% by aerial or ground spray equipment. As an aerial spray, use a spray system capable of producing a uniform spray pattern of medium to fine spray droplets at 10 gallons per acre (GPA). Apply no less than 3 GPA of total spray volume. Use low pressure ground sprayers equipped with boom and flat fan nozzles using 10 to 15 GPA spray volume.

#### **Compatibility with Other Chemicals**

GA<sub>3</sub> 20% can be tank mixed with most commonly used rice herbicides and fungicides. When applying GA<sub>3</sub> 20% in mixtures with Arrosolo<sup>®</sup>, Riverside Propanil 60DF, Stam<sup>®</sup> M4, Stam<sup>®</sup> 80EDF, or Wham!<sup>®</sup> EZ, plus one of their recommended adjuvants, use of an additional surfactant is not necessary. Do not apply GA<sub>3</sub> 20% with products containing fenoxaprop-*p*-ethyl as the active ingredient. When preparing tank mixes, ensure adequate physical compatibility and mixing characteristics. Refer to the Compatibility Test for Tank Mix Components under the GENERAL INFORMATION section of this label for additional information.

### **SEED TREATMENT APPLICATION**

Use GA<sub>3</sub> 20% as a seed treatment for rice. GA<sub>3</sub> 20% stimulates seed germination and promotes faster and more uniform stand establishment. The following table describes GA<sub>3</sub> 20% application and expected benefits.

**GA<sub>3</sub> 20% Seed Treatment Application**

<b>Crop</b>	<b>GA<sub>3</sub> 20% Use Rates</b>	<b>Important Considerations</b>	<b>Benefits</b>
Rice	0.5 to 2.1 grams a.i. in 8-20 fl oz water/100 lbs seed	For use with drill or broadcast seeding systems. Do not apply GA <sub>3</sub> 20% prior to a 24 hour presoak or to water used for the presoak. Do not exceed 2.1 grams a.i./100 lbs of seed.	GA <sub>3</sub> 20% promotes germination and emergence for semi-dwarf and tall varieties. GA <sub>3</sub> 20% helps increase final stand density and uniformity when seed are planted deeper to receive adequate moisture.

**Mixing Instructions**

Apply GA<sub>3</sub> 20% to seed with standard mist treating equipment. For best results, higher treatment volume of 12 to 20 fl oz per 100 pounds of seed ensures complete and uniform coverage.

Fill the treatment tank with half of the final tank mix volume. Add the required amount of GA<sub>3</sub> 20% and mix thoroughly while adding water and other co-applied seed treatment products (see Compatibility with Other Chemicals section) to the desired final volume.

Add an approved dye to distinguish GA<sub>3</sub> 20% treated seed and prevent inadvertent use for food, feed or oil purposes. Treated seed must be labeled in accordance with the requirements of the Federal Seed Act.

**Use Restriction**

Do not use treated seed for food, feed or oil purposes.

**Compatibility with Other Chemicals**

GA<sub>3</sub> 20% is compatible with most commonly used fungicide seed treatments (e.g. Vitavax® CT and Dithane®), standard dyes, and sticker/binding agents. When preparing tank mixes, ensure adequate physical compatibility and mixing characteristics. Refer to the Compatibility Test for Tank Mix Components under the GENERAL INFORMATION section of this label for additional information.

**HYBRID RICE SEED PRODUCTION**

(All States except California)

Apply GA<sub>3</sub> 20% during heading to increase panicle height of hybrid rice. This will facilitate pollination and harvest efficiency in the field, thus maximizing potential seed yield.

**Timing and Rate Instructions**

For hybrid rice, make 1 to 5 applications of 20 to 100 grams a.i./acre at regular intervals during the heading period to promote main culm and tiller panicle extension, thus helping to maximize flower pollination.

**Use Precautions**

Avoid drift or accidental application to other crops. Higher rates of GA<sub>3</sub> 20% application to hybrid rice plants can result in excessive vegetative growth, thus producing a taller plant that is more prone to lodging.

**Compatibility with Other Chemicals**

Most commonly used rice herbicides and fungicides are compatible with GA<sub>3</sub> 20%. When applying GA<sub>3</sub> 20% in mixtures with Arrosolo®, Riverside Propanil 60DF, Stam® M4, Stam® 80EDF, or Wham!® EZ, plus one of their suggested adjuvants, use of an additional surfactant is not necessary. Do not apply GA<sub>3</sub> 20% with products containing fenoxaprop-*p*-ethyl as the active ingredient. When preparing tank mixes, ensure adequate physical compatibility and mixing characteristics. Refer to the Compatibility Test for Tank Mix Components under the GENERAL INFORMATION section of this label for additional information.

**Mixing Instructions**

Fill the treatment tank with half of the final tank mix volume. Add the required amount of GA<sub>3</sub> 20% and mix thoroughly while adding water to the desired final volume. Dispose of any unused spray material at the end of the day.

**Application Equipment**

Make aerial applications of GA<sub>3</sub> 20% with spray systems capable of producing a uniform spray pattern of medium to fine spray droplets. Apply no less than 3 gallons per acre (GPA) of total spray volume. Use low pressure ground sprayers equipped with boom and flat fan nozzles using 10 to 15 GPA spray volume.

**CONVERSION TABLES**

<b>Grams of Gibberellic Acid Per Acre</b> Desired Gibberellic Acid Concentration (Grams Active Ingredient per acre) in Finished Spray	<b>To</b>	<b>Amount of GA<sub>3</sub> 20% Formulation Per Acre</b> GA <sub>3</sub> 20% Contains Approximately 1.0 Gram Active Ingredient in 5 Grams of Formulated Product
0.5		2.5 grams
1.0		5 grams
2.0		10 grams
4.0		20 grams
5.0		25 grams
8.0		40 grams
10.0		50 grams
12.0		60 grams
16.0		80 grams
20.0		100 grams
24.0		120 grams
32.0		160 grams
40.0		200 grams
48.0		240 grams
50.0		250 grams

<b>ppm (parts per million) GA<sub>3</sub></b>	<b>Grams of GA<sub>3</sub> 20% per gallon of spray solution</b>
1	0.02
5	0.09
10	0.19
25	0.47
50	0.94
100	1.9
250	4.7
500	9.4
750	14.2
1,000	18.9

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Keep containers tightly closed when not in use.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL: Non-refillable container.** Do not reuse or refill this container. Triple rinse (or equivalent) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ~ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank and store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.  
**For chemical spill, leak, fire or exposure, call CHEMTREC at 1-800-424-9300.**

#### **LIMITED WARRANTY AND DISCLAIMER**

NOTICE: CH Biotech R&D Co., Ltd. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use. Buyer assumes all risks of use and handling which is a variance in any way with the directions herein. To the extent consistent with applicable law, CH Biotech R&D Co., Ltd. makes no other express or implied warranty of fitness or merchantability. To the extent consistent with applicable law, in no case shall CH Biotech R&D Co., Ltd. or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. CH Biotech R&D Co., Ltd. and Seller offer this product and the Buyer and user accept it, subject to the foregoing Limited Warranty and Disclaimer which may be varied only by agreement in writing signed by a duly authorized representative of CH Biotech R&D Co., Ltd.

Dithane® and Stam® are registered trademarks of Dow AgroSciences LLC.

Wham® is a registered trademark of RiceCo.

Arrosolo® is a registered trademark of Syngenta Crop Protection, Inc.

Vitavax® is a registered trademark of Uniroyal Chemical Company, Inc.

**Produced For  
CH BIOTECH R&D CO., LTD.  
By CH BIOTECH LLC  
601 KETTERING DRIVE  
ONTARIO, CA 91761**