



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

December 9, 2021

Premjit Halarnkar, Ph.D., MPA
Global Head of Regulatory Affairs
CH Biotech R&D Co., Ltd.
601 Kettering Drive
Ontario, CA 91761

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment – Modify application directions; Add “foliar” to description; Add precautionary statement
Product Name: MegaGro 2.0
EPA Registration Number: 90866-34
Application Date: 08/30/2021
OPP Case Number: 00319349

Dear Mr. Halarnkar:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the U.S. Environmental Protection Agency (EPA). If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false


Page 2 of 2
EPA Reg. No. 90866-34
OPP Decision No. 00319349

or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.


If you have any questions, please contact Susannah Powell via email at powell.susannah@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read "James Parker", with a long, sweeping underline that extends to the left.

James Parker, Team Leader
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)
Office of Pesticide Programs

Enclosure



MEGAGRO™ 2.0

A foliar plant growth regulator for stimulation of early and improved root and shoot development, increased vegetative growth, and growth promotion of flowers and fruit of listed crops

ACTIVE INGREDIENTS:	BY WT
Choline Chloride	2.50%
3-Indolebutyric acid (IBA).....	0.80%
Gamma Aminobutyric Acid.....	0.50%
Cytokinin, as Kinetin	0.18%
OTHER INGREDIENTS.....	96.02%
TOTAL	100.00%

Not for use in furrow with liquid fertilizers

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For non-emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 (NPIC Web site: www.npic.orst.edu).</p> <p>For general product information, call CH Biotech LLC at (909) 472-3033 between the hours of 9 a.m. – 4 p.m. Pacific Time.</p>	

EPA REG. NO. 90866-34

EPA EST. NO. 90866-CA-001

NET CONTENTS:

BATCH NUMBER:

Produced For
 CH Biotech R&D., Ltd.
 By CH Biotech LLC
 601 Kettering Drive
 Ontario, CA 91761
 Tel: 909-472-3033
 Email: info@chbio.com

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long sleeved shirt and long pants,
- Chemical resistant gloves in Category A, such as butyl rubber > 14 mils, or natural rubber > 14 mils, or neoprene rubber > 14 mils or nitrile rubber > 14 mils, and
- Shoes plus socks

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during applications. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls,
- Chemical resistant gloves in Category A, such as butyl rubber > 14 mils, or natural rubber >14 mils, or neoprene rubber > 14 mils or nitrile rubber > 14 mils, and
- Shoes plus socks

PRODUCT INFORMATION

MEGAGRO™ 2.0 contains four active ingredients classified as plant growth regulators (PGRs): 3-Indolebutyric acid (IBA), Cytokinin (as Kinetin), Choline Chloride, and Gamma Aminobutyric Acid (GABA). Benefits derived from the use of this product include: stimulation of early and improved root and shoot development, increased vegetative growth, and growth development of flowers and fruit. Read the use instructions for specific details by crop.

A surfactant can be included in the tank mixture if desired based on field experience or further instructions from your local extension service, crop consultant or field representative or if indicated by a tank mix partner.

USE DIRECTIONS FOR CHEMIGATION

Apply MegaGro 2.0 through fixed or standing irrigation systems or through foliar applications. Foliar applications are preferred.

Apply this product only through the following types of irrigation systems:

1. Sprinkler including big gun, solid set or hand move irrigation systems.
2. Calibrated overhead watering booms.

Before applying this product through any type of irrigation system, perform a small-scale trial to determine if product performance and phytotoxicity results are acceptable.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have any questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems), used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively

designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Agitate the pesticide supply tank throughout the application of MegaGro 2.0. Apply MegaGro 2.0 at the rate of 2.0 - 4.0 fluid ounces per acre at the end of the irrigation period in a sufficient amount of water to allow proper coverage of plant or crop. Fill the supply tank one-half full with water, add the appropriate amount of MegaGro 2.0 to the tank and finish filling the tank with water. Application of MegaGro 2.0, through sprinkler irrigation systems, on turf grasses is not recommended.

GROUND AND AERIAL APPLICATION

Minimum Spray Volume (Gal/A)

CROP/CROP GROUP	GROUND		AIR
	Concentrate	Dilute	
Field Crops, Miscellaneous	10.0	15.0	3.0
Berry and Small Fruits, Vegetables, Vines	25.0	100.0	5.0
Pome Fruits, Stone Fruits, Tree Crops and Tree Nuts	50.0	200.0	10.0
Citrus	50.0	300.0	—

Special considerations: MegaGro 2.0 compatibility with other agricultural products has not been fully investigated. Compatibility of MegaGro 2.0 with other products requires testing for crop safety and performance prior to large-scale use. Repeat application may be necessary if it rains within 2 hours after application.

Depending upon the equipment used, and specific crop, spray volume applied per acre will differ. Apply sufficient water volume to ensure thorough coverage.

APPLICATION INSTRUCTIONS		
Crop/Crop Group	Amount of MegaGro 2.0	Application Directions and Timing
Asparagus	13.0 fl oz/100 gallons water	<ul style="list-style-type: none"> Apply at green up to sustain inter-harvest Repeat applications every 14 to 21 days as needed Apply after harvest when asparagus is in fern stage to promote overwinter health
Berry and Small Fruit Group. [Crops such as Blackberry, Blueberry, Caneberry, Kiwi and Raspberry (except Grape and Strawberry)]	13.0 fl oz/100 gallons water	<ul style="list-style-type: none"> 1st: After leaves are fully emerged through bloom Repeat applications every 14 to 21 days as needed through harvest Application 2 to 3 weeks before frost to promote overwinter health
Brassica Vegetables Group. [Crops such as Broccoli, Cabbage, Cauliflower and Mustard greens]	13.0 fl oz/100 gallons water	<ul style="list-style-type: none"> Foliar application: Apply to achieve full coverage 1st: At 2 to 4 true leaf stage Repeat applications every 14 to 21 days as needed Use a non-ionic surfactant for hard to wet crops such as cabbage
Bulb Vegetables Group. [Crops such as Garlic, Leek, and Onion]	13.0 fl oz/100 gallons water	<ul style="list-style-type: none"> 1st: At 2 to 4 leaf stage Repeat application every 14 to 21 days as needed up to 10 days prior to harvest Thorough coverage and leaf wetting are required

APPLICATION INSTRUCTIONS

Crop/Crop Group	Amount of MegaGro 2.0	Application Directions and Timing
Cereal Grains Group. [Crops such as Barley, Corn ⁽¹⁾⁽²⁾ (field, pop, sweet), Millet, Oats, Rice, Sorghum and Wheat]	2.0 to 4.0* fl oz/A	<ul style="list-style-type: none"> • 1st: At 2 to 4 true leaf stage (fall or spring) • 2nd: After tillering through grain set (cereals), up to full tassel (corn)
Citrus Fruit Group. [Crops such as Grapefruit, Lemon, Lime, Sweet Orange and Tangerine]	13.0 fl oz/100 gallons water	<ul style="list-style-type: none"> • 1st application at 1st bloom • Repeat applications at each flush of new growth • Full coverage is necessary
Coffee	13.0 fl oz/100 gallons water	<ul style="list-style-type: none"> • Apply to newly transplanted coffee to assist rooting • 1st: Annual application prior to bloom • 2nd: 14 to 21 days after first application • 3rd: 45 days prior to harvest • 4th: 30 days prior to harvest
Cotton ⁽¹⁾⁽²⁾	2.0 to 4.0* fl oz/A	<ul style="list-style-type: none"> • 1st: At 2 to 4 true leaf stage • Repeat applications at 14 to 21-day interval up to boll development
Cucurbit Vegetables Group. [Crops such as Cantaloupe, Cucumber, Honeydew, Muskmelon, Squash (summer and winter) and Watermelon]	13.0 fl oz/100 gallons water	<ul style="list-style-type: none"> • 1st: At beginning vine run • Repeat applications every 14 to 21 days
Forage, Fodder and Straw of Cereal Grains	2.0 to 4.0* fl oz/A	<ul style="list-style-type: none"> • 1st: At 2 to 4 true leaf stage • 2nd: 14 to 21 days after first application • Applications between cuttings will improve root structure and increase stand vigor
Fruiting Vegetables Group. [Crops such as Eggplant, Pepper and Tomato]	13.0 fl oz/100 gallons water	<ul style="list-style-type: none"> • 1st: At 2 leaves to first bloom • Repeat applications every 14 to 21 days until harvest
Grass Forage, Fodder and Hay	2.0 to 4.0* fl oz/A	<ul style="list-style-type: none"> • 1st: At 2 to 4 true leaf stage • 2nd: 14 to 21 days after first application • Applications between cuttings will improve root structure and increase stand vigor
Grass Grown for Seed such as Perennial Ryegrass, Tall Fescue or Bentgrass	2.0 to 4.0* fl oz/A	<ul style="list-style-type: none"> • Apply when growth resumes in the spring • Applications every 14 to 21 days until harvest
Grape	13.0 fl oz/100 gallons water	<ul style="list-style-type: none"> • 1st: Apply when grapes are 2 to 3 mm in size • 2nd: 14 to 21 days after first application • 3rd: 45 days prior to harvest • 4th: 30 days prior to harvest
Herbs and Spices Group. [Crops such as Basil, Dill, Mustard and Sage]	2.0 to 4.0* fl oz/A or 13.0 fl oz/100 gallons water	<ul style="list-style-type: none"> • 1st: At 2 to 4 true leaf stage • Repeat applications every 14 to 21 days as needed until harvest
Hops	13.0 fl oz/100 gallons water	<ul style="list-style-type: none"> • 1st: At 1 to 2 pairs of leaves on main vines • Repeat applications every 14 to 21 days as needed

APPLICATION INSTRUCTIONS

Crop/Crop Group	Amount of MegaGro 2.0	Application Directions and Timing
Leafy Vegetables Group. [Crops such as Celery, Head and Leaf Lettuce, Kale and Spinach]	13.0 fl oz/100 gallons water	<ul style="list-style-type: none"> • 1st: At 2 to 4 true leaf stage • Repeat applications every 14 to 21 days as needed
Legume Vegetables Group. (Succulent or Dried) [Crops such as Bean (all types), Peas and Soybeans ⁽¹⁾⁽²⁾]	2.0 to 4.0* fl oz/A	<ul style="list-style-type: none"> • 1st: At 2 to 4 true leaf stage • Repeat applications every 14 to 21 days as needed through R3
Mint, Peppermint, Spearmint	2.0 to 4.0* fl oz/A	<ul style="list-style-type: none"> • 1st: At 2 to 4 true leaf stage • 2nd: Sequential application at 5 to 10 days after each harvest
Nongrass Animal Feeds such as Alfalfa, Clover, Hay and Vetch	2.0 to 4.0* fl oz/A	<ul style="list-style-type: none"> • Seedling alfalfa, clover, hay, and vetch: Apply at 2 to 4-trifoliate stage • For established crop, apply at green-up and 5 to 10 days after each cutting
Oil Seed Crops Group. [Crops such as Canola ⁽¹⁾⁽²⁾ , Flax and Sunflower]	2.0 to 4.0* fl oz/A	<ul style="list-style-type: none"> • 1st: At 2 to 4 leaf stage • Repeat applications can be made every 10 to 14 days as needed
Peanut	2.0 to 4.0* fl oz/A	<ul style="list-style-type: none"> • 1st: At 2 to 4 true leaf stage • Repeat applications every 14 to 21 days as needed through seed fill
Pome Fruits Group. [Crops such as Apple and Pear]	13.0 fl oz/100 gallons water	<ul style="list-style-type: none"> • 1st: From green tip to tight cluster • 2nd: Repeat applications as needed (12 mm to 30 mm fruit size)
Root and Tuber Vegetables Group. [Crops such as Carrot, Ginseng, Horseradish, Parsley (turnip-rooted), Potato, Radish, Sugar Beet, Sweet Potato and Turnip]	2.0 to 4.0* fl oz/A or 13.0 fl oz/100 gallons water	<ul style="list-style-type: none"> • 1st: At 2 to 4 true leaf stage • Repeat applications every 14 to 21 days as needed until harvest • Foliar application: thorough spray coverage is necessary
Stone Fruits Group. [Crops such as Apricot, Cherry, Peach and Plumcot]	13.0* fl oz/100 gallons water	<ul style="list-style-type: none"> • 1st: At 5 to 10 mm fruit • Repeat applications every 14 to 21 days as needed
Strawberry	13.0 fl oz/100 gallons water	<ul style="list-style-type: none"> • 1st: Spray immediately after transplant • Repeat applications every 14 to 21 days as needed
Sugarcane	2.0 to 4.0* fl oz/A	<ul style="list-style-type: none"> • 1st: At 2 to 4 true leaf stage • Repeat applications every 14 to 21 days as needed
Tobacco	2.0 to 4.0* fl oz/A	<ul style="list-style-type: none"> • 1st: At 2 to 4 true leaf stage • Repeat applications every 14 to 21 days as needed
Tree Nuts Group. [Crops such as Almonds, Cashews and Pecans]	13.0 fl oz/100 gallons water	<ul style="list-style-type: none"> • 1st: After leaves are fully emerged through bloom • Repeat applications every 14 to 21 days as needed

*If application spray volume is greater than 15.0 gallons per acre, use the dilution rate of 13.0 fluid ounces per 100 gallons water.

⁽¹⁾ This product can be tank mixed with glyphosate products registered for use on Roundup Ready® crops.

⁽²⁾ This product can be tank mixed with products registered for use on LibertyLink® crops.

OTHER CROPS

Crop/Crop Group	Amount of MegaGro 2.0	Application Directions and Timing
Hemp	13.0 fl oz/100 gallons water	<ul style="list-style-type: none"> 1st: At 1 to 2 pairs of leaves on main stem Repeat applications every 14 to 21 days as needed up to bud formation

GREENHOUSE AND NURSERY

Greenhouse Transplant - For Early Root Growth

Crop Group	Use Rate Oz/A
Leafy Vegetables Group. [Crops such as Celery, Head and Leaf Lettuce, Kale and Spinach]	3.0 to 4.0*
Fruiting Vegetables Group. [Crops such as Eggplant, Pepper and Tomato]	3.0 to 4.0*

Greenhouse Transplant – For Early Shoot Growth

Crop Group	Use Rate Oz/A
Leafy Vegetables Group. [Crops such as Celery, Head and Leaf Lettuce, Kale and Spinach]	3.0 to 4.0*
Fruiting Vegetables Group. [Crops such as Eggplant, Pepper and Tomato]	3.0 to 4.0*

*If application spray volume is greater than 15.0 gallons per acre, use the dilution rate of 13.0 fluid ounces per 100 gallons water.

For Greenhouse Establishment or Production

Crop Group	Amount of MegaGro 2.0	Application Directions and Timing
Leafy Vegetables Group. [Crops such as Celery, Head and Leaf Lettuce, Kale and Spinach]	13.0 oz/100 gallons in 30 to 80 gallons per acre	<ul style="list-style-type: none"> 1st: 2 to 4 true leaves Repeat applications every 14 to 21 days as needed
Fruiting Vegetables Group. [Crops such as Eggplant, Pepper and Tomato]	13.0 oz/100 gallons in 30 to 80 gallons per acre	<ul style="list-style-type: none"> 1st: 2 to 4 true leaves Repeat applications every 14 to 21 days as needed

TURFGRASS

For Sod Grass: Apply MegaGro 2.0 by ground using 20.0 to 40.0 gallons of water per acre. Use 2.5 fluid ounces to 6.5 fluid ounces product in 20.0 gallons to 40.0 gallons of water, respectively, at a 1:1000 dilution rate.

For Turfgrass: Apply MegaGro 2.0 by ground according to the table below using 1.0 to 10.0 gallons of water per 1000 square feet. Use MegaGro 2.0 for turf growth suppression at the dilution rate of 1:300 (4.2 fluid ounces product per 10.0 gallons water).

Turf	Amount (MegaGro 2.0 /Gal Water/1000 Sq Ft*)	Application Directions and Timing
Warm climate grasses including Bermuda, Bermuda hybrids,	0.13 to 0.65 fl oz/1.0 to 5.0 gal of water/1000 sq ft	<ul style="list-style-type: none"> Make applications at 2 to 3-week intervals during the growing season

Turf	Amount (MegaGro 2.0 /Gal Water/1000 Sq Ft*)	Application Directions and Timing
Centipede, St. Augustine and similar warm season grasses		
Dichondra	0.65 to 1.3 fl oz/5.0 to 10.0 gal of water/1000 sq ft	• Make applications at 2 to 3-week intervals during the growing season
Cool climate grasses including Bluegrass, Fescue, Rye and similar cool season grasses	0.13 to 0.65 fl oz/1.0 to 5.0 gal of water/1000 sq ft	• Make applications at 2 to 3-week intervals during the growing season

*Apply 0.13 fluid ounce per gallon.

GREENHOUSE AND NURSERY

Differences in responsiveness may vary from one cultivar to another or from one set of growing conditions to another. Unless previous experience dictates otherwise, prior to widespread use, test a small number of plants from each cultivar to verify desired efficacy.

Foliage Plants

Aglaonema	Aphelandra	Dieffenbachia	Fittonia	Maranta	Philodendron	Schefflera	Syngonium
Ajuga	Caladium	Dracaena	Gynura	Palms	Pilea	Schlumbergera	Tradescantia
Anthurium	Cissus	Ficus	Hoya	Peperomia	Pothos	Spathiphyllum	Similar foliage plants

Application Rates and Timing: Dilute 0.85 fluid ounce of MegaGro 2.0 in 10.0 gallons of water (1:1500 dilution rate) for plants less than 2 years old. Dilute 1.3 fluid ounces MegaGro 2.0 in 10.0 gallons of water (1:1000 dilution rate) for mature plants. Repeat applications at 10 to 14-day intervals when required. Apply the last spray 1 to 2 weeks prior to sale. Uniform and thorough spray coverage is necessary for best results.

Bedding and Flowering Plants

Abutilon	Carnation	Coral Bells	Foxglove	Gladiolus	Lily	Osmachus	Salvia	Vinca
Aglais	Champaca	Cyclamen	Fuchsia	Gloxinia	Lupine	Petunia	Scabiosa	Zinnia
Alyssum	Chrysanthemum	Dahlia	Gardenia	Impatiens	Marigold	Poinsettia	Sedum	Similar plants
Calceolaria	Cineraria	Delphinium	Gazania	Iris	Michelia	Portulaca	Sempervivum	
Canna	Columbine	Dianthus	Geranium	Jasminum	Monarda	Roses	Tulips	

Application Rates and Timing: Dilute 0.85 fluid ounce of MegaGro 2.0 in 10.0 gallons of water (1:1500 dilution rate) for plants less than 2 years old. Dilute 1.3 fluid ounces MegaGro 2.0 in 10.0 gallons of water (1:1000 dilution rate) for mature plants. Repeat applications at 10 to 14-day intervals when required. Apply the last spray 1 to 2 weeks prior to sale. Uniform and thorough spray coverage is necessary for best results.

Woody Ornamentals

Arborvitae	Azalea	Carissa	English ivy	Juniper	Pine	Rhododendron	Similar plants
Aucuba	Boxwood	Chinese magnolia	Holly	Maple	Podocarpus	Viburnum	

Application Rates and Timing: Dilute 0.85 fluid ounce of MegaGro 2.0 in 10.0 gallons of water (1:1500 dilution rate) for plants less than 2 years old. Dilute 1.3 fluid ounces MegaGro 2.0 in 10.0 gallons of water (1:1000 dilution rate) for mature plants. Repeat applications at 10 to 14-day intervals when required. Apply the last spray 1 to 2 weeks prior to sale. Uniform and thorough spray coverage is necessary for best results.

Garden Grown Tree Fruits

Apple	Cherry	Grape	Lemon	Mango	Persimmon	Starfruit
Asian pear	Fig	Jujubee	Litchi	Orange	Plum	Similar plants
Apricot	Guava	Kumquat	Longara	Peach	Prunes	

Application Rates and Timing: Dilute 0.85 fluid ounce of MegaGro 2.0 in 10.0 gallons of water (1:1500 dilution rate) for plants less than 2 years old. Dilute 1.3 fluid ounces MegaGro 2.0 in 10.0 gallons of water (1:1000 dilution rate) for

mature plants. Repeat applications at 10 to 14-day intervals when required. Apply the last spray 1 to 2 weeks prior to sale. Uniform and thorough spray coverage is necessary for best results.

PLANT CUTTINGS

- To propagate new plants from cuttings.
- Treated cuttings can be expected to produce uniform roots resulting in beautiful, symmetrical plants.
- For use on most home, tropical, greenhouse and nursery plants.

Type of Cutting	Dilution rate
Soft wood	1:20 dilution rate (0.5 fluid ounce product in 10.0 fluid ounces of water)
Medium wood	1:10 dilution rate (1.0 fluid ounce product in 10.0 fluid ounces of water)
Hard wood	1:5 dilution rate (2.0 fluid ounces product in 10.0 fluid ounces water)

For Rooting House Foliage, Tropical and Hardy Ornamentals, Leaf, Greenwood and Softwood Cuttings, Woody ornamentals, Deciduous hardwoods, Evergreens, Ground Covers, and Perennials including:

Acanthopanax	Catalpa	Dogwood	Heath	Manzanita	Rhododendron
African violet	Chamaecyparis (False Cypress)	Douglas Fir	Heather	Maple	Rose
Apple (ornamental)	Chaste tree	Escallonia	Hemlock	Matrimony vine	Russian Olive
Arborvitae, American	Chestnut	Euonymus	Hibiscus	Minor	Snowball
Arbutus	Chokeberry	Flowering Crab apple	Holly	Myrtle	Sourwood
Azalea (evergreen & semi-evergreen)	Cotoneaster	Flowering Quince	Honeysuckle	Oak	Spirea
Aster	Crape-myrtle	Forsythia	Ivy	Olive (ornamental)	Tulip Tree
Barberry	Clematis	Fuchsia	Japanese quince	Orange, sour (ornamental)	Umbrella Pine
Begonia	Cryptomeria	Gardenia	Jasmine	Pachysandra	Viburnum
Birch	Chrysanthemum	Geranium	Juniper	Pecan (ornamental)	Vinca
Bittersweet	Cypress	Germander	Lilac	Photinia	Yew
Boxwood	Dahlia	Grape (ornamental)	Locust	Pivet	Wriggle
Camellia	Delphinium	Hawthorne	Magnolia	Pyracantha (Firethorn)	Many others

USE INSTRUCTIONS: Obtain cuttings from vigorous, healthy plants. Keep cuttings moist and cool such as in an ice chest. With a sharp knife, trim the cutting (2 to 8 inches long) with a diagonal cut just below a node or leaf. Dip the basal end of cutting, individually or in bunches, into the MegaGro 2.0 solution for 3 to 5 seconds.

Note: Following dipping, place cuttings into planting medium. Depending on the species, rooting will take place in several weeks or months under a moist greenhouse environment. Transplant once the cuttings have rooted.

Shrubs, Flowers, Groundcovers and Houseplants including, Rose, Arborvitae, Gardenias, Flowering Trees and other ornamentals

USE INSTRUCTIONS: In bare root transplant or from containers: Use 2.0 tablespoons of MegaGro 2.0 per 10.0 gallons of water. Apply solution to root area in transplanting hole and then cover roots with soil. After planting, repeat applications biweekly as a drench to thoroughly wet the root area using a solution consisting of 1.0 tablespoon of MegaGro 2.0 per 10.0 gallons of water.

Annual and Perennial Flowers (bedding plants)

USE INSTRUCTIONS: Use 1.0 tablespoon of MegaGro 2.0 per 10.0 gallons of water and apply to thoroughly saturate roots at time of planting. Repeat at weekly intervals until plants are well established.

Groundcovers including, Ivy, Iceplant, Geranium, Cotoneaster, Barberry, and Ajuga

USE INSTRUCTIONS: Use 1.0 tablespoon of MegaGro 2.0 per 10.0 gallons of water and apply thoroughly to saturate the root area at time of planting. Repeat at weekly intervals until plants are well established.

Houseplants (repotting and planting)

USE INSTRUCTIONS: Use 1.0 tablespoon of MegaGro 2.0 per 10.0 gallons of water and water thoroughly at weekly intervals to saturate the root zone until plants are well established.

Established Plants

USE INSTRUCTIONS: To continue new root growth, use 1.0 tablespoon of MegaGro 2.0 per 10.0 gallons of water and water plants with solution once a month.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store product in the original container. Store product in a cool, dry locked place out of the reach of children and out of direct sunlight.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. Triple rinse (or equivalent). Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{3}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **For chemical spill, leak, fire or exposure, call CHEMTREC at 1-800-424-9300.**

LIMITED WARRANTY AND DISCLAIMER

NOTICE: CH Biotech R&D Co., Ltd. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use. Buyer assumes all risks of use and handling which is a variance in any way with the directions herein. To the extent consistent with applicable law, CH Biotech R&D Co., Ltd. makes no other express or implied warranty of fitness or merchantability. To the extent consistent with applicable law, in no case shall CH Biotech R&D Co., Ltd. or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. CH Biotech R&D Co., Ltd. and Seller offer this product and the Buyer and user accept it, subject to the foregoing Limited Warranty and Disclaimer which may be varied only by agreement in writing signed by a duly authorized representative of CH Biotech R&D Co., Ltd.

Roundup Ready® is a registered trademark of the Monsanto Company.

LibertyLink® is a registered trademark of Bayer CropScience.