



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Biopesticides and Pollution Prevention Division (7511P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

90866-37

Date of Issuance:

10/17/2022

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Strong Finish

Name and Address of Registrant (include ZIP Code):

CH Biotech R & D Co., Ltd.
 601 Kettering Drive
 Ontario, CA 91761

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA Registration Number.

On the basis of information furnished by the registrant, the above-named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA or the Act).

Registration is in no way to be construed as an endorsement or recommendation of this product by the U.S. Environmental Protection Agency (EPA). In order to protect health and the environment, the Administrator, on his or her motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under the Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration or registration review of your product when EPA requires all registrants of similar products to submit such data.

Signature of Approving Official:

Andrew Bryceland, Team Leader
 Biochemical Pesticides Branch
 Biopesticides and Pollution Prevention Division (7511M)
 Office of Pesticide Programs

Date:

10/17/2022

2. Make the following labeling change before you release this product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 90866-37.”
3. Submit one (1) copy of the final printed labeling for the record before you release this product for shipment.

Should you wish to add/retain a reference to your company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to EPA’s Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6. A stamped copy of the labeling is enclosed for your records. Please also note that the record for this product currently contains the following acceptable Confidential Statement of Formula (CSF):

- Basic CSF dated 2/25/2022

If you have any questions, please contact Cheryl Greene via email at greene.cheryl@epa.gov.

Sincerely,



Andrew Bryceland, Team Leader
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511M)
Office of Pesticide Programs

ACCEPTED

Oct 17, 2022

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 90866-37

Strong Finish

A Plant Growth Regulator for Listed Commodities

ACTIVE INGREDIENT:	By WT
Cytokinin, as Kinetin	0.5%
Choline Chloride	30.0%
OTHER INGREDIENTS:	<u>69.5%</u>
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

First Aid	
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow Do not induce vomiting unless told to do so by a poison control center or doctor Do not give anything by mouth to an unconscious person
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing Rinse skin immediately with plenty of water for 15 to 20 minutes Call a poison control center or doctor for treatment advice
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 to 20 minutes Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye Call a poison control center or doctor for treatment advice
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For non-emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 (NPIC Web site: www.npic.orst.edu).</p> <p>For general product information, call CH Biotech LLC at (909) 472-3033 between the hours of 9a.m. – 4 p.m. Pacific Time.</p>	

EPA Reg. No. 90866-XX

EPA Est. No. 90866-CA-1

Produced for CH Biotech R&D Co., Ltd.

By CH Biotech, LLC
601 Kettering Drive
Ontario, CA 91761
Tel: 909.472.3033

Net Content: 1.0 gallon
2.5 gallons

Email: info@chbio.com

Batch No.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

For Terrestrial Uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean highwater mark. Do not contaminate water when disposing of equipment wash water or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

This product is not compatible with strong oxidizers.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during applications. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and

restricted-entry interval. The requirements in this box only apply to uses that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves Category A, such as butyl rubber \geq 14 mils, or natural rubber \geq 14 mils, or neoprene rubber \geq 14 mils or nitrile rubber \geq 14 mils
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter without appropriate protective clothing until sprays have dried.

GENERAL INFORMATION

Strong Finish contains two active ingredients classified as plant growth regulators (PGRs): Cytokinin (as Kinetin) and Choline Chloride. Benefits derived from the use of this product include: improved photosynthetic efficiency, stimulation of root and shoot development, enhanced plant growth and development, and growth development of flowers and fruit.

Strong Finish is not a fertilizer; therefore, incorporate good fertilization program practices. Kinetin may delay senescence of the leaves on some crops. Make applications at the proper timing and when the crop is actively growing.

CHEMIGATION

Apply this product only through the following types of irrigation systems:

1. Sprinkler including side (wheel) roll, solid set or hand move irrigation systems.
2. Calibrated overhead watering booms

Do not apply this product through any other types of irrigation systems. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems), used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, discharge the water from the public water system into a reservoir tank prior to pesticide introduction. A complete physical break (air gap) must occur between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must contain a normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment. Agitate the pesticide supply tank throughout the application of Strong Finish. Except for turfgrass, apply Strong Finish at the end of the irrigation period in a sufficient amount of water to allow proper coverage of plant or crop. Fill the supply tank one-half full with water, add the appropriate amount of Strong Finish to the tank and finish filling the tank with water.

SPRINKLER CHEMIGATION

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Agitate the pesticide supply tank throughout the application of Strong Finish. Except for turfgrass, apply Strong Finish at the end of the irrigation period in a sufficient amount of water to allow proper coverage of plant or crop but not to exceed 8 fluid ounces of Strong Finish per acre per application. Fill the supply tank one-half full with water, add the appropriate amount of Strong Finish to the tank and finish filling the tank with water.

APPLICATION DIRECTIONS

For all crops, unless otherwise specified, tank mix Strong Finish by adding 13 fluid ounces per 100 gallons of water and spray crop canopy with sufficient volume to ensure uniform coverage. For more specific instructions and rates, see the following table of crops; the rate is expressed as fluid ounces of Strong Finish per acre in a corresponding volume of water per acre. Water volume is usually determined by the grower and/or the particular circumstances that affect uniform crop coverage; however, sufficient volume to wet leaf is required as opposed to just misting it. Use lower rates with the corresponding lower volumes and higher rates with higher volumes.

If sufficient rain to wash leaf occurs within 2 hours of application, re-apply. Compatibility has not been fully determined for all agrochemicals.

With carrier volumes of 5 gallons per acre or less, the required minimum dose is 2 oz./acre (see footnote ⁽³⁾ below).

SPRAY VOLUME (GAL/ACRE) RATES

CROP/CROP GROUP	GROUND		AIR	
	Adequate Coverage	Best Coverage	Adequate Coverage	Best Coverage
Field Crops	10.0	20.0	2.0	5.0
Berry and Small Fruits, Vegetables, Vines	25.0	100.0	5.0	10.0
Pome Fruits, Stone Fruits, Tree Crops and Tree Nuts	50.0	200.0	10.0	20.0

Crop/Crop Group	Application Timing	Strong Finish Rate/Acre	Water Volume/Acre**
Alfalfa, clover, and vetch (for seed)	Make one application, 5 to 8 days after full bloom	2.0 to 3.0 fl oz	15 to 25 gal
Alfalfa, clover, and vetch (not for seed)	Apply to regrowth after cuttings	2.0 to 3.0 fl oz	15 to 25 gal
Brassica Vegetables Group. [Crops such as Broccoli, Cauliflower, Cabbage, and Mustard greens]	1st application: At flowering stage	2.0 fl oz	15 gal
	2nd application: 10 to 14 days after first application	2.0 to 3.0 fl oz	15 to 25 gal
	3rd application: 7 to 10 days before harvest		
Citrus Fruits Group. [Crops such as Grapefruit, Lemon, Sweet orange, and Tangelos]	1st application: At small fruit stage when fruit size is approx. 6 to 8 mm	6.5 to 13.0 fl oz	50 to 100 gal
	2nd application: 40 days prior to harvest	6.5 to 13.0 fl oz	50 to 100 gal
Corn (Sweet, Field ⁽¹⁾⁽²⁾ , and Popcorn)	Make one application, V5 to VT (Full tassel)	2.0 to 3.0 fl oz	10 to 25 gal
Cotton including short staple, acala & pima varieties	1st application: Apply at the pin head square stage	2.0 to 3.0 fl oz	10 to 25 gal
	2nd application: Apply at first bloom	2.0 to 4.0 fl oz	10 to 30 gal
Cucurbit Vegetables Group. [Crops such as Cucumber, Cantaloupe, Honeydew, Muskmelon, Summer squash, and Watermelon]	Make 1st application at early fruiting stage	2.0 to 3.0 fl oz	15 to 25 gal
	Make 2 or more applications at 10 to 14-day intervals after early fruiting	3.0 to 5.0 fl oz	20 to 40 gal
Fruiting Vegetables Group. [Crops such as Eggplant, Pepper, and Tomato]	Make 1st application at early fruiting stage	2.0 to 3.0 fl oz	15 to 25 gal
	Make 2 or more applications at 10 to 14-day intervals after early fruiting	3.0 to 12.8 fl oz	25 to 100 gal
Grapes including table, wine, and raisin varieties	1st application: 10 to 14 days before bud break	3.0 to 5.0 fl oz	25 to 40 gal
	2nd application: Small berry stage (3 to 5 mm in size)	5.0 to 9.0 fl oz	40 to 60 gal
	3rd application: 40 days prior to		

	harvest		
Herbs Group. [Crops such as Basil, Dill, Mustard, and Sage]	1st application: During mid-season growth	3.0 to 5.0 fl oz	25 to 40 gal
	2nd application: At budding stage		
Hops	1st application: At inflorescence emergence	3.0 to 9.0 fl oz	25 to 75 gal
Leafy Vegetables Group. [Crops such as Celery, Head lettuce, Leaf lettuce, and Spinach]	1st application: During mid-season growth	3.0 to 5.0 fl oz	25 to 40 gal
	2nd application: 10 to 14 days prior to harvest		
Legume Vegetables Group. [Crops such as Dry beans, Green beans, Lentils, Peas]	1st application: Apply at the 3 to 5 trifoliate leaf stage	2.0 to 3.0 fl oz	15 to 25 gal
	2nd application: Apply at 5 to 10% bloom		
Oil Seed Crops including Canola ⁽¹⁾⁽²⁾ , Flax, and Sunflower	1st application: First bloom	2.0 fl oz	10 to 15 gal
	2nd application: 14 to 21 days after 1st application		
Peanuts including all commercial varieties	1st application: Apply at the 3 to 5 leaf stage	2.0 to 3.0 fl oz	15 to 25 gal
	2nd application: Apply at initial pegging		
	3rd application: Apply during pod fill	3.0 to 5.0 fl oz	25 to 40 gal
Pome/Stone Fruits Group. [Crops such as Apple, Apricot, Cherry, Plum, Plumcot, and Peach]	1st application: At small fruit stage when fruit size is approximately 3 to 5 mm	6.5 to 13.0 fl oz	50 to 100 gal
	2nd application: 40 days prior to harvest		
Root Vegetables Group. [Crops such as Carrot, Ginseng, Horseradish, Parsley (turnip-rooted), Radish, Sugar beet, and Turnip]	1st application: 40 to 60 days after plant emergence	2.0 fl oz	15 gal
	2nd application: 21 days after first application	2.0 to 3.0 fl oz	15 to 25 gal
	3rd application: Apply 10 to 14 days prior to anticipated harvest		
Small Fruits Group. [Crops such as Blackberry, Blueberry, Raspberry, and Strawberry]	1st application: At early flowering stage	2.0 to 5.0 fl oz	15 to 40 gal
	2nd application: 10 to 14 days after first application		
	3rd application: Repeat applications every 10 to 14 days		
Small Grains Group. [Crops such as Barley, Rice, Sorghum, Rye, and Wheat]	Make one application: Boot stage to anthesis. Wheat Feekes stage 10 to 10.5	2.0 to 3.0 fl oz	10 to 25 gal
Soybeans ⁽¹⁾⁽²⁾	1st application: First bloom	2.0 fl oz	15 gal
	2nd application: 14 to 21 days after 1st application	2.0 to 3.0 fl oz	10 to 25 gal
Tree Nuts Group. [Crops such as Almond, Pecan,	1st application: 2 weeks prior to bloom	6.0 to 12.0 fl oz	50 to 100 gal

and Pistachio]	2nd application: 2 weeks following petal fall		
	3rd application: 30 days after 2nd application		
Tuber Vegetables Group. [Crops such as Potato, Sweet Potato, and Yam]	1st application: At tuber initiation stage	2.0 to 3.0 fl oz	15 to 25 gal
	2nd application: 14 to 21 days after first application		
	3rd application: At early bloom stage	3.0 to 5.0 fl oz	25 to 40 gal

**Note: If using any other volume of water, use the tank mix dilution rate of 13 fluid ounces of Strong Finish per 100 gallons of water and use sufficient water volume to obtain uniform coverage.

(1) This product can be tank mixed with glyphosate products registered for use on Roundup Ready® crops.

(2) This product can be tank mixed with products registered for use on LibertyLink® crops.

(3) Carrier volumes with 10 gallons per acre or less require the use of 2 oz/acre minimum.

TURFGRASS

For Sod Grass: Apply Strong Finish by ground using 20 to 40 gallons of water per acre. Apply 2.5 to 5.0 fluid ounces product in 20 to 40 gallons of water, respectively, at a 1:1000 dilution rate.

For Turfgrass: Apply Strong Finish by ground according to the table below using 1 to 10 gallons of water per 1000 square feet.

Turf	Application & Timing	Strong Finish Rate/1000 ft. ²	Water Volume/1000 ft. ²
Cool Climate grasses such as Bluegrass, Fescue, Rye, and similar grasses	1st application: When seeded grass becomes established or at the beginning of the season for perennials	0.13 to 0.65 fl oz	1 to 5 gal
	Repeat as necessary		
Dichondra	1st application: When turf greens up in the spring	0.65 to 1.30 fl oz	5 to 10 gal
	Repeat at 14 to 21-day intervals during the growing season		
Warm Climate grasses such as St. Augustine, Bermuda, Bermuda hybrids, Centipede, and similar grasses	1st application: When turf greens up in the spring	0.13 to 0.65 fl oz	1 to 5 gal
	Repeat at 14 to 21-day intervals during the growing season		

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Protect from freezing. Store out of direct sunlight. Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at www.acrecycle.org. If not recycled, then puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

For packages up to 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For packages greater than 56 gallons: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For refillable containers: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For help with any spill, leak, fire or exposure involving this material, call day or night
CHEMTREC- 1-800-424-9300.

LIMITED WARRANTY AND DISCLAIMER

NOTICE: CH Biotech R&D Co., Ltd. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use. Buyer assumes all risks of use and handling which is a variance in any way with the directions herein. To the extent consistent with applicable law, CH Biotech R&D Co., Ltd. makes no other express or implied warranty of fitness or merchantability. To the extent consistent with applicable law, in no case shall CH Biotech R&D Co., Ltd. or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. CH Biotech R&D Co., Ltd. and Seller offer this product and the Buyer and user accept it, subject to the foregoing Limited Warranty and Disclaimer which may be varied only by agreement in writing signed by a duly authorized representative of CH Biotech R&D Co., Ltd.

Roundup Ready® is a registered trademark of the Monsanto Company.
LibertyLink® is a registered trademark of Bayer CropScience.