



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Biopesticides and Pollution Prevention Division (7511P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

90930-8

Date of Issuance:

4/19/2019

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Miller 2361

Name and Address of Registrant (include ZIP Code):

Miller Chemical & Fertilizer, LLC
P.O. Box 333
120 Radio Road
Hanover, PA 17331

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA Registration Number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA or the Act).

Registration is in no way to be construed as an endorsement or recommendation of this product by the U.S. Environmental Protection Agency (EPA). In order to protect health and the environment, the Administrator, on his or her motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under the Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration or registration review of your product when the EPA requires all registrants of similar products to submit such data.
2. Submit storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) data as these data requirements are not satisfied. A one-year study is required to satisfy these data requirements. You have 18 months from the date of this registration to provide these data to the EPA.

Signature of Approving Official:

Gina M. Burnett, Senior Regulatory Advisor
Biochemical Pesticides Branch
Biopesticides and Pollution Prevention Division (7511P)
Office of Pesticide Programs

Date:

4/19/2019

3. Make the following labeling change before you release this product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 90930-8.”
4. Submit one (1) copy of the final printed labeling for the record before you release this product for shipment.

Should you wish to add/retain a reference to your company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6. A stamped copy of the labeling is enclosed for your records. Please also note that the record for this product currently contains the following acceptable Confidential Statement of Formula (CSF):

- Basic CSF dated April 18, 2019

Any CSFs other than the one listed above are superseded.



If you have any questions, please contact Cheryl Greene by phone at (703) 308-0352 or via email at greene.cheryl@epa.gov.

Sincerely,



Gina M Burnett, Senior Regulatory Advisor
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)
Office of Pesticide Programs

Enclosure: Stamped Label



Miller 2361*

A Plant Hormone Supplement

Active Ingredients:

Homobrassinolide	0.0025%
Auxin:	
Indole-3-butyric acid	0.0100%
Gibberellin:	
Gibberellic acid GA3	0.0090%
Other Ingredients	99.9785%
Total	100.000%

*Alternate Brand Name: **ARISE®**

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call poison control center or doctor for treatment advice
Have the product container or label with you when calling a poison control center or doctor, or going for treatment You may also call the Pesticide Control Center at 1-800-222-1212 - 24 hours a day for emergency medical treatment information.	

SEE INSIDE THIS LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Contains 0.10% aminoindole-3-propionic acid

EPA Registration Number: 90930-

EPA Establishment # 90930-PA-0001

Manufactured by:
 Miller Chemical & Fertilizer, LLC
 P.O. Box 333, 120 Radio Road
 Hanover, Pennsylvania 17331
 1-800-233-2040
 Net Contents: ONE GALLON (3.78 Liters)

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin clothing or eyes. Wash thoroughly with soap and water after handling and before eating drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment

Applicators and other handlers must wear:

- A. long-sleeved shirt and long pants**
- B. Shoes plus socks**
- C. Protective eyewear**

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment. If no such instructions for washables, use detergent and hot water. Keep and wash personal protective equipment separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Requirements

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove personal protective equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For Terrestrial Uses: Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment, wash water or rinsate.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Before using **Miller 2361**, read and follow the precautions appearing on the label.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and pets out of the treated area until sprays have dried. Keep animals and unprotected persons out of operational areas during treatment.

Chemigation System

Apply **Miller 2361** only through the following types of systems: sprinkler, including center pivot, lateral move, end tow, side roll, traveler, big gun, solid set, or hand move; or drip (trickle) irrigation systems. Do not apply this product through any other type of system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable about the chemigation systems and responsible for its operation, or under supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Chemigation System Connected to Public Water Systems

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), back flow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Supply pesticide tank agitation, especially if product is to sit in tank for over 6 hours.

Miller 2361 may be applied continuously for the duration of water application or with the first quarter to one-half of the watering period.

Mixing instructions: Fill supply tank to 1/4 to 1/2 full. Add **Miller 2361** and complete filling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of reach of children, preferably in locked storage area.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Use Miller 2361 in combination with a well-balanced fertility program and good management practices. The Company recommends the use of soil and tissue testing, and additional nutrients and micronutrients as needed.

WARRANTY: The manufacturer warrants that this product shall be of its standard quality and shall conform to the label thereon. Recommendations for use of the product are based on tests believed to be reliable. The use of the product being beyond the control of the manufacturer, no guarantee expressed or implied is made as to the effects of such or the results to be obtained, except as expressly set forth on the label hereon. The BUYER must assume all responsibility if not used in accordance with directions or established safe practices, including injury or damage resulting from its misuse as such or in combination with other materials. The manufacturer's liability shall be limited to the replacement without charge, FOB warehouse, of all product shown to be otherwise than as warranted.

Miller 2361
For Use as a Plant Hormone Supplement
APPLICATION INSTRUCTIONS
SHAKE WELL BEFORE USING

Miller 2361 is a plant growth regulator/hormone supplement that provides good growing conditions for obtaining maximum benefits. For maximum gain from the use of **Miller 2361**, a well-balanced plant nutrient program should always be used. **Miller 2361**, in any of its applications, is not intended to replace fertilizer or to supply nutrients that would normally be added in a conventional fertility program. Timing of the foliar spray application is very important. Always follow directions precisely. Do not apply within eight hours of forecast rain. For best results, apply **Miller 2361** in the early morning or late afternoon, especially when temperature exceeds 95°F (36°C).

TRANSPLANTS: For a quick start, dip roots in a solution of 1 tablespoon **Miller 2361** per gallon of water prior to transplanting or drench flats with a solution of 1/2 oz. per gallon of water.

CROP USE GUIDELINES

A blanket exemption from the requirement of a tolerance has been issued by the U.S. EPA for the active ingredients in this product. This product is cleared for use on any and all crops.

For local use recommendations for major and minor crops, contact your PCA or local distributor representative.

For maximum benefit, all foliar applications require Calcium EDTA or other highly available calcium source in the tank mix.

<u>CROP</u>	<u>OZ/ACRE</u> <u>(each application)</u>	<u>TIMING AND FREQUENCY</u>
Cotton	1-2 fl. oz.	Apply in seed furrow.
	2 fl. oz.	Spray in a band at 1 to 3 leaf stage.
	3-4 fl. oz.	Spray at pinhead or match head square.
	6-8 fl. oz.	Spray at early bloom and to late bloom.
Corn (field)	2 fl. oz.	Apply in seed furrow at planting.
	6 fl. oz.	Spray in band at 5 to 7 leaf stage.
	6 fl. oz.	Repeat 2 weeks later.
Corn (sweet, popcorn)	2 fl. oz.	Apply in furrow prior to planting or with seed.
	6 fl. oz.	Spray in a band at 3 to 5 leaf stage (12" to 16").
	6 fl. oz.	Repeat 2 weeks later.
Rice	8 fl. oz.	Spray at 3 to 7 leaf stage.
	8 fl. oz.	Spray at panicle differentiation.
Sorghum (Milo)	2 fl. oz.	Apply in seed furrow at planting.
	4-6 fl. oz.	Banded spray at 5 to 7 leaf stage.
	4-6 fl. oz.	Apply at boot to early bloom.
Soybeans	4 fl. oz.	Broadcast preplant incorporated with herbicide or 1-2 fl. oz./A in furrow with seed.
	4 fl. oz.	Spray at third to fifth trifoliolate.

Sugar Beets	6 fl. oz.	Spray at pod fill.
	4 fl. oz.	Banded spray at the 6 to 8 leaf stage.
	8 fl. oz.	30 days after first application

CROP USE GUIDELINES (CONTINUED)

<u>CROP</u>	<u>OZ/ACRE</u> <u>(each application)</u>	<u>TIMING AND FREQUENCY</u>
Winter wheat, Barley, Rye	For winter grazing: Make a second 4 to 8 fl. oz. application when spring growth begins after vernalization to increase grain production.	Apply 8 fl. oz./Acre two weeks after emergence.
Spring wheat, Barley, Rye, Oats	Spray 8 fl. oz. when plants have 3 to 5 true leaves emerged.	
Forage crops- Legumes or grasses	8-16 fl. oz.	Spray Miller 2361 4 to 6 weeks after emergence and monthly thereafter. Mature Crop: Spray Miller 2361 as spring growth begins, 1 week before harvest and again 2 weeks after cutting.
Seed production	8-16 fl. oz.	On established crops: spray Miller 2361 at the beginning of inflorescence development (early tillering) and again 2 weeks later. Spray 8 to 16 oz/acre at the beginning of bloom. Spray fern 2 weeks after last harvest. Spray monthly during fern growth.
Asparagus	12 fl. oz. 8 fl. oz.	Spray banded at the third trifoliolate. Spray (broadcast) at first bloom.
Beans (all)	3 fl. oz. 6 fl. oz.	Spray at the third trifoliolate.
Field Peas	3 fl. oz.	Spray at first bloom. Band 2 weeks after transplant. Band 4 weeks after transplant. Repeat biweekly.
Broccoli, Cabbage, Cauliflower, Celery Brussels Sprouts	2 fl. oz. 4 fl. oz.	
Canola	4 fl. oz. 6-8 fl. oz.	Apply at 3 leaf stage. Apply again two weeks later
Carrots, Beets, Other root crops	8-12 fl. oz. 4-8 fl. oz.	Apply when seedlings have 3 to 6 leaves. Follow at 2 to 4 week intervals.
Cucurbita: watermelons, Cantaloupe, cucumbers, Muskmelons	2 fl. oz. 4 fl. oz. 6 fl. oz.	Banded at 2 to 4 leaf stage. Banded when plants show first signs of running. Broadcast two weeks after first application.
Grapes	4-8 fl. oz.	General: Apply Miller 2361 at 4 oz. with all foliar nutritional or pesticidal sprays. Sizing: Apply as tank mix with all GA sizing sprays. Harvest: Apply Miller 2361 with high potash fertilizer at 2 to 10 days before harvest to enhance sugar accumulation.
Onions, Garlic	Make first application at bulb initiation at 8 to 16 fl. oz./acre. Repeat at two week intervals for up to 4 applications.	
Peas (Field, Chickpea, etc.)	3 fl. oz.	Spray at the third trifoliolate. Spray at first bloom.

Peanuts	2 fl. oz. 6 fl. oz.	Two weeks after emergence, banded. Apply at bloom and at initial pegging.
Peppers: Bell	2-4 fl. oz.	Banded at the 3 to 5 leaf stage.
Chile, Cayenne, Jalapeno	4-8 fl. oz.	Apply at 7 to 14 day intervals for 4 to 6 applications.
Eggplant		

CROP USE GUIDELINES (CONTINUED)

<u>CROP</u>	<u>OZ/ACRE (each application)</u>	<u>TIMING AND FREQUENCY</u>
Potatoes	4 fl. oz. 6 fl. oz. 8 fl. oz.	Add to fertilizer and incorporate in seed furrow prior to planting. Banded at stolonization. Broadcast 2 to 4 weeks later.
Spinach, lettuce, and other leafy vegetables	2-4 fl. oz. 4-8 fl. oz.	Begin at the 3 leaf stage and apply weekly at 4 to 6 oz. thereafter.
Squash	3 fl. oz. 8 fl. oz. 8 fl. oz.	Band at 2 to 4 leaf stage. Broadcast at early bloom. Broadcast at 14-day intervals.
Sugarcane	16 fl. oz.	Planting: In furrow over newly laid cane. Foliar: 1st – At beginning of raton bud extension. 2nd – At beginning of sugar accumulation. 3rd – One to three weeks before harvest.
Sunflower	2 fl. oz. 6 fl. oz.	Apply two weeks after emergence. Apply at bloom.
Strawberries	8 fl. oz. 8 fl. oz.	Broadcast 2 to 3 weeks prior to coming out of dormancy. Broadcast at early bloom and at 14-day intervals thereafter.
Tomatoes (processing)	2 fl. oz. 8 fl. oz.	Apply in a band (14") 1 week after transplant or at 6 to 8 leaf stage. Broadcast at early bloom and again 2 weeks later.
Tomatoes (fresh market)	2 fl. oz. 4 fl. oz. 8 fl. oz.	Apply in a band (14") 1 week after transplant or 6 to 8 leaf stage. Band 3 weeks later. Broadcast with calcium or foliar fertilizer every 14 days.
Nut crops - mid	16 to 32 fl. oz.	Apply Miller 2361 with 10 lb/acre low-biuret urea at nut fill.
Almonds, Pecans, Pistachios, Filberts,		and again one month later. Add 8 oz. of Miller 2361 per acre to each zinc or calcium spray.
Walnuts, Cashews		Apply 16 to 32 oz/acre prior to flowering. Ask your local PCA for specific regional timing.

All Fruits: Apple, Cherry, Citrus (Orange, Lemon, etc.) Banana, Stonefruits (Peach, Plum, etc.), Pear, Mango, Papaya, Pineapple

Transplants: Follow general transplant instructions.

Fruit trees in production: Spray fruit trees with a solution of 1 oz **Miller 2361** in 4 gallons water (or 1 to 2 pints/A) at the following growth stages:

1. At bud break to increase pollination efficiency. (**Miller 2361** will not harm bees or pollinating insects);
2. At 1 week after petal fall to promote cell division;
3. At 1 to 2 weeks before fruit drop to reduce physiological stress and reduce fruit drop;
4. At 20 to 30 days after petal fall to increase fruit size;
5. Monthly during fruit growth and development to promote nutrient translocation to produce larger and better quality fruit.

Non-Bearing Use for TREES, FRUITS, NUTS, BERRIES, SHRUBS AND WOODY ORNAMENTALS:

To aid in propagation of trees, fruits, berries, soft wood cuttings, shrubs and woody ornamentals and to reduce transplant shock, to promote growth and vigor and reduce stress in non-bearing fruit trees such as apple, peach; berry and vine crops such as cranberries; evergreen trees such as spruce, fir, pine; deciduous trees such as birch, elm, maple; flowering plants and shrubs such as poinsettia, rose, azalea, rhododendron, crepe myrtle; and for other flowering and non-flowering shrubs

New cuttings: Spray **Miller 2361** at 1 to 2 pints per acre on the stems, branches, vines or canes to be propagated from 1 to 7 days before cutting. After planting, spray **Miller 2361** at 1/2 to 1 pint or apply through the irrigation system at weekly intervals until the plants are established.

Replant areas: Spray the plants before cutting. Then spray **Miller 2361** weekly at 1/2 to 1 ounce per 1500 square feet and irrigate in. Continue weekly to biweekly applications until the plants are established.

Established Trees and Shrubs: Spray 1 to 2 pints per acre, or a mixture of 1 oz. **Miller 2361** to 4 gallons water to thoroughly wet the foliage at any or all of the following growth stages:

1. Early spring to promote bud initiation;
2. At bud break;
3. At terminal calyx;
4. Early to mid fall.

For best results apply Miller 2361 with foliar nutrients, micronutrients or secondary nutrient sprays such as calcium, iron, and zinc.

TURF

Spring Application: Make an early application of 1/2 to 1 fl. oz. **Miller 2361** per 1000 sq. ft. to promote tiller, rhizome or stolon growth, to develop a deep root system, and to give the turf a rapid start once winter dormancy is broken and growth begins. Continue monthly 1/2 fl. oz./1000 sq. ft. applications throughout spring and summer. Spring application is important to develop a deep root system which will condition the turf and reduce the stress of disease and summer heat or low rainfall. Apply **Miller 2361** with iron sulfate for maximum root growth response.

Fall Application: Two to three applications of **Miller 2361** (1/2 to 1 fl. oz. per 1000 sq. ft.) should be made in the fall beginning about eight weeks before the turf becomes dormant (eight weeks before first frost date in the northern states) to promote root growth and provide the grass with the vigor to better endure the stress of winter (reduce winter kill) and improve survival of a good healthy turf for the following spring.

Golf Greens, Fairways, Football and Soccer Fields, and Baseball Infields and Outfields: At the beginning of spring growth apply 1 to 2 fl. oz. per 1000 square feet at the breaking of dormancy. Make successive maintenance applications of 1/2 to 1 fl. oz. per 1000 sq. ft. at monthly intervals or as needed to maintain root growth, tillering, appearance and vigor throughout the growing season. During periods of intensive use apply 1/2 to 1 fl. oz. per 1000 sq. ft. weekly to the greens, infield or playing field to maintain root structure and renew growth and vigor between games. Make three applications of 1/2 fl. oz. per 1000sq. ft. at 2 week intervals in the fall beginning about eight (8) weeks before turf becomes dormant to promote root growth and increase winter stamina to reduce winter kill.

Sod: Spray **Miller 2361** to newly laid sod at 1/2 to 1 fl. oz. per 1000 sq. ft. to promote rooting and increase the rate of sod establishment. Maintain growth and vigor with monthly applications of 1/2 to 1 fl. oz. per 1000 square feet.

Sod Farming: Spray **Miller 2361** at 1/2 to 1 pint/acre monthly to sod fields to promote root, tiller and rhizome growth and to bring the crop to harvest more quickly. Maintain accelerated growth with **Miller 2361** applications of 1/2 to 1 pints/acre at 2 to 4 week intervals or as needed. Spray **Miller 2361** at 1/2 to 1 pints per acre to sod 1 to 4 days before harvesting to initiate new root growth and speed up establishment when sod is laid.

Lawns, Playgrounds, Parks, Recreational Areas, Landscaped Roadways and Cemeteries: Apply 1 fl. oz. per 1000 sq. ft. at the beginning of spring growth to promote a deep root system and tillering to fill sparse areas. Apply at 1/2 to 1 fl. oz. per 1000sq. ft. monthly to maintain health and vigor of the turf. Application can be made more frequently as needed to condition the turf for stress or for periods of heavy use.

Nutritional sprays: For better color response from nitrogen, iron, sulfur, zinc and other nutrient sprays use 1/2 to 1 pt. **Miller 2361** per acre with nutrient spray solution. For greens or smaller area, add 1/2 to 1 fl. oz. **Miller 2361** per 3 to 5 gallons spray solution.

NURSERY AND GREENHOUSE USE

To promote bud differentiation, cell division, root induction and growth and to reduce apical dominance. Use **Miller 2361** HMS in your watering program or as foliar spray.

Propagation of Cuttings: Dip cuttings in rooting hormone powder or solution and stick in rooting medium. Spray or mist cuttings with a solution of 1 fl. oz. **Miller 2361** to 4 gallons water (1 qt./100 gallons) at weekly intervals until root buds initiate. Then spray at 2 to 4 week intervals.

Transplanting: Add 1 fl. oz. **Miller 2361** per 4 gallons of transplant solution (fertilizer-water). Drench the root zone. Follow with spray to foliage or add through irrigation system at 2 to 4 week intervals at the rate of 1 quart per 100 gallons.

Production: To increase growth rate, improve quality and resilience of nursery and greenhouse crops, add 1 fl. oz. **Miller 2361** per 4 gallons (1 qt./100 gallons) of fertilizer or water solution and apply through the irrigation system or via foliar spray.

Nutritional Deficiencies: To promote rapid uptake and correction of nutrient deficiencies in ornamentals and turf, add **Miller 2361** to iron, nitrogen fertilizers, zinc or other nutrient solutions at the rate of 1 fl. oz. per 4 gallons (1 qt./128 gallons). Apply as a foliar spray or soil drench.

To promote rapid uptake and correction of nutrient deficiencies in ornamentals and turf, add MILLER 2360 to iron, nitrogen, fertilizers, zinc or other nutrient solutions at the rate of 1 fl. oz per 4 gallons (1 quart/128 gallons). Apply as a foliar spray or soil drench.

LANDSCAPE MANAGEMENT (see Turf uses also)

Bedding Plants: Spray bedding plants at 2 to 4 week intervals with a solution of 1 fl. oz. **Miller 2361** per 4 gallons water (1 qt./128 gallons), fungicide or nutrient spray to promote growth, flowering and maximum color development.

Lawn Care: Spray **Miller 2361** to lawns at the rate of 1/2 fl. oz. per 1000 sq. ft. **Miller 2361** can be added to liquid fertilizer, insecticide, fungicide, or herbicide sprays.

Transplanting of trees, shrubs or bedding plants: See transplanting instructions under nursery use.

Maintenance: To promote growth and reduce stress from drought, disease or nutrient deficiency. Spray **Miller 2361** to foliage at the rate of 1 fl. oz. per 4 gallons of water or fertilizer or pesticide solution (1 qt./128 gallons).

Nutritional Deficiencies: To promote rapid uptake and correction of nutrient deficiencies in ornamentals and turf, add **Miller 2361** to iron, nitrogen fertilizer, zinc or other nutrient spray solutions at the rate of 1 fl. oz. per 4 gallons (1 qt./128 gallons). Apply as a foliar spray or soil drench.

Root Feeding: Mix **Miller 2361** with root feeding solutions at the rate of 1 fl. oz. per 4 gallons of nutrient solution (1 qt./128gallons).

SEED TREATMENT MILLER 2361

As a seed treatment for seeds prior to planting **MILLER 2361** may be applied to seed up to 6 months prior to planting. Dilute the recommended rate with a sufficient amount of water for uniform coverage. Mix thoroughly to coat seed and allow to dry before planting.

MILLER 2361 can be applied with fungicide treatment or to fungicide-treated seed. Do not use treated seed for food, feed or oil purposes. Commercial seed processors must apply with sufficient EPA-approved dye to assure adequate seed coloring. Commercially-treated seed must be labeled in accordance with Federal Seed Act. For seed treated at planting, treat only those seeds needed for immediate use and planting. Do not store excess treated seed beyond planting time. Dispose for excess treated seed by burial away from streams and bodies of water.

Crop	Recommended Rate Ounces/10 lbs.	ml/kg
Cotton, Peanuts	0.5 to 1.5	3.0 to 10.0
Wheat, soybeans, beans, peas	0.1 to 0.15 (1.0 to 1.5/100 lbs.)	0.6 to 0.9
Corn, rice, grain sorghum	0.1 to 0.15 (1.0 to 1.5/100 lbs.)	0.6 to 0.9
Potato seed pieces	1/400 dip for 1 minute	
Sweet corn, popcorn	1.0 to 2.0	6.0 to 12.0
Alfalfa, clover	0.75 to 1.5	5.0 to 10.0
Chiles, peppers, tomatoes	1.5 to 3.0	10.0 to 20.0
Cucumbers, melons, cantaloupes, Honeydews, muskmelons, watermelons, squash (all varieties)	0.5 to 1.0	3.0 to 6.0
Carrot, lettuce cabbage, broccoli	2.5 to 5.0	15.0 to 30.0
Okra, onion, garlic, spinach	1.0 to 3.0	6.0 to 18.0
Turf grasses	1.5 to 2.5	10.0 to 15.0

08/14/18