



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

91234-2

Date of Issuance:

7/22/15

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Kylix SC Lawn Care Insecticide

Name and Address of Registrant (include ZIP Code):

Brooke Hedrick
SynTelus, LLC
1204 Village Market Place, #273
Morrisville, NC 27560

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Venus Eagle, Product Manager 01
Invertebrate and Vertebrate Branch 3
Registration Division (7505P)

Date:

7/22/15

2. You are required to comply with the data requirements described in the DCI or EDSP Order identified below:
 - a. Clothianidin GDCI-044309-1185
 - b. Bifenthrin GDCI 128825-902
 - c. Bifenthrin EDSP-128825

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI or EDSP Order listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division:

http://www.epa.gov/oppsrrd1/contacts_prd.htm

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 91234-2.”
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 03/30/2015

If you have any questions, please contact Marianne Lewis by phone at (703) 308-8043, or via email at lewis.marianne@epa.gov.

Enclosure: Stamped label

KYLIX SC Lawn Care INSECTICIDE

GROUP	4A	INSECTICIDE
GROUP	3	INSECTICIDE

For contact and systemic insect pest control in lawns, landscape ornamentals trees (including nonbearing fruit trees), around residential, institutional, public, commercial and industrial buildings, parks, recreational areas, and athletic fields. Not for use on golf courses and sod farms.

ACTIVE INGREDIENT:

Clothianidin*	24.70%
Bifenthrin**	12.30%
OTHER INGREDIENTS	<u>63.00%</u>
TOTAL:	100.0%

* (E)-1-(2-chloro-1,3-thiazol-5-ylmethyl)-3-methyl-2-nitroguanidine.

** 2-methylbiphenyl-3-ylmethyl (1 RS,3RS)-3-[(Z)-2-chloro-3,3,3-trifluoroprop-1-enyl]-2,2-dimethylcyclopropanecarboxylate (cis isomers 97% minimum, trans isomers 3% maximum)

Contains 2.22 lb of Clothianidin and 1.10 lb of Bifenthrin per gallon

KEEP OUT OF REACH OF CHILDREN WARNING AVISO


Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle
(If you do not understand the label, find someone to explain it to you in detail.)
See inside booklet for additional precautionary statements

FIRST AID	
If Swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow Do not induce vomiting unless told to by the poison control center or doctor Do not give anything by mouth to an unconscious person
If Inhaled	<ul style="list-style-type: none"> Move person to fresh air If person is not breathing, call 911 or ambulance, then give artificial respiration preferably mouth-to-mouth, if possible Call a poison control center or doctor for further treatment advice
If In Eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye Call a poison control center or doctor for treatment advice
If on Skin or Clothing	<ul style="list-style-type: none"> Take off contaminated clothing Rinse skin immediately with plenty of water for 15-20 minutes Call a poison control center or doctor for treatment advice
<p>NOTE TO PHYSICIAN: This pesticide contains a pyrethroid. If significant quantities have been swallowed, promptly eliminate the ingested quantities from the stomach and intestines. Treatment is symptomatic and supportive. Alcohol, milk, cream, and other digestible fats and oils should NOT be administered as they may increase absorption.</p>	
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-844-685-9173 for emergency medical treatment information. FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC at 1-800-434-9300 or 1-703-527-3887 if calling from outside of the U.S.</p>	

EPA Reg. No. 91234-xx

EPA Est. No.

NET CONTENTS:
Manufactured For:
SynTelus, LLC
1204 Village Market Place, #273
Morrisville, NC 27560



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING. May be fatal if swallowed. Harmful if inhaled. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid breathing spray mist or vapor. Avoid contact with skin, eyes, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- long-sleeved shirt and long pants
- shoes plus socks
- chemical-resistant gloves: barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride, or viton

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or runoff into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply where runoff is likely to occur.

This product is toxic to bees exposed to treatment and for more than 5 days following treatment. Do not apply this product to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period, unless the application is made in response to a public health emergency declared by appropriate state or federal authorities.

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at:

<http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>.

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: www.aapco.org/officials.html. Pesticide incidents should also be reported to the

National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Carefully read and understand the Directions for Use and restrictions before applying this product.

Do not apply this product in any way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.



Do not apply KYLIX SC Lawn Care while bees are foraging. Do not apply KYLIX SC Lawn Care to plants that are flowering. Only apply after all flower petals have fallen off..

[Language within brackets is optional text related to State specific information.]

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

- 1) Treatment to soil or vegetation around structures;
- 2) Applications to lawns, turf, and other vegetation;
- 3) Applications to building foundations, up to a maximum height of 3 feet.

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only.

RESTRICTIONS

Do not apply by air.

Do not water the treated area to the point of run-off.

Do not make applications during rain.

Do not mow until after irrigation or rainfall has occurred.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

Do not apply through any type of irrigation system.

Do not apply on the following: Commercial nurseries, commercial greenhouses, grass grown for seed, golf courses and sod farms.

Do not allow children or pets on treated surfaces until the spray has dried.

USE INFORMATION

KYLIX SC Lawn Care is a long lasting contact and systemic insecticide for control of listed insects infesting lawns and ornamentals in landscapes and surrounding areas. **KYLIX SC Lawn Care** can be used around residences, business complexes, industrial complexes, airports, playgrounds, school yards, parks and recreation areas, athletic fields, cemeteries and arboretums.

Container Use Directions

1. Remove the measuring chamber cap and induction seal. Replace cap and securely tighten.- Tip container until liquid fills measuring chamber.
2. Return container to level position. No adjustment is needed.
3. Remove measuring chamber cap and dispense into proper application equipment.

For multiple dose measuring: Remove fill chamber cap and dispense according to markings on side of bottle.

PREPARATION OF SPRAY MIXTURES AND APPLICATION

Begin with clean equipment. Add sufficient clean water to the spray tank for one-half of the mix load. Start tank agitation. Add the necessary amount of **KYLIX SC Lawn Care** before adding any other products. Agitate to ensure thorough mixing while adding the remaining water. Maintain agitation during mixing and application. Apply with properly calibrated spray equipment. If agitation is stopped for any reason the spray solution must be thoroughly remixed prior to any further use.

Use sufficient water volume to provide uniform coverage of the lawn and ornamental plants. For control of listed surface insects, do not water in until at least 48 hours after the application. If the target insects are in the thatch, either apply **KYLIX SC Lawn Care** in enough spray volume to wet the thatch or lightly water in after application with enough water to wet the thatch zone.

For control of existing grub populations, water in after application with enough water to move the insecticide into the root zone where the grubs are feeding.

RESISTANCE MANAGEMENT

KYLIX SC Lawn Care contains two active ingredients with different modes of action and provides a recognized built in insecticide resistance management strategy to minimize the development of insecticide resistance in insect pests. The active ingredients in **KYLIX SC Lawn Care** are clothianidin, a Group 4A insecticide, and bifenthrin a Group 3 insecticide. Insect biotypes with acquired resistance to Group 4A or Group 3 insecticides may eventually dominate the insect population if Group 4A or Group 3 insecticides are used alone or repeatedly in the same site or in successive years as the primary method of control for targeted species.

To delay insecticide resistance in insects that are at risk of developing resistance to insecticides consider:

- Avoid the consecutive use of insecticides on the same insect species.
- Use tank-mixtures or pre-mixes of insecticides from a different target site of action.
- Base insecticide use on a comprehensive IPM program.
- Monitor treated insect populations for loss of field efficacy.
- Contact your State Agricultural Experiment Station, State Extension Specialists, other lawn experts and/or SynTelus, LLC for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems.
- For further information or to report suspected resistance, you may contact SynTelus, LLC at 1-984-664-9804.

APPLICATION INSTRUCTIONS - TURFGRASS AND LAWNS

KYLIX SC Lawn Care can be used for control of listed insects in lawns. **KYLIX SC Lawn Care** can be used around residences, business complexes, industrial complexes, airports, playgrounds, school yards, parks and recreation areas, athletic fields, cemeteries and arboretums.

KYLIX SC Lawn Care has sufficient contact and residual activity to provide high levels of control. The need for an application can be based on historical monitoring of the presence of or anticipated occurrence of individual or multiple insect pest species, the site, previous records or experiences, current season adult trapping or other methods. Optimum control of white grubs will be achieved when applications are made just prior to egg laying or to early instar larvae of target pests, followed by sufficient irrigation or rainfall to move the active ingredient into the zone where the insect pests are active. Multiple applications can be made each growing season but do not use more than 23.3 fl oz of product (0.4 lb of clothianidin active ingredient, 0.2 lb bifenthrin) per acre per year. Note: You may also use additional products containing bifenthrin as long as the maximum amount of bifenthrin does not exceed 0.4 lb active ingredient per acre per year.

Apply KYLIX SC Lawn Care to lawns and perimeters at 11.65 - 23.3 fl oz per acre per year (0.27 - 0.54 fl oz/1,000 sq ft). The rate is dependent on the target pest(s), their stage of development and the desired level of control. Optimum control of white grubs (larvae) will be achieved when applications are made just prior to or just after peak egg laying. Use the higher rates within the rate range when long residual control is desired for control of multiple insect pests or late season applications are made to lawns already showing grub damage or damage by scavenging animals. Multiple applications can be made but do not exceed the maximum amount per year (23.3 fl oz/A or 0.54 fl oz/1,000 sq ft). Consult your local State Agricultural Experiment Station, State Extension Turf Specialists or other lawn experts for specific information concerning the timing of applications.

For early season control of listed insect pests (such as annual bluegrass weevil) and for full season control of white grubs, make the first application of **KYLIX SC Lawn Care** at peak adult activity or when other insect pests reach a damaging population in the spring at a rate of 11.65 - 23.3 fl oz per acre (0.27 - 0.54 fl oz/1,000 sq ft). For control of white grubs at standard application timing, apply 11.65-23.3 fl oz/A (0.27-0.54 fl oz/1,000 sq ft) just prior to peak adult activity or to early instar larvae.

For control of listed surface insects, do not water in until at least 24 hours after the application. For control of the target insects in the thatch, use enough carrier volume to move the insecticide into the thatch or lightly water in after application with enough water to wet the thatch zone. For control of armyworms, cutworms and other surface feeders, apply 11.65-23.3 fl oz/A (0.27-0.54 fl oz/1,000 sq ft) at the first sign of insects or insect damage. Repeat applications up to a total of 23.3 fl oz/A per year (0.54 fl oz/1,000 sq ft) to control additional flushes of insects. Up to a total of 23.3 fl oz/A per year (0.54 fl oz/1,000 sq ft) can be applied.

For control of existing grub populations, irrigate after application with enough water to move the insecticide into the zone where the grubs are feeding.

Do not mow until after irrigation or rainfall has occurred.

TURFGRASS AND LAWNS

APPLICATION RATE	INSECT PEST
0.27 - 0.54 fl oz per 1,000 sq ft (11.65 - 23.3 fl oz/A)	Annual Bluegrass Weevil/Hyperodes (Adults) ¹ Annual Bluegrass Weevil/Hyperodes (Larvae) ¹ Ants ² , Nuisance (Excluding fire, harvester, pharaoh, and carpenter ants) Armyworms Billbugs ³ Black Turfgrass Ataenius ⁴ Chiggers Chinch Bugs Cutworms European Crane Fly Grasshoppers Leafhoppers Mole Crickets ⁵ Pillbugs Sod webworms Sowbugs Spittlebug Sugarcane grub Ticks ⁶ White grubs ⁷ (Asiatic Garden Beetle, European Chafer, Green June Beetle, Japanese Beetle, Northern Masked Chafer, <i>Phyllophaga</i> spp. (May or June Beetle), Oriental Beetle, Southern Masked Chafer)
Control 0.46 fl oz per 1,000 sq ft (20 fl oz/A)	Fire Ants, Red Imported ^{8[*]}
Suppression 0.33 fl oz per 1,000 sq ft (14.4 fl oz/A)	

¹ **Annual Bluegrass Weevil:** Application can be timed for adult weevils as they move from their over-wintering sites into turf. This movement usually begins shortly before or when Forsythia is in full bloom and continues to full bloom of dogwoods. Larvae from adult weevils that escape the application will be controlled with the application directed at adult weevils. Applications made after adult weevil activity will control larvae. Consult your State Cooperative Extension Service for more specific information regarding timing of application. Full season control of white grubs will be obtained with an application of 14 fl oz/A targeted towards Annual Bluegrass Weevil/Hyperodes.

² Applications of **KYLIX LC SC** in the late fall or early spring will reduce populations of nuisance ants on residential lawns. Make an application in the fall before ants stop foraging or after they become active in the spring. Make a broadcast application and lightly water in to wet the turf and thatch area where nuisance ants are foraging.

³ **Billbugs:** Application can be made when adult Billbugs are first observed, usually in April and May in temperate regions. A spring application will also control Billbug larvae. Consult your State Cooperative Extension Service for more specific information regarding timing of application. White grubs will be controlled with an application targeted towards Billbugs.

⁴ **Black Turfgrass Ataenius:** Application can be timed to coincide with the first generation of Black Turfgrass Ataenius that usually appears during May. White grubs will be controlled with an application targeted towards Black Turfgrass Ataenius.

⁵ **Mole Crickets:** Apply during peak adult flight and egg lay. If the soil is dry, irrigate before application to stimulate insect activity and water in lightly after application to move the insecticide into the zone where the mole crickets are active.

⁶ **Ticks (including ticks that may transmit Lyme Disease and Rocky Mountain Spotted Fever):** Treat the entire area where exposure to ticks may occur. Use higher spray volumes when treating areas with dense groundcover or heavy leaf litter. Ticks may be reintroduced from surrounding areas on host animals.

Deer Ticks (*Ixodes* sp.): Apply in late fall and/or early spring to control adult ticks that are usually located on brush or grass above the soil surface and in mid- to late spring to control larvae and nymphs that reside in the soil and leaf litter.

American Dog Ticks (*Dermacentor* sp.): These ticks commonly congregate along paths or roadways where humans are likely to be encountered. Apply from mid-spring to early fall to control American dog tick larvae, nymphs and adults.

⁷ **KYLIX LC SC** will control white grubs anytime applications are made when they are active from early spring to late fall. The residual activity of **KYLIX LC SC** is long enough to provide season long control of white grubs when applied for control of listed early season insects such as Annual Bluegrass Weevil, Billbugs, Nuisance Ants, or overwintering White Grubs.

⁸ **Red Imported Fire Ants[*]:** If the soil is not moist, irrigate before application or use a high volume application. For mound treatments, use one teaspoon **KYLIX LC SC** per gallon of water (this is equivalent to 14.4 fl oz **KYLIX LC SC** per acre applied in 87 gallons of water) and apply 1-2 gallons of finished spray per mound. The solution should be applied in such a manner to allow the solution to move into the mound. Also treat a four foot diameter circle around the mound. Apply in cool weather (65 - 80° F) or in early morning or late evening hours when most of the foraging workers are in the colony.

RESTRICTIONS

- Do not apply more than a total of 23.3 fl oz of **KYLIX LC SC** (0.4 lb of clothianidin active ingredient, 0.2 lb bifenthrin) per acre per year.
- If the maximum season limit of **KYLIX LC SC** has been applied and pest populations require additional treatments, use another registered pesticide labeled for control of the targeted insects that does not contain clothianidin. You may use additional products containing bifenthrin as long as the maximum amount of bifenthrin does not exceed 0.4 lb active ingredient per acre per year
- Do not allow this product to contact plants in bloom if bees are foraging the treatment area.
- Do not allow children and pets to enter the treated area until sprays have dried.
- Not for use on golf courses and sod farms.

[*Not approved in California.]

APPLICATION INSTRUCTIONS - LANDSCAPE ORNAMENTALS

KYLIX SC Lawn Care is a contact and long lasting systemic insect control product for listed insects infesting landscape ornamentals adjacent to turf areas. **KYLIX SC Lawn Care** can be used around residences, business complexes, industrial complexes, airports, playgrounds, school yards, parks and recreation areas, athletic fields, cemeteries and arboretums.

For foliar applications, apply enough spray solution to wet the surface of the leaves to the point of drip.

For control of listed surface insects and insects that crawl along the surface of the soil or mulch under ornamentals, water in lightly after application to move the insecticide into the areas where insects are active. Water in with enough irrigation or rainfall to move the active ingredient into the zone where the target insects are active. Soil surface applications will provide control of foliage feeding insects; however, optimum control will not be achieved until the active ingredient is taken up by the plant roots and translocated to the area where insects are feeding.

LANDSCAPE ORNAMENTALS			
Foliar Application KYLIX SC Lawn Care Rate (fl oz/100 gals)	Foliar Insect Pests Controlled		Application Instructions
3.6-15.5 (5.0 -22.5 fl oz/A based on 150 gallons of spray per acre)	Aphids Caterpillars Leafhoppers Mealybugs White Flies		Target the spray towards the areas where the insects are active. Use enough water carrier to wet the surfaces on the target plants to the point of drip.
5.0-15.5 (7.5 -22.5 fl oz/A based on 150 gallons of spray per acre)	Ants ¹ (Excluding fire, harvester, pharaoh, and carpenter ants) Bagworms Black Vine Weevil (adults) Brown Soft Scale Budworms California Red Scale (crawlers) Chiggers Clover Mites Crickets	Cutworms Earwigs Elm Leaf Beetles Flea Beetles Japanese Beetles (Adults) Plant Bugs Spiders Tent Caterpillars Tip Moths Weevils	Repeat applications may be made if insects reappear but do not exceed a total of 0.54 fl oz KYLIX SC Lawn Care per 1,000 sq ft of landscape.
8.0-15.5 (12 -22.5 fl oz/A based on 150 gallons of spray per acre)	Adelgids Azalea lacebugs Beetles Cicadas (adults) Citrus Thrips Diaprepes (adults) Florida wax scale Glassy winged Sharpshooter	Gypsy moth Lacebug Leaf miners Leaf rollers Mites ² Pine Needle Scale (crawlers) Plant bugs (incl. <i>Lieges</i> spp.) Scales, soft	

¹ **KYLIX SC Lawn Care** will control aphids, scale insects, mealybugs and other sucking pests on ornamentals and limit the honeydew available as a food source for ant populations. Ants (excluding fire, harvester, carpenter and pharaoh ants) present at the time of application will also be controlled.

² **KYLIX SC Lawn Care** will provide suppression of Broad Mites, Clover Mites, European Red Mites and Spider Mites when used for control of other listed insects. Use a surfactant on hard to wet plants. The use of an alternate class of chemistry in a mite treatment program is recommended to prevent or delay development of bifenthrin resistant mite populations.

RESTRICTIONS

- Do not apply more than a total of 23.3 fl oz of **KYLIX SC Lawn Care** (0.4 lb of clothianidin active ingredient, 0.2 lb bifenthrin) per acre per year.
- If the maximum season limit of **KYLIX SC Lawn Care** has been applied and pest populations require additional treatments, use another pesticide labeled for control of the targeted insects that does not contain clothianidin. You may also use additional products containing bifenthrin as long as the maximum amount of bifenthrin does not exceed 0.4 lb active ingredient per acre per year.
- Follow application restrictions on this label, designated by the bee icon, to protect bees and other insect pollinators.

SOIL SURFACE or MULCH APPLICATION	SOIL INSECT PESTS CONTROLLED		APPLICATION INSTRUCTIONS
0.27 - 0.54 fl oz per 1,000sqft (11.65-23.3 fl oz/A)	Ants, Nuisance (Excluding fire, harvester, pharaoh, and carpenter ants) Beet Armyworm Beetles Black Vine Weevils (Larvae) Centipedes Crickets Cutworms Diaprepes (Adults) Earwigs Fall Armyworms Grasshoppers Millipedes	Pillbugs White grubs (Asiatic Garden Beetle, European chafer, Green June beetle, Japanese beetle, Northern Masked Chafer, <i>Phyllophaga</i> spp. (May or June Beetle), Oriental Beetle, Southern Masked Chafer	Apply in enough water to wet the areas where the insects are active or water in after applications with enough irrigation to move the insecticide into the areas where insects are active. For ornamental trees and woody shrubs see below.

SOIL SURFACE or MULCH APPLICATION	ABOVE GROUND INSECT PESTS CONTROLLED		APPLICATION INSTRUCTIONS
0.34 - 0.54 fl oz per 1,000sqft (15-23.3 fl oz/A)	Aphids Azalea lacebug Borers Black Vine Weevils (Adults) Caterpillars Flea Beetles Florida wax scale Glassy winged Sharpshooter Gypsy Moth Caterpillars	Japanese Beetles (Adults) Leafhoppers Leafminers Leafrollers Mealybugs Plant Bugs (incl. <i>Lygus</i> spp.) Root Weevils (incl. Black Vine Weevils) Scales, Soft Whiteflies	Apply in enough water to wet the root zone of the plants. Control of leaf feeding insects will be delayed until the insecticide is taken up by the roots and moved into the areas where insects are feeding, usually 2-4 weeks. For ornamental trees and woody shrubs see below.

RESTRICTIONS

- Do not apply more than a total of 23.3 fl oz of **KYLIX SC Lawn Care** (0.4 lb of clothianidin active ingredient, 0.2 lb bifenthrin) per acre per year.
- If the maximum season limit of **KYLIX SC Lawn Care** has been applied and pest populations require additional treatments, use another pesticide labeled for control of the targeted insects that does not contain clothianidin. You may use additional products containing bifenthrin as long as the maximum amount of bifenthrin does not exceed 0.4 lb active ingredient per acre per year.

Drench application to ornamental trees and woody shrubs adjacent to turf:

KYLIX SC Lawn Care can be used for systemic control of listed insects with an application to trees in landscapes and interior plantscapes. **KYLIX SC Lawn Care** contains clothianidin which is systemic and will be translocated throughout the tree. Treat the area over the root zone of the trees or apply the insecticide in the "root well" of the plant.

Apply the insecticide in enough water to get uniform coverage and water in after application. **KYLIX SC Lawn Care** is a systemic product and will be translocated upward into the plant system from root uptake. It must be placed in an area where the roots can absorb the active ingredient. It may take 2-4 weeks for the active ingredient to move from the root zone to the areas of the trees affected. If insect pests are already present and causing damage, consider making a foliar application and then following up with a drench application.

For ornamental trees and shrubs adjacent to turf areas, use 0.06 - 0.12 fl oz **KYLIX SC Lawn Care** per inch of cumulative trunk diameter (DBH) of the plant(s). For hedges six feet or less in height use 0.6 fl oz **KYLIX SC Lawn Care** per 100 linear feet of hedge. For hedges greater than six feet in height, use an additional 0.15 fl oz per 100 linear feet of hedge for each foot of hedge height over six feet. Before application remove any plastic or other barriers that may prevent the insecticide from reaching the root zone of the plants.

For Bronze Birch Borers, Flatheaded Borers and other borers (excluding Emerald Ash Borer and Asian Longhorned Beetles): Use a drench treatment around the trunk of the trees when the trees are actively growing. Mix 0.5 fl oz **KYLIX SC Lawn Care SC** in 10 gallons of water and apply 2 qt of the mixture per 1 inch DBH in a 1-2 ft area around the trunks of the trees.

For outdoor ornamentals, do not apply more than a total of 23.3 fl oz **KYLIX SC Lawn Care** (0.4 lb of clothianidin active ingredient, 0.2 lb bifenthrin) per acre per year.

NUT TREES and Ornamental Trees (including Non-bearing Fruit Trees) In Landscapes that are NOT for Food Use

Apply **KYLIX SC Lawn Care** when threshold populations are observed. Do not wait until large insect populations have been established. For foliar applications, apply **KYLIX SC Lawn Care** in sufficient water to ensure uniform and thorough coverage of foliage. Thorough coverage is required for listed insect control. To achieve optimum coverage, use proper spray pressures, nozzles, nozzle spacing and water volume per acre. For soil applications, apply as a soil surface spray to the entire root zone or as a banded spray to the tree well area.

APPLICATION RATES (POST-BLOOM APPLICATION)		
PEST	APPLICATION RATE	
Aphids Adelgids Leafhoppers Plum curculio Apple maggot Codling moth ¹ Oriental fruit moth ¹ Spotted tentiform leafminer ² Western tentiform leafminer ² Obliquebanded leafrollers ² Whiteflies (suppression) ²	<u>Soil application: For trees under 6 feet tall</u> .35 - .54 fl oz per 1,000 sq ft	Foliar Application 8.0 -15.5 fl oz per 100 gallons*
	<u>Soil Application: For Trees over 6 feet tall</u> .54 -1.25 fl oz per 1,000 sq ft	

* The amount of **KYLIX SC Lawn Care** per acre will depend on tree size, pest pressure and the volume of foliage present.
¹ To control Codling moth and Oriental fruit moth, apply **KYLIX SC Lawn Care** in a program of rotational sprays with other labeled insecticides. When using the foliar application rates of 8.0- 15.5 fl oz per 100 gallons of **KYLIX SC Lawn Care** to control Codling moth and/or Oriental fruit moth, do not apply treatments less than a 14 day interval.
² For control of leafminers and leafrollers, apply as a soil application in early Spring when the soil is not frozen. A foliar application may result in suppression only and a repeat foliar application may be necessary to maintain control.

It may take 2-4 weeks for the active ingredient to move from the root zone to the areas of the trees affected. If insect pests are already present and causing damage, consider making a foliar application and then following up with a drench application.

RESTRICTIONS

- Do not consume fruit or nuts from these trees at any time.
- Do not apply more than a total of 23.3 fl oz of **KYLIX SC Lawn Care** (0.4 lb of clothianidin active ingredient, 0.2 lb bifenthrin) per acre per year.
- When using the rates for trees larger than 6 feet, apply only to the target root area as a surface spray or as a band treatment, do not apply more than 23.3 ounces per acre on the treatment site or property.
- For foliar applications, apply **KYLIX SC Lawn Care** in a full coverage spray to the point of drip. The spray volume will depend on tree size and density of canopy. Repeat applications as needed to maintain control but do not apply more often than every 10 days.
- Do not apply foliar treatments less than 10 days apart.
- Do not apply **KYLIX SC Lawn Care** through any type of irrigation system.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place away from excessive heat. Do not store near food, feed, fertilizers, seeds, or other pesticides. Keep this product out of reach of children and animals at all times. Store in original container only. Open the product container' carefully and after use, seal tightly.

To contain a spill: Cover the spilled product to prevent the material from spreading. If the product container is damaged, place the entire container along with the product in a secure receptacle and clearly label the contents.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Non-refillable Containers equal to or less than 5 gallons:

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows; Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill.

Non-refillable Containers greater than 5 gallons:

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill.

REFILLABLE CONTAINERS:

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collections system. Repeat this procedure two more times.

FOR 24-HOUR CHEMICAL EMERGENCY (Spill, Leak, Fire or Accident) ASSISTANCE: Call CHEMTREC at 1-800-424-9300.

Warranty and Disclaimer Statement

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Such risks may arise from weather conditions, soil factors, off-target movement, unconventional farming techniques, the presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of SynTelus, LLC, and can cause crop injury, injury to non-target crops or plants, ineffectiveness of the product, or other unintended consequences. All such risks shall be assumed by the user or buyer.

SynTelus warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to SynTelus, and is subject to the inherent risks described above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SYNTELUS DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SYNTELUS, MANUFACTURER, AND SELLER DISCLAIM AND SHALL NOT BE LIABLE FOR ANY SPECIAL, INCIDENTAL, INDIRECT, OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE, HANDLING, APPLICATION, STORAGE, OR DISPOSAL OF THIS PRODUCT OR FOR DAMAGES IN THE NATURE OF PENALTIES, AND THE USER AND BUYER WAIVE ANY RIGHT THAT THEY MAY HAVE TO SUCH DAMAGES. NO AGENT, REPRESENTATIVE OR EMPLOYEE OF SYNTELUS IS AUTHORIZED TO MAKE ANY WARRANTY, GUARANTEE OR REPRESENTATION BEYOND THOSE CONTAINED HEREIN OR TO MODIFY THE WARRANTIES CONTAINED HEREIN.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE TOTAL LIABILITY OF SYNTELUS, MANUFACTURER, AND SELLER, SHALL BE LIMITED TO THE

PURCHASE PRICE PAID, OR AT SYNTELUS 'S ELECTION, THE REPLACEMENT OF THE PRODUCT.