



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (7505P)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

91234-64

Date of Issuance:

3/29/18

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

A216.02

Name and Address of Registrant (include ZIP Code):

Dave G. Bolin, Ph.D.  
Atticus, LLC.  
5000 CentreGreen Way, Suite 100  
Cary, North Carolina 27513

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Hope Johnson, Product Manager 21  
Fungicide Branch, Registration Division (7505P)

Date:

3/29/18

2. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, “EPA Reg. No. 91234-64.”
3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.


If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 10/25/2017
- Alternate CSF 1 dated 10/25/2017

If you have any questions, please contact Maryam K. Muhammad by phone at 703-347-0301, or via email at [Muhammad.maryam@epa.gov](mailto:Muhammad.maryam@epa.gov).

Enclosure

**[Note to reviewer: [Text] in brackets denotes optional text].**  
**[Note to reviewer: {Text} in braces denotes where in the final label text will appear.]**  
**{BOOKLET FRONT PANEL LANGUAGE}**



METALAXYL	GROUP	4	FUNGICIDE
-----------	-------	---	-----------

# A216.02 [TM]

[Alternate Brand Name: ReCon 4 F]

**For the control of certain diseases in various crops caused by the Oomycete class of fungi.  
For use as a seed treatment for control of systemic downy mildew, Pythium seed rot, Pythium  
damping-off, and early season Phytophthora diseases of certain crops.  
For the control of certain diseases in conifers, nonbearing citrus, nonbearing deciduous fruits and  
nuts, ornamentals, and turf.  
Not for residential use.**

<b>Active Ingredient:</b>	(% by weight)
Metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl) alanine methyl ester .....	44.08%
<b>Other Ingredients:</b> .....	<u>55.92%</u>
<b>Total</b> .....	100.00%

Contains 4 lbs. of metalaxyl per gallon.

## KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside label booklet for First Aid, Precautionary Statements and Directions for Use.

**EPA Reg. No.: 91234-XX**

**EPA Est. No.:**

**Net Weight:**

**Manufactured For:**  
Atticus, LLC  
5000 CentreGreen Way, Suite 100  
Cary, NC 27513

<b>FIRST AID</b>	
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor.</li> <li>• Have a person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything to an unconscious person.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-844-685-9173 for emergency medical treatment information.</p>	

**For Chemical Emergency  
Spill, Leak, Fire, Exposure, or Accident  
Call CHEMTREC Day or Night  
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

**PRECAUTIONARY STATEMENTS  
Hazards to Humans and Domestic Animals  
CAUTION**

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves: barrier laminate or nitrile rubber ≥ 14 mils or neoprene rubber ≥ 14 mils or Viton ≥ 14 mils.
- Protective eyewear may be worn, if appropriate.

**User Safety Requirements:**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Control Statements:**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

<b>USER SAFETY RECOMMENDATIONS</b>
<p>Users should:</p> <ul style="list-style-type: none"> <li>• Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.</li> <li>• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.</li> <li>• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li> </ul>

**ENVIRONMENTAL HAZARDS:**

For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Apply only as specified on this label. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

**GROUNDWATER ADVISORY STATEMENT**

This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

**PHYSICAL/CHEMICAL HAZARDS:**

Do not mix or allow contact with oxidizing agents. Hazardous chemical reaction may occur.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Not for Residential Use.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Maximum usage when applying both metalaxyl and mefenoxam containing products to the same crop within the same season: Do not apply more than the maximum seasonal total for the active ingredient as stated on the label of the product containing the lowest seasonal total on that crop.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.** Exception: If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

**For Turf & Ornamental: Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.** Exception: If the product is soil-incorporated, or applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. There is no restricted -entry interval (REI) requirement following soil injection, soil incorporated, or a soil drench application to ornamentals.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Shoes plus socks
- Chemical-resistance gloves made of barrier laminate or nitrile rubber ≥ 14 mils or neoprene ≥ 14 mils or Viton ≥ 14 mils
- Protective eyewear (goggles, face shield or safety glasses)

**Non-Agricultural Use Requirements**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. **Do not**

**enter or allow others or pets to enter the treated area until sprays have dried. For Turf & Ornamental:** Do not enter treated areas without footwear until sprays have dried. In addition to coveralls, socks and shoes, protective eyewear and chemical resistant gloves are mandatory early entry PPE.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE RESTRICTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.**

**USE INFORMATION:**

**A216.02** is a systemic fungicide for use on selected crops to control certain diseases caused by members of the Oomycete class of fungi. Other fungicides must be used to control diseases incited by other classes of fungi.

**Resistance Management Recommendations:**

For resistance management, **A216.02** is a systemic fungicide that contains a Group 4 mode of action fungicide. Any fungal population may contain individuals naturally resistant to **A216.02** and other Group 4 mode of action fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of **A216.02** or other Group 4 fungicides within a growing season sequence with different groups that control the same pathogens. Avoid application of more than 3 and consecutive sprays of **A216.02** or other fungicides in the same group in a season.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Atticus, LLC at 984-664-9804. You can also contact your pesticide distributor or university extension specialist to report resistance.

**RESTRICTIONS:**

- Not for Residential Use.
- THIS PRODUCT IS NOT TO BE USED IN FOLIAR APPLICATIONS UNLESS SPECIFIED ON THIS LABEL OTHERWISE OR IN SOLUTIONS USED TO DIP PLANTS.
- Do not make foliar applications to field grown tobacco or other crops unless specified, since this practice may encourage more rapid development of insensitivity.
- **THIS LABEL IS FOR FIELD USE ONLY AND IS NOT FOR USE ON TRANSPLANT TRAYS, GREENHOUSES, LATH HOUSES, FLOAT HOUSES, HYDROPONIC PRODUCTION, OR IN BEDDING PLANT STRUCTURES.**

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result. Where rate ranges are specified on this label, use the higher specified rate when heavy disease pressure is expected and the lower specified rate when disease pressure is expected to be light unless otherwise noted.

**ATTENTION: UNDER CONDITIONS CONDUCIVE TO EXTENDED INFECTION PERIODS, DIFFERENT FUNGICIDE PRODUCT MODES OF ACTION MAY BE NEEDED.**

Conditions conducive to extension of infection periods include: (1) a cool and wet environment for Pythium seeding disease, (2) long growing seasons, (3) a cool and humid environment for downy mildew and (4) use of susceptible varieties.

**Mixing Instructions:**

Add 1/4 – 1/2 of the required amount of water to the spray tank, add the specified amount of **A216.02** then add the rest of the water. When tank mixing other products with **A216.02**, follow the proper sequence of adding products to the spray tank. Wettable powders or water dispersible granules should be added to the water in the tank first, followed by flowable products, with emulsifiable concentrates, such as **A216.02**, added last. Provide sufficient agitation during mixing and application to maintain a uniform emulsion.

To assure the compatibility of **A216.02** with these and other products, pour the products into a small container of water in the correct proportions. After thorough mixing, let stand for five minutes. If the combination remains mixed, or can be remixed readily, the mixture is compatible.

**BEFORE TANK MIXING A216.02 WITH OTHER REGISTERED PRODUCTS FOR ANY USE ON THIS LABEL, READ THE LABEL OF THE TANK MIX PARTNER TO BE CERTAIN IT IS LABELED FOR USE ON THE PARTICULAR CROP AND THAT USE PATTERNS ARE COMPATIBLE WITH THOSE OF A216.02.** It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Application Instructions:**

Apply **A216.02** by ground or air in sufficient water or liquid fertilizer to provide uniform coverage of the soil surface. Apply in a minimum of 20 gals. per acre for ground applications and 5 gals. per acre by air. Refer to the specific crop directions for use for application directions.

**Banded Applications:**

For banded applications, the area actually is the area covered by the band, not total cropland area planted. Some row-crop directions are based on treating in-the-row and these rates generally are specified as amounts (fl. oz.) of product per certain row length (often 1,000 ft.). Others express rates as amount per treated acre which means the total area treated with the pesticide. If rates are expressed as amount per treated acre and banded applications are used, the amount of pesticide used per acre will be proportionately less. The following formula can be used to calculate the amount of **A216.02** needed per acre of crop when banded applications are made.

Calculate the amount of **A216.02** needed for band treatment by the formula:

$$\frac{\text{Band width in inches}}{\text{Row spacing in inches}} \times \text{Broadcast rate per acre} = \text{amount needed per acre of field}$$

**In-Furrow Applications:**

For in-furrow applications to cotton, soybean and other crops where the use is approved, apply **A216.02** as an in-furrow spray in 3 to 7 gallons of water per acre at planting. Mount the spray nozzle so the spray is directed into the furrow just before the seeds are covered.



**Instructions for Moving A216.02 into the Root or Seed Zone:**

To ensure maximum activity on soilborne pathogens, this product must be moved into the seed or root zone of the plant. Some crop directions specify incorporating the product to move it to the seed or root zone (preplant incorporated application, soil drenches, shank applications) while others place the fungicide into the seed or root zone (in-furrow sprays, soil injections, crown dips). For applications made to the soil surface, rainfall will move the fungicide to the seed or root zone. However, if rainfall is not expected within 24 hours after application, mechanically incorporate before planting or sprinkler-irrigate after planting with 1/2 to 1 inch of water.

**Application Through Irrigation Systems:**

**A216.02** alone or in combination with other pesticides which are registered for application through irrigation systems, may be applied through irrigation systems. Apply this product only through center pivot, solid set, hand move, moving wheel, micro-sprinkler or drip irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety device for the public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

**Dilute A216.02 with water on a 1/10 basis prior to injection into an irrigation system. Proper tank-mix agitation is required during this mixing procedure.**

**Operating Instructions:**

1. The system must contain a functional check valve, vacuum relief valve, and low- pressure drain appropriately located on the irrigation pipeline to prevent water- source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick- closing valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended.

**Application Instructions:**

**A216.02** must be applied on the schedule specified in the specific crop use directions, not according to the irrigation schedule.

With the exception of avocados and citrus, **A216.02** has not been sufficiently tested when applied through irrigation systems to assure consistent product performance for all labeled uses. The following calibration and application techniques are provided for user reference, but do not constitute a warranty of fitness for application through sprinkler or drip irrigation equipment. Users must check the state and local regulatory agencies for potential use directions before applying any agricultural chemical through sprinkler or drip irrigation equipment.



**NOTE:** Do not inject **A216.02** at full strength or deterioration of valves and seals may occur. Use a dilution ratio of at least 10 parts water to 1 part **A216.02** in the mix tank. **A216.02** is corrosive to many seal materials. Leather seals are best. EPDM or silicone rubber seals can be used but should be replaced once a year. Do not use Viton, Buna-N, Neoprene or PVC seals.

#### **Center Pivot Irrigation Equipment:**

**NOTE:** (1) Use only with drive systems which provide uniform water distribution. (2) Do not use end guns when chemigating **A216.02** through center pivot irrigation systems because of non-uniform application. (3) Plug the first nozzle closest to well head to protect water source.

1. Determine the size of the area to be treated.
2. Determine the time required to apply ½ to 1 inch of water over the area to be treated when the system and injection equipment are operated at normal pressures recommended by the equipment manufacturer. Run the system at 80 to 95% of the manufacturer's rated capacity.
3. Using water, determine the injection pump output when operated at normal line pressure.
4. Determine the amount of **A216.02** required to treat the area covered by the irrigation system.
5. Add the required amount of **A216.02** and sufficient water to meet the injection time requirements to the solution tank.
6. Make sure the system is fully charged with water before starting injection of **A216.02** solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
7. Maintain constant solution tank agitation during the injection period.
8. Continue to operate the system until the **A216.02** solution has cleared the sprinkler head.

#### **Solid Set, Hand Move and Moving Wheel Irrigation Equipment:**

1. Determine the acreage covered by the sprinklers.
2. Fill injector solution tank with water and adjust flow rate to use the contents over 20 to 30-minute interval.
3. Determine the amount of **A216.02** required to treat the area covered by the irrigation system.
4. Add the required amount of **A216.02** into the same quantity of water used to calibrate the injection period.
5. Operate the system at the same pressure and time interval established during the calibration.
6. Inject **A216.02** at the end of the irrigation cycle in ½ to 1 inch of water or as a separate application to maximize the effectiveness of the fungicide.
7. Stop injection equipment after treatment is completed. Continue to operate the system until the **A216.02** solution has cleared the last sprinkler head.

#### **Micro Sprinkler or Drip Irrigation Systems Use**

##### **Instructions:**

1. Each run of the irrigation system must be calibrated separately to determine the time it takes water to move through the system and to make sure all emitters in the system are putting out the same amount of water.
2. Only pressure injection or venturi equipment is recommended.
3. Determine the area to be treated in each irrigation run.
4. Measure the output of each of the emitter or drip tubes closest to and farthest from the injector site.
5. For calibration, substitute a concentrated detergent (such as Wisk) or a soluble fertilizer for the **A216.02** in the injector tank. The detergent will bubble as it leaves the emitters. The time period over which bubbles occur should be checked for both the closest and farthest emitters. If these times are not within 2 minutes of each other, adjust the dilution ratio and/or the injector rate.
6. If a soluble fertilizer is used, measure the time intervals with a salt bridge. If a drip system is being calibrated, substitute soluble fertilizer for the **A216.02** in the injector and measure the time intervals with a salt bridge.

**Step-by-Step Instructions:**

1. Before starting to calibrate, operate the system until all the emitters are putting out at equal flow rates or until the system is operating at full pressure.
2. Make up an indicator solution of detergent or fertilizer, using the same ratio to be used when mixing **A216.02**.
3. Set the injector to apply the indicator solution at the injection rate to be used in the actual **A216.02** application.
4. Attach a 5-inch length of flexible tubing over the emitter closest to the injection point, another length over the emitter farthest away. Both emitters should be monitored to determine the time intervals that the indicator solutions are observed.
5. Begin injecting the indicator solution. Direct the flow from the tubes at the emitters into a small container. Begin timing when the indicator solution is first detected, stop timing when the indicator solutions are no longer detected.
6. If the period of detection of the indicator solution between the two emitters are within two minutes of each other, comparable coverage will be obtained. If they are not, make adjustments by increasing the dilution ratio, using more water per part of **A216.02**, or adjust the injector to a slower flow rate.
7. Once the system is calibrated, dilute the needed amount of **A216.02** with water using a minimum of 10 parts water to 1 part **A216.02**.
8. Do not begin to inject **A216.02** into the system until all emitters are producing equal flow rates, or until the system is at full pressure.
9. Inject the **A216.02** into the system at the end of the irrigation set in ½ to 1 inch of irrigation water.

**CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS:**

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days of the year.

Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water systems should be discharged into a reservoir tank prior to pesticide introduction.

There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed solenoid- operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where the pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

## CROPS

Use on the crops listed below is for field use only and is not for use on transplant trays, dips, greenhouses, lath houses, float houses, hydroponic production, or in bedding plant structures. Soil surface sprays of this product will not be effective until the fungicide is moved into the root zone by rainfall or irrigation. For banded applications, use the formula in the “Use Instructions” section of this label to calculate the amount of this product needed per acre for the band width actually used.

- Equivalents: 1 quart = 2 pints.

## ALFALFA\*

\* birdsfoot trefoil

**A216.02** applied to the soil at planting will provide control of damping-off caused by *Pythium* spp. and root rots caused by *Phytophthora* spp.

**Stand Establishment:**

Apply 0.5 to 1 pt. (0.25-0.5 lb. a.i.) product per acre as a broadcast surface spray at planting in a minimum of 20 gals. of water. If inter-seeding alfalfa into existing stands for renovation, apply 0.5 pt. (0.25 lb. a.i.) product per acre as a broadcast surface spray at planting in a minimum of 20 gals. of water. If alfalfa seed was previously treated with a metalaxyl or mefenoxam seed treatment product, use an application rate of **A216.02** at 0.5 pt. (0.25 lb. a.i.) per acre at planting. Use the higher specified rate (1 pt. (0.5 lb. a.i.) per acre) in areas where disease pressure is expected to be heavy unless the alfalfa product was previously treated with a metalaxyl or mefenoxam seed treatment product.

**RESTRICTIONS:** Do not feed green forage or cut for 60 days following application.

## APPLES (Bearing and Nonbearing Trees)

Use of **A216.02** will aid in the control of crown, collar and root rot caused by *Phytophthora* spp. when used in conjunction with good cultural practices and rootstocks that are tolerant to the disease. **A216.02** applications should be made before symptoms appear, especially in areas of the orchard favorable for disease development. **A216.02** will not revitalize trees showing moderate to severe disease symptoms.

**Broadcast Spray or Banded Applications:**

Apply 1 gal. (4 lbs. a.i.) product per treated acre as a broadcast spray (3 fl. oz. (0.094 lbs. a.i.) per 1,000 sq. ft.) in sufficient water to obtain thorough coverage. The treated area is based on the area under the tree canopy or the area of the sprayed row. Soil surface sprays of **A216.02** will not be effective until the fungicide is moved into the root zone by rainfall or irrigation. Applications should be made in early spring before growth starts and in the fall after harvest but before the ground freezes.

**Drench:**

Mix 0.5 qt. (0.5 lbs. a.i.) of **A216.02** with 100 gals. of water. Apply the amount of diluted mixture indicated in the table below around the trunk of each tree. Applications should be made in early spring before growth starts and in the fall after harvest but before the ground freezes. On new plantings, delay the first application until 2 weeks after planting.

To determine trunk diameter, measure the trunk 12 inches above the soil line.

Trunk Diameter 12 Inches Above Soil Line	Pints of Diluted Mixture per Tree
< 1 inch	2
1 to 3 inches	4
3 to 5 inches	6
> 5 inches	8

**RESTRICTIONS:**

1. Do not dip roots of trees in or spray bare roots with solutions containing **A216.02**.
2. Do not graze or feed cover crops in treated orchards.

**ASPARAGUS**

**A216.02** will control crown rot and spear rot caused by *Phytophthora* spp.

Apply 1 qt. (1 lb. a.i.) product per acre as a broadcast spray in a minimum of 10 gals. of water over the beds. **Soil surface sprays of this product will not be effective until the fungicide is moved into the root zone by rainfall or irrigation.**

**Cutting Beds:** Apply 30 to 60 days before the first cutting. For additional control, make another application just before the beginning of harvest.

**New Plantings:** Apply after planting seedlings or after covering one-year old crowns.

**RESTRICTIONS:** Do not apply **A216.02** within 1 day of harvest.

**AVOCADOS**

Root Rot – *Phytophthora cinnamomi*

Begin applications at the start of the growing season or at transplanting or when *Phytophthora* spp. is detected in soils. Two additional applications may be made at three-month intervals. **A216.02** may be applied as a sleeve drench at the time of transplanting, as a soil surface spray under sprinkler irrigation systems, as a directed spray under drip emitters or injected into the irrigation water.

**Sleeve Drench:**

Mix 0.5 fl. oz. (0.016 lb. a.i.) of **A216.02** with 18 gals. of water. At the time of transplanting, drench the roots inside the sleeve with 1 qt. (1 lb. a.i.) of **A216.02** solution per tree.

**NOTE:** The sleeve drench will not replace the soil surface applications for long-term control of root rot.

**Soil Spray:**

Apply to the soil under the drip emitter. Irrigate to incorporate residues into the soil. If there is more than one emitter per tree, distribute the total amount of product needed among the emitters.

**Sprinkler Irrigation:**

Apply as a soil surface spray under the canopy of the tree in sufficient water to obtain uniform coverage. See the following table for the amount of **A216.02** to use based on the diameter of the tree canopy. Start applications at the beginning of the growing season or at transplanting and continue at three-month intervals.

**Drip Irrigation:**

Apply the specified amount of **A216.02** (see table) to the soil directly under the drip emitter at each tree. If there is more than one emitter per tree, distribute the total amount of **A216.02** needed among the emitters.

**Injection into Irrigation Water (Sprinkler or Drip Irrigation Only):**

Inject **A216.02** into the irrigation water at a rate of 1 to 2 fl. oz. (0.031-0.063 lb. a.i.) product per 1,000 gals. (3  $\frac{3}{4}$  to 7  $\frac{1}{2}$  ppm active ingredient) at each irrigation. If **A216.02** is not applied at each irrigation, use the table below to determine how much **A216.02** should be injected into the irrigation water. If **A216.02** is to be used more frequently than every 3 months, adjust the rates so that no more than the specified amount is applied during each 3-month period. See the Use Information section of this label for further instructions and precautions when making applications through irrigation systems.

Diameter of Tree Canopy (Ft.)	Amount of A216.02 per Ten Trees Per 3 Months
2	0.25 to 0.5 fl. oz (0.008-0.016 lbs. a.i.)
5	1.5 to 3.0 fl. oz (0.047-0.094 lbs. a.i.)
10	6.5 to 13 fl. oz (0.203-0.41 lbs. a.i.)
15 or wider	14.5 to 29 fl. oz (0.45-0.91 lbs. a.i.)

**NOTE:**

1. For best results, use **A216.02** as soon as soil tests indicate the presence of Phytophthora.
2. For new plantings, the use of Phytophthora-resistant rot stocks with **A216.02** is recommended. Mature trees in moderate to advanced stages of decline cannot be cured with **A216.02**.

**RESTRICTIONS:** Do not apply more than 3 gals. (12 lbs. a.i.) per acre of **A216.02** per year. Do not make an application within 28 days of harvest. Do not apply during the winter months of November through February.

## BERRIES

### BLUEBERRIES

Use of **A216.02** will aid in the control of root rot caused by Phytophthora spp. when used in conjunction with good cultural practices to minimize disease problems. **A216.02** will not revitalize plants showing moderate to severe root rot symptoms.

#### Established Plantings:

Apply 0.5 pt. (0.25 lb. a.i.) product per 1,000 linear ft. of row (7.25 pts. (3.625 lb. a.i.) product per acre broadcast basis) in a three-foot band over the row before the plants start growth in the spring. One additional application may be made to coincide with periods most favorable for root rot development. **Soil surface sprays of this product will not be effective until the fungicide is moved into the root zone by rainfall or irrigation.**

#### New Plantings:

Apply 1 gal. (4 lbs. a.i.) product per acre broadcast to the soil at or after the time of planting. One or two additional applications should be made to coincide with periods most favorable for root rot development. For banded applications, an 18-inch band over the row is recommended. Use the formula in the **Use Information** section of the label to calculate the amount needed per acre. **Soil surface sprays of this product will not be effective until the fungicide is moved into the root zone by rainfall or irrigation.**

**RESTRICTIONS:** On new plantings, do not apply more than 1.8 gals. (7.2 lbs. a.i.) product per acre broadcast during the 12 months before bearing harvestable fruit.

### CRANBERRIES

Use **A216.02** as a soil application for control of Phytophthora root rot of cranberries caused by Phytophthora spp.

Apply **A216.02** at 2 to 3.5 pts. (1-1.75 lbs. a.i.) product per acre as a broadcast soil application for control of Phytophthora root rot of cranberries. Make the first application in the fall after harvest. Make the second application in the spring followed by a third application up to but no later than 45 days before harvest.

Apply **A216.02** using ground or chemigation equipment. Sufficient water should be used to allow movement of **A216.02** into the root zone. Use a minimum of 20 gals. of water per acre when applying by ground equipment. Refer to the Application Through Irrigation Systems section of the label for instructions and precautions when making applications through irrigation systems.

#### RESTRICTIONS:

1. Do not apply **A216.02** to cranberries by air.
2. Do not apply within 45 days before harvest.

3. Do not apply more than 10.5 pts. (5.25 lbs. a.i.) product per acre per crop season.
4. Do not exceed 3 applications per crop season.

### **RASPBERRIES**

**A216.02** is a soil-applied systemic fungicide for use in the control of *Phytophthora* root rot.

Apply 0.5 pt. (0.25 lb. a.i.) product per 1,000 linear ft. of row to the soil surface in a three-foot band over the row. Make 1 application in the spring and another in the fall after harvest. Use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per acre.

**RESTRICTIONS:** Do not apply **A216.02** within 45 days before harvest.

### **STRAWBERRIES**

**A216.02** provides control of red stele (*Phytophthora fragariae*), vascular collapse (*P. cactorum*) and leather rot (*P. cactorum*) when used as directed.

Applications may be made using ground application equipment or through drip irrigation systems.

#### **New Plantings:**

Apply **A216.02** at 1 qt (1 lb. a.i.) per treated acre in sufficient water to move the fungicide into the root zone of the plants. Make one application after transplanting followed by an additional application 30 days before the beginning of harvest or at fruit set. A third application may be made during the harvest season, depending on environmental conditions and disease pressure. For banded applications, use the formula in the Use Information section of this label to determine the amount of **A216.02** needed per acre. When applying **A216.02** through drip irrigation systems, use the same amount as would be applied in a banded application to cover the root zone of the plants. Observe the precautions and restrictions concerning application of **A216.02** through irrigation systems in the Use Information section of this label.

#### **Established Plantings:**

Apply **A216.02** at 1 qt. (1 lb. a.i.) per treated acre in sufficient water to move the fungicide into the root zone of the plants. Make one application in the spring after the ground thaws and before first bloom. A second application may be applied after harvest in the fall. For supplemental control of leather rot, an application may be made during the growing season at fruit set. For banded applications, use the formula in the Use Information section of this label to determine the amount of **A216.02** needed per acre. When applying **A216.02** through drip irrigation systems, use the same amount as would be applied in a banded application to cover the root zone of the plants. Observe the precautions and restrictions concerning application of **A216.02** through drip irrigation systems in the Use Information section of this label.

#### **RESTRICTIONS:**

1. Do not apply more than a total of 3 qts. (3 lbs. a.i.) of **A216.02** per treated acre per year.
2. For low annual rainfall areas, a surface application of **A216.02** needs to be moved into the root zone by rainfall, overhead irrigation, or mechanical incorporation.

## **CITRUS**

**Calamodin, chironja, grapefruits, lemons, limes, oranges, tangelos, tangerines, citrus citron, kumquats, mandarin orange, satsuma mandarin, pummelo, tangor and hybrids of these.**

Use **A216.02** on citrus for control of brown rot, citrus foot rot, root rot, trunk cankers and gummosis caused by *Phytophthora* spp. **A216.02** can be applied as a topical canker application and as a soil application, as a spray or through sprinkler or drip irrigation systems. If trees are on a drip irrigation system, distribute the amount of **A216.02** needed per tree (see tables) to the soil directly under the drip emitters at each tree. If there is more than one emitter per tree, distribute the total amount of **A216.02** needed among the emitters.



**NOTE:** Where nematodes are a problem, best results can be achieved if effective EPA- registered nematicides are used. Nematicides can be used in combination or in sequence with **A216.02** applications.

**Use Precaution:**

For best Phytophthora control, a combination of cultural practices and resistant varieties is recommended. The use of **A216.02** is not recommended in Florida for use on the highly susceptible sweet orange rootstock.

**Citrus in Nurseries (Arizona, California, Florida and Puerto Rico Only):**

Make the first application of **A216.02** at the time of planting. Make repeat applications at three-month intervals during the period when trees are actively growing. For banded applications, use a band wide enough to cover the root systems of the plants.

**Soil Drench:**

Apply 2 to 3 fl. oz. (0.063-0.094 lbs. a.i.) product per 100 gals. of water as a drench over the row at a rate of 100-250 gals. per 1,000 feet of row. The width of the drench treatment should be wide enough to cover the root systems of the plants. Follow with ½ to 1 inch irrigation over the treated area.

**Soil Surface Spray:**

Apply 0.5 to 1 gals. (2-4 lbs. a.i.) product per treated acre in a broadcast or banded surface spray to seedbeds, liners, or bedded stock in sufficient water to obtain thorough coverage. If applications are banded, the treated area should be wide enough to cover the root systems of the plants. Follow applications with a ½ to 1 inch irrigation over the treated area.

**RESTRICTIONS:** Do not use **A216.02** for disease control in greenhouse nurseries. Do not apply **A216.02** solutions to bare roots.

**Citrus Resets or New Plantings (Arizona, California, Florida and Puerto Rico Only):**

Make the first application of **A216.02** to citrus resets or new plantings at the time of transplanting. Make up to three additional applications per year at three-month intervals or when root growth flushes occur.

**Water Ring Drench:**

Mix 2 to 3 fl. oz. (0.063-0.094 lbs. a.i.) product per 100 gals. of water. Apply 5 gals. of the mix around the base of each tree within the watering ring.

**Soil Surface Spray (Arizona and California Only):**

Apply 0.5 to 1 gals. (2-4 lbs. a.i.) product per treated acre (1.5 to 3 fl. oz. (0.047-0.094 lb. a.i.) per 1,000 sq. ft.) in sufficient water to obtain uniform coverage of the soil surface. Apply spray to the soil surface beneath the tree canopy or apply through irrigation water. If natural rainfall is not expected within three days of a soil surface application, irrigate with ½ to 1 inch water over the treated area. See instructions below for application through irrigation water.

**Soil Surface Spray (Florida and Puerto Rico Only):**

Apply 0.5 gal. (2 lbs. a.i.) product per treated acre (1.5 fl. oz. (0.047 lbs. a.i.) per 1,000 sq. ft.) under the canopy of the tree. Applications may be made through low volume irrigation systems at the rate of 0.5 qt. per grove acre for trees less than 5 years old. Two to three applications per year are recommended. Applications may be made on a spring + summer, summer + fall or spring + summer + fall schedule. Restrictions-do not exceed 3 applications per year.



## Established Plantings

### Soil Application (Florida and Puerto Rico Only):

Apply 0.25 gal. (1 lb. a.i.) product per treated acre to groves that have a Phytophthora propagule count of 10 to 20 per cubic centimeter (cc) of soil as a feeder root rot disease maintenance treatment. Applications may be made through low volume irrigation for trees 5 years or older at the rate of 0.5 qt. (0.5 lb. a.i.) per grove acre. Two to three applications per year are recommended. Restriction: Do not exceed 3 applications per year. Applications may be made on a spring + summer, summer + fall or spring + summer + fall schedule. **For groves with extremely high propagule counts (above 20 per cc of soil), apply 0.5 gal. (2 lbs. a.i.) per treated acre for one year (2 to 3 applications) to reduce the population.**

### Soil Surface Spray (Arizona and California Only):

For best results, begin **A216.02** applications during the spring root-flush period. One or two additional applications per year can be made at three-month intervals or to coincide with flushes of root growth. Restrictions: Do not exceed 3 applications per year. Use the following table to determine the proper rate based on tree size and the number of applications per year. For applications based on broadcast rates, use **A216.02** at 0.5 to 1 gals. (2-4 lbs. a.i.) product per acre (1.5 to 3 fl. oz. (0.047-0.094 lbs. a.i.) per 1,000 sq. ft.) when three applications are planned and at 1.5 gals. (6 lbs. a.i.) per acre (4.5 fl. oz. (0.141 lbs. a.i.) per 1,000 sq. ft.) when two applications are planned. Apply in sufficient water to provide uniform coverage or apply through irrigation water. See instructions below for application through irrigation water.

Diameter of Tree Canopy (Ft.)	Fl. Oz. of A216.02 per Ten Trees	
	2 Applications per Year	3 Applications per Year
5	0.75 (0.023 lbs. a.i.)	0.5 (0.016 lbs. a.i.)
10	3.75 (0.117 lbs. a.i.)	2.5 (0.078 lbs. a.i.)
15	7 (0.219 lbs. a.i.)	5 (0.156 lbs. a.i.)
20	15 (0.469 lbs. a.i.)	10 (0.313 lbs. a.i.)

### Trunk Spray for Control of Gummosis Caused by Phytophthora spp. (Arizona, California and Texas Only):

Add 0.5 gal. (2 lbs. a.i.) of **A216.02** to 3 gallons of water and spray the surface of the trunks using enough spray to thoroughly wet the cankers. In Florida, add 0.5 gal. (2 lbs. a.i.) of **A216.02** to 10 gals. of water and spray the surface of the trunks using enough spray to thoroughly wet the cankers. **A216.02** may be applied up to 3 times per year.

### RESTRICTIONS:

1. Do not make trunk and soil applications to the same tree in the same cropping season.
2. Do not apply more than 3 gals. (12 lbs. a.i.) of **A216.02** per treated acre per year.
3. Do not exceed 3 applications per year.

### Application Through Irrigation Water (Sprinkler or Drip Irrigation Only):

See comments and restrictions, precautions in the Use Information section of this label. Inject **A216.02** into the irrigation water at rates specified in the tables above.

## CLOVER

**A216.02** applied to the soil at planting will provide control of damping-off caused by Pythium spp. and root rots caused by Phytophthora spp.

### Stand Establishment:

Apply 0.5 – 1 pt. (0.25-0.5 lbs. a.i.) product per acre as a broadcast surface spray at planting in a minimum of 20 gals. of water. If seed was previously treated with a metalaxyl or mefenoxam seed treatment product, apply the rate of **A216.02** at 0.5 pt. (0.25 lbs. a.i.) per acre at planting. Use the higher specified rate (1 pt. (0.5 lbs. a.i.) per acre) in areas where disease pressure is expected to be heavy.

**Soil surface sprays of A216.02 will not be effective until the fungicide is moved into the root zone by rainfall or irrigation.**

**RESTRICTIONS:** Do not feed green forage or cut hay for 90 days following application.

## **COLE CROPS**

**Broccoli, Broccoli, Chinese (gai lon), Broccoli raab (rapini), Brussels sprouts, Cabbage, Chinese (bok choy and napa), Cauliflower, Cavalo broccoli, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens, All hybrids and varieties**

**A216.02** applied as a soil application at planting will control damping-off caused by *Pythium* spp. and basal stem rot caused by *Phytophthora* spp. Applications may be made preplant incorporated, as a soil surface spray after planting or injected into drip irrigation.

### **Preplant Incorporated Application:**

For control of basal stem rot, apply 2 to 4 pts. (1-2 lbs. a.i.) product per treated acre as a broadcast soil application in sufficient water or liquid fertilizer to provide uniform coverage and incorporate in the top 2 inches of soil. For control of *Pythium* damping-off only use 0.5 to 1 pt. (0.25-0.5 lbs. a.i.) product per acre. For banded applications, a 7-inch band is recommended. Use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per acre.

### **Surface Application:**

Apply 2 to 4 pts. (1-2 lbs. a.i.) product per treated acre at planting in sufficient water or liquid fertilizer to provide uniform coverage. For banded applications, a 7-inch band is recommended. Use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per acre. If natural rainfall is not expected before the seeds start germinating, **A216.02** should be incorporated mechanically before planting or be moved into the seed zone after planting with ½ to 1 inch sprinkler irrigation. **A216.02** may be tank mixed with a pentachloronitrobenzene (PCNB) 75WP product to control club root (*Plasmodiophora brassicae*) or wirestem/black root (*Corticium solani*).

### **Irrigation (Drip):**

Inject this product into the irrigation water at the labeled rates.

### **RESTRICTIONS:**

1. Do not use **A216.02** for disease control in greenhouse crops or field-grown vegetable bedding plants.
2. Do not dip plants in solutions containing **A216.02**, or crop injury may occur.
3. Do not use **A216.02** as a transplant water treatment.

## **COTTON**

### **Seed Rots and Seedling Diseases of Cotton Caused by *Pythium* spp.:**

Apply 0.125 to 0.25 pt. (0.031-0.125 lbs. a.i.) product per 13,000 linear feet of row (0.15 to 0.3 fl. oz. (0.005-0.009 lbs. a.i.) per 1,000 linear ft.) as an in-furrow spray in 5 to 15 gals. of water or liquid fertilizer at planting. Mount the spray nozzle so the spray is directed into the furrow over the seed just before the seeds are covered.

For control of *Pythium* and *Rhizoctonia* apply with a pentachloronitrobenzene (PCNB) 2E or 75WP product in tank mixture with **A216.02**.

## CUCURBIT VEGETABLES

**Balsam pear (bitter melon), Chinese waxgourd (Chinese preserving melon), citron melon, cucumber, gherkin, edible gourds, cantaloupe, casaba, crenshaw, honeydew melon, honey balls, mango melon, muskmelon, Persian melon, pumpkin, summer squash, winter squash, watermelon, and cucurbit hybrids only.**

**A216.02** applied at planting will provide control of damping-off and cottony leak caused by *Pythium* spp. Applications may be made preplant incorporated or as a soil surface spray after planting.

### **Preplant Incorporated Application:**

Apply 2 to 4 pts. (1-2 lbs. a.i.) product per treated acre as a broadcast soil application in sufficient water or liquid fertilizer to provide uniform coverage and incorporate in the top 2 inches of soil. For banded applications, a 7-inch band is recommended. Use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per acre.

### **Surface Application:**

Apply 2 to 4 pts. (1-2 lbs. a.i.) product per treated acre at planting in sufficient water or liquid fertilizer to provide uniform coverage. For banded applications, a 7-inch band is recommended. Use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per acre. If natural rainfall is not expected before the seeds start germinating, **A216.02** should be incorporated mechanically before planting or be moved into the seed zone after planting with ½ to 1 inch sprinkler irrigation.

### **RESTRICTIONS:**

1. Do not use **A216.02** for disease control in greenhouse or field-grown vegetable bedding plants.
2. Do not dip plants in solutions containing **A216.02**, or crop injury may occur.
3. Do not use **A216.02** as a transplant water treatment.

## **DECIDUOUS FRUITS AND NUTS IN ORCHARDS (Nonbearing)\***

\*For bearing Deciduous Fruits & Nuts, see **Directions for Use** under the **Apples** and **Stone Fruits, Walnuts and Almonds** sections of this label.

### **Phytophthora Diseases:**

Use of **A216.02** will aid in the control of crown, collar, and root rot of deciduous fruit and nut trees caused by *Phytophthora* spp. when used in conjunction with good cultural practices and rootstocks that are most tolerant to the disease. Applications should be made before symptoms appear, especially in areas favorable for disease development. **A216.02** will not revitalize trees showing moderate to severe disease symptoms.

On new plantings, make the first applications at the time of planting (See RESTRICTIONS below). Additional applications should be made at three-month intervals during the time when conditions are favorable for disease development. For established plantings, make the first application in the spring before growth starts.

Apply 1 gal. (4 lbs. a.i.) product per treated acre (3 fl. oz. (0.094 lbs. a.i.) per 1,000 sq. ft.) in sufficient water to obtain uniform coverage of the soil under the canopy of the trees. For banded applications, use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per acre. Soil surface sprays of **A216.02** will not be effective until the fungicide is moved into the root zone by rainfall or irrigation.

### **RESTRICTIONS:**

1. Do not apply to plantings that will bear harvestable fruit within 12 months of the last application.
2. Do not use **A216.02** to dip or spray tree roots. Do not concentrate it around the tree trunks.
3. Do not apply to trees under stress.

4. For intense plantings (2 to 3 times the normal planting rate), make applications on a per area basis, i.e. per acre or 1,000 sq. ft. Do not calculate the amount of **A216.02** on a per tree basis.
5. In California, do not apply **A216.02** to newly planted trees within 90 days of planting.
6. Do not apply more than 3 gals. (12 lbs. a.i.) of **A216.02** per treated acre per year.

## PEPPERS AND EGGPLANT

### Eggplant, Pepper (bell, chili, cooking pepper, pimento, sweet)

Soil applications of **A216.02** will control damping-off caused by *Pythium* spp. and crown rot caused by *Phytophthora capsici*. **A216.02** must be applied to the soil before the plants are infected with *Phytophthora* to obtain satisfactory disease control.

Apply 2 to 4 pts. (1-2 lbs. a.i.) product per treated acre at the time of planting in sufficient water (20 to 50 gals.) or liquid fertilizer to provide uniform coverage. For direct seeded peppers, apply preplant or prior to emergence. If rainfall is not expected before the plants begin growth, **A216.02** should be incorporated mechanically before planting or be moved into the root zone after planting with ½ to 1 inch of sprinkler irrigation water. For banded applications, a 12 to 16 inch band is recommended. After the initial application, two additional post-directed applications at 2 pts. (1 lb. a.i.) product per treated acre should be made at 30- day intervals. The spray should be directed at the base of the plants and cover 6 to 8 inches of soil on either side of the plants. Such applications must be incorporated mechanically or by sprinkler irrigation to move the **A216.02** into the root zone.

**A216.02** may be applied with liquid fertilizer shanked in as a band treatment to either side of the plant. Use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per acre.

Injection (drip irrigation): Inject this product into the irrigation water. Make up to 2 additional applications on a 30 day schedule following initial application at planting.

#### Use Precautions:

1. **A216.02** may cause some yellowing of the pepper leaves.
2. Plants already infected with *Phytophthora* cannot be cured with **A216.02** applications.
3. The foliar blight phase of *Phytophthora* cannot be controlled with foliar applications of **A216.02**.
4. In areas where there is a history of late *Phytophthora* infections, an application of another EPA-registered fungicide labeled for *Phytophthora* control is recommended 17 to 21 days following the last **A216.02** application.

#### RESTRICTIONS:

1. Do not apply within 7 days of harvest.
2. Do not apply more than 6 pts. (3 lbs. a.i.) of **A216.02** per acre of crop per season.
3. Do not use **A216.02** for disease control in greenhouses or field-grown vegetable bedding plants.

## GINSENG

**A216.02** applied to the soil before early growth followed by applications of a metalaxyl or mefenoxam granular product will control *Phytophthora* root rot in ginseng caused by *Phytophthora cactorum*.

Apply **A216.02** at 0.75 qts. (0.75 lbs. a.i.) product per acre as a drench in 100 to 400 gallons of water uniformly to the soil surface in the spring before the plants begin growing.

**RESTRICTIONS:** Do not exceed one application per year.

## GRASSES\*

\*Any grass, Graminae family (either green or cured), except the following. **RESTRICTION:** Do not apply to sugarcane; to any of the following that will be fed or grazed by livestock: barley, buckwheat, corn, millet (pearl or proso), oats, popcorn, rice, rye, sorghum, teosinte, triticale, wheat, or wild rice; or to any enclosed pasture grasses or grasses grown for hay or silage such as bermudagrass, bluegrass, bromegrass, or fescue.

**A216.02** applied to the soil at planting will provide control of seedling diseases caused by *Pythium* spp.

### For Stand Establishment:

Apply up to 2 pts. (1 lb. a.i.) product per acre as a broadcast surface spray at planting in a minimum of 20 gals. of water. Use 0.5 to 1 pts. (0.25-0.5 lbs. a.i.) product per acre if grass seed was previously treated with a metalaxyl or mefenoxam seed treatment product. Use the higher specified rate of 1 to 2 pts. (0.5-1 lbs. a.i.) product per acre in areas where there has been a history of *Pythium* disease. Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation.

### RESTRICTIONS:

1. Do not graze, feed green forage, or cut for hay for 60 days following application.
2. Do not apply to range grasses.

## HOPS

**A216.02** applied as a soil drench followed by foliar applications with a copper fungicide registered for use on hops will control downy mildew caused by *Pseudoperonospora humulii*.

### Soil Drench:

Apply 0.5 qt. (0.5 lbs. a.i.) of **A216.02** per acre of hops in a minimum of 20 gals. of water or liquid fertilizer to the soil surface over the crowns after pruning, but before training. Early application before shoots are 6 inches long is preferable.

### Foliar Spray:

When primary infection (spikes) persist after a soil drench treatment and/or there is the first evidence of secondary (foliar) infection, foliar sprays of **A216.02** in combination with contact copper fungicides may be used. Apply **A216.02** at 0.5 qt. (0.5 lbs. a.i.) per acre in a tank-mix combination with a copper fungicide registered for use on hops). Apply with ground equipment in a minimum of 50 gals. of water per acre.

### RESTRICTIONS:

1. Do not make more than 3 applications of **A216.02** per season (1 soil drench + 2 foliar sprays).
2. Do not make the last application within 45 days before harvest.
3. Do not apply foliar sprays of **A216.02** without a copper fungicide registered for use on hops.

## LEAFY VEGETABLES\*

\* Celery, gardengrass, upland cress, endive, fennel, lettuce (head and leafy), parsley, rhubarb, spinach, and Swiss chard.

**A216.02** applied as a soil application will control damping off caused by *Pythium* spp. in leafy vegetables (*Albugo occidentalis*) and downy mildew in spinach. Applications may be made banded over the row, preplant incorporated, or injected with liquid fertilizer.

**Preplant Incorporated Application:**

Apply 2 to 4 pts. (1-2 lbs. a.i.) product per treated acre as a broadcast soil application in sufficient water or liquid fertilizer to provide uniform coverage and incorporate in the top 2 inches of soil. For banded applications, a 7-inch band is recommended. Use the formula in the **Use Information** section of this label to calculate the amount of **A216.02** needed per acre.

**Surface Application:**

Apply 2 to 4 pts. (1-2 lbs. a.i.) product per treated acre at planting in sufficient water or liquid fertilizer to provide uniform coverage. For banded applications, a 7-inch band is recommended. Use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per acre. If natural rainfall is not expected before the seeds start germinating, **A216.02** should be incorporated mechanically before planting or be moved into the seed zone after planting with ½ to 1 inch sprinkler irrigation.

**White Rust and Downy Mildew Control (Spinach Only):**

In addition to the preplant incorporated or surface application described above, apply **A216.02** at 0.5 pt. (0.25 lbs. a.i.) product per acre of crop, shanked in 21 days after planting or after the first cutting. One other application may be shanked in after the next cutting. A total of 2 additional applications may be used on a 21-day interval. Use sufficient mechanical or bypass agitation to keep the **A216.02** mixed with the water or fertilizer.

**NOTE:** The additional applications of **A216.02** noted above and made after each cutting by shanking the fungicide into the beds along with liquid fertilizer provide continuing control of white rust. However, white rust can only be controlled in a preventative disease control program that begins with an application of **A216.02** to the soil at planting. The use of **A216.02** in curative applications greatly increases the risk of the fungus developing insensitivity to metalaxyl. The development of insensitivity will destroy the effectiveness of **A216.02** in controlling white rust.

**RESTRICTIONS:**

1. If **A216.02** is not used at planting, do not use **A216.02** at any other time throughout the season. Do not apply **A216.02** in foliar applications or in situations where white rust infections are already established.
2. Do not harvest spinach within 21 days of a **A216.02** application.
3. Do not use **A216.02** for disease control in greenhouse or field-grown vegetable bedding plants.
4. Do not use **A216.02** as a transplant water treatment.
5. Do not apply more than 5.5 pts. (2.75 lbs. a.i.) of **A216.02** per acre per growing season in spinach.
6. Do not exceed a total of 2.8 lbs. active ingredient per acre of metalaxyl per growing season when using a combination of **A216.02** and other metalaxyl products in spinach.

**LEGUME VEGETABLES (Succulent or Dried)**

**Field beans, French beans, kidney beans, lima beans, mung beans, navy beans, pinto beans, runner beans, snap beans, wax beans, broad beans (fava beans), chickpeas (garbanzo beans), lentils, lupines (sweet, white sweet, white and grain), garden peas, field peas, sugar peas, southern peas (blackeyed peas, crowder peas, cowpeas, catjang) and edible soybeans.**

**Pythium Damping-Off and Root Rot:**

**A216.02** applied at planting will control damping-off and root rot caused by *Pythium* spp. Applications may be made preplant incorporated, or at a soil surface spray after planting.

**Preplant Incorporated Application:**

Apply 1 to 2 pts. (0.5-1 lbs. a.i.) product per treated acre as a broadcast soil application in sufficient water or liquid fertilizer to provide uniform coverage and incorporate in the top 2 inches of soil. For banded applications, a 7-inch band is recommended. Use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per acre.



**Surface Application:**

Apply 1 to 2 pts. (0.5-1 lbs. a.i.) product per treated acre at planting in sufficient water or liquid fertilizer to provide uniform coverage. For banded applications, a 7-inch band is recommended. Use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per acre. If natural rainfall is not expected before the seeds start germinating, **A216.02** should be incorporated mechanically before planting or be moved into the seed zone after planting with ½ to 1 inch sprinkler irrigation. **A216.02** may be tank mixed with a pentachloronitrobenzene (PCNB) 2E or 75WP product to control Rhizoctonia root and stem rot or white mold (*Sclerotinia sclerotiorum*) in snap and dry beans.

**RESTRICTIONS:** Do not use **A216.02** for disease control in greenhouse or field-grown vegetable bedding plants.

**ONIONS – DRY BULBS\*, GREEN\*\*, AND ONION GROWN FOR SEED**

\*Garlic, onions (dry bulb), and shallots (dry bulb).

\*\* Green onions, leeks, spring onions or scallions, Japanese bunching onions, green shallots, or green eschalots.

**A216.02** applied at planting will control damping-off caused by Pythium spp. Applications may be made preplant incorporated or as a soil surface spray after planting.

**Preplant Incorporated Application:**

Apply 1 to 2 pts. (0.5-1 lbs. a.i.) product per treated acre as a broadcast soil application in sufficient water or liquid fertilizer to provide uniform coverage and incorporate in the top two inches of soil. For banded applications, a 7-inch band is recommended. Use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per acre.

**Surface Application:**

Apply 1 to 2 pts. (0.5-1 lbs. a.i.) product per treated acre at planting in sufficient water or liquid fertilizer to provide uniform coverage. For banded applications, a 7-inch band is recommended. Use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per acre. If natural rainfall is not expected before the seeds start germinating, **A216.02** should be incorporated mechanically before planting or be moved into the seed zone after planting with ½ to 1 inch sprinkler irrigation.

**RESTRICTIONS:** Do not use **A216.02** for disease control in greenhouse or field- grown vegetable bedding plants.

**PAPAYA (Hawaii Only)**

**A216.02** aids in the control of Phytophthora root rot of papaya in new plantings in the field.

**Papaya in the Field – New Plantings:**

Make the first application of **A216.02** at the time of transplanting to the field or within one week of transplanting. Apply 1.75 to 3.5 qts. (1.75-3.5 lbs. a.i.) product per acre\* of soil treated (1.25 to 2.5 fl. oz. (0.039-0.078 lbs. a.i.) per 1,000 sq. ft.) as a soil surface spray in sufficient water to obtain thorough coverage of the soil under the canopy of the trees. After application, immediately irrigate with 1/8 to 1/4 inch of water. Repeat in 2 to 4 weeks if conditions are favorable for disease. Use the table below as a guide for treating individual trees.

**RESTRICTIONS:**

1. Do not apply more than two applications per year in the field to newly transplanted stock.
2. Do not apply within 3 months of harvest.



Diameter of Tree Canopy (Ft.)	Fl. Oz. of A216.02 per Ten Trees
3	0.094-0.19 (0.003-0.006 lbs. a.i.)
4	0.19-0.38 (0.006-0.012 lbs. a.i.)
6	0.38-0.75 (0.012-0.023 lbs. a.i.)
8	0.63-1.25 (0.020-0.039 lbs. a.i.)

\* 1 qt. = 1 lb. a.i.

## PEANUTS

**A216.02** is a soil-applied systemic fungicide for use in control of seedling and pod diseases of peanuts incited by *Pythium* spp.

### Seedling Diseases:

Apply 0.5 pt. (0.25 lb. a.i.) product per acre of crop as an application to the seed in- furrow or in a 7-inch band at the time of planting. For the in-furrow applications, position the spray so the fungicide is mixed with the soil covering the seed. Avoid spraying the seed directly with the spray solution or crop injury may occur.

### Pod Rot:

Apply 0.5 to 1 qts. (0.5-1 lbs. a.i.) product per acre at early pod set or pegging through overhead irrigation systems. See the Use Information section of this label for further instructions and precautions when making applications through irrigation systems.

### NOTE:

1. Where pathogens other than *Pythium* spp. are present, use fungicides that control those diseases in combinations with **A216.02**.
2. Where the predominant pod rot pathogens are *Pythium* spp. and *Rhizoctonia* spp., use **A216.02** at 0.5 to 1 qts. (0.5-1 lbs. a.i.) per acre tank mixed with a pentachloronitrobenzene (PCNB) 2E product.

## PINEAPPLE

**A216.02** applied as a "seed piece" dip, provides effective control of heart rot disease of pineapple caused by *Phytophthora* spp.

Apply **A216.02** as a crown dip before planting at the rate of 0.5 to 1 qts. (0.5-1 lbs. a.i.) per 100 gals. of water. Use 75 to 100 gals. of dip solution per planted acre, depending on crown size, plant density and dipping techniques.

**RESTRICTIONS:** If there is a crop failure within one year of planting treated crowns, do not harvest plant material for animal feed.

## ROOT AND TUBER VEGETABLES\*

**Artichoke (Jerusalem), beet (sugar\*\* and table), carrot, cassava, chicory, dasheen (taro), ginger, ginseng\*\*\*, horseradish, parsnip, potato, radish, rutabaga, salsify, sweet potato, tanager, turnip, and yams.**

\* See Note at end of section.

\*\* See separate section for Sugar Beets.

\*\*\* See separate section for Ginseng.

**A216.02** applied to the soil at planting will provide control of diseases caused by *Pythium* and *Phytophthora* spp. Applications may be made preplant incorporated or as a soil surface spray after planting.

**Preplant Incorporated Application:**

Apply 2 to 4 pts. (1-2 lbs. a.i.) product per treated acre as a broadcast soil application in sufficient water or liquid fertilizer to provide uniform coverage and incorporate in the top two inches of soil. For banded applications, a 7-inch band is recommended. Use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per dose.

**Surface Application:**

Apply 2 to 4 pts. (1-2 lbs. a.i.) product per treated acre at planting in sufficient water or liquid fertilizer to provide uniform coverage. For banded applications, a 7-inch band is recommended. Use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per acre. If natural rainfall is not expected before the seeds start germinating, **A216.02** should be incorporated mechanically before planting or be moved into the seed zone after planting with ½ to 1 inch sprinkler irrigation.

**POTATOES**

**A216.02** will provide effective control of Pythium leak caused by *Pythium* spp., pink rot caused by *Phytophthora erythroseptica*, and Pythium seedling caused by *Pythium* spp.

Apply 0.84 fl. oz. (0.026 lbs. a.i.) of product per 1,000 linear ft. of row on a 6 to 8-inch band at planting in a minimum of 3 gals. of water per acre. Make application directly over the seed piece(s) prior to row closure or use markout method (incorporated).

**A216.02** may be impregnated on dry fertilizer provided the application rate of 0.84 fl. oz. (0.026 lbs. a.i.) per linear ft. of row is observed and placement is in a 6 to 8-inch band incorporated within the planted hill. **A216.02** may also be applied in combination with liquid fertilizers.

**Storage Rots:**

**A216.02** will effectively control storage rots caused by *Pythium* leak and pink rot when used in conjunction with other management practices such as crop rotation. Apply product at 6.4 fl. oz. (0.2 lbs. a.i.) per acre at flowering and repeat with a second application 14 days later. If the field has a history of storage rot problems, make a third application 14 days after the second application. If conditions favor the development of foliar diseases, use **A216.02** in tank mixtures with a companion fungicide such as mancozeb, chlorothalonil, or other approved products.

**RESTRICTIONS:**

1. Do not use **A216.02** for disease control in greenhouse or field-grown vegetable bedding plants.
2. Do not apply this product within 14 days of harvest.

**SOYBEANS**

**A216.02** is a soil-applied systemic fungicide for use in the control of *Phytophthora* root and stem rot and *Pythium* damping-off. **A216.02** may be applied broadcast, banded or in the seed furrow before the seeds are covered. The seed furrow applications will provide more consistent results if rain is not expected before the seeds germinate.

For best results against *Phytophthora* root and stem rot, use **A216.02** with soybean varieties that have some tolerance to the races of *Phytophthora* present in the field. The higher specified rate of **A216.02** should be used in areas with a history of heavy *Phytophthora* damage. Under heavy late season *Phytophthora* pressure, **A216.02** may not provide complete control.

**Surface Application:**

For full season control, apply 2.5 pts. (1.25 lbs. a.i.) product per treated acre in sufficient water or liquid fertilizer to provide uniform coverage at the time of planting. For banded applications, a 7-inch band is recommended. Use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per acre. For early to mid-season control, apply 0.75 to 1.5 pts. (0.375-0.75 lbs. a.i.) product per treated acre.

**In-Furrow Applications:**

For full season control, apply 0.55 fl. oz. (0.017 lbs. a.i.) product per 1,000 linear feet of row as an in-furrow spray in 5 to 10 gals. of water or liquid fertilizer at planting.

Position the spray so the fungicide is mixed with the soil covering the seed. Avoid spraying the seed directly with the spray solution or crop injury may occur. Use sufficient water or liquid fertilizer to provide uniform coverage. Use the following table to determine the amount of **A216.02** needed per acre based on row spacing. For early to mid-season control, apply 0.15 to 0.3 fl. oz. (0.005-0.009 lbs. a.i.) product per 1,000 linear feet of row.

**Rates Per Acre According to Row Spacing**

Row Spacing	Linear Ft. of Soybean Row Per Acre	Fluid Ounces of A216.02 Needed Per Acre Rate Desired		
		0.15 fl. oz. (0.005 lbs. a.i.)	0.3 fl. oz. (0.009 lbs. a.i.)	0.55 fl. oz. (0.017 lbs. a.i.)
38 in.	13,756	2	4	7.5
36 in.	14,520	2.25	4.5	8
30 in.	17,424	2.63	5.5	9.5
24 in.	21,136	3.25	6.5	12
20 in.	26,136	4	8	14

**NOTE:** **A216.02** is specific for Pythium and Phytophthora and will not control other diseases that may attack soybeans.

### STONE FRUITS\*, WALNUTS, AND ALMONDS

**\*Apricots, cherries (sweet, tart), nectarines, peaches, plums (Chickasaw, Damson, Japanese), plumcot, prunes (fresh) and hybrids or cultivars of these.**

Use of **A216.02** will aid in the control of crown, collar and root rot caused by Phytophthora spp. when used in conjunction with good cultural practices and rootstocks that are most tolerant to the disease. **A216.02** applications should be made before symptoms appear, especially in areas favorable for disease development. **A216.02** will not revitalize trees showing moderate to severe disease symptoms.

On new plantings, make the first application of **A216.02** 2 weeks after planting. Additional applications should be made at 2 to 3 months intervals or during periods most favorable for root, crown or collar rot development.

For established plantings, the application should be made in the spring before plants start growth. Additional applications should be made at 2 to 3 months intervals or to coincide with periods most favorable for root, crown or collar rot development.

Apply 1 gals. (4 lbs. a.i.) product per treated acre (3 fl. oz. (0.094 lbs. a.i.) per 1,000 sq. ft.) in sufficient carrier to obtain thorough coverage of the soil under the canopy of the trees. Sufficient surface area should be treated in nurseries to cover the root zone on the plants. Up to 3 applications can be made per year. For banded applications, use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per acre. Soil surface sprays of **A216.02** will not be effective until the fungicide is moved into the root zone by rainfall or irrigation.

For intense plantings (2 to 3 times the normal planting rate) make applications on a per area basis, i.e. per acre or 1,000 sq. ft.

**RESTRICTIONS:**

1. Do not dip the roots of trees in **A216.02** solutions, spray the roots or concentrate it around the tree trunks or injury may occur.
2. Do not apply it to trees under stress.

3. Do not calculate the amount of **A216.02** on a per tree basis.
4. In California, do not apply **A216.02** to newly planted trees within 90 days of planting.
5. Do not graze livestock in treated areas.
6. Do not graze or feed cover crops in treated orchards.

## SUGAR BEETS

**A216.02** will provide control of diseases caused by *Pythium* spp. Applications may be made preplant incorporated or as a surface spray at planting.

### Preplant Incorporated Application:

Apply 2 to 4 pts. (1-2 lbs. a.i.) product per treated acre as a broadcast soil application in sufficient water or liquid fertilizer and incorporate in the top 2 inches of soil. For banded applications, a 7-inch band is recommended. Use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per acre.

### Surface Application:

Apply 2 to 4 pts. (1-2 lbs. a.i.) product per treated acre at planting in sufficient water or liquid fertilizer to provide uniform coverage. For banded applications, a 7-inch band is recommended. Use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per acre. If natural rainfall is not expected before the seeds begin germinating, **A216.02** should be incorporated mechanically before planting or be moved into the seed zone after planting with ½ to 1 inch sprinkler irrigation.

## TOBACCO

**A216.02** is a soil-applied systemic fungicide for use in the field before transplanting for control of damping-off (*Pythium* spp.), black shank (*Phytophthora parasitica* var. *Nicotianae*) and blue mold (*Peronospora tabacina*) on all types of tobacco. For control of anthracnose and other tobacco diseases, use approved fungicides to control those diseases.

### RESTRICTIONS:

1. Do not use Metalaxyl 4lb SC for disease control in greenhouse crops or tobacco plant beds.
2. Do not dip plants in solutions containing Metalaxyl 4lb SC, or crop injury may occur.
3. Do not use Metalaxyl 4lb SC for disease control in floathouse, floatbed production facilities, hydroponic production or greenhouse facility.
4. Do not use Metalaxyl 4lb SC as a transplant water treatment.
5. Do not use Metalaxyl 4lb SC as a foliar spray to field planted tobacco.

### FIELD PLANTED TOBACCO

#### Damping-off:

Apply **A216.02** as a preplant soil application before or at time of transplanting. Apply 0.5 to 1 qts. (0.5-1 lbs. a.i.) product in 50 gallons of water (2 gallons water per 150 square yards). Use the higher specified application rate on broadcast tobacco.

#### Blue Mold:

Apply **A216.02** as a broadcast soil application prior to transplanting and incorporate in the top 2 to 4 inches of soil. For flue-cured tobacco, use 0.5 to 1 qt. (0.5-1 lbs. a.i.) product per treated acre, depending on disease pressure and length of control desired. Under low disease pressure or for early season control, use 0.5 qt. (0.5 lbs. a.i.) product per treated acre. For burley and other tobacco types, use 1 qt. (1 lb. a.i.) product per treated acre.

For prolonged control of blue mold in field planted tobacco, make an additional application of 0.5 qt. (0.5 lbs. a.i.) product per acre of crop as a soil application at lay-by or the last cultivation. Position the nozzles so that the spray is deposited under the plants and is covered by soil by the cultivator.

**Black Shank:**

Use **A216.02** as a broadcast soil application prior to transplanting and incorporate in the top 2 to 4 inches of soil. Apply **A216.02** using conventional ground application equipment in sufficient water or fertilizer to provide uniform coverage. Use the following table to determine the amount of **A216.02** needed per acre depending on the black shank severity.

Type of Tobacco	Disease Level in Field	Rate of A216.02 per Acre
Flue-Cured	Low to moderate (less than 6% disease)	1 qt. (1 lb. a.i.)
	High (more than 6% disease)	2 qts. (2 lbs. a.i.)*
Burley and Other**	Low to moderate (less than 6% disease)	2 qts. (2 lbs. a.i.)
	High (more than 6% disease)	3 qts. (3 lbs. a.i.)

\*Florida and Georgia – Use 3 qts. (3 lbs. a.i.) per treated acre of **A216.02** in fields with heavy black shank levels (greater than 6%).

\*\* Pennsylvania – Do not use **A216.02** for black shank control.

For prolonged control of black shank in field planted tobacco, one of the following is recommended: (1) Make a preplant incorporated and an additional lay-by application (last cultivation). Apply the second application at last cultivation at the rate of 0.5 to 1 qts. (0.5-1 lbs. a.i.) product per acre as a soil treatment. Position the nozzles so that the spray is deposited under the plants and is covered with soil by the cultivator. Apply **A216.02** at 1 qt. (1 lb. a.i.) per acre just prior to transplanting followed by a second application of 1 qt. (1 lb. a.i.) per acre at the first cultivation followed by a third application of 1 qt. (1 lb. a.i.) per acre at lay-by or the last cultivation.

**Resistance Management:**

1. For best results against black shank, use **A216.02** with tobacco varieties that have high resistance to black shank and use crop rotation. In fields where there is a history of severe black shank incidence, use the highest rate and plant variety that is resistant to the race of *Phytophthora* present in the field. (Burley L8 hybrids are only resistant to *Phytophthora* Race O.)
2. **A216.02** is not recommended for use in high black shank areas on highly susceptible flue-cured varieties.
3. Failure to adequately control nematodes in fields treated with **A216.02** may result in poor control of black shank.

**No-Till Tobacco:**

For black shank and blue mold on all types of tobacco, apply **A216.02** to the field before transplanting and incorporate in the top 2 to 4 inches of soil. Apply 0.5 to 1 qts. (0.5-1 lbs. a.i.) product per treated acre as a preplant, broadcast or banded soil application. For banded applications, use the formula in the Use Information section of this label to calculate the amount of **A216.02** needed per acre. A lay-by soil application may be made 30 to 35 days after planting at 0.5 qt. (0.5 lb. a.i.) product per acre.

**RESTRICTIONS:**

1. Do not make this application if more than 1 qt. (1 lb. a.i.) per acre of **A216.02** were applied prior to transplanting or if no **A216.02** was applied prior to transplanting.
2. Do not make this application if more than 1 qt. (1 lb. a.i.) per acre of **A216.02** was applied at transplanting.
3. Make a preplant incorporated plus 2 additional soil applications at first cultivation and last cultivation (lay-by).
4. Do not make the lay-by application if more than 1 qt. (1 lb. a.i.) per acre of **A216.02** were applied at transplanting or if no **A216.02** was applied at transplanting.

**TOMATOES**

Soil applications of **A216.02** at planting will provide control of damping-off caused by *Pythium* spp. Soil applications applied 4 to 12 weeks before harvest under the vines will control fruit and root rot caused by *Pythium* spp. and *Phytophthora* spp.

**Damping-Off (*Pythium* spp.):**

Apply 1 to 2 qts. (1-2 lbs. a.i.) product per treated acre in sufficient water or liquid fertilizer to provide uniform coverage at the time of planting. If rainfall is not expected before the seeds start to germinate, **A216.02** should be incorporated mechanically before planting, during the planting operation, or be moved into the seed zone after planting with ½ to 1 inch sprinkler irrigation. For banded applications, a 7- inch band is recommended. Use the formula in the **Use Information** section of this label to calculate the amount of **A216.02** needed per acre.

**Root and Fruit Rot (*Phytophthora* spp. and *Pythium* spp.):**

To aid in the control of root and fruit rot, 1 or 2 additional applications may be made during the growing season, depending on the severity of the conditions for disease infection.

Apply 1 qt. (1 lb. a.i.) product per treated acre beginning 4 to 6 weeks after planting. A second application may be made as needed up to 4 weeks before harvest, but before the last irrigation. **A216.02** may be applied as a directed soil surface spray under the vines or it may be injected into the beds with liquid fertilizer. If less than the full bed is treated, use the formula in the Use Information section to determine the amount of **A216.02** needed per acre. If **A216.02** is injected into the beds with liquid fertilizer, base calculations on a 7-inch band.

If soil surface sprays are used, the **A216.02** must be incorporated into the soil with ½ to 1 inch of rainfall or sprinkler irrigation.

**Injection (drip irrigation):**

Initiate control of Root and Fruit rot with a soil application as described above. Make subsequent applications through drip irrigation. Make the first drip application 4 to 6 weeks after planting. Apply the second drip application as needed up to 4 weeks before harvest, but before the last irrigation.

For injection applications, base rate calculation on a 7-inch band.

**A216.02** may be applied with water or liquid fertilizer. Use the test in the Use Information section to check for compatibility with various fertilizers.

**RESTRICTIONS:**

1. Do not apply more than 3 qts. (3 lbs. a.i.) product per treated acre per season.
2. Keep **A216.02** suspended in the fertilizer solution with bypass or mechanical agitation. Refer to the Use Information section for drip irrigation instructions.
3. Do not use **A216.02** for disease control in greenhouse or field-grown vegetable bedding plants.
4. Do not use **A216.02** as a transplant water treatment.

**REPLANTING**

If replanting is necessary, an additional application of **A216.02** may be made provided that the total amount of active ingredient in **A216.02** applied does not exceed the maximum allowed for the specific crop and the maximum number of applications per season are not exceeded.

## ROTATION (PLANTBACK) RESTRICTION

Do not plant any crop which is not registered for use with the **A216.02** active ingredient in soil treated with this active ingredient for a period of 12 months, unless a shorter interval is specified on the following list.

Rotation Crops	Planting Time from Last A216.02 Application
Alfalfa (birdsfoot trefoil), Almonds, Apples, Asparagus, Avocados Blueberries Citrus, Clover, Cole Crops, Cotton, Cranberries, Cucurbit Vegetables Deciduous Fruits and Nuts* Eggplant Garlic, Ginseng, Grapes, Grasses** Hops Leafy Vegetables (Excluding Brassica), Legume Vegetables (beans and peas—succulent and dried) Onions (dry bulb, green, and seed) Papaya, Peanuts, Peppers, Pineapples, Potatoes Raspberries, Root and Tuber Vegetables Soybeans, Spinach, Stone Fruits, Strawberries, Sugar Beets Tobacco, Tomatoes	0 days
Cereal Grains (Other than Corn)	14 days
Corn	9 months
Crops <b>NOT INTENDED</b> for Food or Feed	0 days
All Other Crops <b>INTENDED</b> for Food or Feed	12 months

\* These crops and other perennial crops may be planted immediately following last application of **A216.02**, provided they will not bear harvestable fruits within 12 months.

\*\* Any grass, Gramineae family (either green or cured), except the following: Do not apply to sugarcane; to any of the following that will be fed to or grazed by livestock: barley, buckwheat, corn, millet (Pearl or proso), oats, popcorn, rice, rye, sorghum, teosinte, triticale, wheat, or wild rice; or to any enclosed pasture grasses or grasses grown for hay or silage such as bermudagrass, bluegrass, bromegrass, or fescue.

### CONIFERS, NON-BEARING CITRUS, NONBEARING DECIDUOUS FRUITS AND NUTS, ORNAMENTALS AND TURF

**A216.02** is a systemic fungicide for use on ornamentals, turf, nonbearing citrus grown in nurseries and as landscape plantings, conifers grown in nurseries and plantations, including Christmas trees and nonbearing deciduous fruit and nut trees grown in nurseries.

**Restriction:** Not for residential use.

**Resistance Management:**

**A216.02** is a systemic fungicide having a specific mode of action and could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible **A216.02** resistant strains of fungi which may occur.

To help decrease the chance of downy mildew resistance, do not use **A216.02** for the control of downy mildew diseases, except for use in turf. Use **A216.02** only as a soil application for control of soil-borne diseases with the exception of azalea petal blight.

To avoid drift, do not apply under windy conditions. Avoid spray overlap, or crop injury may result.



**Mixing Instructions:**

To assure the compatibility of **A216.02** with these and other products, pour the products into a small container of water in the correct proportions. After thorough mixing, let stand for five minutes. If the combination remains mixed, or can be remixed readily, the mixture should be considered compatible. Prepare no more spray mixture than is required for the immediate operation. Agitate the spray solution continuously during mixing and during application. Rinse the spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to already treated area.

**A216.02 Alone:**

Add 1/4 -1/2 of the required amount of water to the spray tank. With the agitator running, add the **A216.02** to the tank. Continue agitation while adding the remainder of the water. Begin application of the spray solution after the **A216.02** has completely dispersed into the mix water. Maintain agitation until all the mixture has been sprayed.

**A216.02 + Tank Mixtures:**

Add 1/4 – 1/2 of the required amount of water to the spray tank. Start the agitator before adding any tank mix partners. In general, tank mix partners should be added in this order: wettable powders, dry flowable formulations, liquid flowable formulations, microencapsulated formulations, and emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water and the **A216.02** to the spray tank. Allow the **A216.02** to completely disperse into the mix water. Maintain agitation until all of the mixture has been sprayed.

**NOTE:** When using **A216.02** in tank mixtures, all products in water-soluble packaging should be added to the tank before any other tank mix partner, including **A216.02**. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

If using **A216.02** in a tank mixture, observe all directions for use, sites, use rates, dilution ratios, precautions and limitations which appear on the tank mix partner label. No label dosage may be exceeded and the most restrictive label precautions and limitations must be followed. This product may not be mixed with any product which prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the products are registered. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Application Instructions:**

For banded applications, calculate the amount of **A216.02** needed as follows:

Bandwidth in inches X broadcast rate per acre = amount needed per  
acre row width in inches

**Application Through Irrigation Systems:**

**A216.02** alone or in tank mixture with other pesticides registered for application through irrigation systems may be applied in irrigation water at rates recommended on this label. This product may be applied through micro sprinkler or drip irrigation systems. Do not apply this product through any other type of irrigation system. Plant injury or lack of effectiveness may result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label- prescribed safety devices for public water supplies are in place. A person knowledgeable of the chemigation system and responsible for its operation shall shut the system down and make necessary adjustments should the need arise.

**A216.02** can be injected into the irrigation line in concentrated form, or diluted with water or liquid fertilizer solutions with pH levels less than 7.5. If diluted, a pesticide supply tank should be used. Agitation is not needed unless the diluted solution will remain in the supply tank more than 24 hours. **A216.02** is normally diluted at a ratio of 10:1 to 50:1, depending on injection setups. Injecting a larger volume of a more dilute mixture will usually allow a more accurate calibration of the metering equipment. Meter the fungicide into the irrigation water during the first part of the irrigation cycle.

**NOTE:** **A216.02** is highly corrosive to seals and other pump components. Recommended components are Teflon, polyethylene, polypropylene and nylon. When **A216.02** is diluted at least 50:1, silicone rubber and viton can be used. Do not use PVC or EPDM based components.

#### **Safety Devices for Irrigation Systems Connected to Public Water Supplies:**

If the source of water for your irrigation system is a public water supply, follow the instructions below.

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

#### **Safety Devices for Irrigation Systems Not Connected to a Public Water Supply:**

1. The system must contain a functional check-valve, vacuum relief valve and low- pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick- closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where the pesticide distribution is adversely affected.
6. Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

**Application Instructions:**

**A216.02** must be applied on the schedule specified in the use directions, not according to the irrigation schedule.

The following calibration and application techniques are provided for the user reference, but do not constitute a warranty of fitness for application through sprinkler irrigation equipment. Users should check with state and local regulatory agencies for potential use restrictions before applying any agricultural chemical through sprinkler irrigation equipment.

**Use Instructions:**

1. Each run of the irrigation system must be calibrated separately to determine the time it takes water to move through the system and to make sure all emitters in all systems are putting out the same amount of water.
2. Only pressure injection or venturi equipment is recommended.
3. Determine the area to be treated in each irrigation run.
4. Measure the output of each of the emitters or drip tubes closest to and farthest from the injector site.
5. For calibration, substitute a concentrated detergent (such as Wisk) for the **A216.02** in the injector tank. It is important to use the same volume of soap solution as the planned volume of **A216.02** solution when calibrating the system. The detergent will bubble as it leaves the emitters. The time period over which the bubbles occur should be checked for both the closest and farthest emitters. If these times are not within 2 minutes of each other, adjust the dilution ratio and/or the injection rate.

**Step-by-Step Instructions:**

1. Before starting to calibrate, operate the system until all the emitters are putting out at equal flow rates or until the system is operating at full pressure.
2. Make up an indicator solution of detergent or fertilizer, using the same ratio to be used with mixing **A216.02**.
3. Set the injector to apply the indicator solution at the injection rate to be used in the actual **A216.02** application.
4. Attach a 5-inch length of flexible tubing over the emitter closest to the injection point, another length over the emitter farthest away. Both emitters should be monitored to determine the time intervals that the indicator solutions are observed.
5. Begin injecting the indicator solution. Direct the flow from the tubes at the emitters into a small container. Begin timing when the indicator solution is first detected. Stop timing when the indicator solutions are no longer detected.
6. If the period of detection of the indicator solution between the 2 emitters are within 2 minutes of each other, comparable coverage will be obtained. If they are not, make adjustments by increasing the dilution ratio, using more water per part of **A216.02**, or adjust the injector to a slower flow rate.
7. Once the system is calibrated, dilute the needed amount of **A216.02** with water using a minimum of 15 parts water to 1 part of **A216.02** in the solution tank.
8. Do not begin to inject **A216.02** into the system until all emitters are producing equal flow rates, or until the system is at full pressure.
9. Inject the **A216.02** into the system at the beginning of the irrigation set in 1/2 – 1 inch of irrigation water.

## ORNAMENTALS

Use **A216.02** on container, bench, or bed grown ornamentals in greenhouses or outdoor nurseries, and for use on ornamentals grown for indoor and outdoor landscaping, for control of damping-off, and root and stem rot diseases caused by Pythium and Phytophthora. **Restriction:** Not for residential use. **A216.02** may be applied through irrigation systems, as a soil drench or as a soil surface spray, or incorporated into a soil mix for subsequent seeding or transplanting of ornamentals. **A216.02** may be applied as a foliar spray on azaleas. **Within a rate range given for a specific group of ornamentals, use the lower specified rate for the shortest interval listed and the higher specified rate for the longest interval. Under severe disease conditions, use the highest specified rate and the shortest interval specified.**

For drench applications, use enough of the specified **A216.02** water solution to wet the root zone of plants. In general, 1 pt. (0.5 lb. a.i.) product per sq. ft. of this solution is sufficient for ornamentals growing in containers with 4 inches of growth media. Containers with growth media depth greater than 4 inches generally require 1 1/2 – 2 pts. (0.75-1 lbs. a.i.) product per sq. ft. of the solution. If soil surface applications are made, irrigate with at least 1/2 inch of water if rainfall does not occur within 7 days.

<p><b>NOTICE TO USER:</b> Due to the large number of species and varieties of ornamentals and nursery plants, it is impossible to test every one for tolerance to <b>A216.02</b>. Neither the manufacturer nor the seller has determined whether or not <b>A216.02</b> can be used safely on ornamental and nursery plants not specified on this label. The professional user should determine if can be used safely prior to commercial use. In a small area, test the specified rates for a particular group of unlabeled plants, i.e., bedding plants, foliage, etc., for phytotoxicity prior to widespread use.</p>	
<p><b>Foliage Plants</b>                  Aglaonema,                  Aphelandra,                  Dieffenbachia,                  Peperomia,                  Philodendron*,                  Pothos,                  Schefflera,                  Sedum,                  Sempervivum,                  Zycactus</p>	<p><b>Drench:</b> Mix 0.25 – 0.63 fl. oz. (0.008-0.020 lbs. a.i.) product with 100 gals. of water. Apply 1 pt. solution per sq. ft. For growth media depth greater than 4 inches, apply 1 1/2 – 2 pts. solution per sq. ft. Repeat applications at two to three-month intervals if necessary.</p> <p><b>*NOTE:</b> On <b>Philodendron</b>, use 0.5-1 fl. oz. (0.016-0.031 lbs. a.i.) product per 100 gals.</p> <p><b>RESTRICTIONS:</b> To minimize the potential for injury to Pothos, do not use more than 0.38 fl. oz. (0.012 lbs. a.i.) product per 100 gals. and do not apply more frequently than once every 3 months.</p> <p><b>Soil Mix:</b> Thoroughly mix 0.13 –0.25 fl. oz. (0.004-0.008 lbs. a.i.) product with each cu. yd. of soil mixture.</p> <p><b>Soil Surface Spray to Foliage Plants in the Landscape:</b> Apply 1 fl. oz. (0.031 lbs. a.i.) product per 1,000 sq. ft. to the soil surface in a broadcast or banded spray in sufficient water to obtain thorough coverage of the plant root zone. After application irrigate with a minimum of 1/2 inch of water if rainfall does not occur within seven days.</p>

<p><b>Bedding Plants</b>  Ageratum,  Algerian ivy,  Artemisia,  Aster,  Begonia,  Caladium,  Carnation,  Chrysanthemum,  Coleus,  Daisy,  English ivy*,  Foxglove,  Gaillardia,  Geranium,  Impatiens,  Marigold,  Pansy,  Petunia,  Phlox,  Pinks,  Primrose,  Prostrate,  Rosemary,  Salvia,  Snapdragon,  Verbena,  Vinca,  Zinnia</p>	<p><b>Drench at Seeding:</b> (Soil 2-3 inches deep) Mix 0.13 – 0.25 fl. oz. (0.004-0.008 lbs. a.i.) product with 100 gals. of water and apply 1 pt. solution per sq. ft.  <b>Drench at Transplanting:</b> (Soil 2-3 inches deep) Mix 0.25 – 1 fl. oz. (0.008-0.031 lbs. a.i.) product with 100 gals. of water and apply 1 pt. solution per sq. ft. For growth media depth greater than 4 inches, apply 1 1/2 –2 pts. solution per sq. ft. Repeat applications at one to two month intervals if necessary.  <b>Soil Mix at Seeding and at Transplanting:</b> Thoroughly mix 0.13 fl. oz. (0.004 lbs. a.i.) product with each cu. yd. of soil mixture.  <b>Soil Surface Spray to Bedding Plants in the Landscape:</b> Apply 1 fl. oz. (0.031 lbs. a.i.) product per 1,000 sq. ft. to the soil surface in a broadcast or banded spray in sufficient water to obtain thorough coverage of the plant root zone. After application irrigate with a minimum of 1/2 inch of water if rainfall does not occur within seven days.  <b>*RESTRICTIONS:</b> Do not apply to English ivy more than once every 6 months or injury may occur. Do not apply rates of 1 1/2 –2 fl. oz. (0.047-0.063 lbs. a.i.) product per 100 gals. more often than once every six weeks.</p>
<p><b>Flowers</b>  African violet,  Anthurium,  Baby's breath,  Carnation,  Chrysanthemum,  Columbine,  Delphinium,  Easter lily*,  Geranium,  Gloxinia,  Poinsettia,  Rose</p>	<p><b>Drench:</b> Mix 0.13 – 1 fl. oz. (0.004-0.031 lbs. a.i.) product with 100 gals. of water and apply 1 pt. solution per sq. ft. For growth media depth greater than 4 inches, apply 1 1/2 – 2 pts. solution per sq. ft. Repeat applications at one to two month intervals if necessary.  <b>*RESTRICTIONS:</b> Do not apply more than 0.5 fl. oz. (0.016 lbs. a.i.) product per 100 gals. of water to Easter lily and only make one at-planting application. Do not apply rates of 1 1/2 – 2 fl. oz. (0.047-0.063 lbs. a.i.) product per 100 gals. more often than every six weeks.  <b>Soil Surface Spray to Flowers in the Landscape:</b> Apply 1 fl. oz. (0.031 lbs. a.i.) product per 1,000 sq. ft. to the soil surface in a broadcast or banded spray in sufficient water to obtain thorough coverage of the plant root zone. After application irrigate with a minimum of 1/2 inch of water if rainfall does not occur within seven days.</p>

<p><b>Rhododendrons and Azaleas</b></p>	<p><b>Drench:</b> Phytophthora root and crown rot – Mix 0.5 - 1.25 fl. oz. (0.016-0.039 lbs. a.i.) product with 100 gals. of water and apply 1 pt. solution per sq. ft. For growth media depth greater than 4 inches, apply 1 1/2 – 2 pts. solution per sq. ft. Repeat applications at two to four month intervals if necessary.</p> <p><b>Soil Surface Spray:</b> Apply 1 - 2 fl. oz. (0.031-0.063 lbs. a.i.) product per 1,000 sq. ft. to the soil surface in a broadcast or banded spray in sufficient water to obtain thorough coverage of the plant root zone. After application irrigate with a minimum of 1/2 inch of water if rainfall does not occur within seven days.</p> <p><b>Foliar Spray:</b> Phytophthora shoot blight – Mix at 0.63 - 1.25 fl. oz. (0.020-0.039 lbs. a.i.) product with 100 gals. of water. Spray to runoff. Repeat at two to three month intervals if necessary.</p> <p><b>RESTRICTIONS:</b> 1) For azaleas, do not apply repeat soil applications of 1.25 fl. oz. (0.039 lbs. a.i.) product per 100 gals. closer than every three months and do not exceed a total of 2 fl. oz. (0.063 lbs. a.i.) product in six months. 2) Use the lower specified rate for “Coral Bell” variety.</p>
<p><b>Woody Ornamentals other than Azaleas</b>  <i>Aucuba japonica</i>,  Arborvitae,  Boxwood,  Ceanothus,  Cotoneaster,  Dogwood,  Ficus,  “Halls”  Honeysuckles,  Ilex,  <i>Juniperus</i> spp.,  Photinia,  <i>Pieris japonica</i>,  <i>Pinus</i> spp.,  Pittosporum,  White cedar,  White pine,  Yew</p>	<p><b>Drench:</b> Mix 0.5 - 2 fl. oz. (0.016-0.063 lbs. a.i.) product with 100 gals. of water and apply 1 pt. solution/sq. ft. For growth media depth greater than 4 inches, apply 1 1/2 – 2 pts. solution/sq. ft. Repeat applications at two to three month intervals if necessary.</p> <p><b>Soil Surface Spray:</b> Apply 1 - 2.5 fl. oz. (0.031-0.078 lbs. a.i.) product per 1,000 sq. ft. to the soil surface in a broadcast or banded spray in sufficient water to obtain thorough coverage of the plant root zone. After application irrigate with a minimum of 1/2 inch of water if rainfall does not occur within seven days.</p> <p><b>RESTRICTIONS:</b> Do not apply to <b>Euonymus</b> or injury may occur. Do not apply rates greater than 3 1/4 fl. oz. (0.102 lbs. a.i.) product per 100 gals. more often than once every ten weeks.</p>

**INTERIORSCAPES AND INDIVIDUAL PLANT USE**

In situations where water volumes used are much less than 100 gals. and the area treated is small, the following table provides the **A216.02** rates to make small quantities of solution. Refer to the plant type for the correct fl. oz. of product to use when utilizing this table.

Rate of A216.02 (fl. oz.)	Amount of A216.02 to Add to Water to Make the Following Quantities			
	1 gal.	5 gals.	10 gals.	25 gals.
0.25 (0.008 lbs. a.i.)	3.5 drops	18.5 drops/0.38 ml	37.5 drops/0.75 ml	1.88 ml/0.38 tsp.
0.5 (0.016 lbs. a.i.)	7.5 drops	37.5 drops/0.75 ml	1.5 ml/0.25 tsp.	3.75 ml/0.75 tsp./0.25 Tbsp.
1.0 (0.031 lbs. a.i.)	15 drops	1.5 ml/0.25 tsp.	3.0 ml/ 0.625 tsp.	7.5 ml/1.5 tsp./0.5 Tbsp.
1.5 (0.047 lbs. a.i.)	22.5 drops	2.25 ml/0.5 tsp.	4.5 ml/ 1 tsp.	11.25 ml/2.25 tsp./ 0.625 Tbsp.
2.0 (0.063 lbs. a.i.)	30 drops	3.0 ml/ 0.625 tsp.	6.0 ml/1.25 tsp.	15.0 ml/ 1 Tbsp./0.5 oz.



**Soil Drench:** Apply enough solution to wet the root area of the plants; apply at least one pint of solution per square foot.

### **CITRUS IN NURSERIES AND LANDSCAPE PLANTINGS (Nonbearing)**

Use **A216.02** on nonbearing citrus for control of citrus foot rot, root rot, and trunk canker caused by *Phytophthora* spp. Apply to the soil as a drench or as a spray in a banded application. Restriction: Not for residential use.

Make the first application of **A216.02** at the time of planting. Make repeated applications at three- month intervals during the period when trees are actively growing.

**Soil Drench:**

Mix 2 - 3 fl. oz. (0.063-0.094 lbs. a.i.) product per 100 gals. of water and apply as a drench over the row at the rate of 100- 250 gals. /1,000 feet of row. The width of the drench treatment should be wide enough to cover the root systems of the plants.

**Soil Surface Spray:**

Apply 1 gals. (4 lbs. a.i.) product per acre of treated soil in a broadcast or banded surface spray to seedbeds, liners, or bedded stock in sufficient water to obtain uniform coverage. If applications are banded, the treated area should be wide enough to cover the root systems of the plants. Follow with a 1/2-inch irrigation.

Calculate the amount of **A216.02** needed for a banded treatment by using the formula at the end of the Use Information section of this label.

**RESTRICTIONS:** Do not use in greenhouse citrus nursery stock intended for commercial fruit production.

### **CONIFERS IN NURSERIES AND PLANTATIONS (Including Christmas Trees)**

**A216.02** provides control of *Phytophthora* root rot of conifers. For best results, apply ½ to 1 inch of water after application if rain is not expected within three days.

Restriction: Not for residential use.

**Conifers in Nurseries**

Seedbeds and Plug-Planting	Apply 1.25 pts. (0.625 lbs. a.i.) of <b>A216.02</b> in at least 50 gals. of water per acre in the spring and again in the fall.
2-0 Transplants	Apply 2.5 pts. (1.25 lbs. a.i.) <b>A216.02</b> in at least 50 gals. of water per acre in the spring and again in the fall.

**Conifers in Plantations:**

Use of **A216.02** will aid in the control of *Phytophthora* root rot when used in conjunction with good cultural practices. The use of **A216.02** will not overcome poor management practices such as planting on sites that are prone to flooding or are poorly drained. **A216.02** fungicide will not revitalize trees showing moderate to severe disease symptoms.

Apply 0.63 – 1.25 gals. (2.52-5 lbs. a.i.) of **A216.02** in a minimum of 50 gals. of water as a directed soil spray. Do not apply as a foliar spray. Applications should be made in early spring before growth starts and in the fall before the ground freezes. Calculate the amount of **A216.02** needed for a banded treatment by using the formula at the end of the Use Information section of the label.

### **DECIDUOUS FRUITS AND NUTS IN NURSERIES (Nonbearing)**

**A216.02** provides control of *Pythium* root rot and *Phytophthora* root, crown, and collar rot of nonbearing deciduous fruits and nuts.



Apply 3 fl. oz. (0.094 lbs. a.i.) product per 1,000 sq. ft. in sufficient water to obtain thorough coverage of the soil under the canopy of the trees. Treat sufficient surface area in nurseries to cover the root zone of the plants. Additional applications may be made as necessary at three-month intervals during the growing season.

**RESTRICTIONS:**

1. Do not apply to trees that will bear harvestable fruit within 12 months of the last application, or possible illegal residues may result.
2. Do not apply more than 8.8 fl. oz. (0.275 lbs. a.i.) per 1,000 sq. ft. (3 gals. per acre; 12 lbs. a.i.) of **A216.02** per year.
3. Not for residential use.

**TURF**

**(Golf Courses, Lawns, Landscape Areas around Institutional, Public, Commercial and Industrial Buildings, Parks, Recreational Areas, and Athletic Fields, Sod Farms)**

**RESTRICTION: Not for residential use.**

**A216.02** controls Pythium blight and Pythium damping-off in turf, yellow tuft (downy mildew) in bluegrass, and downy mildew in St. Augustinegrass. **Within the rate range given for turf, use the lower specified rate for the shortest interval listed** and the higher specified rate for the longest interval. Under severe disease conditions, use the highest specified rate and shortest interval specified.

<p><b>Established Turf</b> Pythium Blight Yellow Tuft Downy Mildew</p>	<p>Apply as a preventative treatment at 0.5 - 1 fl. oz. (0.016-0.031 lbs. a.i.) product in 3-5 gals. of water per 1,000 sq. ft. Retreat at 10-21 day intervals. During periods of prolonged conditions favorable for disease development, use 1 fl. oz. (0.031 lbs. a.i.) product on a 14-day schedule.</p>
<p><b>Newly Seeded Areas</b> Pythium Damping-Off Pythium Blight Yellow Turf Downy Mildew</p>	<p>Apply 0.5 - 1.0 fl. oz. (0.016-0.031 lbs. a.i.) product in 5-10 gals. of water per 1,000 sq. ft. immediately after seeding. Retreat at 7-14 day intervals if conditions remain favorable for disease.</p> <p><b>Note:</b> For long-term control of Pythium in areas when using seed treated with the active ingredient contained in <b>A216.02</b>, make application of <b>A216.02</b> 7-10 days after seeding.</p>

**NOTE:** For control of other diseases of turf, use propiconazole alone or in tank mix combination with **A216.02**. Refer to the propiconazole label for rates, precautions, restrictions, etc.

**Resistance Management:**

To minimize the potential for resistance:

1. Make no more than three applications of **A216.02** per season.
2. Apply an alternate EPA- registered fungicide for Pythium control at least once during the season.

**ROTATIONAL CROPS—TURF USE ONLY**

Do not plant any crop which is not registered for use with metalaxyl in metalaxyl-treated soil for a period of 12 months.

## SEED TREATMENT

**A216.02** is a systemic fungicide for use as a seed dressing.

### Resistance Management:

**A216.02** is a systemic fungicide having a specific mode of action. **A216.02** could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible **A216.02** resistant strains of fungi which may occur.

**A216.02** may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.

<b>Grain Sorghum</b>	For Pythium damping-off control: Apply <b>A216.02</b> as a seed treatment at the rate of 0.5 - 1 fl. oz. (0.016-0.031 lbs. a.i.) product per 100 lbs. of seed.
<b>Soybeans</b>	For Pythium damping-off and early season Phytophthora control: Apply <b>A216.02</b> as a seed treatment at the rate of 0.5 - 1 fl. oz. (0.016-0.031 lbs. a.i.) product per 100 lbs. of seed.
<b>Sunflower</b>	For control of systemic downy mildew: Apply <b>A216.02</b> as a seed treatment at the rate of 2 fl. oz. (0.063 lbs. a.i.) product per 100 lbs. of seed.

### Seed Bag Label Requirement:

The Federal Seed Act that containers containing treated seeds shall be labeled with the following statements:

- This seed has been treated with **A216.02**, a fungicide containing metalaxyl.
- Do not use treated seed for feed, food, or oil purposes.

The U.S. Environmental Protection Agency requires the following statements on containers containing seed treated with metalaxyl:

- Store treated seed away from food and feedstuff.
- Do not allow children, pets, or livestock to have access to treated seeds.
- Do not allow livestock to graze cotton plants grown from treated seeds.
- Wear long pants, long-sleeved shirt and protective gloves when handling treated seed.
- Treated seeds exposed on the soil surface may be hazardous to wildlife. Cover or collect treated seed spilled during loading and planting (such as in row ends).
- Dispose of all excess treated seed by burying seed away from bodies of water.
- Do not contaminate bodies of water when disposing of panting equipment wash water.
- Dispose of seed packaging or container in accordance with local requirements.
- Excess seed may be used for ethanol production if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.

Seed treated with this product must be visually identifiable from untreated seed by the use of an approved colorant or dye to prevent accidental use of treated seed as food for humans or feed for animals. Refer to 21 CFR, Part 2.25. Any colorant or dye added to treated seed must be cleared for use in accordance with 40 CFR, Part 153.155(c).

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store at temperatures above 40°F. Crystals may form at lower storage temperatures. If this occurs, place the product in a warm room (68°F or above) and roll or shake the container at frequent intervals until all crystals are dissolved.

**Pesticide Disposal:** Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide spray mixture, or rinsate is a violation of Federal law. Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office.

**Container Handling: Nonrefillable container:** Do not reuse or refill this container. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Returnable/Refillable Containers:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For minor spills, leaks, etc. follow all precautions indicated on this label and clean-up immediately. Take special care to avoid contamination of equipment and facilities during clean-up procedures and disposal of wastes.

## LIMITATION OF WARRANTY AND LIABILITY

**IMPORTANT: READ BEFORE USE.** Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label.

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

[A216.02] is a trademark of Atticus, LLC

# {LANGUAGE ON LABEL AFFIXED TO CONTAINER}

METALAXYL	GROUP	4	FUNGICIDE
-----------	-------	---	-----------

**A216.02™**

[Alternate Brand Name: ReCon 4 F]

**Active Ingredient:** (by weight)

Metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl) alanine methyl ester ..... 44.08%

**Other Ingredients:** ..... 55.92%

**Total** ..... 100.00%

Contains 4 lbs. of metalaxyl per gallon.

## KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor.</li> <li>Have a person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything to an unconscious person.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<p><b>HOT LINE NUMBER</b></p> <p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-844-685-9173 for emergency medical treatment information.</p>	

### For Chemical Emergency

**Spill, Leak, Fire, Exposure, or Accident Call CHEMTREC Day or Night**

**Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Cause moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco or using the toilet.

**Environmental Hazards:** For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Apply only as specified on this label. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

**Groundwater Advisory Statement:** This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store at temperatures above 40°F. Crystals may form at lower storage temperatures. If this occurs, place the product in a warm room (68°F or above) and roll or shake the container at frequent intervals until all crystals are dissolved.

**Pesticide Disposal:** Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide spray mixture, or rinsate is a violation of Federal law. Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office.

**Container Handling: Nonrefillable container:** Do not reuse or refill this container. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Returnable/Refillable Containers:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For minor spills, leaks, etc. follow all precautions indicated on this label and clean-up immediately. Take special care to avoid contamination of equipment and facilities during clean-up procedures and disposal of wastes.

See inside label booklet for additional Precautionary Statements and Directions for Use.

### Manufactured for:

Atticus, LLC  
5000 CentreGreen Way, Suite 100  
Cary, NC 27513

**EPA Reg. No. 91234-XX**

**EPA Est. No. \_\_\_\_\_**

**NET WEIGHT: \_\_\_\_\_**