



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

August 16, 2022

Kristen Cianni  
Regulatory Specialist  
Atticus, LLC  
500 CentreGreen Way, Suite 100  
Cary, NC 27513

Subject: Notification per PRN 2007-4 – Updated Storage and Disposal language  
Product Name: A148.01  
EPA Registration Number: 91234-114  
Application Date: August 08, 2022  
Decision Number: 586792

Dear Ms. Cianni:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 2007-4 and finds that the action requested falls within the scope of PRN 2007-4.

The label submitted with the application has been stamped “Notification” and will be placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, you may contact please contact David Gardner at (202) 566-2677 or by email at [gardner.david@epa.gov](mailto:gardner.david@epa.gov).

Sincerely,

A handwritten signature in cursive script that reads "Tamica Cain".

Tamica Cain, Product Manager 10  
Invertebrate & Vertebrate Branch 2  
Registration Division (7505T)

[Note to reviewer: [Text] in brackets denotes optional or explanatory language  
[Note to reviewer: {Text} in braces denotes where in the final label text will appear  
{BOOKLET FRONT PANEL LANGUAGE}

CYROMAZINE	GROUP	17	INSECTICIDE
------------	-------	----	-------------

# A148.01 [™]

[Alternate Brand Name: Trignata]

Contains cyromazine, the active ingredient used in Trigard®.

[For control of leafminers in Abyssinian and seakale cabbage, bulb vegetables, cucurbits, certain varieties of dried beans, Hanover salad, leafy vegetables including Brassica, succulent beans, peppers, tomatoes, and turnip greens and control of Colorado potato beetle and leafminer in potatoes. A148.01 Insecticide is a wettable powder.]

ACTIVE INGREDIENT:	(% by weight)
Cyromazine (CAS No. 66215-27-8) 2,4,6-triamine.....	75.0%
OTHER INGREDIENTS: .....	25.0%
TOTAL .....	100.0%

## KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside label booklet for First Aid, Precautionary Statements and Directions for Use.

[A148.01™] is not manufactured, or distributed by ADAMA, seller of Trigard®.

EPA Reg. No.: 91234-114

EPA Est. No.:

20190329c

### NOTIFICATION

91234-114

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

08/16/2022

Manufactured for:  
**Atticus, LLC**

5000 CentreGreen Way, Suite 100  
Cary, NC 27513

## {LANGUAGE INSIDE BOOKLET}

<b>FIRST AID</b>	
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at <b>1-844-685-9173</b> for emergency medical treatment information.	

**For Chemical Emergency:  
Spill, Leak, Fire, Exposure, or Accident,  
Call CHEMTREC Day or Night  
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

## **PRECAUTIONARY STATEMENTS**

### **Hazards to Humans and Domestic Animals**

#### **CAUTION**

Causes moderate eye irritation. Harmful if swallowed, absorbed through the skin, or inhaled. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

#### **PERSONAL PROTECTIVE EQUIPMENT**

##### **Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material – Category A (e.g., barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyethylene, polyvinylchloride [PVC] or viton)
- Chemical-resistant footwear plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

## ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

#### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

## ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water. Do not apply when weather conditions favor drift from target area.

## GROUND WATER ADVISORY

A metabolite of **A148.01** Insecticide has been found in ground water, possibly as a result of agricultural use. Use of this product in areas where soils are permeable and water tables are shallow could result in leaching to ground water. Irrigation could increase the risk of contamination.

### PROTECTION OF POLLINATORS

**“Cyromazine acts as a growth regulator, affecting the molting process of target pests, and therefore, may have an effect on bee larvae, but not on adult bees. Exposure to adult bees, either through overspray or contact with residues may lead to effects to bee larvae.”**

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications.
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product, Applicators Should Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this products on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: <http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx> .

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: [www.aapco.org/officials.html](http://www.aapco.org/officials.html) . Pesticide incidents should also be reported to the National Pesticide Information Center at: [www.npic.orst.edu](http://www.npic.orst.edu) or directly to EPA at: [beekill@epa.gov](mailto:beekill@epa.gov) .

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**A148.01** Insecticide must be used only as directed in accordance with recommendations on this label.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material – Category A (e.g., barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyethylene, polyvinylchloride [PVC] or viton)
- Chemical-resistant footwear plus socks

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY AND/OR POOR INSECT CONTROL.**

**NOT FOR SALE, SALE INTO, DISTRIBUTION AND/OR USE IN NASSAU AND SUFFOLK COUNTIES, NEW YORK.**

### GENERAL INFORMATION

**A148.01** Insecticide is a 75% wettable powder. When this product is mixed with water and applied according to the directions for use to the specified crops on this label, **A148.01** Insecticide will control the specified insects in the crops that are listed on this label.

### RESISTANCE MANAGEMENT

**A148.01** Insecticide is a Group 17 Insecticide (contains the active ingredient cyromazine). Because of the inherent risk of resistance development to any product, it is strongly advised that **A148.01** Insecticide be used in a sound resistance management program. Treatment may not be effective against labeled pests if **A148.01** Insecticide tolerant strains develop.

For plants that host a labeled pest with multiple generations per crop or year it is recommended to follow the following resistance management practices that may include, but are not limited to:

- Rotating **A148.01** Insecticide with products with different modes of action
- Avoiding treatment of successive generations with **A148.01** Insecticide
- Using labeled rates at specified spray intervals
- Using non-chemical alternatives such as beneficial arthropods
- Using various cultural practices such as rotating susceptible to non-susceptible plants

For additional information about implementing these or other resistance management practices, consult your local agricultural advisor or company representative.

## **MIXING AND APPLICATION INSTRUCTIONS**

1. Thoroughly clean spray equipment before using this product.
2. Prepare no more spray mixture than is needed for the immediate operation.
3. Agitate spray mixture thoroughly before and during application. Good bypass agitation is sufficient to keep **A148.01** Insecticide thoroughly mixed.
4. Flush spray tank thoroughly with water daily after use and
5. Dispose of pesticide rinsate by application to an already treated area.

### **A148.01 Insecticide Alone:**

1. Add 2/3 of the required amount of water to the spray or mixing tank.
2. With the agitator running, drop the required number of unopened soluble packets of **A148.01** Insecticide into the tank. (One 2.66 oz. water-soluble packet of **A148.01** Insecticide will treat one acre).
3. Continue agitation while adding the remainder of the water.

### **Note**

- Do not let spray mixture stand overnight in the spray tank. Rinse spray tank at the end of the day.
- To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, because crop injury may result.
- Use aerial applications only when conditions prevent application with ground equipment.
- When using ground application equipment direct the spray toward crop foliage.

**Chemigation:** Do not apply this product through any type of irrigation system.

**Greenhouses:** Do not use this product in greenhouses.

## **SPRAY DRIFT REQUIREMENTS**

### **Droplet Size**

Use only medium or coarser spray nozzles (for ground and non-ULV aerial application) according to ASAE (S572) definition for standard nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

## CROP USE DIRECTIONS

### BEANS, DRIED VARIETIES EXCEPT COWPEAS AND SOYBEANS\*

\*Adzuki beans, black-eyed peas, broad beans (dry), catjang, chick peas, Crowder peas, field beans, garbanzo beans, grain lupin, guar, kidney beans, lablab beans, lentils, lima beans (dry), moth beans, mung beans, navy beans, pinto beans, rice beans, southern peas, tepary beans, sweet lupin, urd beans, wax beans, white lupin, white sweet lupin

#### For leafminer control:

- Apply 2.66 oz. (one packet) of **A148.01** Insecticide /A as a foliar spray when leafminers first appear.
- Apply with aircraft in a minimum of 5 gal. of water /A or with ground application equipment in a minimum of 10 gal. of water /A, using sufficient carrier to achieve adequate coverage.
- Repeat application, as needed, at 7-day intervals to maintain control.

**Notes:** To avoid possible illegal residues:

- Do not make more than 6 applications per crop of dry beans.
- Do not apply more than 1.0 lb. product per acre (0.75 lb. a.i./A) per season.
- Do not make the last application within 7 days of harvest. (7-day PHI)

### BEANS, SUCCULENT

#### For leafminer control:

- Apply 2.66 oz. (one packet) of **A148.01** Insecticide /A as a foliar spray when leafminers first appear.
- Apply with aircraft in a minimum of 5 gal. of water/A, or apply with ground application equipment in a minimum of 10 gal. of water/A, using sufficient carrier to achieve adequate coverage.
- Repeat application, as needed, at 7-day intervals to maintain control.
- Do not make more than 6 applications to one crop of lima beans.
- Do not make the last application within 7 days of harvest. (7-day PHI).

Crop/Pest	Rate per Acre	Instructions
<b>Succulent Beans</b> <b>Control of Leafminer</b>  Includes only: <b>Bean (<i>Phaseolus</i> spp.)</b> Broad bean, succulent Lima bean, green Runner bean Snap bean (green bean) Wax bean  <b>Bean (<i>Vigna</i> spp.)</b> Asparagus bean Black-eyed pea Chinese longbean Cowpea Jackbean	2.66 oz.(one packet)	Apply <b>A148.01</b> Insecticide as a foliar spray when leafminers first appear.  Repeat application, as needed, to maintain control. <b>(See Use Restrictions)</b>

Moth bean Southern pea Yardlong bean		
--	--	--

**Use Restrictions – Succulent Beans**

- **PHI:** Allow a minimum of 7 days between the last application and harvest.
- **Application Method:** Ground or aerial application equipment is permitted.
- **Amount of Water:** Apply with aircraft in a minimum of 5 gal. of water/A or with ground application equipment in a minimum of 10 gal. of water/A, using sufficient carrier to achieve adequate coverage.
- **Application Interval:** Allow a minimum of 7 days between applications.
- **Maximum Amount:** Do not apply more than 0.125 lb a.i./A per 7-day interval. Do not apply more than 0.75 lb. a.i./A per crop season. Do not apply more than 6 times per crop season.

**BRASSICA LEAFY VEGETABLES CROP GROUP\*, ABYSSINIAN CABBAGE, SEAKALE CABBAGE, HANOVER SALAD, and TURNIP GREENS\*\***

\*Some of the crops in the Brassica Leafy Vegetables are: broccoli, broccoli raab (rapini), Brussels sprouts, cabbage, Chinese (bok choy) cabbage, Chinese broccoli, Chinese mustard cabbage (gai choy), Chinese (napa) cabbage, cauliflower, cavalo broccolo, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, and rape greens

\*\*Turnip Greens (tops, leaves) only: For use on turnip varieties grown for leaves only. Do not use on turnip varieties grown for roots or dual-purpose varieties grown for roots and leaves.

**For leafminer control:**

- Apply 2.66 oz. (one packet) of **A148.01** Insecticide/A as a foliar spray when leafminers first appear.
- Apply with aircraft in a minimum of 5 gal. of water /A, or with ground application equipment in a minimum of 10 gal. of water /A, using sufficient carrier to achieve adequate coverage.
- Repeat application, as needed, at 7-day intervals to maintain control.

**Notes:** To avoid possible illegal residues:

- Do not make more than 6 applications to one crop of Brassica leafy vegetables, Abyssinian cabbage, seakale cabbage, Hanover salad, or turnip greens.
- Do not apply more than 1.0 lb. (0.75 lb. a.i./A) of product per season.
- Do not make the last application within 7 days of harvest. (7-day PHI).

**BULB VEGETABLES CROP GROUP\***

\*Some of the crops in this group are: garlic, great-headed (elephant) garlic, leek, dry bulb onion, green onion, potato onion, tree onion, Welsh onion, rakkyo, and shallot.

**For leafminer control:**

- Apply 2.66 oz. (one packet) of **A148.01** Insecticide per acre as a foliar spray when leafminers first appear.
- Apply with aircraft in a minimum of 5 gal. of water /A or with ground application equipment in a minimum of 10 gal. of water /A, using sufficient carrier to achieve adequate coverage.



- Repeat application, as needed, at 7-day intervals to maintain control.

**Notes:** To avoid possible illegal residues:

- Do not make more than 6 applications to one crop of bulb vegetables.
- Do not make the last application within 7 days of harvest (or cutting). (7-day PHI)

**CUCURBIT VEGETABLES CROP GROUP\***

\* Some of the crops in this group are: balsam pear (bitter melon), chayote (fruit), Chinese waxgourd (Chinese preserving melon), citron melon, cucumber, gherkin, edible gourds, melons (including hybrids: cantaloupe, casaba, crenshaw, honeydew melons, honey balls, mango melon, muskmelon, Persian melon), pumpkin, summer squash, winter squash, and watermelon (including hybrids).

**For leafminer control:**

- Apply 2.66 oz. (one packet) of **A148.01** Insecticide/A as a foliar spray when leafminers first appear.
- Apply with aircraft in a minimum of 5 gal. of water /A, or with ground application equipment in a minimum of 10 gal. of water /A, using sufficient carrier to achieve adequate coverage.
- Repeat application, as needed, at 7-day intervals to maintain control.

**Notes:** To avoid possible illegal residues:

- Do not make more than 6 applications per growing season to cucurbits.
- Cucurbits may be harvested on the same day of the last application. (0-day PHI)

**LEAFY VEGETABLES\* (EXCEPT BRASSICA VEGETABLES) CROP GROUP**

\*Some of the crops in this group are: Amaranth, arugula, cardoon, celery, celuce, chervil, Chinese celery, chrysanthemum (edible-leaved & garland), corn salad, cress (garden & upland), dandelion, dock, endive (escarole), Florence fennel (finocchio), lettuce (head & leaf), New Zealand spinach, orach, parsley, purslane (garden & winter), radicchio (red chicory), rhubarb, spinach, Swiss chard, and vine spinach

**For leafminer control:**

- Apply 2.66 oz. (one packet) of **A148.01** Insecticide /A as a foliar spray when leafminers first appear.
- Apply with aircraft in a minimum of 5 gal. of water /A, or with ground application equipment in a minimum of 10 gal. of water /A, using sufficient carrier to achieve adequate coverage.
- Repeat application, as needed, at 7-day intervals to maintain control.

**Notes:** To avoid possible illegal residues:

- Do not make more than 6 applications to one crop of celery or head lettuce.
- For all other leafy vegetable crops, do not make more than 5 applications per crop.
- Do not make the last application within 7 days of harvest. (7-day PHI)

**PEPPERS**

**For leafminer control:**

- Apply 2.66 oz. (one packet) of **A148.01** Insecticide/A as a foliar spray when leafminers first appear.
- Apply with aircraft in a minimum of 5 gal. of water /A, or with ground application equipment in a minimum of 10 gal. of water /A, using sufficient carrier to achieve adequate coverage.
- Repeat application, as needed, at 7-day intervals to maintain control.

**Notes:** To avoid possible illegal residues:

- Do not make more than 6 applications to one crop of peppers.
- Peppers may be harvested on the same day of the last application. (0-day PHI)

## POTATOES

**For Colorado potato beetle control:**

- **A148.01** Insecticide is an insect growth regulator and is most effective for control of first and second instar larvae of Colorado potato beetles. Although larger larvae can be affected, treatment of early instars will give best control. Affected larvae may remain on the plant for several days after treatment, but immediately stop feeding.
- **A148.01** Insecticide does not control adult beetles.
- Apply 2.66 oz. (one packet) or 5.32 oz. (two packets) per acre as a foliar spray. The first application should be made just after egg hatch to early instar larvae. If a second application for the same generation is needed, apply 2.66 oz. (one packet) or 5.32 oz. (two packets) per acre 7 to 14 days later.
- Apply with aircraft in a minimum of 5 gal. of water /A. With ground application equipment use a sufficient amount of water to ensure thorough coverage of foliage.

**For leafminer control:**

- Apply 2.66 oz. (one packet) per acre as a foliar spray.
- Use sufficient water volume to ensure thorough coverage of foliage when using ground application equipment and a minimum of 5.0 gallons of water when applying with aircraft.
- Repeat application at 7-day intervals, as needed, to maintain good control.

**Notes:** To avoid possible illegal residues:

- Do not exceed a total of 1 lb. of **A148.01** Insecticide per acre during the growing season.
- Allow at least 7 days between applications.
- Do not make the last application within 17 days of harvest (17-day PHI).

## TOMATOES

**For leafminer control and suppression of Colorado potato beetle:**

- Apply 2.66 oz. (one packet) of **A148.01** Insecticide /A as a foliar spray when leafminers first appear. At this application rate, **A148.01** Insecticide will also provide suppression of Colorado Potato Beetle.
- Apply with aircraft in a minimum of 5 gal. of water /A, or with ground application equipment in a minimum of 10 gal. of water /A, using sufficient carrier to achieve adequate coverage.
- Repeat application, as needed, at 7-day intervals to maintain control.

**Notes:** To avoid possible illegal residues:

- Do not make more than 6 applications to one crop of tomatoes.
- Allow at least 7 days between applications.
- Tomatoes may be harvested on the same day of the last application.

## ROTATIONAL CROPS

The following “plant-back intervals” must be observed when planting rotational crops after an application of **A148.01** Insecticide:

Rotational Crop	Planting Time From Last Application of A148.01 Insecticide (Days)
All crops on the <b>A148.01</b> Insecticide label	0 days
Sweet corn, radishes, and cotton	30 days
Any other crop	300 days

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not use in animal feeds.

**PESTICIDE STORAGE:** Store in a tightly closed container in a cool, dry place. Do not store this product under wet conditions. Handle outer container carefully to avoid breakage of water-soluble packets.

**PESTICIDE DISPOSAL:** Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### CONTAINER HANDLING:

Nonrefillable container. Do not reuse or refill this container. Empty residue into application equipment. Offer container and foil bag for recycling, if available, or dispose of container and/or foil bag in a sanitary landfill or by incineration, if allowed by State and local authorities. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

[Nonrefillable Container: Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Offer for recycling, if available.]

## LIMITATION OF WARRANTY AND LIABILITY

**IMPORTANT: READ BEFORE USE.** Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

[**A148.01™**] is a trademark of Atticus, LLC

Trigard® is a registered trademark of an ADAMA group company.

**{LANGUAGE ON LABEL AFFIXED TO CONTAINER}**

CYROMAZINE	GROUP 17	INSECTICIDE
------------	----------	-------------

**A148.01™**

[Alternate Brand Name: *Trignata* ]

Contains cyromazine, the active ingredient used in Trigard®.  
[Statement for Use]

<b>ACTIVE INGREDIENT:</b>	(% by weight)
Cyromazine (CAS No. 66215-27-8) 2,4,6-triamine.....	75.0%
<b>OTHER INGREDIENTS:</b> .....	25.0%
<b>TOTAL</b> .....	100.0%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>● Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>● Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>● Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>● Take off contaminated clothing.</li> <li>● Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>● Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>● Call a poison control center or doctor immediately for treatment advice.</li> <li>● Have person sip a glass of water if able to swallow.</li> <li>● Do not induce vomiting unless told to do so by the poison control center or doctor.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>● Move person to fresh air.</li> <li>● If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>● Call a poison control center or doctor for further treatment advice.</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at <b>1-844-685-9173</b> for emergency medical treatment information.	

**For Chemical Emergency:**

Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC Day or Night  
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Causes moderate eye irritation. Harmful if swallowed, absorbed through the skin, or inhaled. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water. Do not apply when weather conditions favor drift from target area.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. Do not use in animal feeds.

**PESTICIDE STORAGE:** Store in a tightly closed container in a cool, dry place. Do not store this product under wet conditions. Handle outer container carefully to avoid breakage of water-soluble packets.

**PESTICIDE DISPOSAL:** Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING:**

Nonrefillable container. Do not reuse or refill this container. Empty residue into application equipment. Offer container and foil bag for recycling, if available, or dispose of container and/or foil bag in a sanitary landfill or by incineration, if allowed by State and local authorities. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

[Nonrefillable Container: Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Offer for recycling, if available.]

See inside label booklet for additional Precautionary Statements and Directions for Use.

[A148.01™] is not manufactured, or distributed by ADAMA, seller of Trigard®.

Manufactured for:  
**Atticus, LLC**  
5000 CentreGreen Way, Suite 100  
Cary, NC 27513

**EPA Reg. No.: 91234-114**  
**EPA Est. No.:** \_\_\_\_\_  
**NET WEIGHT:** \_\_\_\_\_

20190329c