



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number: **91234-131** Date of Issuance:

8/1/19

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

A326.03

Name and Address of Registrant (include ZIP Code):

Atticus, LLC
5000 CentreGreen Way, Suite 100
Cary, NC 27513

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 91234-131."
3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Signature of Approving Official:

Erik Kraft, Product Manager 24
Fungicide Herbicide Branch, Registration Division (7505P)

Date:

8/1/19

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 02/20/2019

If you have any questions, please contact BeWanda Alexander by phone at (703)347-0313, or via email at alexander.bewanda@epa.gov.

Enclosure

[Note to reviewer: [Text] in brackets denotes optional or explanatory language

[Note to reviewer: {Text} in braces denotes where in the final label text will appear

{BOOKLET FRONT PANEL LANGUAGE}

A C C E P T E D

08/01/2019

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 91234-131

MESOTRIONE	GROUP	27	HERBICIDE
FLUTHIACET METHYL	GROUP	14	HERBICIDE

A326.03 [TM]

[Alternate Brand Name: Cavallo STX]

[Contains mesotrione and fluthiacet, the active ingredients used in Solstice®.]

ACTIVE INGREDIENTS:

(% by weight)

Fluthiacet methyl.....2.20%

Mesotrione.....38.52%

OTHER INGREDIENTS:.....59.28%

TOTAL100.0%

Contains a total of 4.0 lb/gal which include 0.216 lb ai Fluthiacet methyl and 3.784 lb ai mesotrione per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If on skin or clothing:	<ul style="list-style-type: none">Take off contaminated clothing.Rinse skin immediately with plenty of water for 15-20 minutes.Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none">Hold eye open and rinse slowly and gently with water for 15-20 minutes.Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

For Chemical Emergency:

Spill, Leak, Fire, Exposure, or Accident,

Call CHEMTREC Day or Night

Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

See inside label booklet for Precautionary Statements and Directions for Use.

[[A326.03™]] is not manufactured, or distributed by FMC Corporation, seller of Solstice®.]

EPA Reg. No.: 91234-XX

EPA Est. No.:

Net Weight:

Manufactured for:
Atticus, LLC
5000 CentreGreen Way, Suite 100
Cary, NC 27513

{LANGUAGE INSIDE BOOKLET}

ATTENTION

- Although this label may appear similar to the label on a product you may have used, there may be important label differences. Users must read, understand and strictly follow all label directions, precautions and restrictions.
- It is the user's responsibility to be sure the product is approved for sale or use on the intended crop and for use in the specific geographic area.
- It is the user's responsibility to be aware of and to follow all State or local precautions or restrictions not appearing on this product label.
- Prior to purchase or use of this product, read the Terms of Sale or Use and Limitation of Warranty and Liability. If the terms and conditions are unacceptable, return the product immediately in the original and unopened container.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: Coveralls worn over long-sleeved shirt and long pants, protective eyewear (goggles or face shield), chemical-resistant gloves, and chemical-resistant footwear plus socks. When mixing and loading wear a chemical-resistant apron.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

IMPORTANT:

When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, including a spill or equipment breakdown.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240)(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS. The closed system must be used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)]. When using the closed system, the mixers' and loaders' PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

A326.03 is very toxic to algae and moderately toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high-water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

MIXING AND LOADING INSTRUCTIONS

This product may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas.

Operations that involve mixing, loading rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad.

Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

This product must be used in a manner which will prevent back siphoning into wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

1. The spray equipment must be clean before using this product. If it is contaminated with other materials, mixing problems and/or clogging may occur or crop injury may occur.
2. Prepare no more spray mixture than is needed for the immediate application, and do not let the spray mixture stand in the spray tank overnight.
3. Maintain maximum agitation throughout the spraying operation.
4. Flush the spray equipment thoroughly after each use and apply rinsate to an appropriate area.

GROUNDWATER ADVISORY:

This chemical and its degradation products have properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

SURFACE WATER ADVISORY:

This product may impact surface water quality due to spray drift and runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months after application. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential loading of **A326.03** from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

PHYSICAL/CHEMICAL HAZARDS

Do not use or store near heat or open flame.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. These requirements only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Personal Protective Equipment (PPE) required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: Coveralls over short-sleeve shirt and short pants, goggles, face shield or safety glasses, chemical-resistant gloves made of any waterproof material including polyethylene or polyvinyl chloride and chemical-resistant footwear plus socks.

PRODUCT INFORMATION

A326.03 contains two active ingredients possessing both contact and systemic activity that can be applied Post emergence for selective control of broadleaf weeds in field corn, seed corn, yellow pop-corn, sweet corn. **DO NOT** apply to White Popcorn or ornamental (Indian) corn. When applied as POST application it may take 2 to 15 days to kill the weeds on the list. The product is absorbed through the soil and /or by the weed foliage.

For effective control of grasses, **A326.03** can be tank mixed with other POST grass herbicides to provide broader spectrum weed control in corn. **A326.03** application can also be combined with a burndown herbicide, prior to planting, to provide added burndown and residual weed control in field corn, seed corn, yellow popcorn, and sweet corn.

RESISTANCE MANAGEMENT

For resistance management, please note that **A326.03** contains both a Group 27 [Mesotrione] and a Group 14 [Fluthiacet methyl] herbicide. Any weed population may contain plants naturally resistant to Group 27 and/or Group 14 herbicides. The resistant individuals may dominate the weed population if these herbicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of **A326.03** or other Group 27 & 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with

controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method including hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.

- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Atticus, LLC at (984) 465-4754.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

Utilize a sprayer equipped with the appropriate nozzles providing optimum spray distribution and coverage at the appropriate operating pressures. Apply a minimum of 10 gallons of finished spray solution per acre by ground. If a dense crop and or weed canopy is present use a minimum of 15 gallons per acre of finished spray volume by ground application. The sprayer must be properly calibrated to deliver the appropriate volume of herbicide solution. Be aware that overlaps and slower ground speeds while starting, stopping or turning while spraying may result in excessive application and subsequent crop response. Mix the amount which will be used for spraying on that day.

Restrictions

- **DO NOT** exceed 3.15 fl oz. (0.098 lb. a.i./A of **A326.03** contains 0.0053 lb ai fluthiacet methyl + 0.0931 lb ai mesotrione) in a single postemergence application.
- **DO NOT** apply more than a total of 5.25 fl. oz. (0.164 lb a.i./A of **A326.03** contains 0.0089 lb ai fluthiacet methyl + 0.155 lb ai mesotrione) in a twelve month period (year) including preplant burndown and labeled postemerge applications to corn.
- **DO NOT** apply more than 0.0089 lb ai of fluthiacet-methyl or 0.24 lb of mesotrione per year including all preplant or postemergence herbicides containing these active ingredients.
- **DO NOT** make the second application of **A326.03** within 14 days of the first application. **DO NOT** make more than 2 applications per year.
- **DO NOT** harvest or feed forage within 45 days after application.
- **DO NOT** harvest or feed grain or stover (fodder) within 70 days after application.
- **DO NOT** harvest or feed sweet corn forage or ears within 40 days after application.
- **DO NOT** include nitrogen-based adjuvants (UAN or AMS) when making postemergence applications of **A326.03** to yellow popcorn or sweet corn.
- **DO NOT** apply **A326.03** to white popcorn or ornamental (Indian) corn.
- **DO NOT** apply to corn that is more than 12 inches in height if atrazine is mixed with **A326.03**.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** use aerial application to apply **A326.03**.
- **DO NOT** harvest or feed field corn forage until 30 days after the last application.
- **DO NOT** apply this product with suspension fertilizers as the carrier.
- **DO NOT** apply product postemergence in a tank mix with emulsifiable concentrate grass herbicides, unless specifically addressed under one of the tank mix sections of this label, or injury may occur.
- **DO NOT** cultivate corn within 7 days before or after a **A326.03** application as weed control from the **A326.03** application may be reduced.

Use precautions

1. Avoid drift onto adjacent crops.

2. Severe corn injury may occur if **A326.03** is applied postemergence to corn that was treated with Counter® (terbufos, EPA Reg. No. 5481-562) or Lorsban® (chlorpyrifos, EPA Reg. No. 62719-34) in-furrow at planting, which may result in corn crop yield loss.
3. Severe corn injury may occur if **A326.03** is applied foliar postemergence in a tank mix with any organophosphate or carbamate insecticide which may result in corn crop yield loss.
4. Severe corn injury may occur if any organophosphate or carbamate insecticide is applied foliar postemergence within 7 days before or 7 days after **A326.03** application, which may result in corn crop yield loss.

Rainfastness

A326.03 requires a minimum of 1-hour rain-free period after application for best results when applied postemergence.

CULTIVATION

Cultivation immediately prior to postemergence application is not advised. Cultivation may put weeds under stress, reducing weed control. Timely cultivation 1-3 weeks after applying **A326.03** may assist weed control.

POSTEMERGENCE GROUND APPLICATION

Spray nozzles must be uniformly spaced, the same size and type, and must provide accurate and uniform application.

Apply in a spray volume of 10-30 gals./A. Use a pump that can maintain a pressure of at least 35-40 psi at the nozzles and provide proper agitation within the tank to keep the product dispersed. Lower pressures may be used with extended range or drift reduction nozzles. When weed foliage is dense, use a minimum of 15 gals.

Flat fan nozzles of 80° or 110° are advised for optimum postemergence coverage.

DO NOT use flood-jet nozzles, extremely coarse droplet nozzles, or controlled droplet application equipment for postemergence applications.

Nozzles may be angled forward 45° to enhance penetration of the crop and provide better coverage. Ensure that all in-line strainer and nozzle screens in the sprayer are 50-mesh or coarser.

Always ensure that agitation is maintained until spraying is completed, even if stopped for brief periods of time. If the agitation is stopped for more than 5 minutes, re-suspend the spray solution by running on full agitation prior to spraying.

ADJUVANT REQUIREMENTS

An adjuvant or a product containing an adjuvant is required with **A326.03** for maximum consistent performance. Only spray additives cleared for use on growing crops under 40 CFR 180.1001 may be used in spray mixture.

POSTEMERGENCE APPLICATIONS TO FIELD CORN AND SEED CORN

Add crop oil concentrate (COC) to the Postemergence application at the rate of 0.5 to 1.0 gal./100 gals. Of water (0.5% to 1.0% v/v). The use of a nonionic surfactant (NIS) at 1 qt./100 gallons of water (0.25% v/v) instead of COC is allowed, but the weed control achieved with COC is consistently better than NIS. The use of methylated seed oil (MSO) adjuvants or MSO blend adjuvants for postemergence applications of **A326.03** may cause severe crop injury. In addition to COC, always add spray grade UAN (e.g., 28-0-0) to the spray solution at a rate of 2.5% (v/v) or AMS at 8.5 lb./100 gals. of spray solution, except if precluded elsewhere on this label. If **A326.03** is being tank mixed with another registered herbicide in this situation, refer to the tank mix partner label for adjuvant precautions and restrictions and follow the most restrictive requirements. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

DO NOT use liquid fertilizer as the total carrier solution except for preplant burndown applications.

POSTEMERGENCE APPLICATIONS TO SWEET CORN AND YELLOW POPCORN

Do not add UAN or AMS when making postemergence applications of A326.03 to yellow popcorn or sweet corn, or severe crop injury may occur.

For postemergence applications to yellow popcorn and sweet corn, use nonionic surfactant (NIS) at 1 qt./100 gallons of water (0.25% v/v) to minimize the risk of crop injury. A COC may be used, and will increase the level of weed control achieved, especially under dry growing conditions, but the risk of crop injury is increased. For optimum control, the addition of atrazine is specified wherever rotational or local atrazine restrictions allow.

MIXING A326.03 ALONE

1. Add 1/4-1/2 of the required amount of clean water to the spray or mixing tank.
2. With the agitator running, add the required amount of **A326.03** to the spray tank. Continue agitation in the spray tank and allow product to fully and uniformly disperse.
3. Add the spray adjuvant and continue agitation while adding the rest of the water.
4. Maintain agitation until all of the mixture has been applied.

MIXING A326.03 IN TANK MIXTURES WITH OTHER PESTICIDES

A326.03 is compatible with most commonly used herbicides, insecticides, fungicides, and spray adjuvants. Follow WALE (Wettable/dry, Agitate, Liquids, and Emulsifiable Concentrates) mixing guidelines. BEFORE MIXING **A326.03** WITH OTHER REGISTERED PRODUCTS FOR ANY USE ON THIS LABEL, READ THE LABEL OF THE TANK MIX PARTNER TO BE CERTAIN IT IS LABELED FOR THE USE ON THE TARGET CROP AND THAT USE PATTERNS ARE COMPATIBLE WITH THOSE OF **A326.03**. When using **A326.03** in a tank mixture with other pesticides, observe the most restrictive label limitations and precautions for the products being used. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

TANK MIXING STEPS

1. Add 1/4 -1/2 of the required amount of clean water to the spray or mixing tank.
2. While maintaining agitation, continue filling the spray tank. When the tank is 3/4 full, add any dry formulation tank mix partners and allow them to completely and uniformly disperse.
3. Add the required amount of **A326.03** to the spray tank while maintaining agitation. After the product has completely and uniformly dispersed into the tank mix, add any other liquid tank mix partners and allow them to completely and uniformly disperse.
4. Add the proper amount of spray adjuvant and continue agitation while adding the remaining water.
5. Complete filling the tank with clean water and maintain sufficient agitation at all times to insure surface action until the mixture is uniform.
6. After use, thoroughly clean the sprayer according to this label (see Cleaning Spray Equipment) and any tank mix partner labels.

COMPATIBILITY TEST

A jar test is required before mixing to ensure **A326.03** compatibility with tank mix partners and adjuvants. The following test assumes a spray volume of 25 gallons per acre. For other spray volumes, make appropriate changes in the ingredient rates.

1. Add 1.0 pt. of water to each of 2 one-quart jars.

Note: Use the same source of water and the other components in the compatibility test that will actually be tank mixed and applied. It is important that all components are mixed at a temperature similar to the temperature of those used for the actual application.

2. To one of the jars, add 1/4 tsp. or 1.2 milliliters of a compatibility agent approved for this use (1/4 tsp. is equivalent to 2 pt/100 gallons spray). Shake or stir gently to mix.
3. To both jars, add the appropriate amount of herbicide(s). If more than one herbicide is used, add them separately with dry herbicides first, flowables next, and emulsifiable concentrates last. Finally, add the appropriate amount of any adjuvants that will be used. After each addition, shake or stir gently to thoroughly mix. The appropriate amount of herbicides for this test follows:
 - a. **Dry Herbicides and Adjuvants:** For each pound to be applied per acre, add 1.4 tsp. to each jar.
 - b. **Liquid Herbicides and Adjuvants:** For each pint to be applied per acre, add 0.5 tsp. or 2.5 milliliters to each jar.
4. After adding all ingredients, put lids on and tighten, and invert each jar 10 times to mix. Let the mixtures stand 15-30 minutes and look for separation, large flakes, precipitates, gels, heavy oil film on the jar, or other signs of incompatibility. Determine if a compatibility agent is needed in the spray mixture by comparing the two jars. If either mixture separates, but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility.
 - a. Slurry the dry pesticide(s) in water before addition, or
 - b. Add 1/2 the compatibility agent to the water and the other 1/2 to the emulsifiable concentrate or flowable pesticide before addition to the mixture. If incompatibility is still observed, do not use the mixture. After compatibility testing is complete, dispose of any pesticide wastes according to the Storage and Disposal section of this label.

MANDATORY SPRAY DRIFT

Ground Applications

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For all applications, applicators are required to use a coarse or coarser spray droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Boom-less Ground Applications:

- Applicators are required to use a coarse or coarser droplet size (ASABE S572.1) for all applications.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Spray Drift Advisories

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.
- **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.

- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
 - Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.
- **BOOM HEIGHT - Ground Boom**
Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.
- **SHIELDED SPRAYERS**
Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.
- **TEMPERATURE AND HUMIDITY**
When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.
- **TEMPERATURE INVERSIONS**
Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.
- **WIND**
Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.
- **Boom-less Ground Applications:**
Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.
- **Handheld Technology Applications:**
Take precautions to minimize spray drift.

CROP ROTATION RESTRICTIONS

Immediate: Corn (field, seed, sweet corn and pop)

Following harvest of corn, the following crops may be planted immediately: Asparagus, cranberry, flax, millet(pearl), grasses grown for seed (Kentucky bluegrass, perennial ryegrass, and tall fescue), oats, rhubarb, sorghum (grain and sweet), and sugarcane.

Following harvest of corn and not less than 4 months after application of A326.03, the following crops may be planted: Small grains including wheat, barley, and rye.

Following harvest of corn and not less than 10 months after application of A326.03, the following crops may be planted: Alfalfa, blueberry, canola, cotton, lingonberry, peanuts, potatoes, soybeans, sunflowers, tobacco, okra and rice can be planted back the following season. If A326.03 is applied postemergence following a mesotrione-containing preemergence herbicide, only corn (field, seed and pop) or grain sorghum may be replanted the year following application or severe crop injury may occur.

Following harvest of corn and not less than 18 months after application of A326.03, the following crops may be planted: Sugar beets, dry beans, snap beans, cucurbits, red clover, and all other rotational crops may be replanted 18 months after application of A326.03.

REPLANTING INSTRUCTIONS

If replanting is necessary in fields previously treated with **A326.03**, the field may be replanted to Corn (field, seed and pop). If tank-mix combinations were used, refer to product labels for any additional replanting instructions.

SPRAYER EQUIPMENT CLEAN-OUT

Many pesticides are very active at low rates, especially to sensitive crops. Residues left in mixing equipment, spray tanks, hoses, spray booms and nozzles can cause crop effects if they are not properly cleaned. As soon as possible after spraying **A326.03** and before using the sprayer equipment for any other applications, the sprayer equipment must be thoroughly cleaned using the following procedure. In addition, users must take appropriate steps to ensure proper equipment clean-out for any other products mixed with **A326.03** as required on the other product labels. More complete cleaning can be achieved if the spray system is cleaned immediately following the application.

1. Drain sprayer tank, hoses, spray boom and spray nozzles. Use a high-pressure detergent wash to remove physical sediment and residues from the inside of the sprayer tank and thoroughly rinse. Then, thoroughly flush sprayer hoses, spray boom and spray nozzles with a clean water rinse. Remove and clean spray tips and all filters and screens (tank, spray hose and spray tips) separately in the ammonia solution of Step 2.
2. Next, prepare a sprayer cleaning solution by adding three gallons of ammonia (containing at least 3% active) per 100 gallons of clean water. Prepare sufficient cleaning solution to allow the operation of the spray system for a minimum of 15 minutes to thoroughly flush hoses, spray boom and spray nozzles.
3. Convenient and thorough cleaning of the sprayer can be achieved if the ammonia solution or fresh water is left in the spray tank, hoses, spray booms and spray nozzles overnight or during storage.
4. Before using the sprayer, completely drain the sprayer system. Rinse the tank with clean water and flush through the hoses, spray boom, and spray nozzles with clean water. Remove and clean spray tips and all filters and screens (tank, spray hose and spray tip) separately in an ammonia solution.
5. Properly dispose of all cleaning solution and rinsate in accordance with Federal, State, and local regulations and guidelines.
6. **DO NOT** apply sprayer cleaning solutions or rinsate to sensitive crops.
7. If the sprayer sets overnight or for any extended period of time with **A326.03** spray solution, the spray tank needs to be agitated and purge the spray boom and nozzles before beginning any application.
8. If small quantities of **A326.03** remain in inadequately cleaned mixing, loading and/or spray equipment, they may be released during subsequent applications potentially causing effects to certain crops and other vegetation. To the extent consistent with applicable law, Atticus, LLC accepts no liability for any effects due to inadequately cleaned equipment.
9. When **A326.03** has been tank mixed refer to the label of the product used previously or tank mixed with **A326.03** for cleaning instructions.
10. **DO NOT** drain or flush equipment on or near desirable trees or plants.
11. **DO NOT** contaminate any body of water including irrigation water that may be used on other crops.

WEEDS CONTROLLED

A326.03 Application Alone

At the rates and weed size listed, **A326.03** controls or suppresses the weeds listed in Table 1 when the product is applied alone. Weeds larger than the size indicated in Table 1 may only be partially controlled. For best postemergence results, apply **A326.03** to actively growing weeds. Dry weather following application of **A326.03** may reduce residual weed control effectiveness. **A326.03** applied alone or in mixture with atrazine will not provide consistent or effective control of weeds identified as resistant to postemergence group 27 herbicides.

Table 1: Weeds controlled or partially controlled by postemergence activity of A326.03 herbicide

Common Name	Scientific Name	A326.03* 2.5 to 3.15 fl oz/A (0.078 to 0.098 lb ai/A)	A326.03** 2.0 to 3.15 fl oz/A + Atrazine ¹ (0.0625 to 0.098 lb ai/A)
-------------	-----------------	---	---

		Apply to Weeds < 5 Inches Tall ²	
Amaranth, palmer	<i>Amaranthus palmeri</i>	PC	C ³
Amaranth, powell	<i>Amaranthus powellii</i>	C	C
Amaranth, spiny	<i>Amaranthus spinosus</i>	C	C
Atriplex	<i>Chenopodium orach</i>	C	C
Broadleaf signalgrass	<i>Urochloa platyphylla</i>	C ³	C ³
Buckwheat, wild	<i>Polygonum convolvulus</i>	PC	C ³
Buffalobur	<i>Solanum rostratum</i>	C	C
Burcucumber	<i>Sicyos angulatus</i>	C ³	C ³
Carpetweed	<i>Mollugo verticillata</i>	C	C
Carrot, wild	<i>Daucus carota</i>	PC	C
Chickweed, common	<i>Stellaria media</i>	C	C
Cocklebur, common	<i>Xanthium strumarium</i>	C	C
Crabgrass, large	<i>Digitaria sanguinalis</i>	C ³	C ³
Dandelion	<i>Taraxacum officinale</i>	PC	PC
Dock, curly	<i>Rumex crispus</i>	PC	PC
Galinsoga	<i>Galinsoga parviflora</i>	C	C
Hemp	<i>Cannabis sativa</i>	C	C
Horsenettle	<i>Solanum carolinense</i>	PC	C
Jimsonweed	<i>Datura stramonium</i>	C	C
Horseweed (maretail)	<i>Conyza Canadensis</i>	PC	C ³
Knotweed, prostrate	<i>Polygonum aviculare</i>	PC	PC
Kochia	<i>Kochia scoparia</i>	PC	C ³
Lambsquarters, common	<i>Chenopodium album</i>	C	C
Mallow, Venice	<i>Hibiscus trionum</i>	C	C
Morningglory, entireleaf	<i>Ipomoea hederacea</i>	C ³	C
Morningglory, ivyleaf	<i>Ipomoea hederacea</i>	C ³	C
Morningglory, pitted	<i>Ipomoea lacunose</i>	C ³	C
Mustard, wild	<i>Brassica kaber</i>	C	C
Nightshade, black	<i>Solanum nigrum</i>	C	C
Nightshade, Eastern black	<i>Solanum ptycanthum</i>	C	C
Nightshade, hairy	<i>Solanum sarachoides</i>	C	C
Nutsedge, yellow	<i>Cyperus esculentus</i>	PC	PC
Pigweed, redroot	<i>Amaranthus retroflexus</i>	C	C
Pigweed, smooth	<i>Amaranthus hybridus</i>	C	C
Pigweed, tumble	<i>Amaranthus albus</i>	C	C
Pokeweed, common	<i>Phytolacca americana</i>	PC	PC
Potatoes, volunteer	<i>Solanum spp.</i>	C	C
Pusley, Florida	<i>Richardia scabra</i>	C ³	C ³
Ragweed, common	<i>Ambrosia artemisiifolia</i>	PC	C
Ragweed, giant	<i>Ambrosia trifida</i>	C ³	C
Sesbania, hemp	<i>Sesbania exaltata</i>	C	C
Sida, prickly (teaweed)	<i>Sida spinosa</i>	NC	C ³
Smartweed, lady's thumb	<i>Polygonum persicaria</i>	C ³	C
Smartweed, pale	<i>Polygonum lapathifolium</i>	C	C

Smartweed, Pennsylvania	<i>Polygonum pensylvanicum</i>	C ³	C
Sunflower, common	<i>Helianthus annuus</i>	C	C
Thistle, Canada	<i>Cirsium arvense</i>	NC	PC
Thistle, Russian	<i>Salsola kali</i>	C	C
Velvetleaf	<i>Abutilon theophrasti</i>	C	C
Waterhemp, common	<i>Amaranthus rudis</i>	C ³	C
Waterhemp, tall	<i>Amaranthus tuberculatus</i>	C ³	C

¹ A326.03 tank mixture with atrazine at a minimum rate of 0.5lb a.i./acre.

² Under certain situations weeds can be controlled at larger than listed sizes, however to protect crop yield, manage weed resistance and provide consistent control, treat weeds before they exceed 5 inches in height.

³ Apply before weed exceeds 3 inches in height. A326.03 will not provide control of weed biotypes known to be resistant to herbicide MOA group 14 and 27.

C = Control PC = Partial Control NC = Not Controlled

* A326.03 2.5 to 3.15 fl oz/A (0.078 to 0.098 lb ai contains 0.0042 – 0.0053 lb ai fluthiacet methyl + 0.074 – 0.0931 lb ai mesotrione)

** A326.03 2.0 to 3.15 fl oz/A (0.0625 to 0.098 lb ai contains 0.0034 – 0.0053 lb ai fluthiacet methyl + 0.0591 – 0.0931 lb ai mesotrione)

CROP USE DIRECTIONS

CORN

(Includes field corn, seed corn, sweet corn and yellow popcorn)

Timing and Method of Application:

Preplant Burndown

Apply A326.03 from 2 to 3.75 fl oz/acre (0.0625 to 0.117 lb a.i./A contains 0.0034 – 0.0063 lb ai fluthiacet methyl + 0.0591 – 0.111 lb ai mesotrione)) with other registered burndown herbicides in water or liquid fertilizers as a burn-down treatment to control or suppress weeds prior to planting. For improved broadleaf weed control with limited residual control prior to planting corn and before corn emergence, apply A326.03 in tank mixes with Aim® (carfentrazone-ethyl, EPA Reg. No. 279-3241), paraquat, glyphosate, glufosinate, dicamba, Sharpen® (saflufenacil, EPA Reg. No. 7969-278), and 2,4- D. Refer to individual product labels for precautionary statements, restrictions, rates, approved uses, and a list of weeds controlled. Always use the most restrictive label language when applied in a tank mix. For optimum performance make applications to actively growing weeds. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Postemergence Application

Use Rate: Apply A326.03 at 2 to 3.15 fl oz/acre (0.0625 to 0.098 lb ai contains 0.0034 – 0.0053 lb ai fluthiacet methyl + 0.0591 – 0.0931 lb ai mesotrione). Refer to weed table 1 for specific use rate specifications. A326.03 may be applied broadcast postemergence up to the V8 growth stage (or 30 inches tall). The spray boom must be maintained at a minimum of 18 inches above the crop canopy to ensure uniform spray delivery and avoid concentrating spray in corn whorls. For optimum performance, make application to actively growing weeds <5 inches tall and rosettes less than 3 inches across. Application after weeds have reached the listed maximum height for control could result in commercially unacceptable weed control. A326.03 may be tank mixed with other herbicides registered for use in corn to improve weed spectrum or general weed control unless restricted under the corn crop section.

Refer to seed company specifications for use on field corn inbred lines. Special adjuvant restrictions must be followed for postemergence applications of A326.03 in yellow popcorn or sweet corn (see the spray adjuvant specification in the Directions For Use).

Postemergence applications (after crop emergence) of **A326.03** may cause crop bleaching, leaf speckling in field corn, yellow popcorn and sweet corn hybrids. Crop bleaching, leaf speckling is typically transitory and has no effect on final yield or quality. However, herbicide sensitivity in yellow popcorn and sweet corn varies widely, and all yellow popcorn and sweet corn hybrids have not been tested. Contact your popcorn or sweet corn company, Field man, or University Specialist about hybrid specifications before making a postemergence application of **A326.03** to yellow popcorn or sweet corn.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[**Nonrefillable container.** Do not reuse or refill this container. If empty: Offer for recycling if available or discard in a sanitary landfill. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.]

[**For plastic containers ≤ 5 gallons: Nonrefillable Container:** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

[**For plastic containers > 5 gallons: Nonrefillable container.** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS**

OF LIABILITY: To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

[A326.03™] is a trademark of Atticus, LLC

[Solstice®] & Aim® is a registered trademark of FMC Corporation.

Sharpen® is a trademark of BASF.

{LANGUAGE ON LABEL AFFIXED TO CONTAINER}

MESOTRIONE	GROUP	27	HERBICIDE
FLUTHIACET METHYL	GROUP	14	HERBICIDE

A326.03™

[Alternate Brand Name: Cavallo STX]

[Contains mesotrione and fluthiacet, the active ingredients used in Solstice®.]

ACTIVE INGREDIENTS:	(% by weight)
Fluthiacet methyl.....	2.20%
Mesotrione.....	38.52%
OTHER INGREDIENTS:.....	59.28%
TOTAL	100.0%
Contains a total of 4.0 lb/gal which include 0.216 lb ai Fluthiacet methyl and 3.784 lb ai mesotrione per gallon.	

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If on skin or clothing:	<ul style="list-style-type: none"> ● Take off contaminated clothing. ● Rinse skin immediately with plenty of water for 15-20 minutes. ● Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> ● Hold eye open and rinse slowly and gently with water for 15-20 minutes. ● Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ● Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

For Chemical Emergency:

Spill, Leak, Fire, Exposure, or Accident, Call CHEMREC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS: A326.03 is very toxic to algae and moderately toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high-water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS: Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the user according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[**Nonrefillable container.** Do not reuse or refill this container. If empty: Offer for recycling if available or discard in a sanitary landfill. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.]

[**For plastic containers ≤ 5 gallons:** Nonrefillable Container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

[**For plastic containers > 5 gallons:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

See inside label booklet for additional Precautionary Statements and Directions for Use.

[[A326.03™] is not manufactured, or distributed by FMC Corporation, seller of Solstice®.]

Manufactured for:

Atticus, LLC

5000 CentreGreen Way, Suite 100

Cary, NC 27513

EPA Reg. No.: 91234-XX

EPA Est. No.: _____

NET WEIGHT: _____