



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

91234-207

Date of Issuance:

5/24/21

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

A253.01

Name and Address of Registrant (include ZIP Code):

Atticus, LLC
5000 CentreGreen Way, Suite 100
Cary, NC 27513

Katy DeGroot
Agent for Atticus, LLC
c/o Pyxis Regulatory Consulting Inc.
4110 136th St. Ct. NW
Gig Harbor, WA 98332

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Lindsay Roe, Product Manager 22
Fungicide Branch, Registration Division (7505P)

Date:

5/24/21

2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 91234-207.”
3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSF:

- Basic CSF dated 04/01/2020

The alternate brand name Esquire has been added to the registration.

If you have any questions, please contact Yasmin Bowers by phone at 703 347-8634, or via email at bowers.yasmin@epa.gov.

Enclosure

[Note to reviewer: [Text] in brackets denotes optional or explanatory language
[Note to reviewer: {Text} in braces denotes where in the final label text will appear
{BOOKLET FRONT PANEL LANGUAGE}

DIFENOCONAZOLE	GROUP 3	FUNGICIDE
----------------	---------	-----------

A253.01 [™]

[Alternate Brand Name: Esquire]

Contains Difenoconazole, the active ingredient used in [Inspire®] [and] [Top MP].

ACTIVE INGREDIENT:	(% by weight)
Difenoconazole*:	23.2%
OTHER INGREDIENTS**:	76.8%
TOTAL	100.0%

*CAS No. 119446-68-3

**Contains petroleum distillates

A253.01 is formulated as an emulsifiable concentrate (EC) containing 2.10 lb ai/gal of difenoconazole.

KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you DO NOT understand the label, find someone to explain it to you in detail.)

See inside label booklet for First Aid, Precautionary Statements and Directions for Use.


[A253.01™] Esquire is not manufactured, or distributed by Syngenta Crop Protection, LLC, seller of [Inspire®] [and] [Top MP].

EPA Reg. No.: 91234-XX

EPA Est. No.:

Net Weight:

Manufactured for:
Atticus, LLC
5000 CentreGreen Way, Suite 100
Cary, NC 27513



{LANGUAGE INSIDE BOOKLET}

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give any liquid to the person. • DO NOT give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
Contains petroleum distillates - vomiting may cause aspiration pneumonia.	

For Chemical Emergency:
Spill, Leak, Fire, Exposure, or Accident,
Call CHEMTREC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING/AVISO

Causes substantial but temporary eye injury. **DO NOT** get in eyes or on clothing. Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Mixers, loaders, applicators and all other handlers must wear:

- Chemical-resistant gloves such as barrier laminate, or Viton® ≥ 14 mils
- Protective eyewear
- Long-sleeved shirt and long pants
- Socks and shoes

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Environmental Hazards

This pesticide is toxic to fish, mammals, and aquatic invertebrates. Drift and runoff may be hazardous to aquatic estuarine/marine organisms in water adjacent to treated areas. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

DO NOT discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. **DO NOT** discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or regional office of the EPA.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product may contaminate water through drift or spray in wind. This product has a potential for runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is expected to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

Physical or Chemical Hazards

DO NOT use or store near heat or open flame. Do not mix or allow to come in contact with oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Chemical-resistant gloves such as barrier laminate, or Viton ≥ 14 mils
- Protective eyewear
- Coveralls
- Socks and shoes

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.

PRODUCT INFORMATION

A253.01 is a broad spectrum fungicide with systemic and curative properties recommended for the control of many important plant diseases. **A253.01** may be applied as a foliar spray in alternating spray programs or in tank mixes with other crop protection products. All applications must be made according to the use directions that follow.

PRODUCT USE INSTRUCTIONS

Application: Thorough coverage is necessary to provide good disease control. Make up no more spray solution than is needed for application. Avoid spray overlap, as crop injury may occur.

Efficacy: Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if maximum amount of **A253.01** has been used. If resistant isolates to Group 3 fungicides are present, efficacy can be reduced. Under high disease pressure, it is recommended to use the highest rate and shortest interval.

Rotational Crops: Please see the table below for the crop rotational restrictions.

Rotational Crop	Planting Time From Last A253.01 Application
Artichoke, Globe	0 days
Bean and Pea, Dried Shelled Subgroup 6C	
Berry, Bushberry Subgroup 13-07B	
Berry, Low Growing Subgroup 13-07G, [except Cranberry]	
Brassica (Cole) Head and Stem vegetables Group 5-16[*]	
Brassica (Cole) Leafy greens Subgroup 4-16B	
Bulb vegetables, bulb onion Subgroup 3-07A and green onion Subgroup 3-07B	
Carrots	
Chickpeas	
Citrus fruit Crop Group 10-10[*]	
Cotton Subgroup 20C	
Cucurbit vegetables Crop Group 9	
Fruit, small, vine climbing Subgroup 13-07F, except fuzzy kiwifruit[*]	
Fruiting vegetables Crop Group 8-10	
Ginseng	
Guava[*]	
Kohlrabi[*]	
Papaya[*]	
Pome fruit Crop Group 11-10[*]	
Potatoes	
Rapeseed Subgroup 20A[*]	
Rice	
Soybeans	
Stone fruit Crop Group 12-12[*]	
Strawberry	
Sugar beets	
Tomatoes and tomatillos	
Tree nuts Crop Group 14-12[*]	
Tuberous and corm vegetable subgroup 1C	
Watercress[*]	
Wild rice	
Cereals (wheat, barley, triticale, oats and rye) Root and Tuber Vegetables, Crop Group 1 (except Carrot, Sugar Beet, and Tuberous Corm Vegetable Subgroup 1C)	30 days
All other crops intended for food and feed	60 days

[*] [NOT FOR USE IN WASHINGTON]

Crop Tolerance: Plant tolerance has been found acceptable for all crops on the label; however, not all possible tank-mix combinations have been tested under all conditions. When possible, it is recommended to test the combinations on a small portion of the crop to ensure a phytotoxic response will not occur as a result of application.

SPRAY DRIFT

MANDATORY SPRAY DRIFT

Aerial Applications

- **DO NOT** release spray at a height greater than 10 ft above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For all applications, applicators are required to use a medium or coarser spray droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Ground Applications

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For all applications, applicators are required to use a medium or coarser spray droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Boom-less Ground Applications:

- Applicators are required to use a medium or coarser droplet size (ASABE S572.1) for all applications.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Spray Drift Advisories

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.
- IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher rate nozzles instead of increasing pressure.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size - Aircraft

- Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.
- BOOM HEIGHT - Ground Boom
Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.
- RELEASE HEIGHT - Aircraft
Higher release heights increase the potential for spray drift. When applying aurally to crops, **DO NOT** release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.
- SHIELDED SPRAYERS
Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.
- TEMPERATURE AND HUMIDITY
When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.
- TEMPERATURE INVERSIONS
Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.
- WIND
Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.**
Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.
- Boom-less Ground Applications:
Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.
- Handheld Technology Applications:
Take precautions to minimize spray drift.

RESISTANCE MANAGEMENT

DIFENOCONAZOLE	GROUP	3	FUNGICIDE
----------------	-------	---	-----------

For resistance management, **A253.01** contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to **A253.01** and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

DO NOT apply to plants grown for transplanting purposes.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of **A253.01** or other Group 3 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Atticus, LLC at 984-465-4754. You can also contact your pesticide distributor or university extension specialist to report resistance.

SPRAY EQUIPMENT

Nozzles

- Equip sprayers with nozzles that provide accurate and uniform application.
- Use the same size nozzles uniformly spaced across the boom.
- Calibrate sprayer before use.
- Use screens to protect the pump and to prevent nozzles from clogging.
- On suction side of pump use screens that are *16-mesh or coarser*.
- **DO NOT** place a screen in the recirculation line.
- Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles.
- Check nozzle manufacturer's recommendations.

Pump

- Use a pump with capacity to:
 - (1) Maintain sufficient pressure at the nozzle tip to give the required flow rate and droplet size to provide acceptable coverage of the target crop.
 - (2) Provide sufficient agitation in tank to keep mixture in suspension – this requires recirculation of 10% of tank volume per minute.
- Use a jet agitator or liquid sparge tube for agitation.
- **DO NOT** air sparge.

For more information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state agricultural recommendations.

MIXING INSTRUCTIONS

- Prepare no more spray mixture than is required for the immediate operation.
- Thoroughly clean spray equipment before using this product.
- Agitate the spray solution before and during application.
- Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

A253.01 Alone (no tank mix):

- Add 1/2-2/3 of the required amount of water to the spray or mixing tank.
- With the agitator running, add **A253.01** to the tank.

- Continue agitation while adding the remainder of the water.
- Begin application of the spray solution after **A253.01** has completely dispersed into the mix water.
- Maintain agitation until all of the mixture has been sprayed.

A253.01 + Tank Mixtures: **A253.01** is usually compatible with tank-mix partners. To determine the physical compatibility of **A253.01** with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 qt of water. Add wettable powders and water dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Mixing in the Spray Tank

- Add 1/2-2/3 of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank-mix partner(s) into the tank in the same order as described above.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and **A253.01** to the spray tank.
- Allow **A253.01** to completely disperse.
- Spray the mixture with the agitator running.
- Observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label.
- Label dosage rate must not be exceeded, and the most restrictive label precautions and limitations must be followed.
- This product must not be mixed with any product which prohibits such mixing.
- It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

APPLICATION PROCEDURES

Ground Application:

- Apply in a minimum of 10 gal of water per acre, unless specified otherwise.

Aerial Application:

- Use only on crops where aerial applications are indicated.
- Apply in a minimum of 5 gal of water per acre, unless specified otherwise. For pome fruit apply in a minimum of 10 gal of water per acre.
- **DO NOT** apply under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur.
- **DO NOT** apply directly to humans or animals.

Application Through Irrigation Systems (Chemigation)

- Use only on crops for which chemigation is specified on this label.
- Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. **DO NOT** apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- Apply in 0.125-0.25 inches/A of water. Excessive water may reduce efficacy.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

- **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Note: **DO NOT** inject **A253.01** at full strength or deterioration of valves and seals may occur. Use a dilution ratio of at least 10 parts water to 1-part **A253.01**. **A253.01** is corrosive to many seal materials. Leather seals are best. EPDM or silicone rubber seals can be used but should be replaced once a year. **DO NOT** use Viton, Buna-N, Neoprene, or PVC seals.

Operating Instructions

1. The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. **DO NOT** apply when wind speed favors drift beyond the area intended.

Center Pivot Irrigation Equipment

Notes: (1) Use only with drive systems which provide uniform water distribution. (2) **DO NOT** use end guns when chemigating **A253.01** through center pivot systems because of non-uniform application.

- Determine the size of the area to be treated.
- Determine the time required to apply 1/8-1/2 inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as recommended by the equipment manufacturer. When applying **A253.01** through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution. Run the system at 80-95% of the manufacturer's rated capacity.
- Using water, determine the injection pump output when operated at normal line pressure.
- Determine the amount of **A253.01** required to treat the area covered by the irrigation system.
- Add the required amount of **A253.01** and sufficient water to meet the injection time requirements to the solution tank.
- Make sure the system is fully charged with water before starting injection of the **A253.01** solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
- Maintain constant solution tank agitation during the injection period.
- Continue to operate the system until the **A253.01** solution has cleared from the last sprinkler head.

Solid Set, Hand Move, and Moving Wheel Irrigation Equipment

- Determine the acreage covered by the sprinklers.

- Fill injector solution tank with water and adjust flow rate to use the contents over a 20 to 30-minute interval. When applying **A253.01** through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution.
- Determine the amount of **A253.01** required to treat the area covered by the irrigation system.
- Add the required amount of **A253.01** into the same quantity of water used to calibrate the injection period.
- Operate the system at the same pressure and time interval established during the calibration.
- Stop injection equipment after treatment is completed. Continue to operate the system until the **A253.01** solution has cleared the last sprinkler head.

SPECIFIC INSTRUCTIONS FOR PUBLIC WATER SYSTEMS

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

7. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

SPECIFIC USE DIRECTIONS

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Almonds[*]	Alternaria leafspot <i>(A. alternata)</i> Anthracnose <i>(Colletotrichum acutatum)</i> Blossom Blight Brown rot <i>(Monilinia spp.)</i>	5 – 6.95 (0.082 – 0.114) [6.95 (0.114) (CA Only)]	Begin applications prior to disease onset when conditions are conducive for disease. Apply A253.01 on a 14- to 21-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
	Brown rot/hull rot (<i>Monilinia</i> spp.) Powdery mildew (<i>Podosphaera tridactyla</i> , <i>Sphaerotheca pannosa</i>) Scab (<i>Venturia carpophila</i>) Shot hole (<i>Wilsonmyces carpophilus</i>)		<p>If monitoring or history indicates the presence of Alternaria, apply 6.95 fl oz/A (0.114 lb ai) of A253.01 in the late spring (mid-April to beginning of May) and then repeat the treatment 2-3 weeks later.</p> <p>For blossom blight, begin applications at early bloom and continue through petal fall.</p> <p><i>[Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.]</i></p> <p><i>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.]</i></p> <p><i>[Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]</i></p> <p>Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications use a minimum of 10 gal/A of water.</p>

Specific Use Restrictions:

- 1) **DO NOT** apply more than 27.8 fl oz (0.46 lb ai) /A/year of **A253.01**.
- 2) **DO NOT** apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole).
- 3) **DO NOT** make more than 4 applications per year at the maximum application rate.
- 4) Minimum Application Interval: 14 days.
- 5) **DO NOT** apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 6) **DO NOT** apply within 14 days of harvest (14-day PHI).

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Artichoke, Globe[*]	Ramularia leaf spot Ramularia bud spot (<i>R. cynarae</i>)	6.95 (0.114)	<p>Begin applications prior to disease onset when conditions are conducive for disease. Apply A253.01 on a 14-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p> <p>Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by ground, chemigation, or aerial application. For ground applications, apply in 50-200 gallons of water per acre to obtain coverage without excessive runoff. For aerial applications Use a minimum of 10 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.</p> <p>Specific Use Restrictions:</p> <ol style="list-style-type: none"> 1) DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01.

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
2) DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 3) DO NOT make more than 4 applications per year. 4) Minimum Application Interval: 14 days. 5) DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 6) DO NOT apply A253.01 within 3 days of harvest (3-day PHI).			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Bean and Pea, Dried Shelled (except soybean) Subgroup 6C[*] To be grown for bean, dried seed only Phaseolus Vigna Pisum Lupinus See complete list below See specific directions for soybeans and chickpea	Anthranose (<i>Colletotrichum lindemuthianum</i>) Alternaria leaf spot (<i>A. alternata</i>) Alternaria blight (<i>Alternaria</i> spp.) Ascochyta leaf and pod spot (<i>Ascochyta</i> spp.) Ascochyta blight (<i>Mycosphaerella pinodes</i>) Cercospora leaf spot (<i>Cercospora cruenta</i>)	6.95 (0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply A253.01 on a 14-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.

Application: For best results, sufficient water volume must be used to provide thorough coverage. **A253.01** can be applied by ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications use a minimum of 10 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Complete List of Bean and Pea, dried shelled (except soybean) – Subgroup 6C: Dried cultivars of bean (*Lupinus*); bean (*Phaseolus*) (includes field bean, kidney bean, lima bean, navy bean, pinto bean, tepary bean); bean (*Vigna*) (includes adzuki bean, blackeyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean); broad bean; guar; lablab bean; lentil; pea (*Pisum*) (includes field pea); pigeon pea

Specific Use Restrictions:

- DO NOT** apply more than 27.8 fl oz (0.46 lb ai)/A/year of **A253.01**.
- DO NOT** apply more than 14 fl oz/A/year of **A253.01** (0.230 lb ai/A/year) for pea vines and hay.
- DO NOT** apply more than 6.95 fl oz per acre per application. (0.114 lb ai of difenoconazole)
- DO NOT** make more than 4 applications year.
- DO NOT** apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- DO NOT** feed or harvest cowpeas forage and hay.
- DO NOT** apply **A253.01** within 14 days of harvest (14-day PHI)

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Berry, Low Growing Subgroup 13-07G[*] Strawberry[*] See additional crops below See separate Instructions for cranberry	Anthracnose <i>(Colletotrichum acutatum)</i> Leaf spot <i>(Cercospora fragariae)</i> Powdery mildew <i>(Sphaerotheca macularis)</i> Leaf rust <i>(Phragmidium potentillae)</i>	5.5 – 6.95 (0.090 – 0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply A253.01 on a 7- to 14-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. <i>[Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.]</i> <i>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.]</i> <i>[Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]</i>
Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by ground, chemigation or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Additional Low Growing Berries: Bearberry, Bilberry, Cloudberry, Muntries, Partridgeberry, Strawberry including all cultivars and/or hybrids of these.			
Specific Use Restrictions: 1) DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01 . 2) DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 3) DO NOT make more than 4 applications per year at the maximum application rate. 4) Minimum Application Interval: 7 days. 5) DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 6) A253.01 may be applied up to the day of harvest (0-day PHI).			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Brassica Head and Stem Vegetable Crop group 5-16[*] Broccoli	Alternaria diseases <i>(Alternaria spp.)</i> Anthracnose <i>(Colletotrichum higginsianum)</i> Cercospora	5 – 6.95 (0.082 – 0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply A253.01 on a 7- to 10-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Brussels sprouts Cabbage Cauliflower See additional crops below Including all Cultivars and/or hybrids of these	leafspot (<i>C. brassicicola</i>) Powdery mildew (<i>Erysiphe polygoni</i>)		<i>[Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.]</i> <i>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.]</i> <i>[Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]</i>
Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Complete list of Brassica Head and Stem Vegetables Crop group 5-16 including: Broccoli; Brussels sprouts; cabbage; cabbage, Chinese, napa; cauliflower; cultivars, varieties, and hybrids of these commodities			
Specific Use Restrictions: 1) DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01 . 2) DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 3) DO NOT make more than 4 applications per year at the maximum application rate. 4) Minimum Application Interval: 7 days. 5) DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 6) DO NOT apply within 1 day of harvest (1-day PHI).			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Brassica Leafy Greens Subgroup 4-16B[*] Arugula Chinese cabbage Bok Choy Collards Kale Mustard greens Turnip greens See additional crops below See separate	Alternaria diseases (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum higginsianum</i>) Cercospora leafspot (<i>C. brassicicola</i>) Powdery mildew (<i>Erysiphe polygoni</i>)	5 – 6.95 (0.082 – 0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply A253.01 on a 7- to 10-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. <i>[Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.]</i> <i>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.]</i>

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
instructions for watercress			[Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]
	Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.		
Complete list of Brassica Leafy Greens Vegetable subgroup 4-16B including: Arugula; broccoli raab; broccoli, Chinese; cabbage, abyssinian; cabbage, seakale; cabbage, Chinese, bok choy; collards; cress, garden; cress, upland; hanover salad; kale; maca, leaves; mizuna; mustard greens; radish, leaves; rape greens; rocket, wild; shepherd's purse; turnip greens; cultivars, varieties, and hybrids of these commodities.			
Specific Use Restrictions: 1) DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01 . 2) DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 3) DO NOT make more than 4 applications per year at the maximum application rate. 4) Minimum Application Interval: 7 days. 5) DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 6) DO NOT apply within 1 day of harvest (1-day PHI).			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Bulb Vegetables[*]	Cercospora leafspot (<i>C. duddiae</i>)	5 – 6.95 (0.082 – 0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply A253.01 on a 7- to 10-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.
Bulb Onion Subgroup 3-07A Onions, bulb Garlic Shallot	Leaf blotch (<i>Cladosporium allii-cepae</i>) Powdery Mildew (<i>Leveillula taurica</i>) Purple blotch (<i>Alternaria porri</i>)		
Green Onion Subgroup 3-07B Onions, green Leek Welsh onion tops	Stemphyllium leaf blight (<i>S. vesicarium</i>) Cladosporium leaf blotch (<i>C. allii</i>) Rust (<i>Puccinia allii</i>)		[Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.] [Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.] [Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]
Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
<p>Bulb onion subgroup 3-07A: Daylily, bulb; fritillaria, bulb; garlic, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; lily, bulb; onion, bulb; onion, Chinese, bulb; onion, pearl; onion, potato, bulb; shallot, bulb; cultivars, varieties, and/or hybrids of these.</p> <p>Green onion subgroup 3-07B: Chive, fresh leaves; chive, Chinese, fresh leaves; elegans hosta; fritillaria, leaves; kurrat; lady's leek; leek; leek, wild; onion, Beltsville bunching; onion, fresh; onion, green; onion, macrostem; onion, tree, tops; onion, Welsh, tops; shallot, fresh leaves; cultivars, varieties, and/or hybrids of these.</p>			
<p>Specific Use Restrictions:</p> <ol style="list-style-type: none"> 1) For green onions, DO NOT apply more than 20.85 fl oz (0.34 lb ai)/A/year of A253.01. 2) For dry bulb onions, DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01. 3) DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 4) For green onions, DO NOT make more than 3 applications per year at the maximum application rate. 5) For dry bulb onions, DO NOT make more than 4 applications per year at the maximum application rate. 6) Minimum Application Interval: 7 days. 7) DO NOT apply more than 0.34 lb ai/A/year of difenoconazole-containing products to green onions. 8) DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products to dry bulb onions. 9) DO NOT apply within 7 days of harvest (7-day PHI). 			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Carrots[*]	<p>Cercospora leaf spot (<i>Cercospora carotae</i>)</p> <p>Alternaria leaf blight (<i>Alternaria dauci</i>)</p> <p>Powdery mildew (<i>Erysiphe</i> spp.)</p>	5.5 – 6.95 (0.090 – 0.114)	<p>Begin applications prior to disease onset when conditions are conducive for disease. Apply A253.01 on a 7- to 10-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p> <p><i>[Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.]</i></p> <p><i>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.]</i></p> <p><i>[Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]</i></p>
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.</p>			
<p>Specific Use Restrictions:</p> <ol style="list-style-type: none"> 1) DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01. 2) DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 			

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
3) DO NOT make more than 4 applications per year at the maximum application rate. 4) Minimum Application Interval: 7 days 5) DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 6) DO NOT apply within 7 days of harvest (7-day PHI).			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Chickpea[*]	Alternaria blight <i>(A. Alternata)</i> Ascochyta blight <i>(A. Rabiei)</i> Powdery mildew <i>(Leveillula taurica)</i> Rust <i>(Uromyces ciceris-arietini)</i>	5.5 – 6.95 (0.090 – 0.114)	Begin applications prior to disease onset when conditions are conducive for disease. The minimum retreatment interval is 14 days. Make A253.01 applications at least 14 days apart, making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. <i>[Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.]</i>
	Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.		

Specific Use Restrictions:

- 1) **DO NOT** apply more than 27.8 fl oz (0.46 lb ai)/A/year of **A253.01**.
- 2) **DO NOT** apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole).
- 3) **DO NOT** make more than 4 applications per year at the maximum application rate.
- 4) **Minimum Application Interval:** 14 days.
- 5) **DO NOT** apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 6) **DO NOT** apply within 14 days of harvest (14-day PHI).

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Citrus Fruit Crop Group 10-10[*] Grapefruit Lemon Lime Orange (sour and sweet)	Greasy spot <i>(Mycosphaerella citri)</i> Melanose <i>(Diaporthe citri)</i> Scab <i>(Elsinoe fawcettii)</i> Sweet orange scab <i>(Elsinoe australis)</i>	5 – 7.6 (0.082 – 0.125)	A253.01 applications should begin prior to disease development and continue throughout the season on 7- to 21-day intervals following the resistance management guidelines. Applications may be made by ground or air. An adjuvant may be added at recommended rates. A horticultural spray oil should be used to improve control of greasy spot.

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Tangerine Including all cultivars and/or hybrids of these See additional crops in this subgroup below	Albinism (<i>Alternaria alternata</i> pv <i>citri</i>) Post bloom fruit drop (PFD) (<i>Colletotrichum acutatum</i>) Alternaria leaf and fruit spot (<i>Alternaria citri</i>) Diplodia stem-end rot (<i>Diplodia natalensis</i>) Phomopsis stem-end rot (<i>Phomopsis citrii</i>) Anthracnose (<i>Colletotrichum</i> spp.) Black spot (<i>Guignardia citricarpa</i>)		[Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.] [Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.] [Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]
Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.			
Citrus Fruit subgroup: Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citrus citron; citrus hybrids (includes chironja, tangelo, tangor); grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; mount white lime; New Guinea wild lime; mandarin (tangerine); orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangor; trifoliate orange; unqi fruit; cultivars, varieties, and/or hybrids of these.			
Specific Use Restrictions: 1) DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01 . 2) DO NOT apply more than 7.6 fl oz per application (0.125 lb ai of difenoconazole). 3) DO NOT make more than 3 applications per year at the maximum application rate. 4) Minimum Application Interval: 7 days. 5) DO NOT apply more than 0.5 lb ai/A/year of difenoconazole-containing products. 6) A253.01 may be applied up to the day of harvest (0-day PHI).			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Cotton (Subgroup 20C)[*]	Alternaria leafspot (<i>Alternaria</i> spp.) Areolate mildew (<i>Ramularia gossypii</i>) Leafspots and blights (<i>Alternaria</i> spp., <i>Ascochyta gossypii</i> , <i>Cercospora</i> spp., <i>Stemphyllium</i> spp.)	5.0 – 6.95 (0.082 – 0.114)	For best activity, apply A253.01 prior to or early in the disease development. An adjuvant may be added at specified rates. For foliar disease control, the first application should be targeted approximately at pin-head square to first bloom or when conditions are conducive for disease development. For best control of target spot, adjust the GPA to ensure coverage of upper and lower leaves. Subsequent applications may be made on a 14-21 day interval.

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
	Target spot (<i>Corynespora cassicola</i>)		DO NOT apply more than two sequential applications before alternating to a fungicide with a different mode of action.

Application: For best results, sufficient water volume must be used to provide thorough coverage. **A253.01** can be applied by ground, chemigation, or aerial application. For aerial applications, use a minimum of 5 gal/A of water. Applicators must use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) **DO NOT** apply more than 20.7 fl oz (0.34 lb ai)/A/year of **A253.01**.
- 2) **DO NOT** apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole).
- 3) **DO NOT** make more than 3 applications per year at the maximum application rate.
- 4) **Minimum Application Interval:** 14 days.
- 5) **DO NOT** apply more than 0.34 lb ai/A/year of difenoconazole-containing products.
- 6) **DO NOT** apply **A253.01** within 45 days of harvest (45-day PHI).

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Cranberry[*]	Bitter rot (<i>Colletotrichum gloeosporioides</i>) Blotch rot (<i>Physalospora vacciniae</i>) Cottonball (<i>Monilinia oxycocci</i>) Leaf rust (<i>Pucciniastrum vaccinii</i>) Ripe rot (<i>Coleophoma empetri</i>)	6.95 (0.114)	For best activity, apply A253.01 prior to or early in the disease development. An adjuvant may be added at specified rates. Apply on a 7-14 day interval. DO NOT apply more than two sequential applications before alternating to a fungicide with a different mode of action.

Application: For best results, sufficient water volume must be used to provide thorough coverage. **A253.01** can be applied by ground, chemigation, or aerial application. For aerial applications, use a minimum of 5 gal/A of water. Applicators must use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) **DO NOT** apply more than 20.85 fl oz (0.34 lb ai)/A/year of **A253.01**.
- 2) **DO NOT** apply more than 0.34 lb ai/A/year of difenoconazole-containing products.
- 3) **DO NOT** apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole).
- 4) **DO NOT** make more than 3 applications per year.
- 5) **Minimum Application Interval:** 7 days.
- 6) **DO NOT** allow release of irrigation or flood water for at least 7 days after the last application.
- 7) **DO NOT** apply when weather conditions favor drift from treated areas to non-target aquatic habitat.
- 8) **DO NOT** treat fields used for aquaculture of fish or crustacean.
- 9) **DO NOT** drain water from treated fields into ponds used for aquaculture of fish or crustacean.
- 10) **DO NOT** use water drained from treated field to irrigate other crops.
- 11) **DO NOT** apply **A253.01** within 30 days of harvest (30-day PHI).

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate	Application Instructions
------	-----------------	----------	--------------------------

		fl oz product/A (lb ai)	
Cucurbit Vegetables Crop Group 9[*] Cantaloupe Cucumber Honeydew Muskmelon Watermelon Pumpkin Squash Zucchini Including cultivars and/or hybrids of these See additional cucurbit crops below	Powdery mildew (<i>Sphaerotheca fuliginea</i> , <i>Erysiphe cichoracearum</i>)	5 – 6.95 (0.082 – 0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply 253.01 on a 7- to 10-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. [Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.] [Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.]
	Alternaria leaf blight (<i>A. cucumerina</i>) Alternaria leaf spot (<i>A. alternata</i>) Anthracnose (<i>Colletotrichum orbiculare</i>) Cercospora leafspot (<i>C. citrullina</i>) Gummy stem blight (<i>Didymella bryoniae</i>)	fl oz/1000 sq ft	
	Septoria leaf blight (<i>S. cucurbitacearum</i>) Plectosporium blight (<i>P. tabacinum</i>) Phoma blight (<i>P. exigua</i>) Phyllosticta leafspot (<i>P. cucurbitacearum</i>)	0.13 – 0.16	[Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.] Greenhouse Use for Cucumber only: For production in covered areas, use A253.01 for no more than 50% of sprays per crop. Rotate with other registered products with different modes of action (FRAC codes).
Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications (20 for gummy stem blight). For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Complete list of cucurbit crops: Chayote (fruit); Chinese waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); <i>Momordica</i> spp. (includes balsam apple, balsam pear, bittermelon, Chinese cucumber); muskmelon (includes cantaloupe); pumpkin; squash, summer; squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon			
Specific Use Restrictions: 1) Greenhouse use is only for cucumber. <ul style="list-style-type: none"> • DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01 for greenhouse use. • DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 2) Field Use: <ul style="list-style-type: none"> • DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01. • DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 3) DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 4) DO NOT make more than 4 applications per year at the maximum application rate. 5) Minimum Application Interval: 7 days. 6) A253.01 may be applied up to the day of harvest (0-day PHI).			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Filberts[*] (Hazelnuts)	Eastern Filbert Blight (<i>Anisogramma anomala</i>)	5 – 6.95 (0.082 – 0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply A253.01 on a 14- to 21-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. [Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.] [Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.] [Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]
Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.			
Specific Use Restrictions: 1) DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01 . 2) DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 3) DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 4) DO NOT make more than 4 applications per year at the maximum application rate. 5) Minimum Application Interval: 14 days. 6) DO NOT apply within 14 days of harvest (14-day PHI).			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Fruiting Vegetables Crop Group 8-10[*] Eggplant Groundcherry Pepino Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper) Tomatillo Tomato	Early blight (<i>Alternaria solani</i>) Black mold (<i>A. alternata</i>) Gray leafspot (<i>Stemphylium botryosum</i>) Powdery mildew (<i>Leveillula taurica</i>) Septoria leafspot (<i>S. lycopersici</i>) Target spot (<i>Corynespora cassiicola</i>) Anthracnose (<i>Colletotrichum</i> spp.)	4 – 6.95 (0.066 – 0.114)	Begin applications at first sign of disease or when conditions are conducive for disease development. Apply A253.01 on a 7- to 14-day schedule. A253.01 can be used in blocking program using a maximum of two consecutive applications before rotating to fungicides with another mode of action that are registered for these diseases. If disease pressure is high, use the highest rate and shortest interval. [Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.]

See additional crops in this subgroup below	Leaf mold (<i>Fulvia fulva</i>)	[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.] [Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]
	Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 5 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.	
Fruiting Vegetables subgroup: African eggplant; bush tomato; bell pepper; cocona; currant tomato; eggplant; garden huckleberry; goji berry; groundcherry; martynia; naranjilla; okra; pea eggplant; pepino; pepper, bell; pepper, nonbell; roselle; scarlet eggplant; sunberry; tomatillo; tomato; tree tomato; cultivars, varieties, and/or hybrids of these.		
Specific Use Restrictions: 1) DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01 . 2) DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 3) DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 4) DO NOT make more than 4 applications per year at the maximum application rate. 5) Minimum Application Interval: 7 days. 6) A253.01 may be applied up to the day of harvest (0-day PHI).		

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Ginseng[*]	Alternaria blight (<i>A. Panax</i>) Powdery mildew (<i>Erysiphe</i> spp.)	6.95 (0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply A253.01 on a 7- to 14-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.
Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by ground, chemigation, or aerial application. Use a minimum of 15 gal/A for ground applications. For aerial applications, use a minimum of 10 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions: 1) DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01 . 2) DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 3) DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 4) DO NOT make more than 4 applications per year. 5) Minimum Application Interval: 7 days. 6) A253.01 may be applied up to the day of harvest (0-day PHI).			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
<p>Grapes[*] (except Concord, Concord Seedless, and Thomcord. See Precaution under Application Instructions)</p> <p>(Fruit, small, vine climbing, except fuzzy kiwifruit – subgroup 13-07F)[*]</p> <p>See additional crops in this subgroup below</p>	<p>Alternaria rot (<i>A. alternata</i>)</p> <p>Powdery mildew (<i>Uncinula necator</i>)</p> <p>Rotbrenner (<i>Pseudopezicula tracheiphila</i>)</p> <p>Septoria leaf spot (<i>S. ampelina</i>)</p> <p>Black Rot (<i>Guignardia bidwellii</i>)</p> <p>Angular leafspot (<i>Mycosphaerella angulata</i>)</p> <p>Anthraco-nose (<i>Elsinoe ampelina</i>)</p> <p>Leaf Blight (<i>Pseudocercospora vitis</i>)</p>	<p>5 – 6.95 (0.082 – 0.114)</p>	<p>For powdery mildew, begin at bud break and apply on a 10- to 21-day interval, making no more than 2 sequential applications before alternating to a fungicide with a different mode of action.</p> <p>For Black rot - begin when shoot length is 1-3 inches and continue on a 10-day interval making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p> <p>For all other diseases, begin applications prior to disease onset when conditions are conducive for disease. Apply A253.01 on a 10- to 21-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p> <p><i>[Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.]</i></p> <p><i>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.]</i></p> <p><i>[Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]</i></p> <p>PRECAUTION: On <i>V. labrusca</i>, <i>V. labrusca</i> hybrids and other non-vinifera hybrids where sensitivity is not known, the use of A253.01 by itself or in tank mixtures with materials that may increase uptake (adjuvants, foliar fertilizers) may result in leaf burning or other phytotoxic effects.</p>
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.</p>			
<p>Complete list of small fruit vine climbing, except fuzzy kiwifruit, subgroup 13-07F: Amur river grape; gooseberry; grape; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and/or hybrids of these</p>			
<p>Specific Use Restrictions:</p> <ol style="list-style-type: none"> DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01. DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products. DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 			

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
4) DO NOT make more than 4 applications per year at the maximum application rate. 5) Minimum Application Interval: 10 days. 6) DO NOT apply within 7 days of harvest (7-day PHI).			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Guava[*]	Alternaria Fruit Rot <i>(Alternaria spp.)</i> Anthracnose <i>(Colletotrichum gloeosporioides)</i> <u>Suppression</u> Rust <i>(Puccinia psidii)</i>	6.95 (0.114)	For best activity, apply A253.01 prior to or early in the disease development. An adjuvant may be added at specified rates. Apply on a 7-14 day interval. DO NOT apply more than two sequential applications before alternating to a fungicide with a different mode of action.

Application: For best results, sufficient water volume must be used to provide thorough coverage. **A253.01** can be applied by ground, chemigation, or aerial application. For aerial applications, use a minimum of 10 gal/A of water. Applicators must use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) **DO NOT** apply more than 27.8 fl oz (0.46 lb ai)/A/year of **A253.01**.
- 2) **DO NOT** apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) **DO NOT** apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole).
- 4) **DO NOT** make more than 4 applications per year.
- 5) **Minimum Application Interval:** 7 days.
- 6) **A253.01** may be applied the day of harvest. (0-day PHI).

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Kohlrabi[*]	Alternaria diseases <i>(Alternaria spp.)</i> Anthracnose <i>(Colletotrichum higginsianum)</i> Cercospora leafspot <i>(C. brassicicola)</i> Powdery mildew <i>(Erysiphe polygoni)</i>	5 – 6.95 (0.082 – 0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply A253.01 on a 7- to 10-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. <i>[Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.]</i> <i>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.]</i>

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
			[Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]
	<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.</p>		
<p>Specific Use Restrictions:</p> <ol style="list-style-type: none"> 1) DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01. 2) DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 3) DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 4) DO NOT make more than 4 applications per year at the maximum application rate. 5) Minimum Application Interval: 7 days. 6) DO NOT apply within 1 day of harvest (1-day PHI). 			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Papaya[*]	Blossom blight and fruit rot <i>(Colletotrichum gloeosporioides)</i> Alternaria fruit spot <i>(A. alternata)</i> Powdery Mildew <i>(Erysiphe spp.)</i> Brown Spot <i>(Corynespora cassicola)</i>	6.95 (0.114)	For best activity, apply A253.01 prior to or early in the disease development. An adjuvant may be added at specified rates. Apply on a 7-14 day interval. DO NOT apply more than two sequential applications before alternating to a fungicide with a different mode of action.
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by ground, chemigation, or aerial application. For aerial applications, use a minimum of 10 gal/A of water. Applicators must use care in making applications near non-target aquatic habitats.</p>			
<p>Specific Use Restrictions:</p> <ol style="list-style-type: none"> 1) DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01. 2) DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 3) DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 4) DO NOT make more than 4 applications per year. 5) Minimum Application Interval: 7 days. 6) A253.01 may be applied the day of harvest. (0-day PHI). 			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Pecans[*]	Pecan Scab (<i>Cladosporium caryigenum</i>) Downy Spot (<i>Mycosphaerella caryigena</i>) Liver Spot (<i>Gnomonia caryae</i>) Vein Spot (<i>Gnomonia nerviseda</i>) Zonate Leaf Spot (<i>Grovesinia pyramidalis</i>) Powdery Mildew (<i>Microsphaera penicillata</i>)	5 – 6.95 (0.082 – 0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply A253.01 on a 14-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. [Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.]
	Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.		

Specific Use Restrictions:

- 1) **DO NOT** apply more than 27.8 fl oz (0.46 lb ai)/A/year of **A253.01**.
- 2) **DO NOT** apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) **DO NOT** apply more than 6.95 fl oz per application (0.113 lb ai of difenoconazole).
- 4) **DO NOT** make more than 4 applications per year at the maximum application rate.
- 5) **Minimum Application Interval:** 14 days.
- 6) **DO NOT** apply within 14 days of harvest (14-day PHI).

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Pistachios[*]	Panicle and Shoot Blight (<i>Botryosphaeria dothidea</i>) Alternaria late blight (<i>Alternaria</i> spp.) Septoria leaf spot (<i>S. pistaciarum</i>)	5 – 6.95 (0.082 – 0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply A253.01 on a 14- to 21-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. For <i>Botryosphaeria</i> , begin applications when green leaf tissue becomes visible and continue on a 14- to 21-day interval. Make no more than 2 sequential applications before alternating to another fungicide with a different mode of action. [Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
			<p>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.]</p> <p>[Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]</p>
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.</p>			
<p>Specific Use Restrictions:</p> <ol style="list-style-type: none"> 1) DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01. 2) DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 3) DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 4) DO NOT make more than 4 applications per year at the maximum application rate. 5) Minimum Application Interval: 14 days. 6) DO NOT apply within 14 days of harvest (14-day PHI). 			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
<p>Pome Fruit Crop Group 11-10[*]</p> <p>Apple</p> <p>Crabapple</p> <p>Loquat</p> <p>Mayhaw</p> <p>Pear</p> <p>Pear, Oriental</p> <p>Quince</p> <p>See additional crops in this subgroup below</p>	<p>Alternaria blotch (<i>Alternaria</i> spp.)</p> <p>Brooks fruit spot (<i>Mycosphaerella pomii</i>)</p> <p>Cedar apple rust (<i>Gymnosporangium juniperi-virginianae</i>)</p> <p>Flyspeck (<i>Zygophiala jamacaicensis</i> (Formerly known as <i>Schizothyrium pomii</i>)</p> <p>Powdery mildew (<i>Podosphaera leucotricha</i>)</p> <p>Quince rust (<i>Gymnosporangium</i> spp.)</p> <p>Scab (<i>Venturia</i> spp.)</p> <p>Sooty blotch (<i>Gloeodes pomigena</i>)</p>	<p>3.97 (0.065)</p>	<p>Apple Scab - Protective Spray Schedule: Apply every 7-10 days starting at 1/4-1/2 inch green tip or when environmental conditions become conducive for scab. Continue through petal fall until the threat of primary scab is complete. For improved fruit scab control, combine A253.01 with a protectant fungicide beginning at bloom.</p> <p>Apple Scab - Curative Spray Schedule: Use a forecasting system beginning at green tip. Apply within 48 hours of the onset of an infection period. Apply a follow-up spray in 7 days. For improved fruit scab control, combine A253.01 with a protectant fungicide beginning at bloom.</p> <p>Powdery Mildew: Begin applications of A253.01 at tight cluster, and continue on a 7- to 10-day schedule. Follow A253.01 with other fungicides as needed.</p> <p>Cedar-Apple Rust, Quince Rust, and Brooks Fruit Spot: Begin applications preventively. Apply A253.01 alone or in combination with a</p>

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
			<p>protectant fungicide on a 7- to 10-day schedule through the second cover-spray.</p> <p>Sooty Blotch, Flyspeck: For early season control, begin applications preventively. Apply A253.01 alone or in combination with a protectant fungicide on a 7- to 10-day schedule through the second cover spray. Follow with other fungicides as needed.</p>
	<p>Resistance Management: To help prevent resistance, make no more than 2 consecutive applications with A253.01 or another Group 3 fungicide or a Group 3 fungicide mixture before alternating to a different mode of action.</p> <p>Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground or aerial application. Use a minimum of 40 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.</p>		
<p>Pome Fruit subgroup: Apple; azarole; crabapple; loquat; mayhaw; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these.</p>			
<p>Specific Use Restrictions:</p> <ol style="list-style-type: none"> DO NOT apply more than 19.85 fl oz (0.33 lb ai)/A/year of A253.01. DO NOT apply more than 0.33 lb ai/A/year of difenoconazole-containing products. DO NOT apply more than 3.97 fl oz per application (0.065 lb ai of difenoconazole). DO NOT make more than 5 applications per year. Minimum Application Interval: 7 days. DO NOT apply A253.01 within 14 days of harvest (14-day PHI). 			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
<p>Potatoes Vegetable, Tuberous and corm subgroup 1C[*]</p> <p>For other crops in this subgroup, please see below</p>	<p>Black dot (<i>Colletotrichum coccodes</i>)</p> <p>Brown spot (<i>Alternaria alternata</i>)</p> <p>Early blight (<i>Alternaria solani</i>)</p> <p>Powdery mildew (<i>Erysiphe cichoracearum</i>)</p> <p>Septoria leafspot (<i>Septoria</i> spp.)</p>	<p>4 – 6.95 (0.066 – 0.114)</p>	<p>Begin applications at first sign of disease or when conditions are conducive for disease development. Apply A253.01 on a 7- to 14-day schedule. A253.01 can be used in blocking program using a maximum of 2 consecutive applications before rotating to fungicides with another mode of action that are registered for these diseases.</p> <p>[Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.]</p> <p>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.]</p>

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
			[Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]
	<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 5 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.</p>		
<p>Vegetables, tuberous and corm, subgroup: Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna, Cassava (bitter and sweet), Chayote (root), Chufa, Dasheen (Taro), Ginger, Leren, Tanier, Turmeric, and Yam (bean and true).</p>			
<p>Specific Use Restrictions:</p> <ol style="list-style-type: none"> 1) DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01. 2) DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 3) DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 4) DO NOT make more than 4 applications per year at the maximum application rate. 5) Minimum Application Interval: 7 days. 6) DO NOT apply within 14 days of harvest (14-day PHI). 			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
<p>Rapeseed Subgroup 20A[*]</p> <p>Including cultivars, varieties, and/or hybrids of these and others in this group (see below)</p>	<p>Alternaria black spot (<i>Alternaria brassicae</i>)</p> <p>Black leg/Phoma (<i>Leptosphaeria maculans</i>)</p> <p>Cercospora leaf spot (<i>C. brassicicola</i>)</p> <p>Head rot (<i>Rhizoctonia solani</i>)</p> <p>Leaf spot and pod rot (<i>Alternaria alternata</i>)</p> <p>Powdery mildew (<i>Erysiphe polygoni</i>)</p>	<p>4 – 6.95 (0.066 – 0.114)</p>	<p>For Phoma control, apply during the rosette stage between 2nd true leaf and bolting.</p> <p>For Alternaria, make an application at the end of flowering/early pod set. For other foliar diseases, apply at first sign of disease.</p> <p>For head rot, apply at 50% flowering.</p> <p>If disease pressure is high, use the highest rate.</p>
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by ground, air, or chemigation. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.</p>			
<p>Complete List of Rapeseed Subgroup 20A: Borage, crambe, cuphea, echium, flax seed, gold of pleasure, hare’s ear mustard, lesquerella, lunaria, meadowfoam, milkweed, mustard seed, oil radish, poppy seed, rapeseed, sesame, sweet rocket and cultivars and/or hybrids of these.</p>			
<p>Specific Use Restrictions:</p> <ol style="list-style-type: none"> 1) DO NOT apply more than one application of A253.01 per year. 2) DO NOT apply more than 6.95 fl oz/A/year of A253.01. 3) DO NOT apply more than 0.114 lb ai/A/year of difenoconazole-containing products. 4) DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 5) DO NOT apply within 30 days of harvest (30-day PHI). 			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Rice[*]	Sheath Blight (<i>Rhizoctonia solani</i>) Aggregate Sheath Spot (<i>Rhizoctonia oryzaesativae</i>) Black Sheath Rot (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>) Sheath Spot (<i>Rhizoctonia oryzae</i>) Brown Leaf spot (<i>Cochliobolus miyabeanus</i>) Narrow Brown Leaf spot (<i>Cercospora oryzae</i>) <u>Suppression only:</u> False smut (<i>Ustilaginoidea virens</i>) Kernel Smut (<i>Neovossia barclayana</i>) Stem Rot (<i>Sclerotium oryzae</i>)	7.45 (0.122)	Apply 7.45 fl oz when disease is less than 4 inches above water line usually between panicle differentiation (PD) +5 days to PD +10 days or at initial sign of disease. Under heavy disease pressure and conditions favorable for disease development, a second application may be applied. The minimum re-treatment interval is 14 days.

Application: For best results, sufficient water volume must be used to provide thorough coverage. **A253.01** can be applied by ground or aerial application. For aerial applications, use a minimum of 5 gal/A of water. Applicators must use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) **DO NOT** allow release of irrigation or flood water for at least 7 days after the last application.
- 2) **DO NOT** apply when weather conditions favor drift from treated areas to non-target aquatic habitat.
- 3) **DO NOT** apply more than 14.9 fl oz (0.244 lb ai)/A/year of **A253.01**.
- 4) **DO NOT** apply more than 0.244 lb ai/A/year of difenoconazole-containing products.
- 5) **DO NOT** apply more than 7.45 fl oz per application (0.122 lb ai of difenoconazole).
- 6) **DO NOT** make more than 2 applications per year.
- 7) **Minimum Application Interval:** 14 days.
- 8) **DO NOT** apply **A253.01** within 28 days of harvest (28-day PHI).
- 9) **DO NOT** treat fields used for aquaculture of fish or crustacean.
- 10) **DO NOT** drain water from treated rice fields into ponds used for aquaculture of fish or crustacean.
- 11) **DO NOT** use water drained from treated field to irrigate other crops.

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Root Vegetables, Subgroup 1A Radish Sugar beet[*]	Alternaria leafspots and blight (<i>Alternaria</i> spp.) Beet rust (<i>Uromyces betae</i>) Cercospora leafspot	4 – 6.95 (0.066 – 0.114)	Apply A253.01 on a 7- to 28-day schedule, alternating with a non-triazole fungicide (non-Group 3) that is registered for these diseases. <i>[Optional language if label has a rate range]</i>

See additional crops in this subgroup below See specific directions for carrots and ginseng	(<i>Cercospora</i> spp.) Powdery mildew (<i>Erysiphe</i> spp.)		<i>and a single interval:</i> If disease pressure is high, use the highest rate.] <i>[Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval.] <i>[Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.]
	Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 5 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.		
Vegetables, Root Subgroup: Beet, garden; beet, sugar; burdock, edible; celeriac; chervil, turnip- rooted; chicory; horseradish; parsley, turnip-rooted; parsnip; radish; radish, oriental (daikon); rutabaga; salsify; salsify, black; salsify, Spanish; skirret; turnip			
Specific Use Restrictions: 1) DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01 . 2) DO NOT apply more than 0.46 lb ai/A/year of difenoconazole. 3) DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 4) DO NOT make more than 4 applications per year at the maximum application rate. 5) Minimum Application Interval: 7 days 6) DO NOT apply within 7 days of harvest (7-day PHI).			

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Leaves of Root and Tuber Vegetables, Group 2 Turnip Sugar beet[*] See additional crops in this group below	Alternaria leafspots and blight (<i>Alternaria</i> spp.) Beet rust (<i>Uromyces betae</i>) Cercospora leafspot (<i>Cercospora</i> spp.) Powdery mildew (<i>Erysiphe</i> spp.)	4 – 6.95 (0.066 – 0.114)	Apply A253.01 on a 7- to 28-day schedule, alternating with a non-triazole fungicide (non-Group 3) that is registered for these diseases. <i>[Optional language if label has a rate range and a single interval:</i> If disease pressure is high, use the highest rate.] <i>[Optional language if label has a single rate and interval range:</i> If disease pressure is high, use the shortest interval.] <i>[Optional language if label has a rate range and interval range:</i> If disease pressure is high, use the shortest interval and highest rate.]
Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 5 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			

Leaves of Root and Tuber Vegetables Group: Beet, garden; beet, sugar; burdock, edible; carrot; cassava, bitter and sweet; celeriac (celery root); chervil, turnip-rooted; chicory; dasheen (taro); parsnip; radish; radish, oriental (daikon); rutabaga; salsify, black; sweet potato; tanier (cocoyam); turnip; yam, true.

Specific Use Restrictions:

- 1) **DO NOT** apply more than 27.8 fl oz (0.46 lb ai)/A/year of **A253.01**.
- 2) **DO NOT** apply more than 0.46 lb ai/A/year of difenoconazole.
- 3) **DO NOT** apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole).
- 4) **DO NOT** make more than 4 applications per year at the maximum application rate.
- 5) **Minimum Application Interval:** 7 days
- 6) **DO NOT** apply within 7 days of harvest (7-day PHI).

[*Not for use in California]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Soybean[*]	Anthracnose (<i>Colletotrichum truncatum</i>) Alternaria leaf spot (<i>Alternaria</i> spp.) Brown spot (<i>Septoria glycines</i>) Cercospora blight and leaf spot (<i>A. kikuchii</i>) Frogeye leaf spot (<i>Cercospora sojina</i>) Pod and stem blight (<i>Diaporthe phaseolorum</i>) Powdery mildew (<i>Microsphaera diffusa</i>)	5.5 – 6.95 (0.090 – 0.114)	Begin applications prior to disease onset when conditions are conducive for disease. Apply A253.01 on a 7- to 10-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. <i>[Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.]</i> <i>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.]</i> <i>[Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]</i>
Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground, chemigation, or aerial application. For aerial applications, use a minimum of 2 gal/A of water. For chemigation, apply in 0.1- 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			

Specific Use Restrictions:

- 1) **DO NOT** apply more than 13.5 fl oz (0.22 lb ai)/A/year of **A253.01**.
- 2) **DO NOT** apply more than 0.22 lb ai/A/year of difenoconazole-containing products.
- 3) **DO NOT** apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole).
- 4) **DO NOT** make more than 1 application per year at the maximum application rate.
- 5) **Minimum Application Interval:** 7 days.
- 6) **DO NOT** apply within 14 days of harvest (14-day PHI).
- 7) **DO NOT** feed soybean hay, forage and silage.

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
<p>Stone Fruit Crop Group 12-12</p> <p>Apricots[*] Cherries, Sweet Cherries, Tart Nectarines[*] Peaches[*] Plums[*] Plumcot[*] Prunes[*]</p> <p>And cultivars and/or hybrids of these</p>	<p>Brown rot blossom blight and fruit rot (<i>Monilinia fructicola</i>, <i>M. laxa</i>)</p> <p>Scab (<i>Cladosporium carpophilum</i>)</p> <p>Alternaria spot and fruit rot (<i>A. alternata</i>)</p> <p>Anthracnose (<i>Colletotrichum</i> spp.)</p> <p>Leaf rust (<i>Tranzschelia discolor</i>)</p> <p>Powdery mildew (<i>Sphaerotheca pannosa</i>, <i>Podosphaera clandestina</i>)</p> <p>Shot hole (<i>Wilsonomyces carpophilus</i>)</p>	<p>5.5 – 6.95 (0.090 – 0.114)</p> <p>[6.95 (0.114) (CA Only)]</p>	<p>For Brown rot blossom blight, begin applications at early bloom and continue through petal fall. For Brown rot on fruit, apply as needed a maximum of two sprays during the preharvest period up to the day of harvest (minimum of a 7- day retreatment interval). If high inoculum and severe disease conditions persist, apply a registered non-Group 3 fungicide.</p> <p>For all other diseases, follow the Brown rot blossom blight schedule. Make additional applications on a 10- to 14-day interval from the end of petal fall to harvest.</p> <p>[Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.]</p> <p>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.]</p> <p>[Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]</p> <p>Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground or aerial application. Stone fruit diseases are most effectively controlled by ground applications. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.</p>
<p>Stone Fruit Crop Group: Apricot[*]; apricot, Japanese[*]; capulin[*]; cherry, black[*]; cherry, Nanking[*]; cherry, sweet; cherry, tart; Jujube, Chinese[*]; nectarine[*]; peach[*]; plum[*]; plum, American[*]; plum, beach[*]; plum, Canada[*]; plum, cherry[*]; plum, Chickasaw[*]; plum, Damson[*]; plum, Japanese[*]; plum, Klamath[*]; plum, prune[*]; plumcot[*]; sloe[*]; cultivars[*], varieties[*], and/or hybrids[*] of these.</p>			
<p>Specific Use Restrictions:</p> <ol style="list-style-type: none"> DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01. DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products. DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). DO NOT make more than 4 applications per year at the maximum application rate. Minimum Application Interval: 7 days. A253.01 may be applied up to the day of harvest (0-day PHI). 			

[*][NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Sugar Beets[*]	Cercospora leafspot (<i>C. beticola</i>) Powdery mildew (<i>Erysiphe polygoni</i>)	4 – 6.95 (0.066 – 0.114)	<p>Apply A253.01 on a 7- to 28-day schedule, alternating with a non-triazole fungicide (non-Group 3) that is registered on sugar beets for these diseases.</p> <p><i>[Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.]</i></p> <p><i>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.]</i></p> <p><i>[Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]</i></p> <p>Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 5 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.</p>

Specific Use Restrictions:

- 1) **DO NOT** apply more than 27.8 fl oz (0.46 lb ai)/A/year of **A253.01**.
- 2) **DO NOT** apply more than 0.46 lb ai/A/year of difenoconazole-containing products.
- 3) **DO NOT** make more than 4 application per year at the maximum application rate.
- 4) Minimum Application Interval: 7 days.
- 5) **DO NOT** apply within 7 days of harvest (7-day PHI).

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Tree Nuts Crop Group 14-12[*] Beechnut Brazil Nut Butternut Cashew Chestnut Macadamia Walnut, Black Walnut, English	Foliar Diseases	5 – 6.95 (0.082 0.114)	<p>Begin applications prior to disease onset when conditions are conducive for disease. Apply A253.01 on a 14- to 21-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p> <p><i>[Optional language if label has a rate range and a single interval: If disease pressure is high, use the highest rate.]</i></p> <p><i>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.]</i></p>

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
See specific directions for almonds, filberts, pecans, pistachios			[Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]
	Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by either ground, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.		
Tree Nuts Crop Group: African nut-tree; almond; beechnut; Brazil nut; Brazilian pine; bunya; bur oak; butternut; Cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; coquito nut; dika nut; ginkgo; Guiana chestnut; hazelnut (filbert); heartnut; hickory nut; Japanese horse-chestnut; macadamia nut; mongongo nut; monkey-pot; monkey puzzle nut; Okari nut; Pachira nut; peach palm nut; pecan; pequi; Pili nut; pine nut; pistachio; Sapucaia nut; tropical almond; walnut, black; walnut, English; yellowhorn; cultivars, varieties, and/or hybrids of these.			
Specific Use Restrictions: 1) DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01 . 2) DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 3) DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 4) DO NOT make more than 4 applications per year at the maximum application rate. 5) Minimum Application Interval: 14 days. 6) DO NOT apply within 14 days of harvest (14-day PHI).			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Watercress[*]	Cercospora leafspot (<i>Cercospora</i> spp.)	6.95 (0.114)	For best activity, apply A253.01 prior to or early in the disease development. An adjuvant may be added at specified rates. Apply on a 7-14 day interval. For applications made to watercress, production fields must be drained of water at least 24 hours prior to application and water must not be reapplied to the field for a minimum of 24 hours following the application. DO NOT apply more than two sequential applications before alternating to a fungicide with a different mode of action.
Application: For best results, sufficient water volume must be used to provide thorough coverage. A253.01 can be applied by ground, chemigation, or aerial application. For aerial applications, use a minimum of 5 gal/A of water. Applicators must use care in making applications near non-target aquatic habitats.			
Specific Use Restrictions: 1) DO NOT apply directly to water and DO NOT allow water in a treated field for at least 24 hours. 2) DO NOT apply more than 27.8 fl oz (0.46 lb ai)/A/year of A253.01 . 3) DO NOT apply more than 0.46 lb ai/A/year of difenoconazole-containing products. 4) DO NOT apply more than 6.95 fl oz per application (0.114 lb ai of difenoconazole). 5) DO NOT make more than 4 applications per year. 6) Minimum Application Interval: 7 days 7) Production fields must be drained of water at least 24 hours prior to application and water must not be			

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
reapplied to the field for a minimum of 24 hours following the application.			
8) DO NOT apply A253.01 within 30 days of harvest (30-day PHI).			

[*] [NOT FOR USE IN CALIFORNIA]

Crop	Target Diseases	Use Rate fl oz product/A (lb ai)	Application Instructions
Wild Rice[*]	Helminthosporium leaf blight Brown Spot (<i>Bipolaris</i> spp.)	7.45 (0.122)	Apply 7.45 fl oz/A at both booting and heading. Minimum re-treatment interval is 14 days.

Application: For best results, sufficient water volume must be used to provide thorough coverage. **A253.01** can be applied by ground or aerial application. For aerial applications, use a minimum of 5 gal/A of water. Applicators must use care in making applications near non-target aquatic habitats.

Specific Use Restrictions:

- 1) **DO NOT** allow release of irrigation or flood water for at least 7 days after the last application.
- 2) **DO NOT** apply when weather conditions favor drift from treated areas to non-target aquatic habitat.
- 3) **DO NOT** apply more than 14.9 fl oz (0.244 lb ai) /A/year of **A253.01**.
- 4) **DO NOT** apply more than 0.244 lb ai/A/year of difenoconazole-containing products.
- 5) **DO NOT** apply more than 7.45 fl oz per application (0.122 lb ai of difenoconazole).
- 6) **DO NOT** make more than 2 applications per year.
- 7) **Minimum Application Interval:** 14 days.
- 8) **DO NOT** apply **A253.01** within 28 days of harvest (28-day PHI).
- 9) **DO NOT** use water drained from treated field to irrigate other crops.
- 10) **DO NOT** treat fields used for aquaculture of fish or crustacean.
- 11) **DO NOT** drain water from treated rice fields into ponds used for aquaculture of fish or crustacean.

[*] [NOT FOR USE IN CALIFORNIA]

A253.01 Conversion Table

Fl oz product/A	Lb ai/A
2	0.033
3	0.049
3.97	0.065
5	0.082
5.5	0.090
6	0.099
6.95	0.114
7.45	0.122
7.6	0.125
8	0.131

Ornamentals – For outdoor application to established ornamentals in landscaped areas, to include recreational parks, institutional sites, golf courses and residential properties by commercial and non-professional (e.g., homeowner) applicators.

Use the calibrated measuring [device] [syringe] supplied with the product to accurately measure the designated amount of product to be added to each gallon of water.

Note: This use for application to ornamentals does not include crops grown for production, harvest and sale in commercial greenhouse or nursery production sites.

Ornamental	Disease	Rate ml/1 Gal (1 Gal Treats 1000 Sq Ft)	Spray interval days	Timing of application
Carnation [Not for use in California]	Rust	4.76	7-10	Begin applications when flowers are half opened or when conditions are conducive for disease development. Apply A253.01 on a 7- to 10-day schedule. Particular care should be taken to achieve thorough coverage.
Gladiolus [Not for use in California]	Rust	4.76	7-14	Begin applications at the 2- to 3-leaf stage or when conditions are conducive for disease development. Apply A253.01 on a 7- to 10-day schedule. Particular care should be taken to achieve thorough coverage.
Iris [Not for use in California]	Rust	1.4-2.7	7-21	Begin applications at the first sign of disease or when conditions are conducive for disease development. Apply A253.01 on a 7- to 10-day schedule. Particular care should be taken to achieve thorough coverage.
Roses [Not for use in California]	Black spot Rust Powdery mildew	4.76	7-14	Begin applications at the first sign of disease or when conditions are conducive for disease development. Apply A253.01 on a 7- to 10-day schedule. Particular care should be taken to achieve thorough coverage.

Restrictions:

- For ornamentals, **DO NOT** apply more than 4.76 ml/1000 sq ft of **A253.01** (0.115 lb ai/A) in a single application.
- For ornamentals, **DO NOT** apply more than 19.2 ml/1000 sq ft of **A253.01** (0.46 lb ai/A) per year.
- **DO NOT** make more than 4 applications at the maximum application rate.
- Minimum Application Interval: 7 days

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[For Residential uses] **[Nonrefillable container. DO NOT** reuse or refill this container. If empty: Offer for recycling if available or discard in a sanitary landfill. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.]

[For Commercial Uses]

[For plastic containers ≤ 5 gallons: Nonrefillable Container: DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

[For plastic containers > 5 gallons: Nonrefillable container. DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

A253.01 is a trademark of Atticus, LLC

[Inspire®] [and] [Top MP®] [is a] [are] registered trademark[s] of a Syngenta Group Company.

{LANGUAGE ON LABEL AFFIXED TO CONTAINER}

DIFENOCONAZOLE	GROUP 3	FUNGICIDE
----------------	---------	-----------

A253.01 [TM]

[Alternate Brand Name: Esquire]

Contains Difenoconazole, the active ingredient used in [Inspire®] [and] [Top MP].

ACTIVE INGREDIENT:	(% by weight)
Difenoconazole*.....	23.2%
OTHER INGREDIENTS:	76.8%
TOTAL	100.0%

*CAS No. 119446-68-3

**Contains petroleum distillates

A253.01 is formulated as an emulsifiable concentrate (EC) containing 2.10 lb ai/gal of difenoconazole.

KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand the label, find someone to explain it to you in detail.)

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> Immediately call a poison control center or doctor. DO NOT induce vomiting unless told to do so by a poison control center or doctor. DO NOT give any liquid to the person. DO NOT give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

For Chemical Emergency:

Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC Day or Night

Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING/AVISO

Causes substantial but temporary eye injury. **DO NOT** get in eyes or on clothing. Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish, mammals, and aquatic invertebrates. Drift and runoff may be hazardous to aquatic **estuarine/marine** organisms in water adjacent to treated areas. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. **DO NOT** discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. **DO NOT** discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or regional office of the EPA.

Surface Water Advisory: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product may contaminate water through drift or spray in wind. This product has a potential for runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is expected to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

PHYSICAL OR CHEMICAL HAZARDS: DO NOT use or store near heat or open flame. Do not mix or allow to come in contact with oxidizing agents. Hazardous chemical reaction may occur.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[For Residential uses]

[Nonrefillable container. **DO NOT** reuse or refill this container. If empty: Offer for recycling if available or discard in a sanitary landfill. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.]

[For Commercial Uses]

[For plastic containers ≤ 5 gallons: Nonrefillable Container: **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

[For plastic containers > 5 gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

See inside label booklet for additional Precautionary Statements and Directions for Use.

A253.01 is not manufactured, or distributed by Syngenta Group Company, seller of [Inspire®] [and] [Top MP®].

Manufactured for:
Atticus, LLC
5000 CentreGreen Way, Suite 100
Cary, NC 27513

EPA Reg. No.: 91234-XX
EPA Est. No.: _____
NET WEIGHT: _____