



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

91234-245

Date of Issuance:

2/16/22

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

A392.03

Name and Address of Registrant (include ZIP Code):

Atticus, LLC
5000 CentreGreen Way, Suite 100
Cary, NC 27513

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Continued on page 2

Signature of Approving Official:

Mindy Ondish, Product Manager 23
Herbicide Branch, Registration Division (7505P)

Date:

2/16/22

2. You are required to comply with the data requirements described in the Generic Data Call-Ins (GDCIs) identified below:
 - a. Metribuzin GDCI-101101-1304 and GDCI-101101-1825

You must comply with all of the data requirements within the established deadlines. If you have questions about the GDCIs listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division: <http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. You have 18 months from the date of registration to provide these data.
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

Please note that the alternate brand name "**TrovidoR**" has been added to the product record.

Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 02/26/2021
- Alternate CSFs 1-8 dated 02/26/2021

If you have any questions, please contact Endia Blunt by phone at 202-566-2505, or via email at blunt.endia@epa.gov.

Enclosure

ACCEPTED

02/16/2022

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 91234-245

{Note to reviewer: [Text] in brackets denotes optional or explanatory language}
{Note to reviewer: {Text} in braces denotes where in the final label text will appear}
{BOOKLET FRONT PANEL LANGUAGE}

METOLACHLOR	GROUP	15	HERBICIDE
METRIBUZIN	GROUP	5	HERBICIDE
IMAZETHAPYR	GROUP	2	HERBICIDE

A392.03TM

[Alternate Brand Name: TrovidoR]

[Herbicide for preemergent control of certain grasses and broadleaf weeds in soybeans.]

ACTIVE INGREDIENTS:	(% by weight)
Metolachlor*	43.72%
Metribuzin**	6.14%
Imazethapyr***	1.38%
OTHER INGREDIENTS:	48.76%
TOTAL	100.0%

- *contains 4.01 pounds of metolachlor per gallon.
- **contains 0.56 pound of metribuzin per gallon.
- ***contains 0.13 pound of imazethapyr acid per gallon.

KEEP OUT OF REACH OF CHILDREN WARNING-AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See [below] [inside label booklet] for [additional] [First Aid,] [and] [Precautionary Statements] [and] [Directions for Use].

FIRST AID	
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

For Chemical Emergency:
Spill, Leak, Fire, Exposure, or Accident,
Call CHEMTREC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

EPA Reg. No.: 91234-245

EPA Est. No.:

Net Contents:

Manufactured for:
Atticus, LLC
5000 CentreGreen Way, Suite 100
Cary, NC 27513

{LANGUAGE INSIDE BOOKLET}

PRECAUTIONARY STATEMENTS **HAZARDS TO HUMANS AND DOMESTIC ANIMALS** **WARNING**

Causes skin irritation. Causes moderate eye irritation. Do not get on skin or on clothing. Avoid contact with eyes. Wear protective eyewear. This product may cause skin sensitization reactions in some people. Wear coveralls, chemical-resistant gloves and chemical-resistant footwear. A chemical resistant apron plus chemical resistant gloves are required when mixing, loading, cleaning up spills or equipment or when otherwise exposed to this product's concentrate. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment

Mixers, loaders, applicators, flaggers and other handlers must wear:

- Protective eyewear,
- Chemical-resistant headgear for overhead exposure,
- Coveralls worn over short-sleeved shirt and short pants,
- Chemical-resistant gloves made out of butyl rubber ≥ 14 mils or barrier laminate,
- Chemical-resistant footwear plus socks, and
- Chemical-resistant apron when cleaning equipment, mixing or loading.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607 (d-f)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses: do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

NON-TARGET ORGANISM ADVISORY STATEMENT:

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several weeks or months after application. A level, well-maintained vegetative buffer strip

between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of metolachlor and imazethapyr from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Ground Water Advisory and Mixing/Loading Instructions

Metribuzin is a chemical which can travel (seep or leach) through soil and can contaminate ground water which may be used as drinking water. Metribuzin has been found in ground water as a result of agricultural use. Users are advised not to apply metribuzin where the water table (ground water) is close to the surface, and where the soils are very permeable, i.e. well-drained soils such as loamy sands. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

Imazethapyr has properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Metolachlor has the potential to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

DO NOT apply this product through any type of irrigation system.

Product must be used in a manner which will prevent back-siphoning in wells, spills or improper disposal of excess pesticide spray mixture.

This product may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes or reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas.

Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times.

The above specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead set-backs and operational containment.

Reporting Ecological Incidents:

To report ecological incidents, including mortality, injury, or harm to plants and animals, call 984-465-4800.

PHYSICAL CHEMICAL HAZARDS

Do not mix or allow coming in contact with Oxidizing agents & Ammonium phosphate. Hazardous Chemical reaction may occur. When mixed with water, temperature may slightly increase.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Endangered Species Protection Requirements:

It is a Federal offense to use any pesticide in a manner that results in an unauthorized "take" (e.g., kill or otherwise harm) of an endangered species and certain threatened species, under the Endangered Species Act section 9. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the area in which you are applying the product. You must obtain a Bulletin no earlier than six months before using this product. To obtain Bulletins, consult <http://www.epa.gov/espp/>, call 1-844-447-3813, or email ESPP@epa.gov. You must use the Bulletin valid for the month in which you will apply the product.

In New York State - not for sale or use in Nassau and Suffolk Counties.

Observe all precautions and limitations on this label and on the labels of products used in combination with this product.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protections of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protections Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeve shirt and short pants,
- Chemical-resistant gloves, such as butyl rubber \geq 14 mils or barrier laminate,
- Chemical-resistant footwear plus socks.

PRODUCT INFORMATION

A392.03 kills weeds by root and/or foliage uptake and rapid translocation to the growing points. Adequate soil moisture is important for optimum activity of this product. When adequate soil moisture is present, this product will provide residual control of susceptible germinating weeds; activity on established weeds will depend on the weed species and the location of its root system in the soil.

Occasionally, internode shortening and/or temporary yellowing of crop plants may occur following applications of this product. These effects occur infrequently and are temporary. Normal growth and appearance should resume within 1 to 2 weeks.

When organophosphate or carbamate insecticides are tank mixed with this product, temporary injury may result to the treated crops.

Use of this product in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks

associated with the use of this product. Therefore, rotational crop injury is always possible.

Under some conditions (such as heavy texture soil, high organic matter, low pH or low rainfall), this product may cause injury to subsequent planted crops. Vegetable crops (particularly sugar beets) are sensitive to residues of this product in the soil.

Naturally occurring biotypes* of some of the weeds listed on this label may not be effectively controlled by this and/or other products with either the ALS/AHAS enzyme inhibiting mode of action. Other herbicides with the ALS/AHAS enzyme inhibiting mode of action include the sulfonylureas, the sulfonamides and the pyrimidyl benzoates.

*A weed biotype is a naturally occurring plant within a given species that has a slightly different, but distinct, genetic makeup from other plants.

Weed Resistance Management

For resistance management, please note that **A392.03** contains a Group 15 (Metolachlor), a Group 5 (Metribuzin), and a Group 2 (Imazethapyr) herbicide. Any weed population may contain plants naturally resistant to Group 15, Group 5, and/or Group 2 herbicides. The resistant individuals may dominate the weed population if these herbicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of **A392.03** or other Group 15, 5 or 2 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include:
 - 1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - 2) a spreading patch of non-controlled plants of a particular weed species;
 - 3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact an Atticus, LLC representative at 984-465-4800.

Spray Drift Management

Aerial Applications:

- Do not release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select the nozzle and pressure that deliver medium or coarser droplets (ASABE S641).
- If the wind speed is 10 miles per hour or less, applicators must use ½ swath displacement upwind at the downwind edge of the field. When the wind speed is between 11-15 miles per hour, applicators must use ¾ swath displacement upwind at the downwind edge of the field.
- Do not apply when wind speeds exceed 15 mph at the application site. If the wind speed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Do not apply during temperature inversions.

Ground Boom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to select the nozzles and pressure that deliver medium or coarser droplets (ASABE S572).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Boomless Ground Applications:

- Applicators are required to select the nozzle and pressure that deliver medium or coarser droplet size (ASABE S572.3) for all applications.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles – Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT – Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boomless Ground Applications:

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

Take precautions to minimize spray drift.

APPLICATION WITH HERBICIDE SPRAY EQUIPMENT

Use a standard low-pressure (20 to 40 psi) herbicide boom sprayer equipped with suitable nozzles and screens no finer than 50 mesh in nozzle and in-line strainers. Agitate thoroughly before and during application with bypass agitation. Low pressure and high volume hand wand equipment is prohibited.

Ground Application

Apply the proper rate of this product in a minimum of 10.0 to 40.0 gallons of spray mixture per acre broadcast.

Aerial Application

Where permitted, apply specified rate in a minimum of 5.0 gallons of spray mixture per acre. Do not apply aerially when wind speed is greater than 10 mph.

For all Applications

Sprayer must be accurately calibrated before applying this product. Check sprayer during application to be sure it is working properly and delivering a uniform spray pattern. As the volume of spray mixture decreases per acre, the importance of accurate calibration and uniform application increases.

Avoid overapplication, misapplication, and boom and spray swath overlapping that will increase spray dosage. (Crop injury may occur as a result.) Avoid spray skips and gaps which allow weeds to grow in untreated soil. Do not apply when weather conditions favor spray drift and/or when sensitive or cool season crops (such as cole crops, onions, peas, or strawberries) are present in adjacent fields or in areas where wheat is growing in coarse textured soils.

Sprayer Cleanup

Spray equipment must be thoroughly cleaned to remove remaining traces of herbicide that might injure other crops to be sprayed. Drain any remaining spray solution of this product from the spray tank and dispose of according to label disposal instructions. Rinse the spray tank and refill with water, adding a heavy-duty detergent at the rate of 1 cup per 20.0 gallons of water. Recycle this mixture through the equipment for 5 minutes and spray out. Repeat this procedure twice. Fill the spray tank with clean water, recycle for 5 minutes, and spray out. Clean pump and nozzle screens thoroughly. Wash away any spray mixture from the outside of spray tank, nozzles or spray rig. All rinse water must be disposed of in compliance with local, state, and Federal guidelines.

Incorporation and Combination Uses

When this product is to be used in combination with another herbicide, follow the most restrictive directions for use and precautionary statements on all product labels for combinations, rates, crops, incorporation, and special precautions.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.

When using this product, make sure the sprayer is completely clean, free of rust or corrosion which occurs from winter storage. Examine strainers and screens to be sure the sprayer is clean from previously used pesticides.

Any tank mix containing this product must be kept agitated and sprayed out immediately. Do not allow tank mixes to stand for prolonged periods of time.

The proper mixing procedure for this product alone or in tank-mix combinations with other herbicides is:

1. Fill the spray tank 1/4 to 1/3 full with clean water.
2. Add specified rate of this product while recirculating and with agitator running.
3. Mix thoroughly and add clean water to fill spray tank to desired level.
4. Add the other herbicide to tank last and agitate thoroughly.
5. Continue agitation during application and until sprayer tank is empty.

Application of A392.03 in Fluid Fertilizers

This product may be applied in fluid fertilizer solutions by following the appropriate mixing procedures and compatibility check. When using tank mix combinations, be sure all components are compatible.

Tank Mixing Guidelines for Fluid Fertilizer Mixtures

1. Add the required amount of water and compatibility-agent (if required) to the tank. Start agitation-system while adding this product and follow by adding the fluid fertilizer and agitate.
2. If a second herbicide is also to be used, follow as above in Step 1, but use twice the amount of water. Start agitation, add **A392.03**. Follow by adding the second herbicide, then continue filling the tank with fluid fertilizer.
3. Maintain continuous agitation to assure uniform spray mixture until the tank is emptied.

Make compatibility checks of this product plus fluid fertilizers and tank-mix combinations plus fluid fertilizers which include this product for each batch because of the variability of fluid fertilizers.

THE FOLLOWING COMPATIBILITY CHECK SHOULD ONLY BE USED WHEN MIXING WITH FLUID FERTILIZERS.

1. Pre-mix 8.0 teaspoons of water with 2.0 teaspoons of this product (4:1 ratio) in a quart jar by adding the water first and following with this product. Mix thoroughly. If a second herbicide is to be used, double the amount of water (8:1 ratio), mix in this product, and follow with the second herbicide.
2. Then pour 1.0 pint of fluid fertilizer into the quart jar and shake well.
3. Allow to stand for 5 minutes.

Interpretation of Results

If the solution in the jar appears to be uniform, without signs of agglomeration, or without a separation of an oily film on top of the fertilizer, the mixture may be used. If not, repeat the compatibility check using twice the amount of water or add a compatibility agent to the water. If separation occurs, but the mixture can be resuspended by shaking, then application is possible with good agitation in the spray tank.

SOYBEAN APPLICATION DIRECTIONS (Except California)

This product may be applied preplant incorporated, preplant surface or preemergent surface, or as a sequential preemergent application. This product may also be used as an overlay application following a preplant incorporated application of a grass herbicide registered for this same use and in tank mix combinations for burndown weed control. All applications may be applied with ground equipment, and some may be applied with aerial spray equipment.

Restrictions

- Do not apply more than 4.0 pints (2.01 lb Metolachlor + 0.28 lb Metribuzin + 0.07 lb Imazethapyr acid) of **A392.03** per acre per year.
- In North Dakota, and in Minnesota north of Highway #210, do not apply more than 2.9 pints (1.45 lb Metolachlor + 0.2 lb Metribuzin + 0.05 lb Imazethapyr acid) of **A392.03** per acre per year.
- Do not harvest within 85 days of the last application of **A392.03**.
- Do not graze or feed treated soybean forage, hay or straw to livestock.
- Do not rotate to any crop not listed on this label for 40 months following application.
- Do not rotate to food or feed crops other than those listed on this label.
- Do not apply **A392.03** preemergence to soybeans in California.
- Do not incorporate into soil or apply more than once per season except where permitted as part of a sequential application.
- Do not allow sprays to drift onto adjacent desirable plants.
- Do not apply under conditions which favor runoff or wind erosion of soil containing this product to non-target areas.
- To prevent off-site movement due to runoff or wind erosion:
 - Avoid treating powdery dry or light sand soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.
 - Do not apply to impervious substrates such as pavement or highly compacted surfaces.
 - Do not use tailwater from the first flood or furrow irrigation of treated fields to treat non-target crops unless at least 1/2 inch of rainfall has occurred between application and the first irrigation.
- Do not apply using low-pressure and high-volume hand-wand equipment.
- Observe all precautions and limitations on labeling of all products used in mixtures.

Soil Texture and Rate Ranges

As used on this label, "coarse soils" are loamy sand or sandy loam soils. "Medium soils" are loam, silt loam, silt, sandy clay, or sandy clay loam. "Fine soils" are silty clay, silty clay loam, clay, or clay loam. Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.

Where a rate range is shown, use a lower rate on soils that are coarse-textured or low in organic matter. Use a higher rate on soils that are relatively fine-textured or high in organic matter.

Precautions

Injury to soybeans may occur when this product is used under the following conditions:

- When soils have a calcareous surface area or a pH of 7.5 or higher.
- When applied in conjunction with soil-applied organic phosphate pesticides.
- With over-application or boom overlapping, which may result in stand loss and soil residues.

- With uneven application or improper incorporation, which can decrease the level of weed control and/or increase the level of injury.
- When applied to any soil with less than 0.5% organic matter.
- When soil incorporation is deeper than 2 inches.
- When sprayers are not calibrated accurately.
- When heavy rains occur soon after application, especially in poorly drained areas where water may stand for several days.
- When soybeans are planted less than 1.5 inches deep, particularly in preemergence application.
- Where high soil levels of atrazine are present.
- When using poor quality soybean seed.

Certain soybean varieties are sensitive to metribuzin. Prior to use of this product, consult your soybean seed supplier for more information on the tolerance of soybean varieties to **A392.03**.

Activation

A minimum amount of soil moisture is required to activate this product. In areas of low rainfall, preemergence applications to dry soil should be followed with light irrigation of 1/4 acre-inch of water. Do not apply heavy irrigation immediately after application. As with many surface-applied herbicides, weed control and crop tolerance may vary with rainfall and/or soil texture.

Replanting

If replanting is necessary in fields treated with this product as directed -on this label, the field may be replant- ed to soybeans. Rework the soil no deeper than the treated zone. Do not apply more than once per season except where permitted as part of a sequential application as injury to soybeans may occur. Maximum application rate is 4.0 pints of **A392.03** per acre per use season. Do not exceed this amount in any use pattern: single application, replant or sequential application.

TABLE 1: ANNUAL BROADLEAF WEEDS CONTROLLED BY A392.03

C = Control S = Suppression or Erratic Control P = Poor or No Control U= Unknown

Weed Controlled	Level of Control
Bristly starbur (<i>Acanthospermum hispidum</i>)	C
Buffalobur (<i>Solanum rostratum</i>)	C
Carpetweed (<i>Mollugo verticillata</i>)	C
Cocklebur (<i>Xanthium pensylvanicum</i>)	S
Common chickweed (<i>Stellaria media</i>)	C
Copperleaf, Hophornbeam (<i>Acalypha ostryifolia</i>)	C
Field pennycress (<i>Thlaspi arvense</i>)	C
Florida beggarweed (<i>Desmodium tortuosum</i>)	C
Florida pusley (<i>Richardia scabra</i>)	C
Galinsoga (<i>Galinsoga</i> spp.)	C
Henbit (<i>Lamium amplexicaule</i>)	C
Horseweed (Marestail) (<i>Conyza canadensis</i>)	U
Jimsonweed (<i>Datura stramonium</i>)	C
Knotweed (<i>Polygonum</i> spp.)	C
Kochia (<i>Kochia scoparia</i>)	C
Lambsquarters (<i>Chenopodium</i> spp.)	C
Marshelder (<i>Iva Annuua</i>)	C
Morningglory	
Entireleaf (<i>Ipomoea hederacea</i> var. <i>integriuscula</i>)	S
Ivyleaf (<i>Ipomoea hederacea</i>)	S
Pitted (<i>Ipomoea lacunosa</i>)	S
Smallflower (<i>Jacquemontia tamnifolia</i>)	C

Tall (<i>Ipomoea purpurea</i>)	S
Mustard spp.	C
Nightshade	
Black (<i>Solanum nigrum</i>)	C
Eastern black (<i>Solanum ptycanthum</i>)	C
Hairy (<i>Solanum villasum</i>)	C
Pigweed	
Redroot (<i>Amaranthus retroflexus</i>)	C
Smooth (<i>Amaranthus hybridus</i>)	C
Spiny (<i>Amaranthus spinosus</i>)	C
Poinsettia, wild (<i>Euphorbia cyathophora</i>)	C
Prickly lettuce (<i>Lactuca serriola</i>)	C
Prickly sida/Teaweed (<i>Sida spinosa</i>)	C
Puncturevine (<i>Tribulus terrestris</i>)	C
Purslane (<i>Portulaca oleracea</i>)	C
Ragweed	
Common (<i>Ambrosia artemisiifolia</i>)	C
Giant (<i>Ambrosia trifida</i>)	S
Redweed	C
Russian thistle (<i>Salsola kali</i>)	C
Sage, Barnyard	S
Sesbania (<i>Sesbania</i> spp.)	C
Shephard's-purse (<i>Capsella bursa-pastoris</i>)	C
Sicklepod (<i>Cassia obtusifolia</i>) ¹	C
Smartweeds (<i>Polygonum</i> spp.)	
Ladysthumb (<i>Polygonum persicaria</i>)	C
Pennsylvania (<i>Polygonum pennsylvanicum</i>)	C
Spurge	
Prostrate (<i>Euphorbia humistrata</i>)	C
Spotted (<i>Euphorbia maculata</i>)	C
Spurred anoda (<i>Anoda cristata</i>)	C
Sunflower (<i>Helianthus</i> spp.)	C
Velvetleaf (<i>Abutilon theophrasti</i>)	C
Venice mallow (<i>Hibiscus trionum</i>)	C
Virginia pepperweed (<i>Lepidium virginicum</i>)	C
Waterhemp (<i>Amaranthus rudis</i>)	C
Wild mustards (<i>Brassica</i> spp.)	C

¹ For maximum control of sicklepod, use a preemergence application.

TABLE 2: ANNUAL GRASSES AND SEDGES CONTROLLED BY A392.03

C = Control S = Suppression or Erratic Control P = Poor or No Control

Weed Controlled	Level of Control
Barnyardgrass (<i>Echinochloa crus-galli</i>)	C
Bluegrass (<i>Poa annua</i>)	C
Broadleaf signalgrass (<i>Brachiaria platyphylla</i>)	C
Browntop millet (<i>Panicum ramosa</i>)	C
Crabgrass (<i>Digitaria</i> spp.)	C
Crowfootgrass (<i>Dactyloctenium aegyptium</i>)	C
Cupgrass (<i>Eriochloa gracilis</i>)	C

Foxtails (<i>Setaria</i> spp.)	C
Goosegrass (<i>Eleusine indica</i>)	C
Johnsongrass, Seedling (<i>Sorghum halepense</i>)	C
Junglerice (<i>Echinochloa colona</i>)	C
Millet, Wild-proso (<i>Panicum miliaceum</i>)	S
Nutsedge	
Yellow (<i>Cyperus esculentus</i>)	S
Purple (<i>Cyperus rotundus</i>)	S
Panicum, Fall (<i>Panicum dichotomiflorum</i>)	C
Panicum, Texas (<i>Panicum, texanum</i>)	S
Red rice (<i>Oryza sativa</i>)	C
Sandbur (<i>Cenchrus</i> spp.)	S
Shattercane (<i>Sorghum</i> ,bicolor)	S
Sorghum, Volunteer (<i>Sorghum</i> spp.)	S
Sprangletop, (<i>Leptochloa</i> spp.)	p
Stinkgrass (<i>Eragrostis</i> spp.)	p
Wheat, Volunteer (<i>Triticum</i> spp.)	S
Witchgrass (<i>Panicum capillare</i>)	C

A392.03 USE RATES FOR CONVENTIONAL TILLAGE SYSTEMS

A392.03 Used Alone in Preplant Incorporated Application

Incorporate **A392.03** uniformly into the top 2 inches of soil within 14 days before planting using a disk, field cultivator, rolling cultivator or similar equipment. Use incorporated application if furrow irrigation is used or when a period of dry weather after application is expected.

A392.03 Used Alone in Preemergence Application

When used alone, **A392.03** can be applied broadcast by ground or aerially. This application may be made during planting or as a separate operation after planting but must be made before crop emergence. If dry weather follows preemergence application, cultivate uniformly with shallow tilling equipment that will not damage soybeans.

Restrictions

Do not apply to sand soils, or to sandy loam or loamy sand soils containing less than 2% organic matter.

Conversion Table

Pints of A392.03/A	Lbs Metolachlor	Lbs Metribuzin	Lbs Imazethapyr Acid
1.6	0.80	0.11	0.026
2.0	1.003	0.14	0.033
2.4	1.20	0.17	0.039
2.7	1.35	0.19	0.04
3.1	1.55	0.22	0.05
3.5	1.75	0.25	0.057
3.9	1.96	0.27	0.063

TABLE 3: A392.03 RATES WHEN USED ALONE IN PREPLANT OR PREEMERGENCE APPLICATION*

*(In ND, and in MN (north of Highway #210), do not apply more than 2.9 pints of A392.03 per acre per use season.)

Soil Texture	Organic Matter		
	0.5 to 2%	2 to 3%	Over 3% ³
	Pt of A392.03/A		

Coarse Soils ¹	Sandy loam	1.6 to 2.0 ⁴	1.6 to 2.0 ⁴	2.0 to 2.4
	Loamy sand	n/a ¹	1.6 to 2.0 ⁴	2.0 to 2.4
Medium Soils (Loam, Silt loam, Silt, Sandy clay, Sandy clay loam)		2.4 to 2.7		2.7 to 3.1*
Fine Soils (Silty clay, Silty clay loam ² , Clay, Clay loam)		3.1 to 3.5*		3.5 to 3.9*

¹Do not use on sand soils. On coarse-textured soils, do not use on loamy sand with less than 2% organic matter.

²Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.

³For preplant incorporated application, use the lower rate.

⁴For AL, AR, FL, GA, LA, MS, MO, NC, OK, SC, TN, TX, VA, see section below **In Coarse (Light) Soils**.

On soils with pH above 7.0, soybean injury caused by the metribuzin in this product may occur at rates higher than 3.3 pints per acre. To avoid injury; do not use this product at rates greater than 3.3 pints per acre on soils above pH 7.0.

In Coarse (Light) Soils

(Only in AL, AR, FL, GA, LA, MS, MO, NC, OK, SC, TN, TX, VA)

This product may be used at the rates specified in Table 4 as a preplant incorporated or preemergence application in coarse-textured, low organic matter soils in the states listed above. Refer to Table 4 and to the appropriate sections of this label for specific directions on use and restrictions.

TABLE 4: A392.03 RATES WHEN USED ALONE IN PREPLANT OR PREEMERGENCE APPLICATIONS ON COARSE SOILS (Only in AL AR FL. GA LA MS MO NC OK SC TN TX. VA)

Soil Texture		Organic Matter	
		0.5 to 1%	1% or above
		Pt of A392.03/A ²	
Coarse Soils	Sand	n/a ¹	1.6 to 2.7
	Sandy loam, Loamy sand	1.6 to 2.7	1.6 to 2.7

¹Do not use on sand with less than 1% organic matter.

²Use the higher rate under heavy weed pressure and/or soils higher in organic matter.

On soils with pH above 7.0, soybean injury caused by the metribuzin in this product may occur at rates higher than 3.3 pints per acre. To avoid injury, do not use this product at rates greater than 3.3 pints per acre on soils above pH 7.0.

HERBICIDES THAT MAY BE APPLIED POSTEMERGENCE FOLLOWING A392.03

If required, application of this product alone or in tank mixture may be followed by an application of a postemergence herbicide to provide additional control of certain weeds. The following postemergence herbicides may be applied:

Antik EC (Carfentrazone-ethyl, EPA Reg: 91234-247)	glyphosate herbicides ¹
Arrow® (Clethodim, EPA Reg: 66222-60)	Harmony® (Thifensulfuron, EPA Reg: 279-9577)
Assure® II (quizalofop-p-ethyl, EPA Reg: 5481-646)	Intensity® (Clethodim, EPA Reg: 34704-864)
Basagran® (Bentazon, EPA Reg: 66330-413)	Poast® (Sethoxydim, EPA Reg: 7969-58)
Classic® (Chlorimuron, EPA Reg: 5481-681)	Poast Plus® (Sethoxydim, EPA Reg: 7969-88)
Cobra® (Lactofen, EPA Reg: 59639-34)	Reflex® (Fomesafen, EPA Reg: 100-993)
FrontRunner (Cloransulam-methyl, EPA Reg: 91234-84)	Resource® (Flumiclorac, EPA Reg: 59539-82)
Flexstar® (Fomesafen, EPA Reg: 100-1101)	Rezult® (Bentazon, EPA Reg: 7969-112)
Fusilade® DX (Fluazifop-p-butyl, EPA Reg: 100-1070)	Synchrony® XP ² (Chlorimuron-ethyl + Thifensulfuron-methyl, EPA Reg: 352-648)
Fusion® (Fluazifop-p-butyl + Fenoxaprop-p-ethyl, EPA Reg: 100-1059)	Ultra Blazer® (Acifluorfen, EPA Reg: 70506-60)

¹Use on Roundup Ready® or glyphosate tolerant soybean varieties only.

²Use on STS™ soybean varieties only.

Refer to the **Directions for Use** on this label and the individual product labels for use directions, use rates, and special precautions and/or restrictions.

BURNDOWN WEED CONTROL

This product can be used as part of a burndown herbicide program for control of existing vegetation prior to soybean emergence in conservation tillage (reduced-tillage/no-till) systems. This product may be tank mixed with a 2,4-D low volatile ester (LVE) (such as Whiteout® (2-ethylhexyl ester of 2,4-dichlorophenoxyacetic acid, EPA Reg: 34704-1032)) and/or glyphosate herbicides for control of emerged weeds prior to crop emergence. Burndown tank mixes with **A392.03** can be applied before planting or prior to crop emergence.

DO NOT tank mix **A392.03** with clomazone containing herbicides.

Application

This product may be applied up to 30 days before planting or preemergence. Apply only by ground equipment when this product is used for burndown of existing vegetation in conservation tillage systems. Use the high end of the rate range for applications of this product made 14 to 30 days before planting. Refer to Tables 3 and 4 for rates of **A392.03** alone and to Table 5 for rates of tank mix partners.

TABLE 5: RATES OF TANK MIX PARTNERS TO BE USED IN COMBINATION WITH A392.03 FOR BURNDOWN APPLICATIONS

Product	Rate of Tank Mix Partner	Directions and Remarks
2,4-D LVE (Whiteout 2,4-D)	Refer to product label for use rates.	Apply at least 7 days preplant when using Whiteout at 0.25 to 0.5 lb AE ¹ /A and at least 30 days preplant with rates greater than 0.5 lb AE ¹ /A. Include crop oil concentrate (COC) at the rate of 1.0 gal/100 gal of spray solution (1 % v/v).
Glyphosate	Refer to product label for use rates.	Must be applied prior to crop emergence. Use the higher rates within the specified range as weeds approach the maximum weed heights listed in Table 6. Apply in 10.0 to 20.0 gal of water/A. Refer to the label for spray adjuvant instructions. Any glyphosate formulation registered and labeled for use in soybeans may be tank mixed with this product.
Glyphosate (Mad Dog or Makaze brands) + 2,4-D LVE (Whiteout 2,4-D)	Refer to product label for use rates	Follow the Directions and Remarks section above for Whiteout 2,4-D and glyphosate, paying special attention to planting restrictions with Whiteout. Refer to the Glyphosate label for spray adjuvant instructions. Do not use crop oil concentrate (COG).

¹AE = 2,4-D acid equivalent

Precautions

Do not apply these treatments after crop emergence. Observe all precautions and limitations on the labeling of all products used in tank mixtures.

- Apply only 2,4-D LVE formulations (such as Whiteout 2,4-D) that are registered for preplant or burndown use.
- Do not apply tank mixtures containing 2,4-D LVE (Whiteout 2,4-D) if wind is blowing toward desired susceptible plants (i.e. cotton, tobacco, tomato, etc.) or when wind speeds exceed 6 mph.

Observe all precautions and limitations of all products used in tank mixtures.

Follow the most restrictive preharvest interval of all products used in a tank mixture.

Weeds Controlled

A392.03 in tank mixtures with the herbicides listed in Table 6 will provide burndown control of the weeds listed below.

TABLE 6: WEEDS CONTROLLED WITH TANK MIXES OF A392.03 IN BURNDOWN APPLICATION

Weeds Controlled	Whiteout 2,4-D	Glyphosate	Glyphosate + Whiteout 2,4-D
Annual Grasses	Maximum Burndown Height (Inches)		
Barley	Does not improve control of these species.	8	
Barnyardgrass		6	
Crabgrass spp.		6	
Foxtail spp.		8	
Johnsongrass, Seedling		8	
Panicum, Fall		6	
Sandbur, Field		8	
Wheat, Volunteer		6	
Witchgrass		6	
Buffalobur		6	6
Chickweed, Common	6	6	6
Cocklebur, Common	6	6	8
Dandelion, Common	6 dia	2 dia	6 dia
Henbit	4	4	4
Horseweed (Marestail)	6	4	6
Jimsonweed	6	6	6
Kochia	4	4	4
Ladysthumb	6	6	8
Lambsquarters, Common	6	6	8
Lettuce, Prickly	6	6	6
Mallow, Venice	6	6	6
Morningglory spp.	6	2	4
Mustard spp.	6	6	8
Pennycress, Field	6	6	6
Pigweed spp. (annual)	6	6	8
Ragweed, Common	6	6	8
Ragweed, Giant	6	4	6
Shepherd's-purse	6	6	6
Sida, Prickly	6	4	4
Smartweed, Pennsylvania	6	6	8
Sunflower, Common	6	6	6
Thistle, Russian	4	2 - 4	4
Velvetleaf	6	6	8

Waterhemp spp.	6	6	8
----------------	---	---	---

A392.03 USE RATES FOR REDUCED- AND NO-TILL SYSTEMS

Preplant Surface Application

A392.03 may also be used in reduced-till and no-till systems. Applications may be made up to 30 days before planting or after planting, but before soybean emergence. Residual herbicides such as Canopy® (metribuzin + chlorimuron ethyl, EPA Reg: 352-444), FrontRunner, Command (Clomazone, EPA Reg: 279-3054), Python® (Flumetsulam, EPA Reg 5481-677), and Stealth® (Pendimethalin, EPA Reg: 34704-868) may be tank mixed for additional weed control. If weeds are present at time of application, burndown herbicides may be added to the tank mixes (see Burndown Weed Control section). Refer to the tank mix partner product labels for specific rates and use directions.

TABLE 7: A392.03 RATES FOR REDUCED AND NO-TILL SYSTEMS*

*(In ND, and in MN (north of Highway #210), do not apply more than 2.9 pints of A392.03 per acre per use season.

Soil Texture ¹	A392.03 (Pt/A ¹)
Coarse ² (Loamy sand, Sandy loam)	1.6 to 2.7
Medium (Loam, Silt, Sandy clay, Sandy clay loam)	2.7 to 4.0*
Fine (Silty clay, Silty clay loam ³ , Clay, Clay loam)	4.0*

¹Use low rate in specified range for low residue level or soils with less than 3% organic matter. Use the higher rate in specified range for high residue level or soils with greater than 3% organic matter.

²Do not use on sand soils. On coarse-textured soils, do not use on loamy sand soils with less than 2% organic matter.

³Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S. When using **A392.03**, treat this soil as fine-textured.

A392.03 SEQUENTIAL APPLICATION

More consistent control of broadleaf and grass weeds may be obtained by an early preplant (surface-applied or shallow incorporated) application of **A392.03**, followed by a second preemergence application after planting but before soybean emergence. A sequential application will decrease the need for tillage and/or burndown herbicides for the control of existing vegetation before planting, while providing residual control of weeds after planting.

Application

An early preplant application may be made 15 to 30 days before planting soybeans. Follow this application with a preemergence overlay application of **A392.03** after planting but before crop emergence. Follow directions on this label for sequential applications from 0 to 14 days before planting.

Where a rate range is listed, use the higher rates:

- In fields with a history of severe weed pressure.
- When the time between early preplant and preemergence overlay applications approaches the maximum 30 days.
- When the organic matter content of the soil is over 3%.
- When heavy crop residues are present on the soil surface.

When weeds exceed 1 to 1.5 inches in height or diameter at application, use a burndown herbicide, such as Glyphosate, or Whiteout.

Weeds Controlled

In addition to weeds controlled by **A392.03** alone, the sequential application improves control of the following annual broadleaf weeds: Buffalobur, Cocklebur, Common ragweed, Sunflower and Velvetleaf.

TABLE 8: A392.03 SEQUENTIAL USE RATES FOR REDUCED-TILL AND NO-TILL SYSTEMS (BROADCAST RATES)*
 *(In ND, and in MN (north of Highway #210), do not apply more than 2.9 pints of A392.03 per acre per use season.)

Soil Texture ¹	Early Preplant Application A392.03 (Pt/A)	Followed By	Preemergence Overlay Application A392.03 (Pt/A) ²
Coarse ¹ (Sand, Loamy sand, Sandy loam)	1.6 to 2.4	followed by	0.4 to 1.2*
Medium (Loam, Silt loam, Sandy Clay loam, Silt, Sandy clay)	2.0 to 2.7	followed by	0.8 to 1.6*
Fine (Silty clay loam ³ , Clay loam, Silty clay, Clay)	2.4 to 3.5*	followed by	1.2 to 2.0*

¹On coarse-textured soils, do not use on sandy soils with less than 1% organic matter. However, on coarse-textured soils with a calcareous surface area or a pH of 7.5 or higher, do not use on sand soils with less than 2% organic matter, or on loamy sand or sandy loam soils with less than 1% organic matter.

²Total not to exceed 4.0 pints of **A392.03** per acre per use season.

³Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S. When using **A392.03**, treat this soil as "fine-textured."

CROP ROTATION INTERVALS

Full-rate application of products containing chlorimuron-ethyl, chloransulam-methyl, flumetsulam, imazaquin the same year as **A392.03** may increase the risk of injury to sensitive follow crops. Consult the product labels for listed uses of these products in combinations.

Only rotational crops harvested at maturity may be used for feed or food.

TABLE 9: CROP ROTATION INTERVALS

Crop	Crop Rotation Intervals (Months)	Crop	Crop Rotational Intervals (Months)
Alfalfa	4.5	Popcorn ⁵	18
Asparagus	40	Potatoes ⁶	26
Bahiagrass ⁶	40	Rice	40
Barley, Spring (except ND) ²	9.5	Root crops	40
Barley/Winter (except ND) ²	9.5	Rye (except in ND, and in MN north of Hwy #210)	12
Cabbage ⁶	40	Rye in ND, and in MN north of Hwy #210	18
Canola ⁷	40	Safflower	18
Cantaloupe ⁶	40	Sainfoin	40
CLEARFIELD® Corn	8	Sorghum	18
Clover	12	Southern peas	12
Cotton	18	Sweet pepper transplants ⁶	40
Cucumber ⁶	40	Sweet potato transplants ⁶	40
Edible beans	12	Soybeans	0
Field corn ^{3,4}	8.5	Sunflower	18
Field corn (seed) ^{3,4}	8.5	Sugarcane	40
Flax	26	Sweet corn ⁵	18
Forage grasses	40	Tobacco	12

Lentils	40	Tomatoes	40
Lettuce	18	Tomato transplants ⁶	40
Lima	12	Watermelon ⁶	40
Oats	18	Wheat, Spring	8
Onion ⁶	40	Wheat, Winter ¹	4.5
Peanuts	12	Other crops not listed	40
Peas	8	-	-

¹If soybeans are furrow irrigated, till the soil prior to planting winter wheat. The beds should be broken up and the soil mixed with tillage equipment set to cut 4 to 6 inches deep.

²**DE, IN, KY, MD, NJ, OH, PA, and VA only:** Barley may be planted 4 months following a **A392.03** application in these states. **ND only:** Barley may be planted 18 months following a **A392.03** application.

³**Corn inbred lines:** Corn inbred seed lines may be planted the year following an application of **A392.03**. Several seed companies have tested a wide range of inbreds for sensitivity to **A392.03** soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, Loveland Products, Inc. has not been given access to the inbred data. Growers are directed to contact the seed company for information and directions regarding the planting of corn grown for seed in fields treated with **A392.03** the previous year. **Since growing conditions, environmental conditions and grower practices are beyond the control of Loveland Products, Inc. all risks and consequences associated with planting seed corn inbreds into fields treated previously with A392.03 shall be assumed by the user.**

⁴**AZ, HI, ID, MT, NV, OR, UT, WA, and WY only:** Field corn and field corn grown for seed may be planted 9-1/2 months after **A392.03** application.

⁵**IL, IN, IA, MN, OH, TN, and WI only:** Sweet corn and popcorn varieties may be planted the year following an application of **A392.03**. Some sweet corn and popcorn varieties may be injured when planted at less than 18 months following an application of **A392.03**. Before planting sweet corn for processing, contact the processor company for information and directions regarding the tolerance of sweet corn varieties planned for fields treated with **A392.03** the previous year. **DO NOT** plant fresh market sweet corn varieties prior to 18 months after **A392.03** use. Before planting popcorn, contact the popcorn company for information and directions regarding the tolerance of popcorn varieties planned for fields treated with **A392.03** the previous year. **Since growing conditions, environmental conditions and grower practices are beyond the control of Loveland Products, Inc., to the extent consistent with applicable law, all risks and consequences associated with planting sweet corn or popcorn varieties into fields treated previously with A392.03 shall be assumed by the user.** Stunting and maturity delay or other adverse effects may result when sweet corn or popcorn are planted following **A392.03** use.

⁶**AL, DE, FL, GA, IN, KY, MD, NJ, NC, PA, SC, and VA only:** This crop may be planted 18 months following the last application of **A392.03**.

⁷**CLEARFIELD® Canola:** CLEARFIELD varieties of canola may be planted as a rotational crop the 12 months after an application of **A392.03** at specified rates on soybeans.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[For plastic containers ≤ 5 gallons: Nonrefillable Container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

[For plastic containers > 5 gallons: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

[A392.03] is a trademark of Atticus, LLC

{LANGUAGE ON LABEL AFFIXED TO CONTAINER}

METOLACHLOR	GROUP	15	HERBICIDE
METRIBUZIN	GROUP	5	HERBICIDE
IMAZETHAPYR	GROUP	2	HERBICIDE

A392.03TM

[Alternate Brand Name: Trovidor]

[Herbicide for preemergent control of certain grasses and broadleaf weeds in soybeans.]

ACTIVE INGREDIENTS:	(% by weight)
Metolachlor.....	43.72%
Metribuzin**.....	6.14%
Imazethapyr***.....	1.38%
OTHER INGREDIENTS:	48.76%
TOTAL	100.0%

*contains 4.01 pounds of metolachlor per gallon.

**contains 0.56 pound of metribuzin per gallon.

***contains 0.13 pound of imazethapyr acid per gallon.

KEEP OUT OF REACH OF CHILDREN

WARNING-AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

For Chemical Emergency:

Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC Day or Night Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

WARNING

Causes skin irritation. Causes moderate eye irritation. Do not get on skin or on clothing. Avoid contact with eyes. Wear protective eyewear. This product may cause skin sensitization reactions in some people. Wear coveralls, chemical-resistant gloves and chemical-resistant footwear. A chemical resistant apron plus chemical resistant gloves are required when mixing, loading, cleaning up spills or equipment or when otherwise exposed to this product's concentrate. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS: For terrestrial uses: do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Surface Water Advisory: Metolachlor can contaminate surface water through ground spray drift. Under some conditions, metolachlor may also have a high potential for runoff into surface water (primarily via dissolution in runoff water) for several months post-application. These conditions include: poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[For plastic containers ≤ 5 gallons: Nonrefillable Container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

[For plastic containers > 5 gallons: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

See inside label booklet for additional Precautionary Statements and Directions for Use.

Manufactured for:
Atticus, LLC
5000 CentreGreen Way, Suite 100
Cary, NC 27513

EPA Reg. No.: 91234-245
EPA Est. No.: _____
NET CONTENTS: _____