



## OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

January 25, 2024

Kristen Cianni  
Regulatory Manager  
Atticus, LLC  
940 NW Cary Parkway, Suite 200  
Cary, NC 27513

Subject: PRIA Label and CSF Amendment – Added Chemical Hazard Statement; Updated Company Address.  
Product Name: A1101.18  
EPA Registration Number: 91234-264  
Application Date: March 10, 2022  
Decision Number: 582486

Dear Kristen Cianni:

The amended label and Confidential Statement(s) of Formula (CSF) referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, are acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Please note that the record for this product currently contains the following CSF:

- Basic CSF dated 02/23/2023

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to

sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Christopher Taylor at 202-566-2928 or at [taylor.christopher.m@epa.gov](mailto:taylor.christopher.m@epa.gov).

Sincerely,



Michael Walsh  
Product Manager 11  
Invertebrate & Vertebrate Branch 2  
Registration Division  
Office of Pesticide Programs

Attachment

{Note to reviewer: [Text] in brackets denotes optional or explanatory language}  
{Note to reviewer: {Text} in braces denotes where in the final label text will appear}  
{BOOKLET FRONT PANEL LANGUAGE}

SPINOSAD	GROUP	<b>5</b>	INSECTICIDE
----------	-------	----------	-------------

# A1101.18 [™]

[Alternate Brand Name: Fraxel PSP Insecticide]

Control of darkling beetles, hide beetles, house flies (adults and larvae), stable flies, little house flies (*Fannia canicularis*) in and around agricultural animal premises.

Control of Northern fowl mites (*Ornithonyssus sylviarum*) and poultry red mites (*Dermanyssus gallinae*) infesting poultry.

ACTIVE INGREDIENT:	(% by weight)
Spinosad	
(a mixture of spinosyn A and spinosyn D).....	44.2%
<b>OTHER INGREDIENTS:</b> .....	<u>55.8%</u>
<b>TOTAL</b> .....	100.0%

## KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)


[See inside label booklet for Directions for Use.]  
[See below for Directions for Use.]

For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC Day or Night Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

Notice: Shake Well Before Use. Avoid Freezing. Read the entire label before using. Use only according to label directions. See Directions for Use including storage and disposal inside label booklet

EPA Reg. No.: 91234-264  
EPA Est. No.:  
Net Contents:

Manufactured for:  
**Atticus, LLC**  
940 NW Cary Parkway, Suite 100  
Cary, NC 27513



## {LANGUAGE INSIDE BOOKLET}

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates.

This product is highly toxic to mollusks. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning or disposing of equipment washwaters. Apply this product only as specified on this label.

Applying this product when rain is not predicted for the next 24 hours will help reduce potential risk to aquatic invertebrates by reducing pesticide runoff from the treatment area into water bodies.

This product is highly toxic to bees and other pollinating insects exposed to direct treatment, or to residues in/on blooming crops or weeds. Protect pollinating insects by following label directions intended to minimize drift and to reduce risk to these organisms.

### PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow coming into contact with oxidizing agent. Hazardous chemical reaction may occur.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

**Notice: Shake well before use. Avoid freezing.**

### MIXING FOR APPLICATIONS THAT REQUIRE DILUTION

This product is intended to be mixed with water and applied as a coarse or low-pressure spray. Fill the tank of the sprayer with water to about 1/2 of the total spray volume required. Start agitation and add the required amount of product. Continue mixing and agitation while filling the spray tank to the required spray volume. Maintain sufficient agitation during the application to ensure uniformity of the spray. Prepare spray fresh for each use.

### PRODUCT INFORMATION

- This product may be applied as a general surface spray or a crack and crevice application. Applications may be made in and around poultry and livestock premises, including but not limited to poultry houses, beef and dairy barns, feedlots, stables, corrals and housing structures.
- This product may be used as a direct treatment to control Northern Fowl mites infesting poultry.
- Control is enhanced when interior and exterior perimeter applications are made in and around livestock and poultry structures. Normal sanitation practices of the structure consistent with good animal husbandry should be followed to enhance effectiveness of insect control agents.
- Lactating and non-lactating dairy and beef cattle, swine, sheep and poultry may be present at the time of premise treatment. Horses not intended for human consumption may also be present at the time of premise treatment.
- Cover or remove exposed feed and water from the area being treated.

### Restrictions

- Do not use in milking rooms.
- Do not apply as a fog or space spray.

## SPRAY DRIFT ADVISORIES

**THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.**

### IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

- Applicators need to be familiar with local wind patterns and terrain that could affect spray drift

## APPLICATIONS

TO CONTROL	APPLICATION
House flies (adults and larvae) Stable flies Little house flies	<ul style="list-style-type: none"> <li>• 2 fl. oz. (60 ml) <b>A1101.18</b> in 10 gallons of water will treat 5,000 - 10,000 ft<sup>2</sup>. Use enough water in a directed spray to contact flies and their resting surfaces. Spray thoroughly, but do not allow run-off to occur.</li> </ul>
Darkling beetles Hide beetles	<ul style="list-style-type: none"> <li>• Use enough water to ensure thorough coverage of the area treated, especially litter, along feed and water lines, walls and support beams. Use a sprayer of appropriate design that is equipped to effectively apply low to moderate volumes of spray.                             <ul style="list-style-type: none"> <li>○ Low volume spray: Up to 15 gallons of diluted spray applied to 20,000 ft<sup>2</sup></li> <li>○ Moderate volume spray: Up to 40 gallons of diluted spray applied to 20,000 ft<sup>2</sup></li> </ul> </li> <li>• 2 fl. oz. (60 mL) of product treats 5,000 ft<sup>2</sup> (8 fl. oz. of product treats 20,000 ft<sup>2</sup>).</li> <li>• For severe infestations, 4 fl. oz. of product may be used to treat 5,000 ft<sup>2</sup> (16 fl. oz. treats 20,000 ft<sup>2</sup>).</li> </ul>
Northern fowl mite ( <i>Ornithonyssus sylviarum</i> )	<ul style="list-style-type: none"> <li>• Dilute 3 fl. oz. of product in 10 gallons of water.</li> <li>• Apply no more than 1 gallon of coarse spray per 100 birds to ensure adequate coverage, directed toward the vent area.</li> </ul>
Poultry red mite ( <i>Dermanyssus gallinae</i> )	<ul style="list-style-type: none"> <li>• Dilute 2 fl. oz. of product in 5 gallons of water.</li> <li>• This dilution treats 2,500 ft<sup>2</sup>.</li> </ul>

### Egg Producing Poultry

**Northern Fowl Mite:** Spray birds in cages (layers) or on the floor (layers/breeders) with a coarse spray. Use a sprayer of appropriate design that is equipped to effectively apply the spray. Ensure adequate coverage of the vent area of the bird with a direct saturation spray for optimum efficacy. Re-application may occur every 14 days as needed.

**Poultry Red Mites:** Apply to areas where red mites live: this includes areas between joints, cracks and crevices, under bent surfaces, the ends of metal bars, any place where two surfaces connect, and any place where dust can accumulate. Application to the exposed and unexposed areas is required to obtain a complete and thorough treatment. Ensure a uniform spray with enough pressure to penetrate cracks and crevices where red mites are located during the day to achieve the highest efficacy of the treatment.

## Poultry Houses

**House Flies (adults and larvae), Stable Flies and little house flies:** Treat areas where flies rest: pay attention to side walls, the ends of buildings, upper sides of posts, crossbeams, and manure. Spray in early morning when flies are resting. Use enough water in a directed spray to contact flies and their resting surfaces thoroughly, but do not allow run-off to occur. Do not contaminate feedstuffs, water or watering and feeding equipment with premise treatment. Re-application may occur every 7 to 10 days as needed, or when fly populations reach nuisance levels.

**Darkling Beetles and Hide Beetles:** Apply to floor area, especially to litter around food and water lines and to walls and support beams, but do not allow run-off to occur. Application should also be made into cracks and crevices around insulation, and where pests have been seen or can find harborage. Reapply after each grow-out or sanitation procedure. Indoor control can be enhanced by making perimeter treatments to the outside of building foundations to prevent migrating adult beetles.

## Livestock Housing

**House Flies (adults and larvae), Stable Flies and Little house flies:** Apply to vertical and overhead surfaces where flies may congregate. Feeders, waterers and feedcarts should be covered before application to prevent contamination. Do not contaminate feedstuffs or water with premise treatment. Pay attention to animal areas including fence railings, alleyways, manure, stanchions, pipes, window and doors, and other areas where flies congregate. Spray in early morning when flies are resting. Reapplication may occur every 7 to 10 days as needed, or when fly populations reach nuisance levels.

### Resistance Management:

For resistance management, **A1101.18** contains a Group 5 insecticide. Any insect population may contain individuals naturally resistant to **A1101.18** and other Group 5 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same use sites. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of **A1101.18** or other Group 5 insecticides with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
  - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
  - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
  - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pests.
  - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
  - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use and record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor

- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Atticus, LLC at (948-465-4800).

### **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container only, tightly closed in a secure place. Keep out of reach of children and domestic animals. In case of leak or spill, remove with absorbent materials and dispose of as waste.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING:**

**For plastic containers ≤ 5 gallons: Nonrefillable Container:** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

### **LIMITATION OF WARRANTY AND LIABILITY**

**IMPORTANT: READ BEFORE USE.** Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

[A1101.18] is a trademark of Atticus, LLC

{LANGUAGE ON LABEL AFFIXED TO CONTAINER}

SPINOSAD	GROUP 5	INSECTICIDE
----------	---------	-------------

**A1101.18<sup>TM</sup>**

[Alternate Brand Name: Fraxel PSP Insecticide]

Control of darkling beetles, hide beetles, house flies (adults and larvae), stable flies, little house flies (*Fannia canicularis*) in and around agricultural animal premises.

Control of northern fowl mites (*Ornithonyssus sylviarum*) and poultry red mites (*Dermanyssus gallinae*) infesting poultry.

**ACTIVE INGREDIENT:** (% by weight)

Spinosad (a mixture of spinosyn A and spinosyn D) .....	44.2%
<b>OTHER INGREDIENTS:</b> .....	<u>55.8%</u>
<b>TOTAL</b> .....	100.0%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC Day or Night Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

**ENVIRONMENTAL HAZARDS:** This pesticide is toxic to aquatic invertebrates. This product is highly toxic to mollusks. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning or disposing of equipment washwaters. Apply this product only as specified on this label. Applying this product when rain is not predicted for the next 24 hours will help reduce potential risk to aquatic invertebrates by reducing pesticide runoff from the treatment area into water bodies. This product is highly toxic to bees and other pollinating insects exposed to direct treatment, or to residues in/on blooming crops or weeds. Protect pollinating insects by following label directions intended to minimize drift and to reduce risk to these organisms.

**PHYSICAL OR CHEMICAL HAZARDS:** Do not mix or allow coming into contact with oxidizing agent. Hazardous chemical reaction may occur.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container only, tightly closed in a secure place. Keep out of reach of children and domestic animals. In case of leak or spill, remove with absorbent materials and dispose of as waste.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING:**

**For plastic containers ≤ 5 gallons: Nonrefillable Container:** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

See inside label booklet for additional Precautionary Statements and Directions for Use.

Manufactured for:  
**Atticus, LLC**  
940 NW Cary Parkway, Suite 200  
Cary, NC 27513

**EPA Reg. No.: 91234-264**  
**EPA Est. No.:** \_\_\_\_\_  
**NET CONTENTS:** \_\_\_\_\_