



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

91234-277

Date of Issuance:

3/6/23

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

A1100.04

Name and Address of Registrant (include ZIP Code):

Kristen Cianni
Regulatory Manager
Atticus LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Continues page 2

Signature of Approving Official:

Melissa E Bridges, Acting Product Manager 07
Invertebrate & Vertebrate Branch 3
Registration Division (7505T)
Office of Pesticide Programs

Date:

3/6/23

2. You are required to comply with the data requirements described in the generic data call-in (GDCI) identified below:
 - a. Spiromesifen GDCI-024875-1535

You must comply with all of the data requirements within the established deadlines. If you have questions about the GDCI listed above, you may contact the Chemical Review Manager in the Pesticide Re-Evaluation Division: <http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. You have 18 months from the date of registration to provide these data.
4. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 91234-277.”
5. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance. If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 10/10/2021
- Alternate CSF 1 dated 10/10/2021

If you have any questions, please contact Ralph Narain at 202-566-2853 or at Narain.Ralph@epa.gov.

Enclosure: Stamped label

{Note to reviewer: [Text] in brackets denotes optional or explanatory language}
{Note to reviewer: {Text} in braces denotes where in the final label text will appear}
{BOOKLET FRONT PANEL LANGUAGE}

SPIROMESIFEN GROUP 23 INSECTICIDE/MITICIDE

A1100.04^[TM]

[Alternate Brand Name: Asenra]

Contains spiromesifen, the active ingredient used in Savate®.
A1100.04 is not manufactured, or distributed by Bayer Environmental Science, seller of Savate®.

ACTIVE INGREDIENT: (% by weight)

Spiromesifen: 2-oxo-3-(2,4,6-trimethylphenyl)-1-oxaspiro[4.4]non-3-en-4-yl	
3, 3-dimethylbutanoate.....	45.2%
OTHER INGREDIENTS:	<u>54.8%</u>
TOTAL	100.0%

A1100.04 contains 4 pounds of spiromesifen per US gallon (480 grams ai per liter).

**STOP – Read the label before use
KEEP OUT OF REACH OF CHILDREN**

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

[See inside label booklet for First Aid, [additional] Precautionary Statements, and Directions for Use.]
[See below for additional Precautionary Statements]


{Note to reviewer: If applicable commercially, the statement for use and disclaimer will appear on the front panel of the final product packaging.}

EPA Reg. No.: 91234-XX

EPA Est. No.:

Net Contents:

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513



{LANGUAGE INSIDE BOOKLET}

FIRST AID	
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration.• Call a poison control center or doctor for further treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
Note To Physician: No specific antidote is known. Treat symptomatically.	
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

**For Chemical Emergency:
Spill, Leak, Fire, Exposure, or Accident,
Call CHEMTREC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material: Barrier Laminate Butyl Rubber ≥ 14 mils Nitrile Rubber ≥ 14 mils Neoprene Rubber ≥ 14 mils Natural Rubber ≥ 14 mils Polyethylene Polyvinyl Chloride (PVC) ≥ 14 mils Viton ≥ 14 mils.
- Shoes plus socks

Applicators must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove Personal Protective Equipment immediately after handling this product.
- Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Avoid contamination of surface water through runoff or spray drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater.

SURFACE WATER ADVISORY: This product may contaminate water through runoff or through drift of spray in wind. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

GROUNDWATER ADVISORY: Degradates of spiromesifen have properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow to come in contact with oxidizing or reducing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read entire label and Conditions of Sale before using this product.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours following application. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, are:

- Coveralls worn over short sleeved shirt and short pants
- Chemical resistant gloves made of any waterproof material: Barrier Laminate Butyl Rubber ≥ 14 mils Nitrile Rubber ≥ 14 mils Neoprene Rubber ≥ 14 mils Natural Rubber ≥ 14 mils Polyethylene Polyvinyl Chloride (PVC) ≥ 14 mils Viton ≥ 14 mils.
- Shoes plus socks

RESISTANCE MANAGEMENT

For resistance management, **A1100.04** contains a Group 23 insecticide. Any insect population may contain individuals naturally resistant to **A1100.04** and other Group 23 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of **A1100.04** or other Group 23 insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues for the targeted pests between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - o Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
- Rotate the use of **A1100.04** or other Group 23 insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues for the targeted pests between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - o Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - o Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - o When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - o Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - o The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
 - o Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Atticus, LLC at 984-465-4800.

SPRAY DRIFT ADVISORIES

**THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.**

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

WIND

Drift potential generally increases with wind speed. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boomless Ground Applications:

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.

GENERAL INFORMATION

A1100.04 is a Flowable (Suspension Concentrate) formulation that is intended for control of mites and whiteflies on ornamental plants, flowers and foliage plants in the greenhouse and field and container nurseries. The active ingredient contained in **A1100.04** is active on all mite development stages. However, mite juvenile stages are often more susceptible than adults. Application should be timed to coincide with early threshold levels in a developing mite population.

A1100.04 is also highly effective against whitefly nymphs, plus it has a significant effect on the otherwise difficult to control pupal stage. For best results against whitefly, applications should target these life stages.

A1100.04 can be applied by ground equipment. Thorough coverage of all plant parts is required for optimum performance. Performance evaluations should be made 4 – 10 days after application.

Apply specific dosage of **A1100.04** as needed for control. For best results good coverage of the upper and lower leaf surfaces is recommended. An adjuvant may be used to improve coverage on hard to wet foliage. For best results the treatment should be made when whitefly or mite population begins to build and before a damaging population becomes established. Rate range is provided and is generally dependent on size of the plant and density of the foliage. Apply with ground application equipment in adequate water for uniform coverage.

PRODUCT FEATURES

- Controls all stages of whitefly (nymphs, pupae, and adults).
- Provides knockdown and residual control of all developmental stages of mites

PESTS CONTROLLED	USE RATE
Spider mites (including twospotted spider mite, Southern red mite, Lewis mite, tumid mite, maple spider mite, spruce spider mite, honeylocust spider mite, euonymus mite, boxwood spider mite)	1 to 4 fl oz (30 – 120 mL)/100 gallons of spray solution or
Tarsonemid mites (including broad mite,	1.5 – 6 mL/5 gallons of spray solution

<p>cyclamen mite)</p> <p>Tenuipalpid mites (including false spider mite)</p> <p>Rust and blister mites (family <i>Eriophyidae</i>)</p>	
<p>Whiteflies (Including greenhouse, silverleaf and sweetpotato whitefly)</p>	<p>2 to 4 fl oz (60-120 mL)/100 gallons of spray solution</p> <p>or</p> <p>3-6 mL/5 gallons of spray solution</p>
<p>WHERE TO APPLY</p>	<ul style="list-style-type: none"> To shrubs, trees (including non-bearing fruit and nut trees), flowers and foliage plants in greenhouses, shade houses and plants grown in field and container nurseries.
<p>HOW TO APPLY</p>	<ul style="list-style-type: none"> Apply specific dosage of A1100.04 as needed for control. For best results, apply to achieve uniform coverage of the upper and lower leaf surfaces but avoid excess application that results in runoff from plant foliage. An adjuvant may be used to improve coverage on hard-to-wet foliage. For best results the treatment should be made when the whitefly or mite population begins to build and before a damaging population becomes established. A1100.04 is very effective against the egg and nymphal stages of whiteflies and mites. Control should be directed at these stages. Apply as a full coverage spray to the foliage using a minimum volume of 1 - 2 qts of final solution per 100 sq ft (or 100 to 200 gal per acre). Actual spray volume will vary depending on the size of plants. For maximum efficacy against adult whiteflies, use the 4 fl oz rate of A1100.04. For heavy infestations, two applications may be necessary to control adult whiteflies. Rate range is provided and is generally dependent on size of the plant and density of the foliage. Apply when pests first appear and prior to leaf damage or discoloration. Apply with ground application equipment in adequate water for uniform coverage. Do not apply via chemigation. Do not apply A1100.04 more than three times per season.
<p>PLANT TOLERANCE</p>	<ul style="list-style-type: none"> A1100.04 has been evaluated for phytotoxicity on a wide range of ornamental plants. However, due to the large number of species and varieties of ornamental plants, it is impossible to test every one for tolerance to A1100.04. The professional user should determine if A1100.04 could be used safely prior to commercial use. In a small area, test the recommended rates on a small number of plants for phytotoxicity prior to widespread use. Before using A1100.04 in tank mixture with other products and adjuvants, test the mixture on a small number of plants for phytotoxicity prior to widespread use.

	<ul style="list-style-type: none"> Do not use A1100.04 on geraniums (<i>Pelargonium</i> sp.), <i>Peperomia</i>, <i>Dracaena</i>, and 'Classy', 'Attache', or 'Vogue' varieties of rose. Insufficient information on tolerance is available for the following ornamentals: lily-of-the-Incas (<i>Alstromoeria</i>), New Guinea impatiens, <i>Bacopa</i>, ferns, phlox, English ivy (<i>Hedera helix</i>), cyclamen, orchids[*], hoyo[*], <i>Agyranthemum</i>, hydrangea, schefflera, <i>Matthiola</i> (stock), Mexican heather, <i>Lobelia</i>, fuschia, <i>Cordyline</i>, croton, neanthebella palm, and primula. It is recommended that A1100.04 not be used on these plants. Do not apply more than 2 fl oz per 100 gal per application or 4 fl oz per cropping cycle to chrysanthemum, Shasta daisy, snapdragon, impatiens, verbena, lantana, gerbera daisy, or marigolds. <p>[*Not for Use in California]</p>
IRRIGATION TIMING	<ul style="list-style-type: none"> If irrigation is used, conduct irrigation efficiently to prevent excessive loss of irrigation water through runoff. Time applications to allow sprays to dry prior to rain or sprinkler irrigation. Allow at least 24 hours between applications of product and any irrigation that results in surface water runoff into lakes, reservoirs, rivers, permanent streams, marshes, potholes, vernal pools, natural ponds, estuaries, or commercial fish farm ponds.
ENDANGERED SPECIES	<ul style="list-style-type: none"> The use of any pesticide in a manner that may kill or otherwise harm endangered species or adversely modify their habitat is a violation of Federal law.
RESTRICTIONS	<ul style="list-style-type: none"> Do not use through irrigation systems. Do not use on vegetable gardens. Do not use on plants intended for use as feed or forage. Do not use on bearing fruit or nut trees. Maximum Single Application Rate for ornamentals for commercial production, residential trees, and greenhouses: 0.125 lbs. ai/A per application. Do not use more than 0.35 lbs. spiromesifen per acre per calendar year.
PACKAGE HANDLING	<ul style="list-style-type: none"> A1100.04 is packaged in polyethylene containers. Do not allow product or containers to freeze.
COMPATIBILITY / MIXING / ORDER-OF-MIXING	
<ul style="list-style-type: none"> A1100.04 is physically and biologically compatible with many registered pesticides and fertilizers or micronutrients. When considering mixing A1100.04 with other pesticides, or additives, first contact your supplier for advice. For further information, contact your local Atticus, LLC representative. If your supplier and Atticus, LLC representative have no experience with the combination you are considering, you should conduct a test to determine physical compatibility. To determine physical compatibility, add the recommended proportions of each chemical with the same proportion of water as will be present in the chemical supply tank, into a suitable container, mix thoroughly and allow to stand for five minutes. If the combination remains mixed, or can be readily re-mixed, the mixture is considered physically compatible. A1100.04 may be used with other recommended pesticides, fertilizers and micronutrients. The proper mixing procedure for A1100.04 alone or in tank mix combinations with other pesticides is: <ol style="list-style-type: none"> Fill the spray tank 1/4 to 1/3 full with clean water; 	

2. While recirculating and with the agitator running, add any products in PVA bags. Allow time for thorough mixing;
 3. Continue to fill spray tank with water until 1/2 full;
 4. Add any other wettable powder (WP) or wettable granules (WG) products;
 5. Add the required amount of **A1100.04**, and any other “flowable” (FL or SC) type products;
 6. Allow enough time for thorough mixing of each product added to tank;
 7. If applicable, add any remaining tank mix components: emulsifiable concentrates (EC), fertilizers and micronutrients.
 8. Fill spray tank to desired level and maintain constant agitation to ensure uniformity of spray mixture.
- Mix pesticides in areas not prone to runoff such as concrete mixing/loading pads, disked soil in flat terrain or graveled mix pads, or use a suitable method to contain spills and/or rinsate. Properly empty and triple-rinse pesticide containers at time of use.

For Use in California:

Allow growth of a vegetative filter strip within 25-ft (on which this product should not be applied) along lakes, reservoirs, rivers, permanent streams, marshes, potholes, vernal pools, natural ponds, estuaries or commercial fish farm ponds. This requirement does not apply to retention or holding ponds used to collect and recycle water for irrigation.

[Optional Marketing Claims]

- For Greenhouse and Nursery Use
- Miticide/Insecticide

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[For containers ≤ 5 gallons: Nonrefillable Container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

[For containers > 5 gallons: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

A1101.04 is a trademark of Atticus, LLC
Savate® is a registered trademark of Bayer.

{LANGUAGE ON LABEL AFFIXED TO CONTAINER}

SPIROMESIFEN	GROUP	23	INSECTICIDE/MITICIDE
--------------	-------	----	----------------------

A1100.04^[TM]

[Alternate Brand Name: *Asebra*]

Contains spiromesifen, the active ingredient used in Savate®.

A1100.04 is not manufactured, or distributed by Bayer Environmental Science, seller of Savate®.

ACTIVE INGREDIENT:	(% by weight)
Spiromesifen: 2-oxo-3-(2,4,6-trimethylphenyl)-1-oxaspiro[4.4]non-3-en-4-yl 3, 3-dimethylbutanoate.....	45.2%
OTHER INGREDIENTS:	54.8%
TOTAL	100.0%

A1100.04 contains 4 pounds of spiromesifen per US gallon (480 grams ai per liter).

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If inhaled:	<ul style="list-style-type: none"> ● Move person to fresh air. ● If person is not breathing, call 911 or an ambulance, then give artificial respiration. ● Call a poison control center or doctor for further treatment advice.
If swallowed:	<ul style="list-style-type: none"> ● Call a poison control center or doctor immediately for treatment advice. ● Have person sip a glass of water if able to swallow. ● Do not induce vomiting unless told to do so by the poison control center or doctor. ● Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> ● Take off contaminated clothing. ● Rinse skin immediately with plenty of water for 15-20 minutes. ● Call a poison control center or doctor for treatment advice.
Note To Physician: No specific antidote is known. Treat symptomatically.	
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

For Chemical Emergency:

Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish and aquatic invertebrates. Avoid contamination of surface water through runoff or spray drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not

contaminate water when cleaning equipment or disposing of equipment washwater.

SURFACE WATER ADVISORY: This product may contaminate water through runoff or through drift of spray in wind. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

GROUNDWATER ADVISORY: Degradates of spiromesifen have properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow to come in contact with oxidizing or reducing agents. Hazardous chemical reaction may occur.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[For containers ≤ 5 gallons: Nonrefillable Container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

[For containers > 5 gallons: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

See inside label booklet for additional Precautionary Statements and Directions for Use.

{Note to reviewer: If applicable commercially, the statement for use and disclaimer will appear on the front panel of the final product packaging.}

Manufactured for:
Atticus, LLC
 940 NW Cary Parkway, Suite 200
 Cary, NC 27513

EPA Reg. No.: 91234-XX
EPA Est. No.: _____
NET CONTENTS: _____

{Optional Marketing graphics}

