



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

91234-291

Date of Issuance:

3/14/25

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration

(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

A177.03

Name and Address of Registrant (include ZIP Code):

Beth Anderson
Atticus, LLC
5000 CentreGreen Way, Suite 100
Cary, NC 27513

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. You have 18 months from the date of registration to provide these data.

Continues page 2

Signature of Approving Official:

Jacquelyn Herrick, Product Manager 3
Invertebrate and Vertebrate Branch 1, Registration Division
(7505T)

Date:

3/14/25

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 91234-291."
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 4/3/2023

If you have any questions, please contact Elizabeth Andrews at 202-566-2467 or at Andrews.Elizabeth@epa.gov.

Enclosure

{Note to reviewer: [Text] in brackets denotes optional or explanatory language}
{Note to reviewer: {Text} in braces denotes where in the final label text will appear}
{BOOKLET FRONT PANEL LANGUAGE}

BETA-CYFLUTHRIN	GROUP 3A	INSECTICIDE
-----------------	----------	-------------

A177.03 [™]

[Alternate Brand Name: Tirade Ultra WSP]

Contains beta-cyfluthrin, the active ingredient used in [Tempo® Ultra WSP].

For Pest Management Professionals and commercial use only [For broad-spectrum control of crawling, flying, and wood-infesting insect pests on indoor and outdoor surfaces, as well as insect pests of trees, landscape ornamentals, residential, and commercial lawns]

ACTIVE INGREDIENT: (by weight)

Beta-Cyfluthrin, cyano(4-fluoro-3-phenoxyphenyl)methyl 3-(2,2-dichloroethenyl)- 2,2-dimethylcyclopropane-carboxylate.....10.0%

OTHER INGREDIENTS:90.0%

TOTAL100.0%

**STOP – READ THE LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN**

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you DO NOT understand the label, find someone to explain it to you in detail.)

See [below] [inside label booklet] for [additional] [First Aid,] [and] [Precautionary Statements] [and] [Directions for Use].

[A177.03] is not manufactured, or distributed by Bayer Environmental Science, seller of [Tempo® Ultra WSP].

{Note to reviewer: If applicable commercially, the statement for use and disclaimer will appear on the front panel of the final product packaging.}

EPA Reg. No.: 91234-XX

EPA Est. No.:

Net Contents:

Manufactured for:

Atticus, LLC

940 NW Cary Parkway, Suite 200

Cary, NC 27513

1

ACCEPTED

03/14/2025

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under

EPA Reg. No. 91234-291

{LANGUAGE INSIDE BOOKLET}

FIRST AID	
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• DO NOT induce vomiting unless told to do so by the poison control center or doctor.• DO NOT give anything to an unconscious person.
NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.	
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

For Chemical Emergency:
Spill, Leak, Fire, Exposure, or Accident,
Call CHEMTREC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Harmful if swallowed, inhaled, or absorbed through the skin. **DO NOT** get in eyes, on skin, or on clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and warm water after handling. Remove contaminated clothing and wash before reuse. Wear safety glasses, goggles, or face shield when handling the undiluted material, and wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters when making general surface overhead treatments indoors. **DO NOT** allow children or pets to enter treated areas until surfaces are dry.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Remove from premises or tightly cover fish tanks and disconnect aerators when applying indoors where such containers are present. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. To protect the environment, **DO NOT** allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters.

Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or residues on crops or weeds. **DO NOT** apply **A177.03** or allow it to drift onto crops or weeds on which bees are actively foraging. Additional information may be obtained by consulting your Cooperative Extension Service.

Physical Or Chemical Hazards

Do not mix or allow to come into contact with any oxidizing agent. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
For both indoor and outdoor use.

PRO FACTS <ul style="list-style-type: none"> • Quick Knockdown & Residual Control • Low Odor • Non-staining • Ideal for Power Spray Applications • No phytotoxicity on a wide range of ornamentals 	
A177.03	<ul style="list-style-type: none"> • Water soluble packets containing 10% beta-cyfluthrin wettable powder.
CONTROLS	<ul style="list-style-type: none"> • Indoor Pests <ul style="list-style-type: none"> ○ General Household Pests ○ Pantry and Stored Product Pests ○ Food Processing Pests • Wood-Infesting Pests • Turf and Ornamental Pests <p><i>For list of specific insects, see inside</i></p>
WHERE TO APPLY	<p>Apply where pests have been seen or found, or can find shelter.</p> <ul style="list-style-type: none"> • In and around buildings and structures. • On residential, commercial, and recreational areas of turf. • On ornamentals in landscapes and interior plantscapes.
COVERAGE	<ul style="list-style-type: none"> • 1 packet treats up to 10,000 square feet outdoors. • 1 packet makes up to 5 gallons of finished spray solution for indoors.
MIXING INFORMATION	<ul style="list-style-type: none"> • The inner packet contains A177.03 and dissolves in water. • Remove the inner packet and drop into spray tank while adding water
HOW TO APPLY	<ul style="list-style-type: none"> • General surface, spot, mist, or crack & crevice application. • Broadcast application to turf and foliar application to ornamentals.
RE-APPLY	<ul style="list-style-type: none"> • Every 7 to 10 days, if needed. • For turf, reapply if needed, but not more than 6 times per year.
RE-ENTRY	<ul style="list-style-type: none"> • Avoid contact with treated surfaces until dry. • People and pets may re-enter after the treated area is dry.
APPLICATION RESTRICTIONS	<p>All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:</p>

	<ol style="list-style-type: none"> 1. Application to pervious surfaces such as soil, lawn, turf, and other vegetation; 2. Perimeter band treatments of 7 feet wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or lawn); 3. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure; 4. Applications around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a band not to exceed one inch; 5. Applications to vertical surfaces (such as the side of a man-made structure) directly above impervious surfaces (e.g., driveways, sidewalks, etc.), up to 2 feet above ground level; 6. Applications to vertical surfaces directly above pervious surfaces, such as soil, lawn, turf, mulch or other vegetation) only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters. <ul style="list-style-type: none"> • DO NOT apply or irrigate to the point of run-off. • DO NOT apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment. • Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.). • For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds. • DO NOT spray the product into fish pools, ponds, streams, or lakes. DO NOT apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. • DO NOT allow the product to enter any drain during or after application • DO NOT make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours). • Rainfall within 24 hours after application may cause unintended runoff of pesticide application. • DO NOT apply when the wind speed is greater than 15 mph. • Treat surfaces to ensure thorough coverage but avoid runoff. • To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.
QUESTIONS	<ul style="list-style-type: none"> • For questions or comments 984-465-4800

Resistance Management

For resistance management, A177.03 contains a group 3A insecticide. Any insect population may contain individuals naturally resistant to A177.03 and other Group 3A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of A177.03 or other group 3A insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the insecticide Resistance Action Committee (IRAC):

- Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
- Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
- When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pests.
- Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
- The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological, and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Atticus, LLC at 984-465-4800.

STRUCTURAL PESTS			
FOR RESIDUAL PEST CONTROL IN AND AROUND BUILDINGS, STRUCTURES, FOOD/FEED HANDLING ESTABLISHMENTS, AND MODES OF TRANSPORT. Use A177.03 as a general surface, spot, mist, or crack and crevice treatment in and around buildings and structures. This includes schools, factories, stores, warehouses, hospitals, food processing facilities, and modes of transport. For a list of common facilities where this product may be used, see Appendix.			
STRUCTURAL PESTS			
OUTDOOR/PERIMETER PESTS			
Ants Bees Boxelder bugs Carpenter ants Centipedes Cockroaches Cluster flies	Crickets Elm leaf beetles Fire ants Firebrats Fleas Flies Gnats	Ground beetles Hornets Midges Millipedes Moths Pillbugs Scorpions	Silverfish Sowbugs Spiders Ticks Wasps Yellowjackets
WHERE TO APPLY	<ul style="list-style-type: none"> ● Apply where pests have been seen or found, or can find shelter. ● Apply to buildings, garages, fences, and other areas where pests are found. ● Apply to porches and patios as a spot or crack and crevice treatment. ● To prevent infestation, treat the building foundation, walls, and around doors, windows, and soffits where pests are active or may enter or hide. Also treat the soil, turf, or other substrates adjacent to buildings. 		
USE RATE	<ul style="list-style-type: none"> ● One 50 g packet covers 2,500 square feet (20 g/1,000 square feet) to 5,000 square feet (10 g/1000 square feet). See Table 1 for mixing directions. ● For fire ants and flying pests: <ul style="list-style-type: none"> ○ 0.025% -- one 50 g packet / 5 gallons water ○ 0.05% -- one 50 g packet / 2.5 gallons water ● Always use the high rate for fire ants, bees, wasps, yellowjackets, and hornets. ● Use high rate for severe infestations, fast knockdown, and long residual. 		

<p>HOW TO APPLY</p> <p>(also see APPLICATIONS FOR OUTDOOR/PERIMETER PESTS)</p>	<ul style="list-style-type: none"> • Perimeter Treatment: To help prevent infestation of structures, treat area adjacent to buildings in a band up to 7 feet wide. Also treat the building foundation to a height of 2 feet in areas where pests are active or can find entrance. • Apply just enough dilution to adequately cover the area without excessive dripping or runoff. Volume can vary depending on the surface type treated. For example, mulch requires more spray than bare soil. If mulch is heavy, rake it back to expose soil before treating. • Avoid contact with treated surfaces until dry.
WHEN TO APPLY	<ul style="list-style-type: none"> • When pests are seen or found. • Re-apply every 7 to 10 days if needed.
MIXING INFORMATION	<ul style="list-style-type: none"> • The inner packet contains A177.03 and dissolves in water. • To prepare the spray mixture, drop the unopened inner packet into spray tank while adding water. • Operate the agitator while mixing. Depending on water temperature and agitation, the packets will dissolve within a few minutes from the time they are added to the water. <p>NOTE: DO NOT mix A177.03 with products that contain boron or release free chlorine. Mixing of these products prevents the breakdown of the water soluble packets. To avoid this problem, pre- dissolve packet(s) in a separate container of water and then pour mixture into the spray tank.</p> <ul style="list-style-type: none"> • Diluted spray mixture may be stored overnight. Agitate before using. • A177.03 can be tank mixed with pesticides currently registered for similar uses. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

TABLE 1: MIXING DIRECTIONS FOR OUTDOOR/PERIMETER APPLICATIONS

Application Volume: Gallons/1,000 square feet	Use Rate: Grams/1,000 square feet	Packets of A177.03 Diluted to These Volumes of Finished Spray	
		25 gal	50 gal
2.5	10	2	4
	20	4	8
5	10	1	2
	20	2	4
10	10	-	1
	20	1	2

Formula for Determining the Percent Concentration in the Finished Spray

(Packets of A177.03)			
(Gallons of finished spray)	x	7.57	= Percentage Concentration

APPLICATION FOR OUTDOOR/PERIMETER PESTS	
ANTS	Locate and treat ant nests, if possible. Follow trails of foraging workers, check under objects near the structure, and inspect other likely nest sites. If ants are nesting in the soil, thoroughly drench or inject the nest and surrounding soil. Treat ant trails around windows and doors, along sidewalks, or other areas where seen. Apply a perimeter treatment as described in the HOW TO APPLY box.
BEEES, WASPS, YELLOWJACKETS, HORNETS	Use Rate: 0.05% concentration. Make pin stream applications directly to the nests. To help avoid stings treat nests in the evening when the insects are inside the nest. For best results, check nests within one or two days to ensure complete kill. Remove and destroy treated nests to prevent emergence of newly hatched insects, secondary pest problems, and possible staining from honey. If removal is not possible, retreat nests if necessary
BOXELDER BUGS	Apply directly to boxelder bugs that have congregated around doors and windows and similar areas where they may enter structures. Apply perimeter treatment as described in the HOW TO APPLY box. For best results, treat infested trees early in the season. Tree spraying may require ornamental license - check with your local regulator.
CENTIPEDES, MILLIPEDES, PILLBUGS, SOWBUGS	Apply perimeter treatment to reduce the number of insects from entering the structure. Thoroughly treat mulched areas. Reduce moisture areas from around the structure.
FIRE ANTS	Use Rate: 0.05% concentration. Thoroughly drench or inject the entire mound. When using injection treatment, you must treat the bottom of the mound. Apply dilution or foam using a downward and upward motion of the injector tool. For complete distribution, inject in several points throughout the mound. Treat new mounds as they appear. DO NOT use in pastures or areas of food crop production.
FLEAS & TICKS	Make a perimeter treatment, concentrating on areas where pets rest or run. For ticks, treat likely harborage areas like along paths, tall grass, and other vegetation. For best results, treat the entire area where ticks could be present. Begin treating in the spring. Ticks may be re-introduced from surrounding areas by host animals. Continue treating until frost to control both larvae and adult ticks.
FLIES	Use Rate: 0.025% to 0.05% concentration. Apply low-volume mist or spray to outside surfaces of buildings, garages, fences, and other areas where flies rest. Apply to porches and patios as a spot or crack and crevice treatment. CLUSTER FLY: Treat in late summer or fall, just before flies begin looking for winter shelter. Siding, eaves, around windows and doors, window screens, and attics are limited to spot or crack and crevice treatments. See APPLICATION RESTRICTIONS for exceptions and permissible uses. To help reduce infestations, treat cracks and crevices, attics, and other indoor winter hiding places. Note: Cluster flies may appear during winter warm spells, while snow is still on the ground.
MOSQUITOES	Spray around landscape plants, turf and ground cover, under decks, around building foundations, and other places where mosquitoes may rest.
SCORPIONS	Treat around piles of firewood and other areas where scorpions may hide. When possible, remove piles of materials that can harbor the pests. Apply a perimeter treatment.

STRUCTURAL PESTS					
INDOOR PESTS					
General Household Pests		Pantry & Stores Product Pests		Food Processing Pests	
<u>Crawling Pests</u> Ants (except Pharaoh) Bed bugs Carpet beetles Centipedes Clothes moths Cluster flies Cockroaches Crickets Darkling beetles Earwigs Firebrats Millipedes	Pillbugs Silverfish Sowbugs Spiders <u>Flying Pests</u> Flies Gnats Hornets Midges Moths Wasps Yellowjackets	<u>Beetles</u> (exposed adults & immature stages) Cadelle Cigarette Confused flour Dermeestid Drugstore Granary weevil Hide Larder Leather Lesser grain borer	Lesser mealworm Merchant grain Mealworm Red flour Rice weevil Sawtoothed grain Warehouse <u>Moths</u> Indian meal Mediterranean flour	Fruit flies Phorid flies Sciarid flies	Also includes: General Household Pests Pantry & Stored Product Pests
WHERE TO APPLY	Apply where pests have been seen or found, or can find shelter. Use a general surface, spot, mist, or crack and crevice application.				
USE RATE	0.025% -- one 50 g packet / 5 gallons water 0.05% -- one 50 g packet / 2.5 gallons water <ul style="list-style-type: none"> Use high rate for severe infestations, fast knockdown, and long residual. Always use the high rate for bees, wasps, yellowjackets, and hornets. 				
HOW TO APPLY (Also see APPLICATION FOR INDOOR PESTS)	<ul style="list-style-type: none"> Remove or tightly cover fish tanks and disconnect aerators during application. Remove pets during application and until treated surfaces are completely dry. Avoid contact with treated surfaces until dry. Wear safety glasses, goggles, or face shield and wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters when treating overhead areas. <u>General Surface Applications</u> <ul style="list-style-type: none"> Use a low-pressure system (DO NOT exceed 50 psi at nozzle tip), with a fan-type nozzle to apply the dilution uniformly. In the home, cover all food processing surfaces and utensils or thoroughly wash following treatment. Cover exposed food or remove from area being treated. <u>Spot, Mist, or Crack and Crevice Applications</u> <ul style="list-style-type: none"> Apply directly into voids, cracks and crevices using spray or foam equipment. Spray: use a low-pressure system (DO NOT exceed 50 psi at nozzle tip), with pinpoint or variable pattern nozzle or applicator equipment that delivers low volume treatments, such as the Micro-Injector® or Actisol®. Foam: Use appropriate foam generating equipment to supply a sufficient volume of foam. Mix the dilution with manufacturer's recommended volume of foaming agent. 				
WHEN TO APPLY	<ul style="list-style-type: none"> When pests are seen or found. Re-apply every 7 to 10 days if needed. 				
MIXING INFORMATION	<ul style="list-style-type: none"> The inner packet contains A177.03 and dissolves in water. To prepare the spray mixture, drop the unopened inner packet into spray tank while adding water. Operate the agitator while mixing. Depending on water temperature and agitation, the packets will dissolve within a few minutes from the time they are added to the water. 				

	<p>NOTE: DO NOT mix A177.03 with products that contain boron or release free chlorine. Mixing of these products prevents the breakdown of the water soluble packets. To avoid this problem, pre- dissolve packet(s) in a separate container of water and then pour mixture into the spray tank.</p> <ul style="list-style-type: none"> • Diluted spray mixture may be stored overnight. Agitate before using. • A177.03 can be tank mixed with pesticides currently registered for similar uses. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.
RESTRICTIONS	<ul style="list-style-type: none"> • DO NOT treat entire area of floor or floor coverings. • DO NOT apply where electrical short circuits can occur. Use a dust or dry bait in these areas. • DO NOT apply in occupied hospital and/or nursing home rooms. Occupants may return when surfaces have dried. • DO NOT apply in occupied classrooms. • DO NOT apply to furniture or upholstery where prolonged contact by humans will occur. • DO NOT apply as a space spray.

ADDITIONAL DIRECTIONS FOR FOOD AND FEED HANDLING ESTABLISHMENTS	
WHERE TO APPLY	<ul style="list-style-type: none"> • Applications may be made to both food/feed and non-food areas of food and feed handling establishments. • Food/feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared, or served: <ul style="list-style-type: none"> ○ receiving, storing, and packing (canning, bottling, wrapping, boxing) food ○ preparing food ○ edible waste storage ○ enclosed processing systems (mills, dairies, edible oils, syrups) of food. • Serving areas where food is exposed and the facility is in operation are also considered food areas.
USE RATE	<ul style="list-style-type: none"> • Apply as a general surface, spot, mist, or crack & crevice application. 0.025% -- one 50 g packet / 5 gallons water 0.05% -- one 50 g packet / 2.5 gallons water • Use high rate for severe infestations, fast knockdown, and long residual.
HOW TO APPLY (Also see INDOOR PESTS)	<p><u>General Surface Applications</u> Cover or remove all food processing and handling equipment before treatment. After applying where food is commercially prepared or processed, wash all handling and processing equipment, benches, shelving, and other surfaces which food will contact. Rinse thoroughly with clean, fresh water. DO NOT apply to surfaces in food-handling areas when facility is in use or when foods are exposed. DO NOT apply directly to food.</p> <p><u>Spot or Crack & Crevice Applications</u> Spot or crack and crevice application is permitted while facility is in use. Cover or remove food from area being treated. DO NOT apply directly to food or food handling surfaces</p>

APPLICATION FOR INDOOR PESTS	
<p>GENERAL HOUSEHOLD PESTS</p> <p>Cockroaches, Crickets, Silverfish, Spiders, and Others</p>	<ul style="list-style-type: none"> • Use a coarse, low-pressure spray. • Treat entry points around water pipes, doors, windows, and eaves. Treat areas where pests normally feed or hide. This may also include: baseboards, corners, around water pipes, under refrigerators, cabinets, sinks, and stoves, being sure to spray cracks & crevices. Spot treat floor or rugs beneath furniture, in closets, and storage areas, but DO NOT apply to entire floor area. • DO NOT use as a space spray. <p>IGR NOTE: This product can be tank mixed with an insect growth regulator (IGR) for extended cockroach control</p>
ANTS	<ul style="list-style-type: none"> • Apply to ant trails on walls, tile, baseboards, around pipes, under cabinets, and other places where ants are active. • Treat entry points such as windows and doors and the outdoor perimeter. • For best results, treat nests when possible.
PANTRY AND STORED PRODUCT PESTS	<ul style="list-style-type: none"> • For best results, clean area before treating. Remove and destroy infested food. Discard food accidentally contaminated with spray. • For cupboards, shelving, or food storage areas: Remove utensils, exposed food, shelf paper, and other items before spraying. When dried, cover shelves with clean paper before replacing items.
COMMERCIAL STORED PRODUCT PESTS	<ul style="list-style-type: none"> • Commercial storage or transportation: Clean and treat warehouses, production facilities, storage areas, rail cars, truck beds, and other areas where products are stored before filling with the product.
PESTS IN AIRCRAFT	<ul style="list-style-type: none"> • DO NOT use in aircraft cabins. • May be applied to carpet or upholstery in the cargo area only where there is no prolonged human contact.
PESTS IN LIVESTOCK HOUSING STRUCTURES Including poultry houses & pet kennels	<ul style="list-style-type: none"> • Apply as a general surface and/or a crack and crevice spray. • For best results, clean facilities before applying and treat outdoor perimeter areas. • DO NOT apply indoors where animals other than cattle or horses are present. • DO NOT apply to animal feedstuffs or watering equipment.
FLYING INSECTS	<ul style="list-style-type: none"> • Apply to surfaces where pests collect or rest
STRUCTURAL PESTS	
WOOD-INFESTING PESTS	
<p><u>Beetles</u></p> <p>Ambrosia</p> <p>Deathwatch</p> <p>False powderpost</p> <p>Old house borer</p> <p>Powderpost</p>	<p>Carpenter ants</p> <p>Carpenter bees</p> <p><u>Termites</u></p> <p>Subterranean</p> <p>Formosan</p> <p>Drywood</p>
WHERE TO APPLY	<ul style="list-style-type: none"> • Apply where pests have been seen or found, or can find shelter.
USE RATE	<ul style="list-style-type: none"> • 0.05% -- one 50 g packet / 2.5 gal water
HOW TO APPLY (Also see Application section)	<ul style="list-style-type: none"> • Apply spray or foam as described in application section. • Avoid contact with treated surfaces until dry.
WHEN TO APPLY	<ul style="list-style-type: none"> • When pests are seen or found. • Re-apply every 7 to 10 days, if needed.

MIXING INFORMATION	<ul style="list-style-type: none"> • The inner packet contains A177.03 and dissolves in water. • To prepare the spray mixture, drop the unopened packet into the spray tank while filling with water. Operate the agitator while mixing. Depending on water temperature and agitation, the packets will dissolve within a few minutes from the time they are added to the water. <p>NOTE: DO NOT mix A177.03 with products that contain boron or release free chlorine. Mixing of these products prevents the breakdown of the water soluble packets. To avoid this problem, pre- dissolve packet(s) in a separate container of water and then pour mixture into the spray tank.</p> <ul style="list-style-type: none"> • Diluted spray mixture may be stored overnight. Agitate before using. • A177.03 can be tank mixed with pesticides currently registered for similar uses. Follow the most restrictive of labeling directions and precautions. • <u>Foam applications</u>: Mix end-use dilution with manufacturer's recommended volume of foaming agent for 0.05% concentration
APPLICATION FOR WOOD-INFESTING PESTS	
CARPENTER ANTS	<p><u>Around houses and other structures</u>: Apply around doors, windows, and other places where carpenter ants may enter. Spray or foam into cracks & crevices, or spray, mist or foam through small drilled holes into voids where ants or nests are present. Use enough spray or foam to cover the area. DO NOT exceed 1 gallon of dilution per 1,000 square feet.</p> <p><u>Inside trees, utility poles, fencing, and decking materials and similar structural members</u>: Drill to find the interior infested cavity and inject dilution or foam using an appropriate treatment tool with a splash back guard.</p> <p><u>Tunneling in soil</u>: Apply as a soil drench or inject the dilution or foam in the soil at intervals of 8 to 12 inches. Use 1 gallon of dilution per 8 square feet. Establish a uniform vertical barrier at the edge of walls, driveways, or other hard surfaces where ants are tunneling beneath the surfaces.</p> <p><u>Protecting firewood from carpenter ants</u>: Treat soil beneath firewood before stacking at 1 gallon of dilution per 8 square feet.</p> <p>DO NOT treat firewood with this product.</p>
CARPENTER BEES	<p><u>In structures</u>: Spray or foam directly into gallery entrance holes. Leave entrance open for 24 hours to be sure that returning adult bees are killed. When there is no activity, the hole may be sealed with a suitable sealant.</p>
WOOD-INFESTING BEETLES	<p><u>On surfaces in homes, apartments, and other structures</u>: For small areas, brush or spray evenly on wood surfaces.</p> <p>For large or overhead areas, thoroughly cover wood surfaces with a coarse spray. Protect surfaces below the treated area from accidental dripping by covering with disposable plastic sheeting or other material that can be thrown away after treatment.</p> <p><u>In localized areas</u>: Apply dilution or foam to voids and galleries in damaged wood, in spaces between wooden structural members, and between the sill plate and foundation. Apply to inaccessible areas by drilling, and then injecting, spraying, or misting the dilution or foam into the damaged wood or wall voids.</p>
DRYWOOD AND ABOVE-GROUND SUBTERRANEAN TERMITES	<p>Do not use this product as the sole source of control for active, structural infestations by subterranean termites; the purpose of this application is to kill workers or winged reproductive forms of termites which are present at the time of treatment. It is not intended to provide structural pest control. It is not a substitute for mechanical alteration, soil and foundation treatment, but merely a supplement. For active, structural infestations by subterranean termites, this product can only be used to supplement a federally registered conventional product that is registered as a sole source for termite control. This product will not eliminate termite infestations or provide protection against future infestation.</p> <p><u>In localized areas</u>: Apply dilution or foam to voids and galleries in damaged wood, to spaces</p>

	<p>between wooden structural members, and between the sill plate and foundation. Apply to inaccessible areas by drilling, and then injecting, spraying, or misting the dilution or foam into damaged wood or wall voids.</p> <p><u>Swarming or reproductive stages:</u> Apply dilution to the swarm or area on which the termites collect. Make applications to attics, crawl spaces, unfinished basements, or man-made voids with a coarse fan spray to control exposed reproductive forms of termites.</p> <p>Above-ground termites inside trees, utility poles, fencing, and decking materials and similar structural members: Drill to find the infested cavity and inject dilution or foam using an appropriate tool with a splash back guard.</p> <p><u>Termite carton nests:</u> Inject dilution or foam into building voids or tree cavities using a pointed injection tool. Several injection points to varying depths may be needed. For best results, remove carton nest material when possible.</p> <p><u>Protecting firewood from termites:</u> Treat soil beneath firewood before stacking at 1 gallon of dilution per 8 square feet.</p> <p>DO NOT treat firewood with this product.</p>
--	--

TURF PESTS		
A177.03 controls designated pests on landscape and recreational turf.		
TURF PESTS		USE RATE
Ants	Grasshoppers	50 grams (1 packet) per 5,000 to 10,000 square feet Use high rate for severe infestations, fast knockdown, and long residual.
Armyworms	Japanese beetles (adult)	
Crickets	Sod webworms	
Cutworms	Ticks (including deer ticks)	
Earwigs		
Black turfgrass ataenius (adult)	Fleas	50 grams (1 packet) per 5,000 square feet
Bluegrass billbugs (adult)	Hyperodes weevils (adult)	
Chinch bugs	Mole crickets	
Chiggers	Scorpions	
Fire ants (aids in control)		
WHERE TO APPLY	<ul style="list-style-type: none">• For use on landscape and recreational turfgrass. Not for use on turf grown for research, commercial seed production, sale, or other commercial use.• Not for use on golf course turf.• Apply where pests have been seen or found, or can find shelter. Permitted areas are home lawns, business complexes, apartment buildings, airports, cemeteries, parks, playgrounds, recreational fields, institutional grounds, athletic fields, and other landscape turfgrass.	
HOW TO APPLY (also see APPLICATION FOR TURF PESTS)	<ul style="list-style-type: none">• Use an accurately calibrated sprayer. Apply uniformly using spray nozzles with a coarse droplet size such as flood jet nozzles.• Keep children and pets off treated areas until lawn has dried.	
WHEN TO APPLY	Apply when pests first appear. Reapply as needed based on pest reinfestation. DO NOT treat more than 6 times per year. Consult your state Cooperative Extension Office for the specific timing in your area.	
MIXING INFORMATION	<ul style="list-style-type: none">• The inner packet contains A177.03 and dissolves in water.• To prepare the spray mixture, drop the unopened packet into the spray tank while filling with water. Operate the agitator while mixing. Depending on water temperature and agitation, the packets will dissolve within a few minutes from the time they are added to the water. <p>NOTE: DO NOT mix A177.03 with products that contain boron or release free chlorine.</p>	

	<p>Mixing of these products prevents the breakdown of the water soluble packets. To avoid this problem, pre-dissolve packet(s) in a separate container of water and then pour mixture into the spray tank.</p> <ul style="list-style-type: none"> • Diluted spray mixture may be stored overnight. Agitate before using. • A177.03 can be tank mixed with pesticides currently registered for similar uses. Follow the most restrictive of labeling directions and precautions. • COMPATABILITY: Compatible with commonly used fungicides, miticides, liquid fertilizers, and other insecticides.
RESTRICTIONS	<ul style="list-style-type: none"> • DO NOT apply this product by aerial application or by any type of chemigation system. • DO NOT apply when the average wind speed is greater than 15 mph. • DO NOT allow drift onto ponds, streams, or lakes. • DO NOT apply when turfgrass areas are waterlogged or the soil is saturated with water (i.e., will not accept irrigation).

APPLICATION FOR TURF PESTS	
ARMYWORMS, CUTWORMS, SOD WEBWORMS	DO NOT water or mow for at least 24 hours following application.
BILLBUGS	Apply when adults first appear in the spring. In temperate climates, this application will also control overwintering chinch bugs.
CHINCH BUGS	For best results, water the turf immediately after application to aid in the penetration of the active ingredient into the thatch layer where chinch bugs dwell.
TICKS (Including ticks that may transmit Lyme Disease and Rocky Mountain Spotted Fever)	For best results, treat the entire area where ticks could be present. Begin treating in the spring. Ticks may be re-introduced from surrounding areas by host animals. Continue treating until frost to control both larvae and adult ticks.

ORNAMENTAL PESTS				
A177.03 controls designated pests on trees, shrubs, foliage plants and flowers in outdoor landscaped areas and interior plantscapes where these plants are grown.				
Ants	Clover mites	June beetles (adult)	Peppertree psyllids	Striped oakworms
Aphids	Cockroaches	Lace bugs	Pillbugs	Tent caterpillars
Armyworms	Crickets	Leaf-feeding caterpillars	Pine shoot moths	Thrips
Azalea caterpillars	Cutworms	Leafhoppers	Pine tip moths	Ticks (including deer ticks)
Bagworms	Elm leaf beetles	Leafrollers	Plant bugs	Tussock moth larvae
Black vine weevil (adult)	Elm spanworms	Leaf Skeletonizers	Redhumped caterpillars	Walnut caterpillars
Boxelder bugs	Emerald ash borers (adult)*	Mealybugs	Sawfly larvae	Wasps
Bristly rose slugs	Flea beetles	Midges	Scale insects (crawler stages)	Webworms
Budworms	Flies	Mosquitoes	Sowbugs	Whiteflies
California oakworms	Fungus gnats	Oleander moth larvae	Spiders	Yellownecked caterpillars
Cankerworms	Grasshoppers	Orchid weevils	Spittlebugs	
Casebearers	Gypsy moth larvae	Pear psylla	Striped cucumber beetles	
	Japanese beetles (adult)			
*For emerald ash borer, make a minimum of two applications per season. Make the first application in late- May/early June or when adult beetles begin flying. Make subsequent applications approximately one month after the previous application.				

WHERE TO APPLY	<ul style="list-style-type: none"> • Use in landscapes and interiorscapes. Permitted areas are parks, recreational areas, athletic fields, institutional grounds, hotels, malls, office buildings, and airports. Not for use on plants or turf grown for research, commercial seed production, sale, or other commercial use. Not for use in greenhouses or nurseries. • Apply where pests are seen, have been found, or can find shelter among trees, shrubs, evergreens, ground covers, flowers, and foliage plants. • No phytotoxicity has been observed on a wide range of ornamentals tested under various environmental conditions. If in doubt, first try this product on a small sample.
USE RATE	<ul style="list-style-type: none"> • 1 to 3 packets A177.03 per 100 gallons of water. • Use high rate for severe infestations, fast knockdown, and long residual.
HOW TO APPLY (Also see APPLICATION FOR ORNAMENTAL PESTS)	<p>Apply as a dilute spray just to the point of run-off but DO NOT allow excess run-off. For hard-to-wet foliage such as holly, pine, or ivy, adding a spreader-sticker may enhance knockdown and residual activity. For best results, spray thoroughly for good coverage including the underside of leaves for pests that feed or rest there.</p> <ul style="list-style-type: none"> • Spray plants in bloom at times when pollinating insects are not present: early morning, late evening, or in temperatures below 50°F. • Remove any animal feeding dishes prior to treatment. <p>Avoid contact with treated surfaces until dry.</p>
WHEN TO APPLY	<ul style="list-style-type: none"> • Apply when pests first appear. Reapply as needed based on pest infestation. • Consult your state Cooperative Extension Office for the specific timing in your area.
MIXING INFORMATION	<ul style="list-style-type: none"> • The inner packet contains A177.03 and dissolves in water. • To prepare the spray mixture, drop the unopened packet into the spray tank while filling with water. Operate the agitator while mixing. Depending on water temperature and agitation, the packets will dissolve within a few minutes from the time they are added to the water. <p>NOTE: DO NOT mix A177.03 with products that contain boron or release free chlorine. Mixing of these products prevents the breakdown of the water soluble packets. To avoid this problem, pre-dissolve packet(s) in a separate container of water and then pour mixture into the spray tank.</p> <ul style="list-style-type: none"> • Diluted spray mixture may be stored overnight. Agitate before using. • A177.03 can be tank mixed with pesticides currently registered for similar uses. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. • COMPATABILITY: Compatible with commonly used fungicides, miticides, liquid fertilizers, and other insecticides
RESTRICTIONS	<ul style="list-style-type: none"> • DO NOT apply to any food crop. • DO NOT allow drift onto ponds, streams, or lakes.

APPLICATION FOR ORNAMENTAL PESTS	
ANTS, CRICKETS, SPIDERS, MIDGES, WASPS, AND FLIES	Apply as a general spray, concentrating on damp areas beneath shrubbery, around foundations and in areas of tall grass or weeds in ornamental areas and areas adjacent to building.
BAGWORMS	Spray when bagworms first begin to hatch. If possible, apply directly to the larvae. Young larvae are easier to control.
BLACK VINE WEEVILS (adult)	<p>Use rate: 2-3 packets per 100 gallons of water.</p> <p>Apply as needed when adults and damage are first noticed in spring and at 3 to 4 week intervals</p>

	while weevils and fresh leaf notching are observed. Addition of a spreader/sticker at recommended rates may enhance control of insects on certain species of ornamentals having difficult to wet foliage.
SCALE (CRAWLERS)	Spray thoroughly, including foliage, trunk, limbs, and twigs.

APPENDIX		
Use A177.03 as a general surface, spot, mist, or crack and crevice treatment. Permitted areas of use in buildings and structures, and modes of transport:		
<ul style="list-style-type: none"> • Apartment buildings • Bakeries • Bottling facilities • Breweries • Buses • Cafeterias • Candy plants • Cereal processing and manufacturing plants • Correctional facilities • Dairies and dairy product processing plants • Frozen food plants • Grain mills 	<ul style="list-style-type: none"> • Granaries • Greenhouses (structures only) (non-commercial) • Hospitals • Hotels • Houses • Industrial buildings • Kitchens • Laboratories • Livestock housing • Pet kennels • Manufacturing establishments • Mausoleums 	<ul style="list-style-type: none"> • Meat and produce canneries • Nursing homes • Pickle factories • Poultry houses • Processing and packing plants • Restaurants • Schools • Spice plants • Stores • Warehouses • Wineries and similar structures • Transportation vessels: aircraft (cargo areas only), buses, trucks, trailers, rail cars, and marine vessels.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Do not pour or dispose down-the-drain or sewer. Call your local solid waste agency for local disposal options.

CONTAINER HANDLING:

Nonrefillable container. Do not reuse or refill this container. Empty residue into application equipment. Offer container and foil bag for recycling, if available, or dispose of container and/or foil bag in a sanitary landfill or by incineration, if allowed by State and local authorities.

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

[A177.03] is a trademark of Atticus, LLC

[Tempo®] is a registered trademark of [Bayer].

{LANGUAGE ON LABEL AFFIXED TO CONTAINER}

BETA-CYFLUTHRIN

GROUP

3A

INSECTICIDE

A177.03TM

[Alternate Brand Name: *Tirade Ultra WSP*]

Contains beta-cyfluthrin, the active ingredient used in [Tempo® Ultra WSP].

[For Pest Management Professionals and commercial use only] [For broad-spectrum control of crawling, flying, and wood-infesting insect pests on indoor and outdoor surfaces, as well as insect pests of trees, landscape ornamentals, residential, and commercial lawns]

ACTIVE INGREDIENT: (% by weight)

Beta-Cyfluthrin, cyano(4-fluoro-3-phenoxyphenyl)methyl 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropane-carboxylate.....10.0%

OTHER INGREDIENTS:.....90.0%

TOTAL100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand the label, find someone to explain it to you in detail.)

FIRST AID	
If on skin or clothing:	<ul style="list-style-type: none">Take off contaminated clothing.Rinse skin immediately with plenty of water for 15-20 minutes.Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">Move person to fresh air.If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.Call a poison control center or doctor for further treatment
If in eyes:	<ul style="list-style-type: none">Hold eye open and rinse slowly and gently with water for 15-20 minutes.Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">Call a poison control center or doctor immediately for treatment advice.Have person sip a glass of water if able to swallow.DO NOT induce vomiting unless told to do so by the poison control center or doctor.DO NOT give anything to an unconscious person.
NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.	
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

For Chemical Emergency:

Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Harmful if swallowed, inhaled, or absorbed through the skin. DO NOT get in eyes, on skin, or on clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and warm water after handling. Remove contaminated clothing and wash before reuse. Wear safety glasses, goggles, or face shield when handling the undiluted material, and wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters when making general surface overhead treatments indoors. **DO NOT** allow children or pets to enter treated areas until surfaces are dry.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Remove from premises or tightly cover fish tanks and disconnect aerators when applying indoors where such containers are present. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. To protect the environment, **DO NOT** allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters.

Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. Apply this product only as specified on this label. This pesticide is highly toxic to bees exposed to direct treatment or residues on crops or weeds. **DO NOT** apply **A177.03** or allow it to drift onto crops or weeds on which bees are actively foraging. Additional information may be obtained by consulting your Cooperative Extension Service.

Physical Or Chemical Hazards: Do not mix or allow to come into contact with any oxidizing agent. Hazardous chemical reaction may occur.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Do not pour or dispose down-the-drain or sewer. Call your local solid waste agency for local disposal options.

CONTAINER HANDLING:

Nonrefillable container. Do not reuse or refill this container. Empty residue into application equipment. Offer container and foil bag for recycling, if available, or dispose of container and/or foil bag in a sanitary landfill or by incineration, if allowed by State and local authorities.

See inside label booklet for additional Precautionary Statements and Directions for Use.

[A177.03] is not manufactured, or distributed by Bayer Environmental Science, seller of [Tempo® Ultra WSP].

Manufactured for:

Atticus, LLC

940 NW Cary Parkway, Suite 200
Cary, NC 27513

EPA Reg. No.: 91234-XX

EPA Est. No.: _____

NET CONTENTS: _____