



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505T)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

91234-321

Date of Issuance:

10/20/25

NOTICE OF PESTICIDE:

☒ Registration  
☐ Reregistration  
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

A1101.07

Name and Address of Registrant (include ZIP Code):

Atticus, LLC  
5000 CentreGreen Way, Suite 100  
Cary, NC 27513

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. You have 18 months from the date of registration to provide these data.

*Continues page 2*

Signature of Approving Official:

Digitally signed by Debra Rate  
Date: 2025.10.20 15:49:07  
-04'00'

Date:

10/20/25

Debra Rate, Acting Product Manager 11  
Invertebrate-Vertebrate Branch 2, Registration Division (7505T)

EPA Form 8570-6

3. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, "EPA Reg. No. 91234-321."
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is attached for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 10/16/2025

If you have any questions, please contact Christopher Taylor at 202-566-2928 or at [taylor.christopher.m@epa.gov](mailto:taylor.christopher.m@epa.gov).

Attachment

{Note to reviewer: [Text] in brackets denotes optional language}

{Note to reviewer: {Text} in braces denotes explanatory language and where in the final label text will appear}

**{BOOKLET FRONT PANEL LANGUAGE}**

SPINOSAD	GROUP	<b>5</b>	INSECTICIDE
----------	-------	----------	-------------

**A1101.07™**

[*Alternate Brand Name: Cachera Fruit Fly Bait*]

[Contains spinosad, the active ingredient used in GF-120® NF Naturalyte® Fruit Fly Bait.]

[**A1101.07** is not manufactured, or distributed by Corteva Agriscience United States, seller of GF-120® NF Naturalyte® Fruit Fly Bait.]

**[INSECTICIDAL BAIT]**

[For Selective Attractance and Control of Multiple Species of Tephritid Fruit Flies Infesting Any Tree, Fruit, Nut, Vine, Vegetable or Food Crop and Ornamentals, and on Vegetation which May Serve as Resting Sites for Adult Flies]

<b>ACTIVE INGREDIENT:</b>	(% by weight)
Spinosad (a mixture of spinosyn A and spinosyn D).....	0.02%
<b>OTHER INGREDIENTS:</b> .....	99.98%
<b>TOTAL</b> .....	100.0%

Contains [0.02% active ingredient on a weight basis -] 0.002 lb. ai per gallon.



[Listed by the Organic Materials Review Institute (OMRI) for use in organic production.]



**For Organic Production**

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION / PRECAUCIÓN**

**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)**

[See below for additional Precautionary Statements.]

See inside label booklet for First Aid, Precautionary Statements and Directions for Use.


**AVOID FREEZING**

**EPA Reg. No.: 91234-XXX**

**EPA Est. No.:**

**Net Contents:**

Manufactured for:  
**Atticus, LLC**  
940 NW Cary Parkway, Suite 200  
Cary, NC 27513



## {LANGUAGE INSIDE BOOKLET}

FIRST AID	
If swallowed:	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li><li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li></ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at <b>1-844-685-9173</b> for emergency medical treatment information.	

**For Chemical Emergency:**  
**Spill, Leak, Fire, Exposure, or Accident,**  
**Call CHEMTREC Day or Night**  
**Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

## **PRECAUTIONARY STATEMENTS**

### **HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

#### **CAUTION**

Harmful if swallowed. Wash thoroughly with soap and water after handling.

#### **PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### **USER SAFETY RECOMMENDATIONS**

**Users Should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

#### **ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates. This product is highly toxic to bees and other pollinating insects exposed to direct treatment, or to residues in/on blooming crops or weeds. Protect pollinating insects by following label directions intended to minimize drift and to reduce risk to these organisms. **DO NOT** apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when cleaning equipment or when disposing of equipment washwaters. **DO NOT** apply where runoff is likely to occur. **DO NOT** apply when weather conditions favor drift from treated area. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Applying this product when rain is not predicted for the next 24 hours will help reduce potential risk to aquatic invertebrates by reducing pesticide runoff from the treatment area into water bodies. Apply this product only as specified on the label.

#### **PHYSICAL OR CHEMICAL HAZARDS**

Do not mix or allow coming in contact with Oxidizing agents. Hazardous Chemical reaction may occur.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**Entry Restrictions for Non-WPS Uses:** There are no specific entry requirements following application of this product when applied by government agencies in area-wide eradication and prevention programs.

### PRODUCT INFORMATION

**A1101.07** attracts and controls multiple species of tephritid fruit flies infesting any tree, fruit, nut, vine, vegetable or food crop and ornamentals, and on vegetation which may serve as resting sites for adult flies. Aerial or ground applications in production agriculture or directed ground applications to individual plants are permitted. **DO NOT** make aerial applications in immediate proximity of residential, commercial, government, institutional or other structures where people may be present including homes, apartments, offices, churches, schools, and businesses. Aerial applicators should evaluate conditions existing at the time of application and make appropriate adjustments to reduce drift. In urban areas, however, use is limited to directed ground applications.

There is no preharvest interval when **A1101.07** is applied to any tree, fruit, nut, vine, vegetable or any other food crop and ornamentals to control multiple species of tephritid fruit flies. **DO NOT** enter or allow worker entry into treated areas during the restricted entry interval of 4 hours except as allowed under the Agricultural Use Requirements section when the specified PPE is worn.

**Chemigation:** **DO NOT** apply through any type of irrigation equipment.

## RESISTANCE MANAGEMENT

For resistance management, **A1101.07** contains a Group 5 insecticide. Insect population may contain individuals naturally resistant to **A1101.07** and other Group 5 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of **A1101.07** or other Group 5 insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
  - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
  - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
  - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
  - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
  - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance, you may contact your local Atticus, LLC representative or by calling 984-465-4800.

Currently, only spinetoram and spinosad active ingredients are classified as Group 5 insecticides. These two insecticide active ingredients share a common mode of action and must not be rotated with each other for control of pests listed on this label. Spinetoram and spinosad may be rotated with all other labeled insecticide active ingredients.

### SPRAY DRIFT

#### Aerial Applications:

- **DO NOT** release spray at a height greater than 10 feet above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- If the wind speed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- **DO NOT** apply during temperature inversions.

**Ground Boom Applications:**

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

**SPRAY DRIFT ADVISORIES**

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

**IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

**Controlling Droplet Size – Ground Boom**

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

**Controlling Droplet Size – Aircraft**

- **Adjust Nozzles** - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

**BOOM HEIGHT – Ground Boom**

For ground equipment, the boom should remain level with the crop and have minimal bounce.

**RELEASE HEIGHT - Aircraft**

Higher release heights increase the potential for spray drift.

**SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

**TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

**TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

## WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

## MIXING DIRECTIONS

Dilute **A1101.07**, a bait concentrate, with water. The most effective dilution for aerial and most other applications is a 1:1.5 (**A1101.07**:water) dilution (e.g., to make 10 liters of spray solution, add 6 liters of water to 4 liters of **A1101.07**). For ground applications and applications in low relative humidity, dilutions of up to 1:5 (**A1101.07**:water) can be made. First add water (one-half of the volume to be mixed) to the spray tank or premixing tank and start the agitation system. Then add the full amount of **A1101.07** followed by an equal amount of water. If a full container of **A1101.07** is used, triple rinse the empty container by filling it one-third full with water, shake well and add the rinsate to the spray tank. Repeat two more times so the container is triple rinsed and then complete filling the spray tank until the proper dilution is obtained. Maintain constant agitation of the spray solution to ensure uniformity of spray mixture. Allow agitation system to operate for at least 5 minutes before applying. Once diluted, use **A1101.07** within 24 hours. Concentrated **A1101.07** will not settle and does not need to be shaken before mixing.

## APPLICATION DIRECTIONS

Proper application techniques help ensure adequate coverage and correct dosage necessary to obtain optimum control of insect pests. Use a large spray droplet size of 4000 - 6000  $\mu\text{m}$  (4 - 6 mm) to optimize length of bait attractance. Fruit flies can detect the bait from several yards away.

When aerially applying use ULV applications but with coarse nozzles that will produce the desired droplet size and target 20 - 80 droplets per square meter.

By ground, spot or strip spray several areas on the inner canopy of fruiting plants. Avoid weather conditions that could result in drift to nontarget areas. Direct spray application to bottoms of leaves and leaves inside the foliage canopy to reduce direct exposure to sun and rain. This product resists wash off but will lose effectiveness if exposed to rain and overhead irrigation. When possible, potential for rain or irrigation schedules should be considered when planning applications.

Begin applications as soon as monitoring traps indicate flies are present or 2 - 3 weeks before fruit begins to ripen. Repeat applications every 7 - 14 days, shortening the application interval during rainy periods and as fruit ripens. Remove fruit as soon as ripe, particularly any overly ripe fruit on the tree or ground.

## PESTS AND APPLICATION RATES

Pests Controlled or Suppressed	Amount of Undiluted A1101.07 Spray Solution <sup>1</sup>			
	Broadcast Application		Spray of Individual Plants <sup>2</sup>	
	Fl. oz./Acre	Liters/Hectare	Fl. oz./Tree	mL/Tree
Tephritid fruit flies including: Apple maggot Caribbean fruit fly Cherry fruit fly Mediterranean fruit fly Melon fly Mexican fruit fly Olive fly Oriental fruit fly Walnut husk fly	10 – 20	0.8 – 1.6	1 – 3	30 – 90

<sup>1</sup>Choose rate based on pest pressure and amount of foliage needed to cover.

<sup>2</sup>Avoid leaving large spots of product on foliage or fruit.



**AMOUNT OF FINAL SPRAY SOLUTION FOR DIFFERENT RATES AND DILUTION RATIOS<sup>1</sup>**

<b>Rates of A1101.07</b>					
<b>Dilution Ratio (A1101.07:Water)</b>	<b>10 fl. oz./acre</b>	<b>12 fl. oz./acre</b>	<b>15 fl. oz./acre</b>	<b>18 fl. oz./acre</b>	<b>20 fl. oz./acre</b>
1:1.5	25	30	37.5	45	50
1:2	30	36	45	54	60
1:3	40	48	60	72	80
1:4	50	60	75	90	100
1:5	60	72	90	108	120

<sup>1</sup>Entries in the table are the amount of final diluted spray solution per acre.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container only. In case of leak or spill, contain material with absorbent materials and dispose as waste.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### CONTAINER HANDLING:

**[Nonrefillable containers ≤ 5 gallons: Nonrefillable Container:** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

**[Nonrefillable containers > 5 gallons: Nonrefillable container.** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

## LIMITATION OF WARRANTY AND LIABILITY

**IMPORTANT: READ BEFORE USE.** Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

**A1101.07** is a trademark of Atticus, LLC.

[GF-120® and Naturalyte® are registered trademarks of Corteva Agriscience and its affiliated companies.]

## {LANGUAGE ON LABEL AFFIXED TO CONTAINER}

SPINOSAD	GROUP	5	INSECTICIDE
----------	-------	---	-------------

**A1101.07™**

[Alternate Brand Name: Cachera Fruit Fly Bait]

[Contains spinosad, the active ingredient used in GF-120® NF Naturalyte® Fruit Fly Bait®.]

[A1101.07 is not manufactured, or distributed by Corteva Agriscience United States, seller of GF-120®NF Naturalyte® Fruit Fly Bait.]

[INSECTICIDAL BAIT]

[For Selective Attractance and Control of Multiple Species of Tephritid Fruit Flies Infesting Any Tree, Fruit, Nut, Vine and Vegetable Crops and Ornamentals, and on Non-Crop Vegetation which May Serve as Resting Sites]

**ACTIVE INGREDIENT:** (% by weight)

Spinosad (a mixture of spinosyn A and spinosyn D).....0.02%

**OTHER INGREDIENTS:**.....99.98%

**TOTAL** .....100.0%

Contains [0.02% active ingredient on a weight basis -] 0.002 lb. ai per gallon.



[Listed by the Organic Materials Review Institute (OMRI) for use in organic production.]



For Organic Production

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION / PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If swallowed:	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li><li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li></ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at <b>1-844-685-9173</b> for emergency medical treatment information.	

**For Chemical Emergency:**

**Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC Day or Night**

**Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

### PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

#### CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling.

**ENVIRONMENTAL HAZARDS:** This product is toxic to aquatic invertebrates. This product is highly toxic to bees and other pollinating insects exposed to direct treatment, or to residues in/on blooming crops or weeds. Protect pollinating insects by following label directions intended to minimize drift and to reduce risk to these organisms. **DO NOT** apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when cleaning equipment or when disposing of equipment washwaters. **DO NOT** apply where runoff is likely to occur. **DO NOT** apply when weather conditions favor drift from treated area. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Applying this product when rain is not predicted for the next 24 hours will help reduce potential risk to aquatic invertebrates by reducing pesticide runoff from the treatment area into water bodies. Apply this product only as specified on the label.

**PHYSICAL OR CHEMICAL HAZARDS:** Do not mix or allow coming in contact with Oxidizing agents. Hazardous Chemical reaction may occur.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container only. In case of leak or spill, contain material with absorbent materials and dispose as waste.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

#### CONTAINER HANDLING:

**[Nonrefillable containers ≤ 5 gallons:** Nonrefillable Container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

**[Nonrefillable containers > 5 gallons:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

See inside label booklet for additional Precautionary Statements and Directions for Use.

### AVOID FREEZING

Manufactured for:

**Atticus, LLC**

940 NW Cary Parkway, Suite 200

Cary, NC 27513

**EPA Reg. No.: 91234-XXX**

**EPA Est. No.: \_\_\_\_\_**

**NET CONTENTS: \_\_\_\_\_**