



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

91234-348

Date of Issuance:

11/5/25

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

A2114.02

Name and Address of Registrant (include ZIP Code):

Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. You have 18 months from the date of registration to provide these data.

Continues page 2

Signature of Approving Official:

Stephanie Suarez, Ph.D., Acting Product Manager 22
Fungicide Branch, Registration Division (7505T)

Date:

11/5/25

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 91234-348."
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The alternate brand name "Centyk" has been added to the product record.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 08/20/2025

If you have any questions, please contact James Orrock by phone at 202-566-2862 or by email at orrock.james@epa.gov.

Enclosure

11/5/2025

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 91234-348

{Note to reviewer: [Text] in brackets denotes optional or explanatory language}

{Note to reviewer: {Text} in braces denotes where in the final label text will appear}

{BOOKLET FRONT PANEL LANGUAGE}

TRIFLOXYSTROBIN

GROUP

11

FUNGICIDE

A2114.02 [™]

[Alternate Brand Name: Centyk]

[Contains trifloxystrobin , the active ingredient used in Compass®.]

[For Turf, Landscape, Greenhouse and Nursery Use]

[FUNGICIDE]

[Turf and Ornamental Fungicide]

[For Control of Certain Foliar, Stem, and Root Diseases of Turfgrass Including Golf Courses, Institutional, Commercial and Residential Lawns, Sod Farms, Sports Fields, Parks, Municipal Grounds and Cemeteries, and of Ornamentals Grown in Interiorscapes, Field Nursery Plantings, Forest Nurseries, Residential and Commercial Landscapes, Greenhouses, Lath and Shadehouses, Containers, and Other Enclosed Structures.]

ACTIVE INGREDIENT: (% by weight)

Trifloxystrobin (CAS No. 141517-21-7).....50.0%

OTHER INGREDIENTS:50.0%

TOTAL 100.0%

A2114.02 is a water-dispersible granule.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

[See [Back][Side] Panel for First Aid and [Leaflet] for Precautionary Statements and Directions for Use.]

[See inside label booklet for First Aid, [additional] Precautionary Statements, and Directions for Use.]

[See below for additional Precautionary Statements]

[A2114.02 is not manufactured, or distributed by Bayer Environmental Science, seller of Compass®.]

EPA Reg. No.: 91234-XX

EPA Est. No.:

Net Contents:

Manufactured for:

Atticus, LLC

940 NW Cary Parkway, Suite 200

Cary, NC 27513

{LANGUAGE INSIDE BOOKLET}

FIRST AID	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	
NOTE TO PHYSICIAN: If ingested, induce emesis or lavage stomach. Treat symptomatically.	

For Chemical Emergency:
Spill, Leak, Fire, Exposure, or Accident,
Call CHEMTREC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Harmful if absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils., nitrile rubber ≥ 14 mils., neoprene rubber ≥ 14 mils., natural rubber ≥ 14 mils., polyethylene, polyvinyl chloride (PVC) ≥ 14 mils., or viton ≥ 14 mils.
- Shoes plus socks

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves removing. As soon possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** contaminate water when disposing of equipment wash water or rinsate. Applying this product when rain is not predicted for the next 24 hours will help reduce potential risk to aquatic invertebrates by reducing pesticide runoff from the treatment area into water bodies.

GROUND WATER ADVISORY

Several trifloxystrobin degradates have properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT use, pour, spill, or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Residential Use Sites: Intended for use by professional applicators.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the **restricted-entry interval (REI) of 12 hours**.

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, butyl rubber \geq 14 mils., nitrile rubber \geq 14 mils., neoprene rubber \geq 14 mils., natural rubber \geq 14 mils., polyethylene, polyvinyl chloride (PVC) \geq 14 mils., or viton \geq 14 mils.
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

DO NOT enter treated areas without protective clothing until sprays have dried.

PRODUCT INFORMATION

A2114.02 is a modern site-specific fungicide for use on turfgrass and ornamentals with protective and curative activity. **A2114.02** penetrates the plant and provides translaminar activity via a high affinity for the waxy layer of the plant surface, localized vapor movement and re-deposition on the plant.

CHEMIGATION

DO NOT apply this product through any type of irrigation system.

AERIAL APPLICATION

DO NOT apply by aerial application.

SPRAY DRIFT MANAGEMENT

Airblast Applications:

- Sprays must be directed into the canopy
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer row.
- **DO NOT** apply during temperature inversions.

Ground Boom Applications:

- Apply with nozzle height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- **Volume** – Increasing the spray volume so that larger droplets are produced that will reduce drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATION DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Handheld Technology Applications

Take precautions to minimize spray drift.

FUNGICIDE RESISTANCE MANAGEMENT

For resistance management, **A2114.02** contains a Group 11 fungicide. Any fungal population may contain individuals naturally resistant to **A2114.02** and other Group 11 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of **A2114.02** or other Group 11 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Atticus, LLC at 984-465-4800. You can also contact your pesticide distributor or university extension specialist to report resistance.

A2114.02 belongs to the strobilurin class of chemistry which exhibits no known cross-resistance to other chemical classes including sterol inhibitors, dicarboximides, benzimidazoles, anilinopyrimidines, or phenylamides. However, certain fungal pathogens are known to develop resistance to products used repeatedly. Because resistance development cannot be predicted, the use of this product must conform to resistance management strategies. Such strategies may include rotating and/or tank mixing with products having different modes of action; or limiting the total number of applications per year. Atticus, LLC encourages responsible product stewardship to ensure effective long-term control of the fungal diseases on this label. See specific recommendations in the **Turfgrass** section and the **Ornamentals** section.

MIXING PROCEDURES

WATER DISPERSIBLE GRANULES

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Agitation is necessary for proper dispersal of the product. Maintain agitation throughout the spraying operation. **DO NOT** let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

A2114.02 Alone

Add ½ of the required amount of water to the mix tank. With the agitator running, add the **A2114.02** to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after **A2114.02** has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

A2114.02 Tank Mixtures

Add 1/2 of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. Add tank mix partners in this order:

- (1) products packaged in water-soluble packaging, wettable powders, wettable granules (dry flowables) such as **A2114.02**;
- (2) liquid flowables, liquids; and
- (3) emulsifiable concentrates.

Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all of the mixture has been applied.

NOTE: When using **A2114.02** in tank mixtures, add all products in water-soluble packaging to the tank before any other tank mix partner, including **A2114.02**. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

If using **A2114.02** in a tank mixture, observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations that appear on the tank mix product label. Label dosage rate must not be exceeded, and the most restrictive label precautions and limitations must be followed. **DO NOT** mix this product with any product that prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

A2114.02 is compatible with most insecticide, fungicide, and foliar nutrient products. However, the compatibility of **A2114.02** with tank mix partners must be tested before use.

To determine biological compatibility with other products, mix the products in the desired proportions, spray on target plants, and observe for phytotoxicity seven days after the application.

To determine the physical compatibility of **A2114.02** with other products, use a jar test, as described below. Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Use with Additives

Use of spray additives are not required. Evaluate any spray additive prior to use. **DO NOT** use in conjunction with organosilicate-based products, or plant injury may occur. Label directions are based on data with no additives.

WATER SOLUBLE PACKETS

The enclosed packets containing **A2114.02** are water soluble. **DO NOT** allow packets to become wet prior to adding to the spray tank. **DO NOT** handle with wet hands. Reseal outer carton/container to protect remaining packets.

To prepare the spray mixture, drop the required number of unopened packets, as determined under the **Turfgrass** and **Ornamentals** sections, into the spray tank while filling the water to the desired level. Operate the agitator while mixing. Depending on the water temperature and the degree of agitation, the packets should be completely dissolved within a few minutes from the time they were added to the water.

ATTENTION: DO NOT use **A2114.02** water-soluble packets in a tank-mix with products that contain boron or release free chlorine. The resultant reaction of PVA and boron or free chlorine is a plastic which is not soluble in water or solvents such as diesel oils, kerosene, gasoline or alcohol. Use of chlorinated water is acceptable. **DO NOT** attempt to use the PVA packets directly in liquid fertilizer, diesel oils, or summer spray type oils as in ULV or LV uses. PVA packets are water soluble not fertilizer or oil soluble. If liquid fertilizer or oils are to be added to the mix water, add **A2114.02** packets first and wait until they are dissolved before adding liquid fertilizer or oil.

A2114.02 Alone

Add 1/2 of the required water to the mix tank. With the agitator running, add the required number of **A2114.02** packets to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after **A2114.02** packets have completely dissolved and dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

A2114.02 + Tank Mixtures

Add 1/2 of the required amount of water to the tank mix. Start the agitation running before adding **A2114.02** water-soluble packets and any tank mix partner(s). Add tank mix partners in this order:

- (1) products packaged in water-soluble packaging (such as **A2114.02**), wettable powders, wettable granules (dry flowables);
- (2) liquid flowables, liquids; and
- (3) emulsifiable concentrates.

Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all of the mixture has been applied.

NOTE: When using **A2114.02** in tank mixes, add all products in water-soluble packaging (including **A2114.02**) to the tank before any other tank mix partner. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

If using **A2114.02** in a tank mixture, observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations that appear on the tank mix product label. Label dosage rate must not be exceeded, and the most restrictive label precautions and limitations must be followed. **DO NOT** mix this product with any product that prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

A2114.02 is compatible with most insecticide, fungicide, and foliar nutrient products. However, the compatibility of **A2114.02** with tank mix partners must be tested before use.

To determine biological compatibility with other products, mix the products in the desired proportions, spray on target plants and observe for phytotoxicity seven days after the application.

Observe all directions, precautions, and limitations on labeling of all products used in tank mixes. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

Use with Additives

Use of spray additives are not required. Evaluate any spray additive prior to use. **DO NOT** use in conjunction with organosilicate-based products, or plant injury may occur. Label directions are based on data with no additives.

{**NOTE TO REVIEWER:** Registrant may add the following state-driven statements as required throughout.

[Not Registered for Use by (Insert State)] [Not Registered for Use on (Insert Commodity) by (Insert State)]
[Not Registered for Sale, Sale Into, Distribution and/or Use in (Insert County Name(s)) Counties of (Insert State)]]

TURFGRASS

A2114.02 is a preventive and curative fungicide that may be applied to turf sites including golf courses, institutional, commercial and residential* lawns, sod farms, sports fields, parks, municipal grounds and cemeteries.

*Intended for use by professional applicators.

A2114.02 is a broad spectrum fungicide for the control of brown patch (*Rhizoctonia solani*), anthracnose (*Colletotrichum graminicola*), red thread (*Corticium fuciforme*), pink patch (*Limonomyces roseipellis*), Leaf spot (*Bipolaris* spp., *Drechslera* spp.), gray leaf spot (*Pyricularia grisea*), rust (*Puccinia* spp.), pink snowmold (*Microdochium nivale*), fusarium patch (*Fusarium nivale*), rapid blight, and summer patch (*Magnaporthe poae*).

For optimum control of pink snowmold, fusarium patch, and summer patch, mix or rotate **A2114.02** with Triadimefon [Bayleton® 50 Turf and Ornamental Fungicide in Water Soluble Packets Systemic Fungicide (BAYLETON 50) (EPA Reg No. 432-1360)]. When conditions are favorable for Pythium blight (*Pythium* spp.), mix a Pythium control fungicide with **A2114.02**.

A2114.02 will provide suppression of dollar spot (*Sclerotinia homoeocarpa*) and will not flare this disease. However, during periods of dollar spot pressure, mix **A2114.02** with triadimefon [(BAYLETON® 50)] or other dollar spot control fungicides.

- STROBILURIN FUNGICIDES SUCH AS **A2114.02** ARE BEST USED IN A PREVENTIVE DISEASE CONTROL PROGRAM. CURATIVE APPLICATIONS CAN BE MADE FOR CERTAIN DISEASES.
- Apply in 1 - 2 gals. of water per 1,000 sq. ft. to ensure thorough coverage. For soil-borne diseases, use sufficient water to reach the crown and upper root zone.
- Apply after mowing OR allow sprayed area to completely dry before mowing.
- For control of both foliar and soil-borne diseases, allow sprayed area to completely dry before irrigation.
- Under conditions optimum for high disease pressure, use the higher rate and the shorter interval.
- For optimum turf quality and disease control, use **A2114.02** in conjunction with turf management practices that promote good health and optimum disease control.
- Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.

TANK MIXES

For longer and more broad-spectrum control including dollar spot, **A2114.02** can be tank mixed with other fungicides. Check compatibility before tank mixing.

A2114.02 CAN BE APPLIED TO ALL MAJOR TURFGRASS SPECIES. RATES ARE DEPENDENT UPON DISEASE AND THE ORGANISM THAT IT IS INCITED BY.

RESISTANCE MANAGEMENT FOR TURFGRASS

DO NOT apply more than 2 sequential **A2114.02** applications for gray leaf spot control. **DO NOT** apply more than 3 sequential applications of **A2114.02** for all other diseases. Alternate with a fungicide having a different mode of action.

RESTRICTIONS

- **Maximum Use Rates**
 - For turfgrass, up to 34.50 oz. (1.08 lb. ai) of **A2114.02** can be applied per acre per year.
 - For plants grown in outdoor nurseries, outdoor seedbeds, field plantings, and landscapes, up to 34.50 oz. (1.08 lb. ai) of **A2114.02** per acre of production or acre of landscape per year can be used.
 - For seedlings and plants grown in greenhouses, containers, and other enclosed structures, up to 120.0 oz. (3.75 lb. ai) of **A2114.02** per acre per year can be used. In California only, **DO NOT** apply more than 34.50 oz. (1.08 lb. ai) of **A2114.02** per acre per year to seedlings or plants grown in greenhouses, containers, and other enclosed structures.

TURFGRASS DISEASES CONTROLLED WITH A2114.02

DISEASE	RATE OF PRODUCT (Oz./1,000 Sq. Ft.)	APPLICATION INTERVAL/TIMING	APPLICATION INSTRUCTIONS
Brown Patch	0.10 - 0.20	14 days	Apply when conditions are favorable for disease development.
	0.15 - 0.25	21 days	For curative applications, apply 0.20 oz. of A2114.02 and repeat applications on a 21-day interval.
	A2114.02: 0.10 - 0.15 + Triadimefon [(Bayleton 50)]: See Tank Mix Partner Product Label	14 - 21 days	For longer and more broad-spectrum disease control including dollar spot, mix A2114.02 with Triadimefon [(Bayleton 50)].
Leaf Spot	0.10 - 0.15	14 days	Apply when conditions are favorable for disease development.
	0.15 - 0.25	21 - 28 days	For curative applications, apply A2114.02 at labeled rates and repeat applications according to the application interval timing.
Anthracnose	0.15 - 0.20	14 days	Apply when conditions are favorable for disease development.
	0.25	21 days	
	A2114.02: 0.10 - 0.15 + Triadimefon [(Bayleton 50)]: See Tank Mix Partner Product Label	14 - 21 days	For longer and more broad-spectrum disease control including dollar spot, mix A2114.02 with Triadimefon [(Bayleton 50)].
Summer Patch	A2114.02: 0.20 - 0.25 and/or Triadimefon [(Bayleton 50)]	21 - 28 days	Under heavy disease pressure, apply Triadimefon [(Bayleton 50)] at labeled rates when the soil temperatures reach 65°F. AFTER 21-28 days, apply either:

	See Tank Mix Partner Product Label		<p>A2114.02 at 0.20 oz. every 14 days OR a combination of Triadimefon [(Bayleton 50)] at labeled rates + A2114.02 at 0.20 - 0.25 oz. every 21 - 28 days.</p> <p>DO NOT make more than 3 sequential applications of A2114.02 before rotating back to Triadimefon [(Bayleton 50)].</p> <p>Under light disease pressure, make 2 - 3 applications of A2114.02 at 0.20 - 0.25 oz. + Triadimefon [(Bayleton 50)] at labeled rates every 21 - 28 days.</p>
Gray Leaf Spot Rapid Blight	0.15 - 0.20	14 days	<p>Apply when conditions are favorable for disease development. For best results on Rapid Blight apply A2114.02 at the highest rate.</p> <p>For longer and more broad-spectrum disease control including dollar spot, mix A2114.02 with Triadimefon [(Bayleton 50)].</p>
	0.25	21 days	
	A2114.02: 0.10 - 0.20 + Triadimefon [(Bayleton 50)] See Tank Mix Partner Product Label	14 - 21 days	
Red Thread Pink Patch	0.10 - 0.15	14 days	Apply when conditions are favorable for disease development.
	0.20 - 0.25	21 days	For curative applications, apply A2114.02 at 0.15 - 0.25 oz. and repeat applications according to the application interval timing.
	A2114.02: 0.10 - 0.15 + Triadimefon [(Bayleton 50)] See Tank Mix Partner Product Label	14 - 21 days	For longer and more broad-spectrum disease control including dollar spot, mix A2114.02 with Triadimefon [(Bayleton 50)].
Rust	0.10 - 0.15	14 days	Apply when conditions are favorable for disease development.
	0.20 - 0.25	21 days	For curative applications, apply A2114.02 at 0.15 - 0.25 oz. and repeat applications according to the application interval timing.
	A2114.02: 0.10 - 0.15 + Triadimefon [(Bayleton 50)] See Tank Mix Partner Product Label	14 - 21 days	For longer and more broad-spectrum disease control including dollar spot, mix A2114.02 with Triadimefon [(Bayleton 50)].
Fusarium Patch	0.20 - 0.25	Fall-Early Spring	Apply when conditions are favorable for disease development.
	A2114.02: 0.15 - 0.25 + Triadimefon [(Bayleton 50)]	Fall-Early Spring	Under conditions of heavy disease pressure, tank mix with Triadimefon [(Bayleton 50)].

	See Tank Mix Partner Product Label		
Pink Snow Mold	0.25	Late Fall	Apply one application in late fall before snow cover or early spring after snow melts. DO NOT apply on top of snow.
	A2114.02: 0.15 - 0.25 + Triadimefon [(Bayleton 50)] See Tank Mix Partner Product Label	Late Fall	Under conditions of heavy disease pressure, tank mix with Triadimefon [(Bayleton 50)].

CONVERSION TABLE FOR A2114.02 TURF USES IN WATER SOLUBLE PACKETS

Oz./1,000 sq. ft.	Oz./A	Each 2 Oz. Soluble Packet Treats	Each 8 oz. Soluble Packet Treats	Number of 8 oz. Soluble Packets Needed to Treat 20 Acres of Turfgrass
0.10 (0.003 lb. ai)	4.40 (0.138 lb. ai)	0.46 acres	1.82 acres	11
0.15 (0.005 lb. ai)	6.50 (0.203 lb. ai)	0.31 acres	1.22 acres	16
0.185 (0.006 lb. ai)	8.00 (0.250 lb. ai)	0.25 acres	1.00 acres	20
0.20 (0.007 lb. ai)	8.70 (0.272 lb. ai)	0.23 acres	0.92 acres	22
0.25 (0.008 lb. ai)	10.90 (0.341 lb. ai)	0.18 acres	0.73 acres	27

ORNAMENTAL DISEASE CONTROL

A2114.02 is a broad-spectrum fungicide for the control of certain foliar, stem, and root diseases of ornamentals grown in interiorscapes, field nursery plantings, forest nurseries, residential* and commercial landscapes, greenhouses, lath and shadehouses, containers, and other enclosed structures.

*Intended for use by professional applicators

FOLIAR DISEASES

A2114.02 will control foliar diseases of ornamentals when applied as a foliar spray. Apply **A2114.02** at 1.0 – 4.0 oz./100 gals. (0.032 – 0.125 lb. ai/100 gals.) [(0.5 – 2.0 oz./50 gals.)] to the point of drip and repeat at 7 - 14 day intervals until the threat of disease is over. Start applications when conditions are favorable for disease development and continue until the threat of disease is over.

DAMPING OFF OF NEW SEEDLINGS

A2114.02 will control damping off of new seedlings caused by *Rhizoctonia solani* when applied as a drench to seedlings and transplants. Drench the growth media at a rate of 0.50 oz./100 gals. (0.016 lb. ai/100 gals.). Repeat every 21 - 28 days. If *Pythium* spp. are also present, mix **A2114.02** with a *Pythium* control fungicide.

The plants that **A2114.02** has been tested on, diseases that are controlled, and specific directions for use are listed in **Tables 1, 2, and 3**. Refer to **Table 1** for information on ornamentals and diseases that have been evaluated, **Table 2** for specific pathogens controlled, and to **Table 3** for specific guidelines on the rates and timing of application.

RESISTANCE MANAGEMENT FOR ORNAMENTALS

A2114.02 is a modern, site-specific fungicide belonging to the strobilurin class of chemistry. Fungal pathogens are known to develop resistance to fungicides with a specific mode of action. When site-specific fungicides are introduced without a clear resistance management strategy, resistance development may be rapid, particularly with greenhouse use.

A2114.02 exhibits cross-resistance to other strobilurins and fungicides within the Strobilurin Type Action and Resistance group (STAR compounds), but there is no known cross-resistance to fungicides of other classes including sterol inhibitors, dicarboximides, benzimidazoles, anilinopyrimidines, phenylpyrroles, or phenylamides.

Many fungi which attack ornamentals and flowering plants including Botrytis and powdery mildews have a history of fungicide resistance development. Because resistance development cannot be predicted, implementation of suitable strategies to manage the resistance risk to **A2114.02** is needed. To minimize the risk of resistance development to **A2114.02**, the following practices are recommended:

1. Use **A2114.02** preventively.
 2. **For Leaf Spots and diseases other than Powdery Mildew, Downy Mildew, and Botrytis:**
 - a. Use no more than two (2) applications of **A2114.02** before rotating to another effective product that is not in the strobilurin class of chemistry for two (2) applications before rotating back to **A2114.02**.

OR

 - b. Rotate to another fungicide of non-strobilurin chemistry after each **A2114.02** application.
3. **For Powdery Mildew, Downy Mildew, and Botrytis:**
 - c. Between each **A2114.02** application, make two (2) applications of a fungicide of non-strobilurin chemistry before rotating back to **A2114.02**.

OR

- d. Rotate to another fungicide of non-strobilurin chemistry after each **A2114.02** application.
4. Make no more than four (4) foliar applications of **A2114.02** per year for each at risk pathogen. Soil applications are independent of this limit.
5. **DO NOT** use **A2114.02** for disease control in vegetables grown in greenhouses for crop production or in vegetable production of transplants for outdoor use.

RESTRICTIONS

- For ground application, use a minimum of 50 gals./A.
- To avoid spray drift, **DO NOT** apply when conditions favor drift beyond the target area. Avoid spray overlap.
- For information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state agricultural experiment station recommendations.
- Use of spray additives are not required. Evaluate any spray additive prior to use. **DO NOT** use in conjunction with organosilicate based products, or plant injury may occur. Label directions are based on data with no additives.
- **Maximum Use Rates in Ornamentals**
 - For plants grown in outdoor nurseries, outdoor seedbeds, field plantings, and landscapes, up to 34.50 oz. (1.078 lb. ai) of **A2114.02** per acre of production or acre of landscape per year can be used.
 - For seedlings and plants grown in greenhouses, containers, and other enclosed structures, up to 120.0 oz. (3.75 lb. ai) of **A2114.02** per acre per year can be used. In California only, **DO NOT** apply more than 34.50 oz. (1.078 lb. ai) of **A2114.02** per acre per year to seedlings and plants grown in greenhouses, containers, and other enclosed structures.
 - For foliar applications, **DO NOT** apply more than 8.0 oz. (0.25 lb. ai) of **A2114.02** per acre per application.

TABLE 1.

A2114.02 has been tested for phytotoxicity and been found safe to the following plants*. The numbers in () indicate the diseases listed in Table 2. For plants not listed and for use of A2114.02 in tank mixtures, see Notice To Users.

Ajuga (2,11,12) Aloe Vera (12) Alyssum (7,12) Apple, nonbearing** (11,13,14, 17) Aptenia (12) Aster (11,12,13) Azalea (2,5,11,12,13) Azalea (Rhododendron)(2,10,11,12, 13) Bamboo (12,14) Barberry, Japanese (4) Begonia (5,11,12) Bellis (1,2,5) Betula (11,13) Blue Daze (12) Bottle Brush (14) Brachycome (12) Buddleia (butterfly bush) (8) Camellia (2,5,12) Campanula (Bell Flower) (11,12,13) Caladium (12) Cast Iron Plant (12) Catnip (5,8,12) Cedar (8,13) Celosia (5,12) Cherry, nonbearing** (11,14,17) Chrysanthemum (5,8,12,13) Citrus, nonbearing** (11) Coleus (5,12) Coontie Palm (12) Coreopsis (1,5,7,11,12) Cosmos (11,12) Crabapple, nonbearing** (8,11,13,14) Daisy (12) Dianthus (1,5,8,12,13) Day Lily (8,12,13) Delphinium (5,8,11,12) Dieffenbachia spp. (9,10,12) Digitalis (Foxglove) (7) Dogwood (Cornus spp.) (2,11) Dracaena (8)	Dusty Miller (1,12) Dwarf Ivy (12) Euonymus (2,11) Exacum (5) Ficus (2,12) Geranium (1,5,7,11,13) Gerbera (caution on open flower) (2,5,11) Grasses list (2,8,11,13) <i>Avena</i> : Blue Grass <i>Festuca glauca</i> : Blue Fescue <i>Festuca glauca</i> : Sea Urchin <i>Festuca glauca</i> : Elijah Blue <i>Lagurus</i> : Rabbit Tail Pampas Grass: Pink Pampas Grass: White <i>Pennisetum rubrum</i> (Crimson Fountain Grass) Silver Banner Grass <i>Phalaris picta</i> (Varigated Ribbon Grass) Hawthorn (5,8,14) Hawthorn, Indian (8,12) Heather, Mexican (12) <i>Hedera</i> spp. (2,5,10,12) Hen and Chickens, flowering (12) Hibiscus (2,5,10,12) Holly (Ilex) (4) Hosta (5,12) Hydrangea (11) Hypericum (13) Hypoestes (12) Impatiens (1,7,12) Indian Hawthorne (<i>Raphiolepis</i>) (8) Iris (dwarf, japanese, siberian)(8,12) Jasmine (2,12) <i>Juniperus tortulosum</i> (12) Lantana (12,13) Lagerstroemia (Crape myrtle) (11) Ligustrum (1,2,8,12)	Lilac (5,11,12) Liriope (12) Lupines (2,5,11,12) Marigold (1,5,11,12) Mint (11,13) Monarda (bee Balm) (5,11) Moonflower (12) Nandina (2,11) Nectarine, nonbearing** (8,11,14,17) Pansy* (1,2,5,7,8,11,12) Peach, nonbearing** (5,12, 17) Penstemon (2,8,11) Petunia (5,12) Phlox (5,7,11) Photinia (4,8,11) Pittosporum (1,8,12) Plum, nonbearing** (5,11,14) Poinsettia (1,5,11,12,14) Poppy (5) Primula (Primrose) (5) Prunus (2,5) Rabbit's Foot Fern*** (5,12) Ranunculus (7,11) Photinia (Red Tip) (8) Rose (3,5,6,7,11,12,13,14) Pothos (9,10,12) Salvia (1,5,7,11,12,13) Snapdragon (2,5,7,8,11,12,13) Spathiphyllum (6,9,10) Spirea (11) Syngonium (9,12) Verbena (5,7,11,12) Veronica (11) <i>Viburnum</i> spp.(2,8,12) Vinca (Catharanthus) (1,4,5,10,12) <i>Vinca Minor</i> (2,10) Zinnia (1,11)
--	---	---

RESTRICTIONS:

* **DO NOT** drench pansy

** **DO NOT** apply **A2114.02** to fruit trees that will bear harvestable fruit within 12 months of the last application.

*** **DO NOT** use **A2114.02** on leatherleaf fern.

TABLE 2.

Common and scientific names of diseases controlled by A2114.02.	
Common Name	Scientific Name
(1) Alternaria (B)	<i>Alternaria</i> spp.
(2) Anthracnose (B)	<i>Colletotrichum</i> spp.
(3) Black Root (D)	<i>Triclaviosis</i> spp.
(4) Black spot (B)	<i>Diplocarpon rosae</i>
(5) Botrytis (B)	<i>Botrytis</i> spp.
(6) Cylindrocladium (D)	<i>Cylindrocladium</i> spp.
(7) Downy Mildew (A)	<i>Peronospora</i> spp.
(8) Leaf spot (B)	<i>Septoria</i> spp.
(9) Myrothecium (A)	<i>Myrothecium</i> spp.
(10) Phytophthora aerial (A) Phytophthora root (D)	<i>Phytophthora parasitica</i>
(11) Powdery mildew (A)	<i>Erysiphe</i> spp. <i>Microsphaera</i> spp. <i>Oidium</i> spp. <i>Podosphaera</i> spp. <i>Sphaerotheca</i> spp.
(12) Rhizoctonia stem /root rot	<i>Rhizoctonia solani</i>
(13) Rust (B)	<i>Gymnosporangium</i> spp. <i>Phragmidium</i> spp. <i>Puccinia uromyces</i>
(14) Scab (B)	<i>Cladosporium</i> spp. <i>Sphaceloma Venturia inaequalis</i> .
(15) Myrothecium (A)	<i>Myrothecium</i> spp.
(16) Rhizoctonia root rot (C)	<i>Rhizoctonia solani</i>
(17) Blossom Blight (E)	<i>Monilinia</i> spp.

TABLE 3.

Specific use directions for selected pathogens
(A) Apply A2114.02 as a foliar spray at 1.0 – 2.0 oz./100 gals. (0.032 – 0.063 lb. ai/100 gals.) to the point of drip before disease is detected or when conditions are favorable for disease development. Continue at 7 - 14 day intervals until the threat of disease is over.
(B) Apply A2114.02 as a foliar spray at 2.0 – 4.0 oz./100 gals. (0.063 – 0.125 lb. ai/100 gals.) to the point of drip before disease is detected or when conditions are favorable for disease development. Continue at 7 - 14 day intervals until the threat of disease is over. Under heavy pressure, use the highest rate and the shortest interval. Under light disease pressure, the application interval may be extended.
(C) Apply 0.50 oz./100 gals. (0.016 lb. ai/100 gals.) as a drench to wet the upper 1/2 of the growing media. Start the application at the time of seeding, again at transplanting and at 21 - 28 day intervals thereafter.
(D) Apply 1 - 2 oz./100 gals. (0.032 – 0.063 lb. ai/100 gals.) as a drench to wet the upper 1/2 of the growing media. Start the application at the time of planting and at 14 - 28 days depending on disease pressure.
(E) Apply 2 - 3 oz./100 gals. (0.063 – 0.094 lb. ai/100 gals.) starting at bud break on non-bearing stone and pome fruit listed in Table 1 . If conditions are favorable for disease development, apply again at full bloom and at petal fall or on a 14 - 21 day spray schedule.

NOTICE TO USER: Plant tolerance to **A2114.02** has been found to be acceptable on all ornamentals that it has been tested on. Due to the large number of species and varieties of ornamentals and nursery plants, it is impossible to test every one for tolerance to **A2114.02**. Neither the Manufacturer nor the Seller has determined whether or not **A2114.02** can be used safely on ornamental plants not specified on this label. The professional user must determine if **A2114.02** can be used safely prior to commercial use. In a small area test the prescribed rates on a small number of plants for phytotoxicity prior to widespread use. Before using **A2114.02** in tank mixture with other products, test the mixture on a small number of plants for phytotoxicity prior to widespread use. Before using **A2114.02** on plants for diseases that are not listed in the Directions for Use, test **A2114.02** on a small scale first.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[Bag: Nonrefillable outer bag. **DO NOT** reuse or refill the outer bag. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.]

[Plastic Container: Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.]

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

[A2114.02 is a trademark of Atticus, LLC]

[Compass® is a registered trademark of Bayer.]

[Bayleton® is a registered trademark of Bayer.]

{LANGUAGE ON LABEL AFFIXED TO CONTAINER}

TRIFLOXYSTROBIN	GROUP	11	FUNGICIDE
-----------------	-------	----	-----------

A2114.02TM

[Alternate Brand Name: Centyk]

[Contains trifloxystrobin, the active ingredient used in Compass®.]

[For Turf, Landscape, Greenhouse and Nursery Use]

[FUNGICIDE]

[Turf and Ornamental Fungicide]

[For Control of Certain Foliar, Stem, and Root Diseases of Turfgrass Including Golf Courses, Institutional, Commercial and Residential Lawns, Sod Farms, Sports Fields, Parks, Municipal Grounds and Cemeteries, and of Ornamentals Grown in Interiorscapes, Field Nursery Plantings, Forest Nurseries, Residential and Commercial Landscapes, Greenhouses, Lath and Shadehouses, Containers, and Other Enclosed Structures.]

ACTIVE INGREDIENT:	(% by weight)
Trifloxystrobin (CAS No. 141517-21-7).....	50.0%
OTHER INGREDIENTS:.....	50.0%
TOTAL	100.0%

A2114.02 is a water-dispersible granule.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If in eyes:	<ul style="list-style-type: none">Hold eye open and rinse slowly and gently with water for 15-20 minutes.Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">Take off contaminated clothing.Rinse skin immediately with plenty of water for 15-20 minutes.Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	
NOTE TO PHYSICIAN: If ingested, induce emesis or lavage stomach. Treat symptomatically.	

For Chemical Emergency:

Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Harmful if absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff

may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** contaminate water when disposing of equipment wash water or rinsate. Applying this product when rain is not predicted for the next 24 hours will help reduce potential risk to aquatic invertebrates by reducing pesticide runoff from the treatment area into water bodies.

GROUND WATER ADVISORY: Several trifloxystrobin degradates have properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

PHYSICAL OR CHEMICAL HAZARDS: **DO NOT** use, pour, spill, or store near heat or open flame.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[Bag: Nonrefillable outer bag. **DO NOT** reuse or refill the outer bag. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.]

[Plastic Container: Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.]

See inside label booklet for additional Precautionary Statements and Directions for Use.

[A2114.02 is not manufactured, or distributed by Bayer Environmental Science, seller of Compass®.]

Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

EPA Reg. No.: 91234-XX
EPA Est. No.: _____
NET CONTENTS: _____

{Optional Marketing graphics}

