



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Biopesticides and Pollution Prevention Division (7511P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

91473-3

Date of Issuance:

7/12/2019

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

GEMSEI

Name and Address of Registrant (include ZIP Code):

Seipasa, S.A.
 C/ Almudevar No. 2
 Tardienta (Huesca), 22240
 Spain

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA Registration Number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA or the Act).

Registration is in no way to be construed as an endorsement or recommendation of this product by the U.S. Environmental Protection Agency (EPA). In order to protect health and the environment, the Administrator, on his or her motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under the Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration or registration review of your product when the EPA requires all registrants of similar products to submit such data.
2. Make the following labeling change before you release this product for shipment:
 Revise the EPA Registration Number to read, "EPA Reg. No. 91473-3".
3. Submit one (1) copy of the final printed labeling for the record before you release this product for shipment.

Signature of Approving Official:

Andrew Bryceland, Team Leader
 Biochemical Pesticides Branch
 Biopesticides and Pollution Prevention Division (7511P)
 Office of Pesticide Programs

Date:

7/12/2019

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6. A stamped copy of the labeling is enclosed for your records.

Please also note that the record for this product currently contains the following acceptable Confidential Statement of Formula (CSF):

- Basic CSF dated 5/25/18

If you have any questions, please contact Chris Pfeifer of my team by phone at 703-308-0031 or via email at pfeifer.chris@epa.gov.

Sincerely,

A handwritten signature in blue ink that reads "Andrew C. Bryceland". The signature is fluid and cursive.

Andrew Bryceland, Team Leader
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)
Office of Pesticide Programs

Enclosure

GEMSEI


GLOBAL BIOPESTICIDE / BROAD SPECTRUM BIOPESTICIDE /INSECTICIDE-MITICIDE

ACTIVE INGREDIENT

Garlic oil 6%

OTHER INGREDIENTS94%

TOTAL.....100%



WARNING/AVISO

KEEP OUT OF REACH OF CHILDREN

READ ALL THE DIRECTIONS BEFORE USING THIS PRODUCT

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If in eyes	<ul style="list-style-type: none">•Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.•Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none">•Call a poison control center or doctor immediately for treatment advice.•Have person sip a glass of water if able to swallow.•Do not induce vomiting unless told to do so by a poison control center or a doctor.•Do not give anything to an unconscious person.
If on skin	<ul style="list-style-type: none">•Take off contaminated clothing.•Rinse skin immediately with plenty of water for 15-20 minutes.•Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For non-emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 (www.npic.orst.edu). For medical emergencies, call the poison control center at 1-800-222-1222.	

{Optional statement for use on small container labels} [See (booklet) (pamphlet) (back panel) (side panel) for (Precautionary Statements) (First Aid) (and) (Directions for Use)]

Net Content:

EPA Reg. No. 91473-G

EPA Est. No. 91473-ESP-1

Manufacturer:

Seipasa, S.A.

C/ Almudevar nº 2

22240 Tardienta (Huesca) Spain

www.seipasa.com

{Container Sizes} [4 fl. oz.; 8 fl. oz.; 1 pint; 1 quart; 1, 2.5, 3, 50 and 250 gal.]

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING. Causes substantial but temporary eye injury. Causes skin irritation. Harmful if swallowed or absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Do not get in eyes, skin or on clothing. Avoid contact with skin, eyes or clothing. Wear waterproof gloves and protective eyewear (goggles, face shield, or safety glasses). Wash with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before use.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirts and long pants
- Shoes plus socks
- Protective eyewear
- When mixing, loading and cleaning equipment wear a chemical-resistant apron
- For overhead exposure wear chemical-resistant headgear

USER SAFETY REQUIREMENTS

- Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.
- Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labelling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State and Tribe, consult the State/Tribe agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labelling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and involves contact with anything that has been treated such as plants, soils, or water, is:

- Coveralls.
- Waterproof gloves.
- Shoes plus socks.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter until sprays have dried.

USE INFORMATION

GEMSEI is an innovative formulation with a high efficacy against pests. GEMSEI exerts a powerful effect as an insecticide/miticide, repellent, odor disguising and antifeedant deterrent.

GEMSEI is a green formula which contains a high amount of 4 molecules belonging to the organosulfur family. This complexity fights against pests by four different modes of action:

- Allyl polysulfides (DAPs I, II, III and IV), contained in this botanical extract, act against insects and mites because they cause a disruption of cellular redox balance and enzyme function. Finally, anticholinesterase activity (neurotoxicity) is provoked.
- GEMSEI also works on contact softening exoskeletons and promoting fast dehydration. Its lipophilic nature allows interaction and perturbation of membrane structures.
- GEMSEI's repellency affects pests through interference with the sensing faculties of pests, disguising the odor signal emitted by crops, thus, preventing females from laying eggs, and diminishing pest establishment on crops.
- GEMSEI also has a direct effect on pest food consumption impeding ingestion and causing insect death.

The multiple modes of action of GEMSEI limit the likelihood of resistance developing in target pests.

GEMSEI does not leave a garlic taste on food crops.

GEMSEI can be applied alone or in combination and/or rotation with chemical insecticides as a tool for IPM (Integrated Pest Management) in agricultural crops, ornamentals and turf-grass. GEMSEI offers a valuable tool for management of resistance to chemical insecticides through its multiple and unique modes of action.

TANK MIXING

Shake properly before using. GEMSEI has been found compatible with most commonly used insecticides, and miticides. Prior to use it, in order to avoid problems, conduct a compatibility test before using this product in a tank mixed with other pesticides. Check physical compatibility first, and then mix the correct proportion of products in a small jar. Once this is done, test tank mix combinations on a representative sample of plants, checking if phytotoxicity symptoms appear. Due to the wide variation in climatic conditions, cultural practices, and other factors, the user assumes full responsibility for any crop damage or other liability resulting from the use of GEMSEI in a tank mix combination. Add required amount of GEMSEI to a clean spray tank with at least one-half of the water to be sprayed. Agitate the mixture thoroughly and then fill the tank with remaining water and continue agitation. Always use this product promptly after mixing with water and do not let tank mix sit for any extended period. Preferably do not

mix GEMSEI with Captan, Bordeaux mixtures, oxidizing agents such as bleach or highly alkaline or acid products as they will destabilize the product and can cause unacceptable phytotoxicity and/or reduced effectiveness on target pests.

pH: It is advisable to use GEMSEI in a pH water solution adjusted to neutral or slightly acid (around 6 - 6.5).

APPLICATION INSTRUCTIONS

Read all the directions before using this product. Apply GEMSEI directly to any food or non-food crop growing in fields and greenhouses up to and including the day of harvest, at a maximum rate of 4 pints per acre.

Begin applications at the appearance of the first signs of insects and mites. Repeat applications as required to maintain an effective control (5 days as a minimum interval application).

Pay special attention under moderate to severe pest pressure, or when environmental conditions and plant stage are conducive to rapid pest/mite reproduction rate. Use higher label rates, apply more frequently and rotate GEMSEI with other insecticides/miticides for improved performance.

Refer to the "Crop Uses" section for a complete list of crops. For foliar application, apply GEMSEI in a sufficient spray volume and with adequate spray pressure to ensure complete and thorough coverage of all plant surfaces including both the top and bottom of leaves.

GEMSEI has been found to be compatible when used in conjunction with most beneficial insects.

For greenhouse and outdoor use.

RATE CHART

	<i>First presence*</i>	<i>Common rate*</i>	<i>High pressure*</i>	
GEMSEI (**)	1 pint per acre	2 pints per acre	3 pints per acre	4 pints per acre
	16 fl. oz. per acre	32 fl. oz. per acre	48 fl. oz. per acre	56 fl. oz. per acre
Frequency	5 - 7 days	5 - 10 days	5 - 7 days	

*Rate/100 gallons of water

** in those crops where water volume applications are up to 120 gallons per acre or more, use a Dilution Rate between 40-55 fl. oz. per 100 gallons.

GROUND APPLICATIONS: GEMSEI can be applied in most commonly-used ground application equipment, such as tractor-mounted boom, air-blast high clearance, hose-end, backpack, and other pressurized sprayers; hose-end or hand-help sprayers; foggers or mist blowers; water wheel and other drench applicators; and shank or other soil injection methods.

To achieve better results coverage typically requires a minimum of 30 gallons of total spray volume per acre. Higher volumes will be required for larger perennial crops. Use lower rates for light to moderate pest invasion pressure and higher rates for high infestations.

When pest pressure is intense or plant canopy is dense, use higher rates and increase spray frequency. Repeat application if measurable rain occurs within two to three hours of spraying.

AERIAL APPLICATIONS: GEMSEI can be applied by fixed or rotary winged aircraft in a minimum of 30 gallons per acre (depending on the crop) of total spray volume gallons per acre. Standard precautions should be taken to minimize spray drift.

CHEMIGATION: GEMSEI can be applied through drip (trickle) and sprinkler (including center pivot, lateral move, end tow, side [wheel] row, traveler, big gun, solid set or hand move); flood; furrow; or border irrigation equipment. For detailed instructions refer to the section entitled “Chemigation Instructions”.

PREHARVEST INTERVAL: 0 days

REI (restricted entry interval): 4 hours.

TARGETED PESTS: Use GEMSEI in a foliar spray to control, suppress or repel soft bodied pests, piercing and sucking pests, and mites such as:

- **Mites** (such as two spotted spider mites, *Tetranychus* spp., *Oligonychus* spp., european red mite *Panonychus* spp., Willamette spider mite, *Polyphagotarsonemus latus*, *Phyllocoptura oleivora*, *Aculops lycopersici*, *Brevipalpus phoenicis*, *Steneotarsonemus spinki*, among others).
- **Aphids** (such as cotton aphid, green peach aphid among others).
- **Thrips** (such as *Thrips* spp., *Frankliniella* spp., *Heliethrips* spp., *Scirtothrips citri* among others).
- **White flies** (such as *Dialeurodes* spp., *Trialeurodes* spp. *Bemisia* spp., among others).
- **Armyworms, tuberworms, caterpillars and loopers** (such as cabbage looper, tent caterpillar, fruit worms, alfalfa caterpillar, caterpillar larvae among others, tomato pin worm, peach twig worm, dark sword-grass among others).
- **Moth** (such as codling moth, grape berry moth among others).
- **Flies** (such as walnut husk fly, fungus gnats, *Delia radicum*, and others fruit flies).
- **Leafhoppers** (such as *Empoasca* spp., *Macrostelus* spp., and *Graphocephala* spp., among others).
- **Psyllids** (such as *Bactericera* spp., *Diaphorina citri*, *Cacopsylla* spp., among others).
- **Beetles** (Japanese beetles, carpet beetles, among others).
- **Scales** (such as California red scale, san Jose scale, and other scales).
- **Weevils:** *Cacopsylla chinensis* among others.
- Other pests such as **borers** (Peach twig borer, *Hypothenemus hampei*), **mealybugs** (*Planococcus* spp., *Pseudococcus* spp., *Dysmicoccus brevipes*), **plant bugs** (squash bug, apple maggots, lygus bugs, blueberry maggot, leafrollers), **chinch bugs** and **grasshoppers**.

CROPS USES:

Fruits, Citrus and Nut Trees			
Crops			
Apricots, Cherries, Nectarines, Peaches, Plums, Prunes	Apples, Crabapples, Pears, Quinces	Oranges, Lemons, Limes, Grapefruits, Kumquats, Mandarins	Almonds, Pecans, Pistachio, Walnuts

Vegetables: Leafy, Fruiting, Cucurbit and oriental vegetables		
Crops		
Celery, Endive, Lettuce, Parsley, Radicchio, Mustard greens, Turnip greens	Eggplant, Ground cherry, Okra, Pepper, Tomato	Chayote, Chinese cucumber, Citron melon, Cucumber, Melons, Pumpkin, Squash, Watermelon

Vegetables: Brassica (Cole), Bulb and Root and Tuber Vegetables		
Crops		
Broccoli, Cabbage, Chinese Cabbage, Cauliflower, Kale	Garlic, Leeks, Onions, Shallots	Artichokes, Sugar beet, Chicory, Ginger, Ginseng, Horseradish, Potatoes

Small Fruits, Berries and Grapes

Crops

Blackberry, Blueberry, Cranberry, Currant, Grapes, Guava, Raspberry, Strawberry

Field Crops: Legume Vegetables, Cereal Grains and Forage and Grasses

Crops

Beans, Lentils, Peas, Soybeans	Barley, Corn, Millet, Oat, Rice, Sorghum, Triticale, Wheat	Any grass that will be fed to or grazed by livestock
--------------------------------	--	--

Herbs, Spices and Other Crops

Crops

Alfalfa, Basil, Chives, Cilantro, Dill, Mint, Oregano, Parsley, Pasture, Thyme, Sesame.	Avocados, Bananas, Dates, Figs, Mangoes, Papaya, Pomegranate, Pineapple, Acerola, Artichoke, Cacao, Canistel, Carob, Cherimoya, Cocoa, Coffee, Hops, Lychee, Mamey sapote, Mushroom, Okra, Olives, Palm, Passion fruit, Peanuts, Persimon, Sugarcane, Sunflower, Tea, Tobacco, Sunflower, Cotton, Jojova, Asparagus
---	---

Greenhouse Uses

Crops

Vegetables, Herbs, Spices	Ornamental plants, shrubs and trees; bedding plants; flowers
----------------------------------	---

Note to User: Unintentional consequences such crop injury may result due to environmental or growing conditions, manner of use or application. Because plant tolerance to pesticides varies as conditions vary, treat a few plants under conditions unexpected to be encountered and observe for plant damage prior to full-scale application to large number of plants.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store product in the original container. Store away from food, pet food, feed, seed, fertilizers and veterinary supplies. Protect pesticide containers from extreme heat and cold.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container.

For plastic containers, less than or equal to 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For plastic containers, greater than 5 gallons: Triple rinse as follows: Empty the remain contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip

container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CHEMIGATION INSTRUCTIONS

Apply this product only through drip (trickle), sprinkler (solid set, lateral move, end tow, sideroll, center pivot, or hand move), (basin), furrow, or border irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, discharge the water from the public water system into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

DRIP TRICKLE CHEMIGATION:

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump such as a positive displacement injection pump (i.e., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Dilute the product in water following the label mixing directions. It may be premixed in a supply tank with water, fertilizer, or other appropriate tank-mixed agricultural chemicals. Agitation is necessary. Apply to moderately moist soils. Use volumes that thoroughly wet the soil but that do not cause significant runoff or excessive drip from pots. Application should be continuous in sufficient water to apply the application rate evenly to the entire treated area.

SPRINKLER CHEMIGATION:

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must also contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump such as a positive displacement injection pump (i.e., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Dilute the product in water following the label mixing directions. It may be premixed in a supply tank with water, fertilizer, or other appropriate tank-mixed agricultural chemicals. Agitation is necessary. Apply when soils are moderately moist. Use volumes that thoroughly wet the foliage and/or soil but that do not cause significant runoff or excessive drip from pots. Application should be continuous in sufficient water to apply the application rate evenly to the entire treated area.
8. Do not apply when wind speed favors drift beyond the area intended for treatment.

FLOOD (BASIN), FURROW AND BORDER CHEMIGATION:

1. Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential of water source contamination from the backflow if water flow stops.
2. Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:
 - a. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
 - b. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
 - c. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
 - d. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
 - e. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
 - f. Systems must use a metering pump, such as a positive displacement injection pump (i.e., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
3. Dilute the product in water following the label mixing directions. It may be premixed in a supply tank with water, fertilizer, or other appropriate tank-mixed agricultural chemicals. Agitation is necessary. Apply to moderately moist soils. Use volumes that thoroughly wet the soil but that do not cause significant runoff. Application should be continuous in sufficient water to apply the application rate evenly to the entire treated area.

WARRANTY STATEMENT

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following conditions, Disclaimer of Warranties and Limitations of Liability.

SEIPASA S.A. warrants that this product conforms to its description and is reasonably fit for the purposes stated on the label when used in accordance with directions for use. Buyers and users of this product assume the risk of any use contrary to such directions. Timing and method of application, weather, watering practices, nature of soil, the disease problem, condition of the crop, incompatibility with other influencing factors in the use of this product are beyond the control of the seller. To the extent consistent with applicable law, the Seller's liability for any breach of warranty shall not exceed the purchase price of the material as to which a claim is made. To the extent consistent with applicable law, Buyers and users of this product are responsible for all loss or damage from use or handling of this product which results from conditions beyond the control of Seller, or without the fault or negligence of the Seller, or from failure to follow the label.