



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

February 26, 2019

Kevin Spruce
Regulatory Product Manager
Belchim Crop Protection N.V./S.A.
c/o KNOELL USA, LLC
3070 McCann Farm Drive, Suite 112
Grant Valley, PA 19060

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment –
Correction of Registration Number and the addition of a distributor name and address
Product Name: Beloukha Agricultural Herbicide
EPA Registration Number: 91746-7
Application Date: 11/30/2018
OPP Decision Number: 546917

Dear Mr. Spruce:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the U.S. Environmental Protection Agency (EPA). If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims

Page 2 of 2
EPA Reg. No. 91746-7
OPP Decision No. 546917

made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Sandra Owusu-Ansah by phone at (703) 347-0350 or via email at owusu-ansah.sandra@epa.gov.

Sincerely,

A handwritten signature in blue ink that reads "Andrew C. Bryceland". The signature is fluid and cursive, with the first name being the most prominent.

Andrew Bryceland, Team Leader
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)
Office of Pesticide Programs

Enclosure

ACCEPTED

02/26/2019

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under

EPA Reg. No. 91746-7

GROUP

27

HERBICIDE

BELOUKHA

Agricultural Herbicide

ACTIVE INGREDIENTS:

Pelargonic acid* 71.96%

OTHER INGREDIENTS: 28.04%

TOTAL: 100.0%

* Contains 5.73 lbs of pelargonic acid per gallon of product

KEEP OUT OF REACH OF CHILDREN

DANGER / PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

READ THE LABEL AND ATTACHED BROCHURE BEFORE USING

EPA REG. NO. 91746-7

EPA Establishment No. – 92762-ITA-001

Net Contents – 1 gal to bulk

Produced for:

Belchim Crop Protection N.V./S.A.

Technologielaan 7

B -1840 Londerzeel

Belgium



Distributed by:

Belchim Crop Protection US

Corporation

2751 Centerville Road, Suite 100

Wilmington, DE 19808

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
IF INHALED	<ul style="list-style-type: none"> • Remove victim to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOTLINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact (800) 424-9300 for emergency medical treatment information.</p>	

For Chemical Emergency
 Spill, Leak, Fire, Exposure, or Accident
 Call CHEMTREC Day or Night
 1-800-424-9300

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER / PELIGRO

Causes severe skin burns and eye damage. Harmful if inhaled. Avoid breathing vapor or spray mist. Harmful if swallowed or absorbed through the skin. Avoid contact with eyes, skin and clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Causes irreversible eye damage. Do not get in eyes or on clothing. Wear protective eyewear such as goggles or face shield. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Coverall worn over short-sleeved shirt and short pants
- Waterproof or chemical-resistant gloves such as barrier laminate, nitrile, rubber or neoprene rubber
- Chemical resistant footwear plus socks
- Protective eyewear (goggles or face shield)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Controls Statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-5)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wash waters or rinsate. **DO NOT** apply to water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the WPS.

DO NOT enter or allow worker entry into treated areas during the **REI of 12 hours**.

For early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil, or water, wear: coveralls, chemical-resistant gloves made of any waterproof material, chemical resistant shoes and socks, and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep unprotected persons out of the treated areas until the sprayed area is dry

GENERAL INFORMATION

This product is a contact, non-selective, broad spectrum, foliar-applied herbicide. This product will only control actively growing emerged green vegetation. It provides burndown of both annual and perennial broadleaf and grass weeds, as well as most mosses and other cryptogams. The degree of burndown and the longevity of control are less when the plants are inactive, mature, or biennial/perennial types. This product is not translocated. It will burn only those plant parts that are coated with spray solution. This product is a non-volatile, emulsifiable concentrate. It can be applied through most standard or field type sprayers after dilution and mixing with water in accordance with label instructions. For best results, spray coverage should be uniform and complete. Visible effects on most weeds occur within hours. This product does not damage mature, non-green, woody parts of plants. Cool weather following treatment may slow the activity of this product and delay or reduce visual effects. This product provides no residual weed control. Repeat treatments will be necessary for new plants emerging from

seed or regrowth of treated vegetation. Should residual control be desired, use a product recommended for the use situation.

USE METHODS

See the "Use Sites" section of this label to match the method of use with the crop or use situation.

1	Vegetative Burndown - General control of weeds for seedbed or site preparation, non-crop and around aquatic sites. Spot treatments may be used in crop and pasture situations
2	Directed and Shielded Sprays - Applications may be made in and around desirable plants when contact of foliage and green bark is avoided.
3	Prior to Emergence of Plants from Seed, Perennial Rootstocks, Corms and Bulbs - Ensure applications are made before new growth or crop emerges from soil or damage will occur.
4	Dormant or Post Harvest Spray - For control of weeds growing in dormant turf or fields after this commodity has been harvested. Partially green growth will be killed or stunted
5	Sucker Control, Pruning and Trimming - To burn back unwanted basal sucker growth on woody trees and foliage growth on vines, and excessive cane growth in brambles. Apply only to unwanted vegetative parts. Apply before suckers become woody.
6	Harvest Aid and Desiccation - To remove leaves of plants prior to harvest and/or burn-down of weeds to facilitate harvest. Harvest aid and desiccation uses include applications to root and tuber vegetables, bulb vegetables and cotton only. Applications must be made no later than twenty-four hours prior to harvest (pre-harvest interval = 24 hours).
7	Structural and Building Applications - Apply to unwanted vegetation in and around buildings and structures. Application to walks, benches, walls, floors, roofs, cooling pads for the control of moss and certain algae. A temporary residue or precipitate may result when used on some types of concrete, masonry, brick or stone.

MIXING AND APPLICATION PROCEDURES

APPLY SPRAY SOLUTION IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES. APPLICATIONS SHOULD BE PROPERLY DIRECTED TO AVOID SPRAYING OR ALLOWING DRIFT TO DESIRABLE PLANTS. ALWAYS CLEAN TANK, PUMP, AND LINE THOROUGHLY WITH WATER AFTER USE.

Do not apply this product through any type of irrigation system.

This product may not be aerially applied. Spray droplet size should be adjusted to minimize drift and allow application to the intended exposure area only.

This product mixes readily with water. Prior to application, prepare the spray mixture by filling the spray or mix tank with $\frac{3}{4}$ the required amount of water then add the proper amount of product. Fill remainder of tank with water to the total amount of spray solution needed. Remove hose promptly from tank to avoid spray solution siphoning back into hose or water source. Mix well.

Foaming may occur during mixing and/or application. If needed, a defoaming agent may be added to prevent excessive foaming. Ensure that foaming agent is approved for use around crops if area to be sprayed includes edible crops.

For best results, ensure that the product is thoroughly mixed at filling and during spray application. Agitation should be maintained during operation. For sprayers without agitation, mix or shake solution regularly to maintain suspension. Without agitation, product separates quickly from spray solution.

NOTE: To assure compatibility of this product with other products, pour the products into a small container of water in the correct proportions. After thorough mixing, let stand for 5 minutes. If the combination remains mixed, or can be re-mixed readily, the mixture is compatible.

Make foliar applications by ground in sufficient water for complete coverage of the foliage. Unless otherwise specified elsewhere in this label, use a minimum of 20 gallons of water/Acre. **DO NOT** apply this product aerially. **DO NOT** apply this product through any type of irrigation system. Adjust spray droplet size to minimize drift and

contain application to intended exposure areas only.

For best results with backpack or other small tank applicators ensure thorough mixing of herbicidal solution at filling and during the spray operation. Sprayers without continuous agitation should be shaken or mixed regularly to maintain suspension. Without agitation, this product separates quickly out of the spray solution.

DO NOT apply to weeds when wet from dew, rain or irrigation. **DO NOT** irrigate within 2 hours after application. **DO NOT** apply if rainfall is expected within 2 hours.

BOOM EQUIPMENT: For best control of annual, biennial, or perennial weeds using conventional boom equipment, use the product rate indicated for the intended weed in 75 – 200 gallons of spray solution or water/acre as a broadcast spray. When tank mixed with other products, use lower delivery rates of 10 – 75 gallons/acre spray solution. Increase spray volume within the specified range as weed density increases. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

The pesticide supply tank should be equipped with a means for continuous agitation either by recirculation or a mechanical agitator. Charge the supply tank with the appropriate amount of water and add the pesticide slowly followed by any sticker-spreaders, insecticides, nutrients, etc. Observe all cautions and limitations on the labels of all products used in the mixtures.

HAND-HELD AND HIGH-VOLUME EQUIPMENT: For use with knapsack and backpack sprayers, pump-up pressure sprayers, handguns, hand wands, and other hand-held spray equipment and vehicle mounted high volume spray equipment in spray-to-wet applications. Apply spray solutions of this product to foliage of vegetation to be controlled. For most efficient use of spray mixture, spray all leaf surfaces uniformly and completely to wetness, but not to the point of runoff.

SELECTIVE PLACEMENT EQUIPMENT: Direct spray of the herbicide solution to the weeds using a shielded applicator that uses a physical barrier to protect desirable vegetation from the herbicide spray. Follow spray volume instructions of the equipment or nozzle manufacturer when using this application method.

USE RATES

Mix this product in the proportions as listed in the rates table below and deliver the spray solution through boom or high volume and hand held equipment.

SOLO USE

Use the following amounts of product (volume/volume) in a total spray volume of 75 to 200 gallons / acre using boom, hand-held or high volume equipment.

- Use a 2.5-4.0% solution for control of annual weeds, mosses and cryptogam. Use the lower end of the rate range for young, succulent and actively growing weeds and the higher end of the rate range for weeds taller than 6 inches in height or weeds in the flowering stage. Use the higher rate for herbicidal control mosses, lichens and other cryptogams on structures and surfaces.
- Use a 4.0-6.0% solution for burndown of perennial herbaceous plants, established weeds, and control of sucker growth. Use the high end of the rate range for perennial weeds at or past the flower stage or “hardened” weeds.
- Use a 6.0-8.5% solution for maximum vegetative burndown, edging or foliar trimming.

Repeat application as required to maintain desired level of weed control and to control emerging seedlings and underground growth.

TANK MIX USE

Tank mixing must be done in accordance with the most restrictive label limitations and precautions. **DO NOT** exceed any labelled dose rates. **DO NOT** mix this product with any product containing a label prohibition against such mixing.

Always predetermine tank mix capability of this product with other products prior to application. Mix proportional amounts of all products and water in a container and observe for adverse reactions. When tank mixing, add components in the following sequence (if using): (1) wettable powders, (2) flowable liquids, (3) emulsifiable liquids, (4) this product, (5) water soluble liquids, and (6) adjuvants.

Beloukha and Residual Herbicides: For burndown of vegetation followed by residual control of weeds emerging from seed or underground parts, mix this product with a recommended soil-active herbicide such as all formulations consisting of sulfonylurea, imidazilinone, triazine, dinitroaniline and acylurea herbicides. Follow the use instructions and rates of the residual companion product plus:

- 3 - 5% for annual weeds and vegetation
- 5 - 7% for perennial herbaceous and late stage annuals
- 7 - 10% for maximum vegetation burndown

RATE TABLE

Desired volume of spray solution (gallons)	Amount of Beloukha for Following % Solution (volume/volume)				
	1%	2.5%	4%	6%	8.5%
1	1 1/3 oz	3 1/3 oz	5 1/3 oz	8 oz	11 1/3 oz
5	6 2/3 oz	1 pt	1 2/3 pt	2.5 pt	1 1/2 qt
10	13 oz	1 qt	1 2/3 qt	2 1/2 qt	3 1/2 qt
50	2 qt	1 1/4 gal	2 gal	3 gal	4 1/4 gal
75	3 qt	1 3/4 gal	3 gal	4 1/2 gal	6 1/3 gal
100	1 gal	2 1/2 gal	4 gal	6 gal	8.5 gal

USE SITES

CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE OF DESIRABLE TURFGRASSES, TREES, SHRUBS, OR OTHER DESIRABLE VEGETATION SINCE DAMAGE MAY RESULT.

Best results are obtained when applications are made to young succulent weeds and when spray solutions cover all leaf surfaces. Mature, woody weeds are less susceptible. Repeat applications as needed to give desirable levels of weed control.

Select nozzles/pressure combinations that deliver large coarse droplets such as solid cones or flat fans at low pressures, and avoid nozzle/pressure combinations that generate fine particles or mist. If spraying areas adjacent to desirable plants, use a shield to help prevent spray from contacting foliage of desirable plants.

Treated areas may be reseeded or transplanted into as soon as desirable levels of weed control are obtained.

Crop Group	Representative Crops Such As	Use Method*
Root, tuber and perennial vegetables	Asparagus, artichoke, beet, carrot, ginger, ginseng, horseradish parsnip, potato, radish, rutabaga, sweet potato, turnip and yam. ¹ Harvest aid and desiccation uses are only approved for root and tuber vegetables in this crop group.	1, 2, 3, 4, 6 ¹
Bulb Vegetables	Garlic, leek, onion, and shallot.	1, 2, 3, 6
Cole or Brassica Crops	Broccoli, Brussels sprouts, cabbage, cauliflower, collards, kale, kohlrabi, greens (mustard and turnip)	1, 2, 3
Legume Vegetable	Beans (Phaseolus spp. such as: field green, kidney, lima, mung, navy, pinto, snap,	1, 2, 3

	and wax beans) (<i>Vigna</i> spp. such as: black-eyed, Chinese longbean, cowpea, and southern pea) peas (<i>Pisum</i> spp. such as: garden, green, sugar and snow pea), lentil, and soybean	
Fruiting Vegetables	Eggplant, okra, pepper (bell, chili, sweet), pimento, and tomato.	1, 2, 3
Cucurbits and Melons	Cucumber, gourd, muskmelon, cantaloupe, pumpkin, squash, and watermelon	1, 2, 3
Citrus	Grapefruit, kumquat, lemon, lime, orange, tangerine, and tangelo.	1, 2
Pome Fruit	Apple, crabapple, pear, and quince	1, 2, 5
Stone Fruit	Apricot, cherry, nectarine, peach, plum, and prune	1, 2, 5
Small Fruit, Berries, and Grape	Blackberry, blueberry, boysenberry, cranberry, currant, dewberry, grape (all types), loganberry, olallieberry, raspberry, and strawberry.	1, 2, 3, 5
Nuts	Almond, brazil nut, chestnut, filbert, macadamia, pecan, pistachio, and walnut	1, 2, 5
Tropical and Other Fruit	Avocado, banana, coconut, date, fig, guava, kiwi, mango, olive, persimmon, papaya and pineapple.	1, 2, 5
Field Crops and Cereal Grains	Barley, buckwheat, canola, corn (field, popcorn, sweetcorn), cotton ^a , cowpea, millet, oat, peanut, rice, rye, safflower, sorghum, sugarcane, sunflower, and wheat. ^a Harvest aid and desiccation uses are only approved for cotton in this crop group.	1, 2, 3, 6 ^a
Forages and Pasture Grasses (Forage or Seed)	Alfalfa, clovers, trefoil, vetch, bromegrass, fescue, bluegrass, lespedeza, ryegrass, sudangrass, timothy, range grasses, and crops grown for livestock feed.	1, 3, 4
Herbs & Spices	Anise, basil, caraway, chive, cumin, curry, dill, fennel, oregano, mints, rosemary, sage, savory, sweet bay, tarragon, thyme and wintergreen	1, 3, 2, 4
Beverage and Specialty Crops	Cocoa, coffee, hops, tea, tobacco, and jojoba.	1, 2, 5

*See the corresponding numbers in the "Use Methods" section of this label to determine the use and precautions.

Non-Crop Group	Representative Non-Crops Uses Such As	Use Method*
Turf, Flowers, Bedding and Landscape Plants	Turfgrass (maintenance, sod or seed production), bedding plants, flowers, and ornamentals.	1, 2, 3, 4, 5, 6
Trees and Shrubs	Christmas trees, forest and commercial trees, landscape trees, and nursery trees or shrubs	1, 2, 5
Greenhouse and Indoor Use	All crops, plants, and structures.	1, 2, 3, 7
Noncrop, Industrial, and Public Areas	Farmstead, homestead, fallow land, storage areas, schools, paved areas, rights-of-way (e.g., road, railroad, utilities), parking lots, recreation areas (e.g., athletic fields, campgrounds, golf courses, playgrounds), walks, industrial sites (e.g., lumberyard, tank farms, buildings).	1, 2, 7
Structures, Buildings, and Walkways	Bench, deck, equipment, floor, roof, walls, walks, and evaporative cooling pads	7
Dry Aquatic Sites, Dry Drainage Systems and Around Aquatic Sites	Applications must be made 72 hours prior to reflooding of dry aquatic sites. Dry ditches, dry canals, ditch banks, and for use above the water line or after draw-down of agricultural irrigation water and ditch systems, industrial ponds and disposal systems, and impounded water area	1, 7

*See the corresponding numbers in the "Use Methods" section of this label to determine the use and precautions.

USE PRECAUTIONS

AVOID DRIFT. CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS.

This product is a contact herbicide that kills or damages all green plant tissue. Do not allow spray solution to mist, drip, drift, or splash onto desirable vegetation since any green vegetation contacted by this herbicide may be damaged.

Keep people, pets and livestock off treated areas until spray solution has dried to prevent transfer of this product onto desirable vegetation.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Keep container tightly sealed when not in use. This product should be stored in its original container in a cool, dry locked place out of reach of children and pets.

PESTICIDE DISPOSAL: To avoid waste, use all material in this container by application according to label directions. If waste cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments).

CONTAINER HANDLING:

NOTE: This product is available in multiple containers. Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable" or "Refillable" designation. Follow the container handling instructions below that apply to your container type/size.

Nonrefillable Containers 5 Gallons or Less:

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration. Do not burn unless allowed by state and local ordinance. If burned stay out of smoke.

Nonrefillable containers larger than 5 gallons:

Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration. Do not burn unless allowed by state and local ordinance. If burned stay out of smoke.

Refillable containers larger than 5 gallons:

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration. Do not burn unless allowed by state and local ordinance. If burned stay out of smoke.

NOTICE TO BUYER

Seller warrants that this product conforms to the chemical description on this label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. To the extent consistent with applicable law, this warranty does not extend to use of this product contrary to label directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller. To the extent consistent with applicable law, the Seller shall not be liable for any such use. To the extent consistent with applicable law, the Seller makes no other warranties, either expressed or implied.

All trademarks appearing on this label are the property of their respective owners.