



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

December 19, 2019

Ms. Tracey Angel  
Regulatory Analyst  
Hawthorne Hydroponics LLC d/b/a General Hydroponics  
2877 Giffen Avenue  
Santa Rosa, CA 95407

Subject: Label Amendment – Adding Hemp Use and Updating Label  
Product Name: GH MPMT  
EPA Registration Number: 91865-2  
Application Date: May 7, 2019  
Decision Number: 551594

Dear Ms. Angel:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2  
EPA Reg. No. 91865-2  
Decision No. 551594

with FIFRA section 6. If you have any questions, please contact me by phone at (703) 308-2972 or via email at [walsh.michael@epa.gov](mailto:walsh.michael@epa.gov).

Sincerely,


A handwritten signature in black ink, appearing to read 'Michael Walsh', written in a cursive style.

Michael Walsh  
Product Manager 11  
Invertebrate & Vertebrate Branch 2  
Registration Division  
Office of Pesticide Programs

Attachment

**Note to PM:**  
***Bold, italicized text is information for the reader and is not part of the label.***  
[Bracketed information is optional.]

EPA Reg. No. 91865-2



# GH MPMT

INSECTICIDE – FUNGICIDE - MITICIDE  
CONCENTRATE

For foliar application [to control soft-bodied insects, mites, and powdery mildew]

ACTIVE INGREDIENT:	
Potassium salts of fatty acids .....	49.0%
OTHER INGREDIENTS: .....	51.0%
Total .....	100.0%

1 gallon of GH MPMT weighs 8.50 lb at 68°F and contains 4.16 lb active ingredient per gallon (498 g ai/L)

## KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

EPA Reg. No. 91865-2  
EPA Est. No. XXX-XX-X (*insert EPA Registered Establishment Number(s)*)  
[Superscript used is first letter of lot code]

[Sold by:] General Hydroponics, 2877 Griffen Ave., Santa Rosa, CA 95407

NET CONTENTS XX.XX fl oz (X.XX L)

***{Note to Reviewer: The final printed label will meet the print size requirements outlined in 40 CFR 156.10 for specific package sizes. This text in this note will not appear on the final printed label}***

**Note to PM:**

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional.]

EPA Reg. No. 91865-2

***Alternate Brand Names:***

- GENERAL HYDROPONICS EXILE
- GENERAL HYDROPONICS EXILE INSECTICIDE / FUNGICIDE / MITICIDE
- GENERAL HYDROPONICS M-BOLD INSECTICIDE / FUNGICIDE / MITICIDE
- GENERAL HYDROPONICS BANE INSECTICIDE / FUNGICIDE / MITICIDE

***Alternate Brand Names Sublabel A COMMERCIAL USE:***

- GENERAL HYDROPONICS EXILE PRO INSECTICIDE / FUNGICIDE / MITICIDE
- GENERAL HYDROPONICS EXILE INSECTICIDE / FUNGICIDE / MITICIDE PRO

***Alternate Brand Names Sublabel B HOMEOWNER USE:***

- GENERAL HYDROPONICS EXILE INSECTICIDE/FUNGICIDE/MITICIDE 1

**Note to PM:**  
***Bold, italicized text is information for the reader and is not part of the label.***  
[Bracketed information is optional.]

EPA Reg. No. 91865-2

SUBLABEL A COMMERCIAL USE

# GH MPMT

INSECTICIDE – FUNGICIDE - MITICIDE  
CONCENTRATE

For foliar application [to control soft-bodied insects, mites, and powdery mildew]

ACTIVE INGREDIENT:  
Potassium salts of fatty acids .....49.0%  
OTHER INGREDIENTS: .....51.0%  
Total .....100.0%

1 gallon of GH MPMT weighs 8.50 lb at 68°F and contains 4.16 lb active ingredient per gallon (498 g ai/L)

## KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

EPA Reg. No. 91865-2  
EPA Est. No. XXX-XX-X (*insert EPA Registered Establishment Number(s)*)  
[Superscript used is first letter of lot code]

[Sold by:] General Hydroponics, 2877 Griffen Ave., Santa Rosa, CA 95407

NET CONTENTS XX.XX fl oz (X.XX L)

**Note to PM:**

EPA Reg. No. 91865-2

**Bold, italicized text is information for the reader and is not part of the label.**

[Bracketed information is optional.]

<b>FIRST AID</b>	
If swallowed	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
If on skin or clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
If in eyes	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-465-5161 for emergency medical treatment information.	
<b>NOTE TO PHYSICIAN:</b> Probable mucosal damage may contraindicate the use of gastric lavage.	

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS & DOMESTIC ANIMALS****WARNING**

Causes substantial but temporary eye injury and skin irritation. Do not get in eyes, on skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)****Handlers who may be exposed to the diluted product through application or other tasks must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks

**Handlers who may be exposed to the concentrate through mixing, loading, application, or other tasks must wear:**

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils or viton ≥ 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear (goggles or face shield)
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing or loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

Users should:

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ENVIRONMENTAL HAZARDS**

This product may be hazardous to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

**Note to PM:**

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional.]

## DIRECTIONS FOR USE

It is violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and involves contact with anything that has been treated such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils or viton ≥ 14 mils
- Shoes plus socks

### Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

## USE INFORMATION

Apply GH MPMT to foliage as a contact insecticide, miticide and fungicide for control of soft-bodied insects, mites and powdery mildew. The formulation is based on potassium salts of naturally derived fatty acids. This product may be used to control targeted pests on crops which include vegetables, grapes and other small fruits, tree fruits, tree nuts, cotton, tobacco, hemp, shrubs, shade and ornamental trees, turf, foliage and flowering plants growing outdoors, in greenhouses and in interiorscapes. This product can be applied up to harvest.

## PESTS CONTROLLED

This product may be used only as a foliar spray to control or suppress soft bodied pests which include: adelgids, aphids, caterpillars, earwigs, lace bugs, leafhoppers, leafminers, mealybugs, mole crickets, plant bugs, psyllids, sawfly larvae, scales, spider mites, tent caterpillars, thrips, whiteflies, gypsy moth, and chinch bugs. GH MPMT is most effective on mites, aphids and whiteflies. Effects on other pests are limited to early growth stages and eggs. This product provides curative control of powdery mildew.

## MIXING AND APPLICATION

### RATES

**Stand Alone Rate:** GH MPMT rates as low as 0.25% v/v or as high as 4% v/v may be used. The 2% v/v rate is the most frequently used stand alone rate.

- Refer to individual sections of this label for specific use rates for control of pests on specific crops.
- Rates are provided as a % volume/volume (v/v) solution. A 2% solution is prepared by adding 2 gallons of GH MPMT to 98 gallons of spray water.
- The activity of this product is directly related to the solution concentration and to contact with the pest, not the amount of product applied per acre.

**Tank Mix Rate:** To achieve enhanced and residual pest control, use a 1 to 2% v/v solution of this product in combination with a specified product.

**Note to PM:**

EPA Reg. No. 91865-2

**Bold, italicized text is information for the reader and is not part of the label.**

[Bracketed information is optional.]

**RATE TABLE**

Total Mix Volume	% v/v Solution	Amount of GH MPMT		Tablespoons
		English	Metric	
1 gallon (3.785 liters)	0.25	0.32 fl oz	9.46 ml	0.64
	0.5	0.64 fl oz	18.93 ml	1.28
	1.0	1.28 fl oz	37.85 ml	2.56
	1.5	1.92 fl oz	56.78 ml	3.84
	2.0	2.56 fl oz	75.71 ml	5.12
100 gallon (378.5 liters)	4.0	5.12 fl oz	151.42 ml	10.24
	0.25	0.25 gal	0.95 liters	-
	0.5	0.50 gal	1.89 liters	-
	1.0	1.00 gal	3.79 liters	-
	1.5	1.50 gal	5.68 liters	-
	2.0	2.00 gal	7.57 liters	-
	4.0	4.00 gal	15.14 liters	-

**Mixing**

- Always determine the compatibility of water, pesticides and other tank additives with this product prior to addition to the spray tank. Determine compatibility by conducting a jar test using proportional quantities of each material.
- Use freshly mixed spray made with soft water. If water has high mineral content (hardness >300 ppm or 17.5 grains/gallon), check for compatibility as described above. If needed, add an approved compatibility agent to the tank prior to the addition of this product. Do not lower the pH of the final spray mixture below 8.0. See notes to User: Understanding Plant Sensitivity.
- A defoaming agent may be needed for use in sprayers equipped with an agitator.
- This product is compatible with other labeled pesticides. However, when tank mixing GH MPMT with other products, a compatibility test (jar test) using relative proportions of the tank mix ingredients should be conducted prior to mixing ingredients in the spray tank.
- **It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.**
- Unless otherwise noted, do not tank mix this product with adjuvants such as penetrators, spreader stickers or activators, gibberellic acid, calcium nitrate, or diatomaceous earth, foliar nutrients, alkaline based chelating agents (such as EDTA), chlorothalonil and pesticides containing sulfur or metallic ions (such as manganese, magnesium, iron, zinc, etc.) as they may be physically incompatible and/or phytotoxic.
- **DO NOT** mix GH MPMT with Carzol® insecticide (EPA Reg. No. 10163-265) as they are not physically compatible.
- **DO NOT** mix GH MPMT with sulfur. Do not use within 3 days of a sulfur application.

**When using this product in a tank mix, read and follow all product labels. The order of mixing is:**

- Compatibility agent (if needed)
- Wettable and soluble powders
- Flowable liquids
- Emulsifiable oils and concentrates
- GH MPMT

Application of spray, mixtures should conform to use precautions and directions for all products included in the tank mix. Combinations should be kept agitated and sprayed promptly after mixing.

**Timing of Application**

- Initiate applications as soon as pest is observed, or as needed to avoid economic injury. Repeat applications at one-week intervals, or as needed, to control infesting or increasing pest populations.
- Do not make repeat applications at less than 7-day intervals, or make more than 3 sequential applications at 7- to 14-day intervals unless your experience has shown the use pattern to be non-injurious to the type and the development stage of plant(s) to be treated under local growing conditions.

**Application Guidelines**

- This product may be applied with most types of ground or aerial spray equipment. **Do not** apply through any type of hydroponic systems. **Do not** use with pulse or thermal fogger applicators. **Do not** apply through any type of irrigation system.
- This product must come into contact with the targeted soft-bodied pests or powdery mildew to be effective. Complete coverage of pests is essential for maximum control. Direct sprays to contact the pests. In general, adequate coverage requires a minimum of 50 gallons of total spray volume per acre. Higher volumes may be required on larger perennial crops.
- Apply GH MPMT solutions to wet (minimize run-off) to decrease the potential for injury on foliage, fruit and flowers of sensitive plants.
- Apply promptly after mixing with other pesticides to avoid or reduce alkaline hydrolysis of certain pesticides.
- Avoid applications when wind conditions impair thorough coverage.
- Avoid applications on new transplants and unrooted cuttings.

**Note to PM:**

EPA Reg. No. 91865-2

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional.]

**Notes to User: Understanding Plant Sensitivity**

- Sensitivity of plants to this product can be influenced by several factors which include: pest and disease pressure, cultivar, plant vigor, environmental conditions (temperature, relative humidity, moisture availability, light intensity, etc.), spray concentration, companion products, spray additives, pH of spray mixtures, delivery volume as well as the timing, number and frequency of applications.
- Potential for plant injury increases when this product is used under the following conditions:
  - Plants are stressed, such as under hot (>90°F), humid and/or drought conditions.
  - More than 3 sequential applications are made at 7-day intervals.
  - The pH of the final spray mixture is lowered below 8.0.
  - High volume applications cause the spray to collect on the bottom of fruit such as apple, pear, nectarine, orange and grape.
  - Tender new foliage is present on narrow leaf evergreen trees and shrubs.
  - When existing insect, mite or disease pressure has already stressed plant foliage or damaged fruit (for example, mite burn on leaves).
- Application of this product may alter the appearance of the glaucous bloom on spruces and the waxy bloom on grapes.
- Evaluate the effect of this product alone or in combination with desired companion product(s) on a few test plants prior to full scale applications when sensitivity of plant species or cultivars under local use conditions is unknown.
- **Do not** apply to very sensitive plants such as: horse chestnut, Japanese maple, mountain ash, Cherimoya, bleeding heart or sweetpeas.
- The following plants may be sensitive to this product: bald cypress, begonia, chrysanthemum, Crown of Thorns and other Euphorbia, cucumber, delicate ferns, narrow leaf evergreens (especially when stressed or when tender new growth is present), Dieffenbachia, fuschia, gardenia, impatiens, jade plant, lantana, ornamental ivy, palms, poinsettia, redbud, river birch, schefflera, Zebra plant and some succulents. Test first on a few plants prior to broad scale applications when sensitivity under local use conditions is unknown.
- Flowers of plants including African violet, ageratum, azalea, begonia, camellia, chrysanthemum, dahlia, geranium, gloxinia, impatiens, lily, marigold, orchid, pansy, petunia, poinsettia bracts, rose, salvia, snapdragon, vinca and zinnia may be injured when sprayed with this product.
- Safety to blooms should be determined for each individual cultivar of plant to be treated by conducting a small test under local use conditions.

**Note to PM:**

EPA Reg. No. 91865-2

**Bold, italicized text is information for the reader and is not part of the label.**

[Bracketed information is optional.]

**Insecticidal / Miticidal and Fungicidal Uses**

This product may be used as a foliar spray to control or suppress soft bodied pests which include: adelgid, aphid, earwig, lace bug, leafhopper, mealybug, mole cricket, plant bug, psyllid, sawfly larva, scale, spider mite, tent caterpillar, thrips, whitefly, gypsy moth, and chinch bug.

**Preharvest Interval**

GH MPMT has a 0-Day Preharvest interval for ALL crops contained on this label.

<b>Vegetables, Herbs And Spices (Greenhouse and Outdoor)</b>	
<b>Vegetables:</b> Artichoke*, Asparagus, Brassica (cole), (such as: bok choy, broccoli, broccoli raab, Brussels sprout, cabbage, cauliflower, Chinese broccoli, Chinese cabbage, collard greens, kale, kohlrabi, mustard greens, napa cabbage and Swiss chard), Bulb (such as: green onion, leeks, garlic and onion), Cucurbit (such as: cucumber, melon, pumpkin and summer squash), Fruiting (such as: eggplant, pepper and tomato), Leafy (such as: celery, endive, escarole, lettuce and spinach), Legume (such as: bean, pea and soybean), Root and tuber (such as: beet, carrot, horseradish, potato, radish, rutabaga, shallot, sugarbeet and yam), Herbs and spices: (such as: basil, chive, dill, marjoram, parsley and sage). Watercress* * Not approved for use in California	
<b>Use Rates:</b> Stand Alone Rate: Apply a 1-2% v/v solution of this product, unless otherwise noted. Refer to the Mixing and Application section of this label for rates, timing and application information except as noted below.	
<b>Pest</b>	<b>Requirements</b>
Leafminers and Green peach aphids	Apply this product only in tank mix combination with a labeled companion insecticide.
Other aphids	To achieve enhanced and residual pest control, use a 1-2% v/v solution of this product in combination with a specified rate of another labeled product. Consult your local extension specialist, certified crop advisor, and/or manufacturer representative for specific product recommendations.
Whiteflies	Apply a 1-2% v/v solution of this product for stand-alone use. To achieve enhanced and residual pest control, use a 1 -2% v/v solution of this product in combination with a specified rate of another labeled product. Consult your local extension specialist, certified crop advisor, and/or manufacturer representative for specific product recommendations.
Thrips, Plant bugs, Spider mites, Broad mites, Russet mites and Leafhoppers	See Mixing and Application section of this label for information on rates, mixing and application.
Powdery mildew	<b>Cucurbits, Fruiting Vegetables, Herbs and Spices, Leafy Vegetables, Legumes, Root and Tuber Vegetables and Greenhouse Cucumber:</b> <b>Stand Alone Rate:</b> Apply a 1.5-2% v/v solution of this product when the first signs of powdery mildew appear and repeat at 7 to 10 day intervals, or as needed. Schedule a follow-up fungicide application within 7-10 days after each application when this product is used alone. <b>Tank Mix Rate:</b> Apply a 1-2% v/v solution of this product with GH DNMT [or another specified fungicide] to achieve protectant/residual control of powdery mildew. For additional instructions, see Application Guidelines and Notes to User: Understanding Plant Sensitivity sections of this label.

**Note to PM:**

**Bold, italicized text is information for the reader and is not part of the label.**

[Bracketed information is optional.]

<b>Fruit, Nut And Vine Crops</b>	
<p><b>Citrus fruits</b> (such as: citron, grapefruit, kumquat, lemon lime, tangelo, and tangerine), <b>Pome fruits</b> (such as: apple, crabapple, pear and quince), <b>Stone Fruits</b> (such as: apricot, cherry, nectarine, peach, plum and prune), <b>Small Fruits and Berries</b> (such as: blackberry, blueberry, cane berries, cranberry, currants, grape, raspberry, strawberry, boysenberry, and olallieberry) and <b>Tree Nuts</b> (such as: almond, chestnut, filbert, pecan, pistachio, black and English walnut) and <b>Pomegranates*</b>.                      * Not approved for use in California</p>	
<p><b>Use Rates:</b>  <b>Stand-alone Rate:</b> Apply a 1-2% v/v solution of this product, unless otherwise noted. Refer to the Mixing and Application section of this label for rates, timing and application information except as noted below.</p> <ul style="list-style-type: none"> <li>• <b>For all Fruit and Nut Crops:</b> Apply to wet (minimize run off) with an air blast or other appropriate sprayer. Application of excessive water volumes will increase the potential for fruit injury by causing collection of spray on the bottom of fruit, especially when poor drying conditions exist.</li> <li>• <b>Grapes:</b> This product may alter the waxy bloom and or cause marking on the surface of grapes which can affect the quality rating of table grapes at harvest. To reduce potential for fruit injury, do not apply in greater than 75 gallons of water per acre on table grapes after berries are 6 -7 mm in diameter. <b>Do not apply this product to Calmeria or Italia varieties of grapes.</b></li> <li>• <b>Citrus (Precaution):</b> Fruit marking may occur when sooty mold is present.</li> <li>• <b>Pears: Do not apply</b> this product after the delayed dormant stage on fresh market pear varieties including d'Anjou and Asian pears. Applications after fruit set may cause marking which can reduce market value of the fruit.</li> <li>• <b>Stone Fruits:</b> Concentrate (air blast) applications are specified with a maximum volume of 100 gallons per acre on smooth skinned fruits and 175 gallons per acre on peaches. Do not make dilute spray applications on smooth skinned fruit after fruit formation. This product is not specified for use on yellow skinned nectarine varieties. <b>Do not apply GH MPMT to sweet cherries between fruit formation and harvest.</b></li> </ul> <p>Nursery and non-bearing trees can be sprayed to runoff with no more than 3 consecutive applications with 7-day intervals.</p>	
<b>Pest</b>	<b>Requirements</b>
Citrus rust mites	Apply a 1-2% v/v solution of this product. Use of a suitable citrus spray oil in combination with this product applied to wet (minimize run-off) with an air blast or other appropriate sprayer may help improve performance.
European red mites, Pacific mites, two spotted mites, Willamette mites	Apply a 1-2% v/v solution of this product to control motile stages.
Pear psylla	Apply a 1-2% v/v solution of this product at delayed dormant to control winterform and newly hatched nymphs. Make follow up applications on processing pears at petal fall and turn down for control of 1st and 2nd generation nymphs before they mature into fourth instars.
Rosy apple aphids	Apply a 1-2% v/v solution of this product. Addition of an appropriate spray oil to this product may help improve performance. Time sprays to coincide with pre-pink or pink stages of apples.
White apple leafhoppers	To achieve enhanced and residual control, use a 1 -2% v/v solution of this product in combination with products such as Thiodan. Target 1st generation nymphs before they mature into flyers.
Other aphids; campylomas, mealybugs, rust mites, pear slug (sawfly larvae) scale crawlers, leafhoppers and whiteflies	See Mixing and Application section for information on rates, mixing and application.
Powdery mildew	<p><b>Grapes:</b> Apply a 1-2% v/v solution of this product to control powdery mildew spores and hyphae. Initiate sprays at bud break or as appropriate based on local powdery mildew control requirements. Repeat at 7 to 10 day intervals, or as needed through veraison (development of variety-specific color). Do not apply past veraison. Schedule a follow-up fungicide application within 7 to 10 days after each application when GH MPMT is used alone. Use this product in a scheduled preventive spray program with other labeled fungicides. <b>Do not mix with sulfur. Do not use within 3 days of sulfur application.</b> This product may be applied in tank mixes with other labeled fungicides. Consult your local extension specialist, certified crop advisor, and/or manufacturer representative for specific product recommendations.</p> <p><b>Tank Mix Rate:</b> Apply a 1-2% v/v with labeled rates of companion pesticides. Rates of GH MPMT below 1.5% v/v will not be fungicidal against powdery mildew. Complete coverage of grape clusters as well as leaf surfaces is essential for effective control. For additional instructions on use with sulfur, see Application Guidelines and Notes to User: Understanding Plant Sensitivity sections of this label.</p>
Powdery mildew	<p><b>Citrus fruit, Pome fruit, Stone fruit, Small fruits and Berries, and Tree Nuts:</b> Apply a 1 - 2% v/v solution of this product to control powdery mildew spores and hyphae.</p> <p><b>Stand Alone Rate:</b> Apply a 1 - 2% v/v solution of this product when first signs of powdery mildew appear or as appropriate based on local powdery mildew requirements and repeat at 7 to 14-day intervals, or as needed. Schedule a follow-up fungicide application within 7 to 14-days after each application when this product is applied alone.</p> <p><b>Tank Mix Rate:</b> Apply a 1 - 2% v/v solution of this product with a fungicide to achieve protectant/residual control of powdery mildew. <b>Do not mix with sulfur. Do not use within 3 days of a sulfur application.</b></p>

**Note to PM:**

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional.]

<b>Field Crops</b>
<p><b>Alfalfa*, Canola*, Cotton, Peanut*, Tobacco*, and Wheat*</b>                      * Not approved for use in California</p>
<p><b>Pests Controlled:</b> Aphids, caterpillars (in tank mix combination only), exposed thrips, greenbugs, fleahoppers, plant bugs (such as lygus and stink bugs), spider mites and whiteflies.</p>
<p><b>Use Rates:</b> Refer to the Mixing and Application section of this label for rates, timing and application information except as noted below.</p>
<p><b>Stand Alone Rate:</b> Use 4 pints of this product per acre when applications are made in 3 - 25 gallons of water per acre. Use a 2% v/v spray solution of this product when 25 or more gallons of water are applied per acre.</p>
<p><b>Tank Mixtures:</b></p> <ul style="list-style-type: none"> <li>• When used in combination with other insecticides, use 0.75 to 4 pints of this product per acre when applied in 3 - 25 gallons of water per acre. Use a 1 - 2% v/v solution of this product when greater than 25 gallons of water are applied per acre.</li> <li>• To achieve enhanced and residual control of caterpillars, use this product in combination with other labeled insecticides.</li> <li>• To achieve enhanced and residual control of other insects (aphids, thrips, greenbug, fleahopper, plant bug, spider mite, white fly), use this product in combination with other labeled insecticides.</li> <li>• In situations where thorough coverage of pests is not possible or practical, this product may be tank mixed with one of the companion pesticides listed above.</li> <li>• This product may be tank mixed with penetrator type adjuvants for application to cotton, peanuts and tobacco only if prior use under local use conditions has demonstrated plant safety.</li> </ul>

<b>Hops*, Coffee*, Banana* and Pineapple*</b> * Not approved for use in California
<p><b>Pests controlled:</b> Aphids, leafhoppers, mealybugs, spider mites, plant bugs, scales, exposed thrips and whiteflies.</p>
<p><b>Use Rates:</b>                      Stand Alone Rate: Apply a 1 - 2% v/v solution of this product, unless otherwise noted. Refer to the Mixing and Application section of this label for rates, timing and application information except as noted below.</p>
<p><b>Tank Mix Rates:</b> To achieve enhanced and residual pest control, use a 1 to 2 % v/v solution of this product in combination with a companion product.</p> <ul style="list-style-type: none"> <li>• Use a 1% v/v rate with most companion pesticides targeted at the same pest(s) as GH MPMT.</li> <li>• Lower rates (0.25 -1.0 % v/v solution) of this product maybe used in tank mixes for knockdown/suppression of adult whiteflies.</li> </ul> <p>See Application Guidelines and Notes to User: Understanding Plant Sensitivity for additional comments.</p>

<b>HEMP (Indoor, Greenhouse and Outdoor)</b>	
<b>Pest</b>	<b>Requirements</b>
<p>Aphids, leafhoppers, mealybugs, spider mites, plant bugs, scales, exposed thrips, whiteflies and powdery mildew.</p>	<p><b>Use Rates:</b>                      Stand Alone Rate: Apply a 1 - 2% v/v solution of this product, unless otherwise noted. Refer to the Mixing and Application section of this label for rates, timing and application information except as noted below.  <b>Tank Mix Rates:</b> To achieve enhanced and residual pest control, use a 1 to 2 % v/v solution of this product in combination with a companion product.</p> <ul style="list-style-type: none"> <li>• Use a 1% v/v rate with most companion pesticides targeted at the same pest(s) as GH MPMT.</li> <li>• Lower rates (0.25 -1.0 % v/v solution) of this product maybe used in tank mixes for knockdown/suppression of adult whiteflies.</li> </ul> <p>See Application Guidelines and Notes to User: Understanding Plant Sensitivity for additional comments.</p>

<b>Turf</b>	
<b>Pest</b>	<b>Requirements</b>
<p>Chinch bugs, cutworms, fleas, greenbugs, grubs, mites, mole crickets and sod webworms</p>	<p>Stand Alone Rate: Apply a 1 - 2% v/v solution of this product, unless otherwise noted. Refer to the Mixing and Application section of this label for rates, timing and application information, except as noted below.                      Tank Mix Rate: To achieve enhanced and residual pest control, use a 1 to 2% v/v solution of this product in combination with products such as: malathion or acephate. For use on mole crickets, irrigate turf prior to application of this product to bring mole crickets to the surface.</p>
<p>Powdery Mildew</p>	<p>Stand-alone Rate: Apply a 1 .5 - 2% v/v solution of this product when the powdery mildew is first observed and repeat every 7 to 14-days, or as needed.                      Tank Mix Rate: This product can also be tank mixed with other fungicide products suitable for turf powdery mildew use. Tank mixes with products containing chlorothalonil should be avoided as they can cause severe phytotoxicity.</p>

**Bold, italicized text is information for the reader and is not part of the label.**

[Bracketed information is optional.]

<b>Ornamental Landscape Trees and Shrubs, Christmas Trees, Flowers, Bedding and Indoor Plants</b>	
<b>Pests controlled:</b> Adelgids, ants, aphids, buck moth larvae, earwigs, elm leaf beetle larvae, sawfly larvae, gypsy moth eggs and larvae, Japanese beetles, lace bugs, leafhoppers, mealybugs, rust mites, phylloxera, spider mites, shore fly, plant bugs, psyllid, scales, tent caterpillars, exposed thrips, whitefly and powdery mildew.	
<b>Use Rates:</b> <b>Stand Alone Rate:</b> Apply a 1 -2% v/v solution of this product, unless otherwise noted. Refer to the Mixing and Application section of this label for rates, timing and application information except as noted below. <b>Tank Mix Rate:</b> To achieve enhanced and residual pest control, use a 1 to 2 % v/v solution of this product in combination with a recommended product. Tank mixes with products containing Chlorothalonil should be avoided as they can cause severe phytotoxicity. Lower rates (0.25 - 1.0 % v/v solution) of this product may be used in tank mixes for knockdown/suppression of adult whiteflies. <b>Note:</b> Unless otherwise noted, apply this product to wet (minimize run off). If wilting occurs a few hours after application, rinse with a clean water spray. Since applications may cause injury to flowers of some plant cultivars, test product on a few flowering plants under local use conditions prior to large scale use.	
<b>Pest</b>	<b>Requirements</b>
Adelgid, psyllid, scale on trees and shrubs	Spray to wet infested plant surface.
Balsam woolly adelgid	Spray trees thoroughly when nymphs are abundant in early spring and/or fall. Make a second application 10 days later.
Cooley Spruce gall adelgid	Apply a 4% v/v solution to spruce in fall or spring when females begin egg laying. On Douglas fir, control exposed "woolly aphids" in early spring immediately before bud break (April) or in the fall (Sept.-Oct.) after foliage has hardened off.
Hemlock woolly adelgid	Apply this product immediately after detecting an adelgid infestation to avoid the spread of the pest. In regions with heavy infestations and on trees with a history of infestation, a total of four applications made in April/May, June, August/September and October may be necessary. Because it is a contact insecticide, do not use this product as a preventative control measure.
Conifer and pine needle scale	When insects first appear, apply a 1 to 2% v/v spray to thoroughly wet crawlers.
Tent caterpillar	Apply a 3% v/v solution. Spray directly on young larvae when congregated on outside of tent or dismantled tents and spray to thoroughly wet larvae.
Spider mite	Wet underside of leaves and needle sheaths. Wait 7-10 days to allow viable eggs to hatch and make a second application before mites reach egg laying maturity. For residual control, apply in a tank mix combination with a labeled companion product.
Earwig	Spray directly on insects after disturbing their hiding place.
Leafminer (Dipteran)	Apply in tank mix combination with a labeled companion insecticide which has residual activity.
Whiteflies	Scout plants regularly and apply a 1 to 2% v/v solution when pests first appear. Make a second application 7-days later to target newly hatched nymphs. Make additional applications at weekly intervals, but avoid more than three consecutive weekly applications. Do not apply to poinsettia bracts. Lower rates (0.25 - 1% v/v solution) of this product may be used in tank mixes for knockdown/suppression of adult whiteflies.
Gypsy moth	To suppress gypsy moth populations, apply a 3% v/v solution directly to the larvae.
Blossom thrips on African violets	Spray to contact thrips on blooms.
Euonymous scale	Thoroughly wet crawlers with a 2% solution. Delayed dormant applications of this product (1 % v/v solution) plus summer oil (1 % v/v solution) will also be effective.
Spruce spider mite on conifers	Apply 1-2% solution of this product to thoroughly wet infested needles. Repeat application after 7-days if needed.
Powdery mildew on greenhouse grown roses	Apply a 1-2% solution of this product to control powdery mildew spores and hyphae. Initiate sprays when first signs of powdery mildew appear and repeat at 7 to 10 day intervals, or as needed. Alternate applications of this product with other fungicides to avoid making more than three successive applications of this product at intervals of 7-days or less. Roses may be treated with sublimated sulfur using sulfur-heating pots or similar devices before or after treatment with GH MPMT. To minimize the potential for plant injury, allow 24 hours to elapse between GH MPMT and sublimated sulfur applications. Mixing other forms of sulfur (sprayable or dusting) or applying this product within up to 5 days of a sulfur application may also increase the potential for plant injury, especially during hot weather (>90°F).

**Bold, italicized text is information for the reader and is not part of the label.**

[Bracketed information is optional.]

#### Storage and Disposal

Do not contaminate water, feed or foodstuff by storage or disposal.

#### For rigid nonrefillable containers 5 gallons or less:

**Pesticide Storage:** Store only in original container; in a dry place; inaccessible to children and pets. Keep container tightly sealed when not in use.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

**Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

#### For refillable rigid containers larger than 5 gallons:

**Pesticide Storage:** Store only in original container, in a dry place; inaccessible to children and pets. Keep container tightly sealed when not in use.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in sanitary landfill or by other procedures approved by state and local authorities.

#### For nonrefillable rigid containers larger than 5 gallons:

**Pesticide Storage:** Store only in original container; in a dry place; inaccessible to children and pets. Keep container tightly sealed when not in use.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

**Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank to collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

### OPTIONAL MARKETING CLAIMS

**Text separated by / denotes and/or options and bracketed information is optional**

1. For organic gardening
2. For organic production
3. OMRI Logo
4. [Questions/Comments [Call]] **insert company phone number**
5. [Visit [Our website]] hawthorne-gardening.com
6. For use on vegetables, fruits, tree nuts, flowers, foliage, hemp, trees and shrubs
7. Can be used on [the following plants:] vegetables, fruits, nuts, ornamental trees, hemp, shrubs and flowering plants
8. [GH MPMT is a] contact insecticide/miticide/fungicide for control of soft-bodied insects/mites/powdery mildew.
9. [This product may be used to] control[s] targeted pests on [crops which include] vegetables/grapes/small fruits/tree fruits/tree nuts/cotton/tobacco/ hemp/shrubs/shade/ornamental trees/turf/foliage/flowering plants [growing outdoors/in greenhouses/in interiorscapes].

**Note to PM:**

EPA Reg. No. 91865-2

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional.]

10. This product can be applied up to harvest.
11. Use to control or suppress soft bodied pests
12. This product may be used only as a foliar spray to control or suppress soft bodied pests
13. This product may be used only as a foliar spray to control or suppress soft bodied pests which include:  
adelgids/aphids/caterpillars/earwigs/lace bugs/leafhoppers/leafminers/mealybugs/mole crickets/plant bugs/psyllids/sawfly larvae/scales/spider mites/tent caterpillars/thrips/whiteflies/gypsy moth/chinch bugs.
14. GH MPMT is most effective on mites, aphids and whiteflies.
15. This product provides curative control of powdery mildew.
16. See back/side/top/bottom/booklet [panel/label] for [additional] directions for use/precautions/precautionary statements/first aid [statements].

**WARRANTY INFORMATION**

General Hydroponics warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purpose referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the disease problem, condition of the plant, incompatibility with other influencing factors in the use of this product are beyond the control of the seller. To the extent consistent with applicable law, buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herein, with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OR MERCHANTABILITY IS MADE.

**NOTICE ON CONDITIONS OF SALE**

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of General Hydroponics. To the extent consistent with applicable law, all such risks are assumed by the Buyer and User.

General Hydroponics warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. General Hydroponics MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, General Hydroponics' EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT General Hydroponics' SOLE DISCRETION.

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified. Country: United States of America.

U.S. Patent No. X,XXX,XXX

© 20XX [Licensed by] *insert company name*. [World rights reserved.]

Sold by: *insert company name and address*

Made in/Product of \_\_\_\_\_

Carzol® is registered trademarks of Gowan Company LLC

**Note to PM:**  
***Bold, italicized text is information for the reader and is not part of the label.***  
[Bracketed information is optional.]

EPA Reg. No. 91865-2

SUBLABEL B HOMEOWNER USE

# GH MPMT

INSECTICIDE – FUNGICIDE - MITICIDE  
CONCENTRATE

For foliar application [to control soft-bodied insects, mites, and powdery mildew]

**ACTIVE INGREDIENT:**

Potassium salts of fatty acids .....	49.0%
<b>OTHER INGREDIENTS: .....</b>	<b>51.0%</b>
Total .....	100.0%

1 gallon of GH MPMT weighs 8.50 lb at 68°F and contains 3.8 lb active ingredient per gallon (456 g ai/L)

**KEEP OUT OF REACH OF CHILDREN**  
**WARNING - AVISO**

EPA Reg. No. 91865-2  
EPA Est. No. XXX-XX-X (*insert EPA Registered Establishment Number(s)*)  
[Superscript used is first letter of lot code]

[Sold by:] General Hydroponics, 2877 Griffen Ave., Santa Rosa, CA 95407

NET CONTENTS XX.XX fl oz (X.XX L)

**Note to PM:**

EPA Reg. No. 91865-2

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional.]

<b>FIRST AID</b>	
If swallowed	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
If on skin or clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
If in eyes	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-465-5161 for emergency medical treatment information.	
<b>NOTE TO PHYSICIAN:</b> Probable mucosal damage may contraindicate the use of gastric lavage.	

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS & DOMESTIC ANIMALS****WARNING**

Causes substantial but temporary eye injury and skin irritation. Do not get in eyes, on skin or clothing. Wear goggles or face shield. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

**ENVIRONMENTAL HAZARDS**

This product may be hazardous to aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

**DIRECTIONS FOR USE**

It is violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

**USE INFORMATION**

**Use of this product on agricultural establishments covered by the Worker Protection Standard is prohibited.**

Apply GH MPMT to foliage as a contact insecticide, miticide and fungicide for control of soft-bodied insects, mites and powdery mildew. The formulation is based on potassium salts of naturally derived fatty acids. This product may be used to control targeted pests on crops which include vegetables, grapes and other small fruits, tree fruits, tree nuts, hemp, shrubs, shade and ornamental trees, turf, foliage and flowering plants growing outdoors, in greenhouses and in interiorscapes. This product can be applied up to harvest.

**PESTS CONTROLLED**

This product may be used only as a foliar spray to control or suppress soft bodied pests which include: adelgids, aphids, caterpillars, earwigs, lace bugs, leafhoppers, leafminers, mealybugs, mole crickets, plant bugs, psyllids, sawfly larvae, scales, spider mites, tent caterpillars, thrips, whiteflies, gypsy moth, and chinch bugs. GH MPMT is most effective on mites, aphids and whiteflies. Effects on other pests are limited to early growth stages and eggs. This product provides curative control of powdery mildew.

**MIXING AND APPLICATION****WHEN TO USE:**

- Apply as soon as pest problems are noticed. Repeat applications at one-week intervals, or as needed, to control infesting or increasing pest populations.
- Do not make repeat applications at less than 7-day intervals, or make more than 3 sequential applications at 7- to 14-day intervals unless your experience has shown the use pattern to be non-injurious to the type and the development stage of plant(s) to be treated under local growing conditions.

**BEFORE YOU USE:**

Read and follow these directions when using:

- Do not allow spray to drift from application site.
- Use only with tank sprayers, hand-held sprayer, trigger spray bottles, or hose end sprayers.
- Do not allow spray mixture to stand overnight or for prolonged periods.

**Note to PM:**

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional.]

**Mixing**

- Unless otherwise noted, do not tank mix this product with adjuvants such as penetrators, spreader stickers or activators, gibberellic acid, calcium nitrate, or diatomaceous earth, foliar nutrients, alkaline based chelating agents (such as EDTA, chlorothalonil and pesticides containing sulfur or metallic ions (such as manganese, magnesium, iron, zinc, etc.) as they may be physically incompatible and/or cause damage to plants.
- **DO NOT** mix GH MPMT with Carzol® insecticide (EPA Reg. No. 10163-265) as they are not physically compatible.
- **DO NOT** mix GH MPMT with sulfur. **Do not** use within 3 days of a sulfur application.
- **It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.**
- Combinations should be kept agitated and sprayed promptly after mixing.

**HOW TO USE FOR TANK SPRAYER, HAND-HELD SPRAYERS, AND TRIGGER SPRAYER BOTTLES**

- Fill sprayer or bottle with appropriate amount of water and GH MPMT.
- Mix the spray solution thoroughly.
- Use product promptly after mixing and shake gently periodically during use.

**HOW TO USE FOR HOSE-END SPRAYERS**

Follow hose-end sprayer specific instructions to determine how to fill, set dial, clean and disconnect from hose.

Ready-To-Spray Hose End Applicator

Shake container gently before and during use.

Connect:

- Connect/Attach Sprayer/bottle to end of hose
- Turn on water

Spray:

- Hold/Direct/Point sprayer towards the area to be treated making sure that the end of the nozzle is 8 to 12 inches away from the plants.
- Turn the sprayer to the on position
- Direct spray toward the upper and lower leaf surfaces and stems where pests appear.
- Spray until the surface of the foliage is wet

Finish:

- To stop spraying, turn the knob to the OFF position.
- Turn off water.
- To relieve water pressure, turn knob to ON position until water slows to a drip.

Turn the knob back to the OFF position.

- Disconnect the sprayer from hose.

Do not use hose end applicator indoors

**RATES:** GH MPMT rates from 0.25% v/v to 4% v/v may be used. Refer to individual sections of this label for specific use rates for control of pests on specific plants.

- Rates are provided as a % volume/volume (v/v) solution. For a 2% solution add 2.5 fl oz of concentrate to the desired container and dilute to 1 gallon with water.
- The activity of this product is directly related to the solution concentration and to contact with the pest. This product must come into contact with the targeted pests to be effective. **Tank Mix Rate:** To achieve enhanced and residual pest control, use a 1 to 2% v/v solution of this product in combination with a specified product. Always refer to specified products instructions for correct use rates of that product.

**RATE TABLE**

Total Mix Volume	% v/v Solution	Amount of GH MPMT		Tablespoons
		English	Metric	
1 gallon (3.785 liters)	0.25	0.32 fl oz	9.46 ml	0.64
	0.5	0.64 fl oz	18.93 ml	1.28
	1.0	1.28 fl oz	37.85 ml	2.56
	1.5	1.92 fl oz	56.78 ml	3.84
	2.0	2.56 fl oz	75.71 ml	5.12
5 gallons (18.93 Liters)	4.0	5.12 fl oz	151.42 ml	10.24
	0.25	1.60 fl oz	47.32 ml	-
	0.5	3.20 fl oz	94.64 ml	-
	1.0	6.40 fl oz	189.27 ml	-
	1.5	9.60 fl oz	283.91 ml	-
	2.0	12.80 fl oz	378.54 ml	-
	4.0	25.60 fl oz	757.08 ml	--

**Note to PM:**

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional.]

**Timing of Application**

- Initiate applications as soon as pest is observed, or as needed.
- Repeat applications at one-week intervals, to control infesting or increasing pest populations.

**Application Guidelines**

- Do not apply through any type of hydroponic systems. Do not use with pulse or thermal fogger applicators. Do not apply through any type of irrigation system.
- This product must come into contact with the targeted pests to be effective. Complete coverage of pests is essential for maximum control. Direct sprays to contact the pests. In general, one gallon of total spray volume will treat 870 sq ft.
- Apply GH MPMT solutions to upper and lower leaf surfaces and stems where pests appear. Spray until the surface of the foliage is wet, but minimize run-off (to decrease the potential for injury on foliage, fruit and flowers of sensitive plants.)
- Avoid applications on new transplants and uprooted cuttings.

**Notes to User: Understanding Plant Sensitivity**

- Sensitivity of plants to this product can be influenced by several factors which include: pest and disease pressure, cultivar, plant vigor, environmental conditions (temperature, relative humidity, moisture availability, light intensity, etc.), spray concentration, companion products, spray additives, pH of spray mixtures, delivery volume as well as the timing, number and frequency of applications.
- Potential for plant injury increases when this product is used under the following conditions:
  - Plants are stressed, such as under hot (>90°F), humid and/or drought conditions.
  - More than 3 sequential applications are made at 7-day intervals.
  - The pH of the final spray mixture is lowered below 8.0.
  - High volume applications cause the spray to collect on the bottom of fruit such as apple, pear, nectarine, orange and grape.
  - Tender new foliage is present on narrow leaf evergreen trees and shrubs.
  - When existing insect, mite or disease pressure has already stressed plant foliage or damaged fruit (for example, mite burn on leaves).
- Application of this product may alter the appearance of the glaucous bloom on spruces and the waxy bloom on grapes.
- Evaluate the effect of this product alone or in combination with desired companion product(s) on a few test plants prior to full scale applications when sensitivity of plant species or cultivars under local use conditions is unknown.
- Do not apply to very sensitive plants such as: horse chestnut, Japanese maple, mountain ash, Cherimoya, bleeding heart or sweetpeas.
- The following plants may be sensitive to this product: bald cypress, begonia, chrysanthemum, Crown of Thorns and other Euphorbia, cucumber, delicate ferns, narrow leaf evergreens (especially when stressed or when tender new growth is present), Dieffenbachia, fuschia, gardenia, impatiens, jade plant, lantana, ornamental ivy, palms, poinsettia, redbud, river birch, schefflera, Zebra plant and some succulents. Test first on a few plants prior to broad scale applications when sensitivity under local use conditions is unknown.
- Flowers of plants including African violet, ageratum, azalea, begonia, camellia, chrysanthemum, dahlia, geranium, gloxinia, impatiens, lily, marigold, orchid, pansy, petunia, poinsettia bracts, rose, salvia, snapdragon, vinca and zinnia may be injured when sprayed with this product.
- Safety to blooms should be determined for each individual cultivar of plant to be treated by conducting a small test under local use conditions.

**Insecticidal / Miticidal and Fungicidal Uses**

This product may be used as a foliar spray to control or suppress soft bodied pests which include: adelgid, aphid, earwig, lace bug, leafhopper, mealybug, mole cricket, plant bug, psyllid, sawfly larva, scale, spider mite, tent caterpillar, thrips, whitefly, gypsy moth, and chinch bug.

**Preharvest Interval**

GH MPMT can be used up to harvest

<b>Vegetables, Herbs And Spices (Greenhouse and Outdoor)</b>	
<b>Vegetables:</b> Artichoke*, Asparagus, Brassica (cole), (such as: bok choy, broccoli, broccoli raab, Brussels sprout, cabbage, cauliflower, Chinese broccoli, Chinese cabbage, collard greens, kale, kohlrabi, mustard greens, napa cabbage and Swiss chard), Bulb (such as: green onion, leeks, garlic and onion), Cucurbit (such as: cucumber, melon, pumpkin and summer squash), Fruiting (such as: eggplant, pepper and tomato), Leafy (such as: celery, endive, escarole, lettuce and spinach), Legume (such as: bean, pea and soybean), Root and tuber (such as: beet, carrot, horseradish, potato, radish, rutabaga, shallot, sugarbeet and yam), Herbs and spices: (such as: basil, chive, dill, marjoram, parsley and sage). Watercress* * Not approved for use in California	
<b>Use Rates:</b> Apply a 1-2% v/v solution of this product, unless otherwise noted. Refer to the Mixing and Application section of this label for rates, timing and application information except as noted below.	
<b>Pest</b>	<b>Requirements</b>
Leafminers and Green peach aphids	Apply this product only in tank mix combination with a labeled companion insecticide.
Other aphids	To achieve enhanced and residual pest control, use a 1-2% v/v solution of this

**Note to PM:**

EPA Reg. No. 91865-2

**Bold, italicized text is information for the reader and is not part of the label.**

[Bracketed information is optional.]

	product in tank mix combination with a specified rate of another labeled product. .
Whiteflies	Apply a 1-2% v/v solution of this product.
Thrips, Plant bugs, Spider mites, Broad mites, Russet mites and Leafhoppers	See Mixing and Application section of this label for information on rates, mixing and application.
Powdery mildew	<b>Cucurbits, Fruiting Vegetables, Herbs and Spices, Leafy Vegetables, Legumes, Root and Tuber Vegetables and Greenhouse Cucumber:</b> Apply a 1.5-2% v/v solution of this product when the first signs of powdery mildew appear and repeat at 7 to 10 day intervals, or as needed. <b>Tank Mix Rate:</b> Apply a 1-2% v/v solution of this product with a fungicide to achieve protectant/residual control of powdery mildew. For additional instructions, see Application Guidelines and Notes to User: Understanding Plant Sensitivity sections of this label.

<b>Fruit, Nut And Vine Crops</b>	
<p><b>Citrus fruits</b> (such as: citron, grapefruit, kumquat, lemon lime, tangelo, and tangerine), <b>Pome fruits</b> (such as: apple, crabapple, pear and quince), <b>Stone Fruits</b> (such as: apricot, cherry, nectarine, peach, plum and prune), <b>Small Fruits and Berries</b> (such as: blackberry, blueberry, cane berries, cranberry, currants, grape, raspberry, strawberry, boysenberry, and olallieberry) and <b>Tree Nuts</b> (such as: almond, chestnut, filbert, pecan, pistachio, black and English walnut) and <b>Pomegranates*</b>.</p> <p>* Not approved for use in California</p>	
<p><b>Use Rates:</b> Apply a 1-2% v/v solution of this product, unless otherwise noted. Refer to the Mixing and Application section of this label for rates, timing and application information except as noted below.</p> <ul style="list-style-type: none"> <li>• <b>For all Fruit and Nut Crops:</b> Spray until the surface of the foliage is wet, but minimize run-off to decrease the potential for fruit injury by causing collection of spray on the bottom of fruit, especially when poor drying conditions exist.</li> <li>• <b>Grapes:</b> This product may alter the waxy bloom and or cause marking on the surface of grapes which can affect the quality rating of table grapes at harvest. To reduce potential for fruit injury, do not apply in greater than 1 gallon of water per 581 sq.ft. on table grapes after berries are 6 -7 mm in diameter. <b>Do not apply this product to Calmeria or Italia varieties of grapes.</b></li> <li>• <b>Citrus (Precaution):</b> Fruit marking may occur when sooty mold is present.</li> <li>• <b>Pears:</b> <b>Do not apply</b> this product after the delayed dormant stage on fresh market pear varieties including d'Anjou and Asian pears. Applications after fruit set may cause marking which can reduce market value of the fruit.</li> <li>• <b>Stone Fruits:</b> Do not make dilute spray applications on smooth skinned fruit after fruit formation. This product is not specified for use on yellow skinned nectarine varieties. <b>Do not apply GH MPMT to sweet cherries between fruit formation and harvest.</b></li> </ul> <p>Nursery and non-bearing trees can be sprayed to runoff with no more than 3 consecutive applications with 7-day intervals.</p>	
<b>Pest</b>	<b>Requirements</b>
Citrus rust mites	Apply a 1-2% v/v solution of this product.
European red mites, Pacific mites, two spotted mites, Willamette mites	Apply a 1-2% v/v solution of this product to control motile stages.
Pear psylla	Apply a 1-2% v/v solution of this product at delayed dormant to control winterform and newly hatched nymphs. Make follow up applications on processing pears at petal fall and turn down for control of 1st and 2nd generation nymphs before they mature into fourth instars.
Rosy apple aphids	Apply a 1-2% v/v solution of this product. Time sprays to coincide with pre-pink or pink stages of apples.
White apple leafhoppers	Use a 1 -2% v/v solution of this product. Target 1st generation nymphs before they mature into flyers.
Other aphids; campylomas, mealybugs, rust mites, pear slug (sawfly larvae) scale crawlers, leafhoppers and whiteflies	See Mixing and Application section for information on rates, mixing and application.
Powdery mildew	<b>Grapes:</b> Apply a 1-2% v/v solution of this product to control powdery mildew spores and hyphae. Initiate sprays at bud break or as appropriate based on local powdery mildew control requirements. Repeat at 7 to 10 day intervals, or as needed through veraison (development of variety-specific color). Do not apply past veraison. Schedule a follow-up fungicide application within 7 to 10 days after each application when GH MPMT is used alone. Use this product in a scheduled preventive spray program with other labeled fungicides. Do not mix with sulfur. Do not use within 3 days of sulfur application. This product may be applied in tank mixes with other labeled fungicides. Consult your local extension specialist, certified crop advisor, and/or manufacturer representative for specific product recommendations. <b>Tank Mix Rate:</b> Apply a 1-2% v/v with labeled rates of companion pesticides. Rates of GH MPMT below 1.5% v/v will not be fungicidal against powdery mildew. Complete coverage of grape clusters as well as leaf surfaces is essential for effective control. For additional instructions on use with sulfur, see Application Guidelines and Notes to User: Understanding Plant Sensitivity sections of this label.
Powdery mildew	<b>Citrus fruit, Pome fruit, Stone fruit, Small fruits and Berries, and Tree Nuts:</b> Apply a 1 - 2% v/v solution of this product to control powdery mildew spores and hyphae.

**Note to PM:**

EPA Reg. No. 91865-2

**Bold, italicized text is information for the reader and is not part of the label.**

[Bracketed information is optional.]

	<p><b>Stand Alone Rate:</b> Apply a 1 - 2% v/v solution of this product when first signs of powdery mildew appear or as appropriate based on local powdery mildew requirements and repeat at 7 to 14-day intervals, or as needed. Schedule a follow-up fungicide application within 7 to 14-days after each application when this product is applied alone.</p> <p><b>Tank Mix Rate:</b> Apply a 1 - 2% v/v solution of this product with a fungicide to achieve protectant/residual control of powdery mildew. Do not mix with sulfur. Do not use within 3 days of a sulfur application.</p>
--	---

**Hops\*, Coffee\*, Banana\* and Pineapple\***

\* Not approved for use in California

**Pests controlled:** Aphids, leafhoppers, mealybugs, spider mites, plant bugs, scales, exposed thrips and whiteflies.**Use Rates:**

Stand Alone Rate: Apply a 1 - 2% v/v solution of this product, unless otherwise noted. Refer to the Mixing and Application section of this label for rates, timing and application information except as noted below.

**Tank Mix Rates:** To achieve enhanced and residual pest control, use a 1 to 2 % v/v solution of this product in combination with a companion product.

• Use a 1% v/v rate with most companion pesticides targeted at the same pest(s) as GH MPMT.

• Lower rates (0.25 -1.0 % v/v solution) of this product maybe used in tank mixes for knockdown/suppression of adult whiteflies.

See Application Guidelines and Notes to User: Understanding Plant Sensitivity for additional comments.

**Ornamental Landscape Trees and Shrubs, Christmas Trees, Flowers, Bedding and Indoor Plants****Pests controlled:** Adelgids, ants, aphids, buck moth larvae, earwigs, elm leaf beetle larvae, sawfly larvae, gypsy moth eggs and larvae, Japanese beetles, lace bugs, leafhoppers, mealybugs, rust mites, phylloxera, spider mites, shore fly, plant bugs, psyllid, scales, tent caterpillars, exposed thrips, whitefly and powdery mildew.**Use Rates:**

Apply a 1 -2% v/v solution of this product, unless otherwise noted. Refer to the Mixing and Application section of this label for rates, timing and application information except as noted below.

**Tank Mix Rate:** To achieve enhanced and residual pest control, use a 1 to 2 % v/v solution of this product in combination with a recommended product. Tank mixes with products containing Chlorothalonil should be avoided as they can cause severe phytotoxicity. Lower rates (0.25 - 1 .0 % v/v solution) of this product may be used in tank mixes for knockdown/suppression of adult whiteflies.**Note:** Spray until the surface of the foliage is wet, but minimize run-off wilting occurs a few hours after application, rinse with a clean water spray. Since applications may cause injury to flowers of some plant species , test product on a few flowering plants under local use conditions prior to large scale use.

<b>Pest</b>	<b>Requirements</b>
Adelgid, psyllid, scale on trees and shrubs	Spray until the surface of the foliage is wet but minimize run off.
Balsam woolly adelgid	Spray trees thoroughly when nymphs are abundant in early spring and/or fall. Make a second application 10 days later.
Cooley Spruce gall adelgid	Apply a 4% v/v solution to spruce in fall or spring when females begin egg laying. On Douglas fir, control exposed "woolly aphids" in early spring immediately before bud break (April) or in the fall (Sept.-Oct.) after foliage has hardened off.
Hemlock woolly adelgid	Apply this product immediately after detecting an adelgid infestation to avoid the spread of the pest. In regions with heavy infestations and on trees with a history of infestation, a total of four applications made in April/May, June, August/September and October may be necessary. Because it is a contact insecticide, do not use this product as a preventative control measure.
Conifer and pine needle scale	When insects first appear, apply a 1 to 2% v/v spray to thoroughly wet crawlers.
Tent caterpillar	Apply a 3% v/v solution. Spray directly on young larvae when congregated on outside of tent or dismantle tents and spray to thoroughly wet larvae.
Spider mite	Wet underside of leaves and needle sheaths. Wait 7-10 days to allow viable eggs to hatch and make a second application before mites reach egg laying maturity.
Earwig	Spray directly on insects after disturbing their hiding place.
Leafminer (Dipteran)	Apply in tank mix combination with a labeled companion insecticide which has residual activity.
Whiteflies	Scout plants regularly and apply a 1 to 2% v/v solution when pests first appear. Make a second application 7-days later to target newly hatched nymphs. Make additional applications at weekly intervals, but avoid more than three consecutive weekly applications. Do not apply to poinsettia bracts. Lower rates (0.25 - 1% v/v solution) of this product may be used in tank mixes for knockdown/suppression of adult whiteflies.
Gypsy moth	To suppress gypsy moth populations, apply a 3% v/v solution directly to the larvae.
Blossom thrips on African violets	Spray to contact thrips on blooms.
Euonymous scale	Thoroughly wet crawlers with a 2% solution.
Spruce spider mite on conifers	Apply 1-2% solution of this product to thoroughly wet infested needles. Repeat application after 7-days if needed.

**Note to PM:**

EPA Reg. No. 91865-2

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional.]

Powdery mildew on greenhouse grown roses	Apply a 1-2% solution of this product to control powdery mildew spores and hyphae. Initiate sprays when first signs of powdery mildew appear and repeat at 7 to 10 day intervals, or as needed. Avoid making more than three successive applications of this product at intervals of 7-days or less.
--	--

<b>Hemp (Indoor, Greenhouse and Outdoor)</b>	
<b>Pest</b>	<b>Requirements</b>
Aphids, leafhoppers, mealybugs, spider mites, plant bugs, scales, exposed thrips, whiteflies and powdery mildew.	<p><b>Use Rates:</b> Stand Alone Rate: Apply a 1 - 2% v/v solution of this product, unless otherwise noted. Refer to the Mixing and Application section of this label for rates, timing and application information except as noted below.</p> <p><b>Tank Mix Rates:</b> To achieve enhanced and residual pest control, use a 1 to 2 % v/v solution of this product in combination with a companion product.</p> <ul style="list-style-type: none"> <li>• Use a 1% v/v rate with most companion pesticides targeted at the same pest(s) as GH MPMT.</li> <li>• Lower rates (0.25 -1.0 % v/v solution) of this product maybe used in tank mixes for knockdown/suppression of adult whiteflies.</li> </ul> <p>See Application Guidelines and Notes to User: Understanding Plant Sensitivity for additional comments.</p>

<b>STORAGE AND DISPOSAL</b>
<p><b>PESTICIDE STORAGE:</b> Keep from freezing. To be stored in original container and placed in areas inaccessible to children.</p>
<p><b>PESTICIDE DISPOSAL AND CONTAINER HANDLING:</b> If empty: Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.</p>

**Note to PM:**

EPA Reg. No. 91865-2

***Bold, italicized text is information for the reader and is not part of the label.***

[Bracketed information is optional.]

**OPTIONAL MARKETING CLAIMS**

***Text separated by / denotes and/or options and bracketed information is optional***

1. For organic gardening
2. OMRI Logo
3. [Questions/Comments [Call]] ***insert company phone number***
4. [Visit [Our website]] hawthorne-gardening.com
5. For use on vegetables, fruits, tree nuts, flowers, foliage, hemp, trees and shrubs
6. Can be used on [the following plants:] vegetables, fruits, nuts, ornamental trees, hemp, shrubs and flowering plants
7. [GH MPMT is a] contact insecticide/miticide/fungicide for control of soft-bodied insects/mites/powdery mildew.
8. [This product may be used to] control[s] targeted pests on [crops which include] vegetables/grapes/small fruits/tree fruits/tree nuts/ hemp/shrubs/shade/ornamental trees/foliage/flowering plants [growing outdoors/in greenhouses/in interiorscapes].
9. This product can be applied up to harvest
10. Use to control or suppress soft bodied pests
11. This product may be used only as a foliar spray to control or suppress soft bodied pests
12. This product may be used only as a foliar spray to control or suppress soft bodied pests which include:  
adelgids/aphids/caterpillars/earwigs/lace bugs/leafhoppers/leafminers/mealybugs/mole crickets/plant bugs/psyllids/sawfly larvae/scales/spider mites/tent caterpillars/thrips/whiteflies/gypsy moth/chinch bugs.
13. GH MPMT is most effective on mites, aphids and whiteflies.
14. This product provides curative control of powdery mildew.
15. See back/side/top/bottom/booklet [panel/label] for [additional] directions for use/precautions/precautionary statements/first aid [statements].

**WARRANTY INFORMATION**

General Hydroponics warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purpose referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the disease problem, condition of the plant, incompatibility with other influencing factors in the use of this product are beyond the control of the seller. To the extent consistent with applicable law, buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herein, with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OR MERCHANTABILITY IS MADE.

**NOTICE ON CONDITIONS OF SALE**

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of General Hydroponics. To the extent consistent with applicable law, all such risks are assumed by the Buyer and User.

General Hydroponics warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. General Hydroponics MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, General Hydroponics' EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT General Hydroponics' SOLE DISCRETION.

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified. Country: United States of America.

U.S. Patent No. X,XXX,XXX

© 20XX [Licensed by] ***insert company name***. [World rights reserved.]

Sold by: ***insert company name and address***


Made in/Product of \_\_\_\_\_

Carzol® is registered trademarks of Gowan Company LLC

## Supplemental Label

## GH MPMT

## For use on hemp



This supplemental labeling expires on 12/19/2022 and must not be used or distributed after this date

**KEEP OUT OF REACH OF CHILDREN****WARNING - AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

See affixed label for additional directions for use, precautionary statements and first aid.

**Directions for Use**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the full EPA approved label attached to the container must be in the possession of the user at the time of application. Read the label affixed to the container for this product before applying. Use of this product according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for the product.

**USE INFORMATION**

Apply GH MPMT to foliage as a contact insecticide, miticide and fungicide for control of soft-bodied insects, mites and powdery mildew. The formulation is based on potassium salts of naturally derived fatty acids. This product may be used to control targeted pests on hemp growing outdoors, in greenhouses and in interiorscapes. This product can be applied up to harvest.

<b>Hemp (Indoor, Greenhouse and Outdoor)</b>	
<b>Pest</b>	<b>Requirements</b>
Aphids, leafhoppers, mealybugs, spider mites, plant bugs, scales, exposed thrips, whiteflies and powdery mildew.	<b>Use Rates:</b> Stand Alone Rate: Apply a 1 - 2% v/v solution of this product, unless otherwise noted. Refer to the Mixing and Application section of the affixed label for rates, timing and application information except as noted below. <b>Tank Mix Rates:</b> To achieve enhanced and residual pest control, use a 1 to 2 % v/v solution of this product in combination with a companion product. • Use a 1% v/v rate with most companion pesticides targeted at the same pest(s) as this product. • Lower rates (0.25 -1.0 % v/v solution) of this product may be used in tank mixes for knockdown/suppression of adult whiteflies. See Application Guidelines and Notes to User: Understanding Plant Sensitivity for additional comments.

**RATE TABLE**

Total Mix Volume	% v/v Solution	Amount of GH MPMT		Tablespoons
		English	Metric	
1 gallon (3.785 liters)	0.25	0.32 fl oz	9.46 ml	0.64
	0.5	0.64 fl oz	18.93 ml	1.28
	1.0	1.28 fl oz	37.85 ml	2.56
	1.5	1.92 fl oz	56.78 ml	3.84
	2.0	2.56 fl oz	75.71 ml	5.12
	4.0	5.12 fl oz	151.42 ml	10.24
100 gallon (378.5 liters)	0.25	0.25 gal	0.95 liters	-
	0.5	0.50 gal	1.89 liters	-
	1.0	1.00 gal	3.79 liters	-
	1.5	1.50 gal	5.68 liters	-
	2.0	2.00 gal	7.57 liters	-
	4.0	4.00 gal	15.14 liters	-

EPA Reg. No. 91865-2

©2019 General Hydroponics. World Rights Reserved.

[OMRI logo]