



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

December 19, 2019

Tracey Angel
Regulatory Analyst
Hawthorne Hydroponics LLC
d/b/a General Hydroponics
2877 Giffen Avenue
Santa Rosa, CA 95407

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment – Adding Crop (Hemp) to Sublabels 1 through 4; Adding New Sublabel 5 for Homeowner Use; Adding Telephone Number to First Aide Box; and Adding Alternate Brand Names “General Hydroponics Azamax Pro Botanical Insecticide/Miticide/Nematicide” and “General Hydroponics Azamax Botanical Insecticide/Miticide/Nematicide Pro” to the Product Registration

Product Name: GH NAMT
EPA Registration Number: 91865-4
Application Date: May 07, 2019
OPP Decision Number: 551122

Dear Ms. Angel:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

The alternate brand names “General Hydroponics Azamax Pro Botanical Insecticide/Miticide/Nematicide” and “General Hydroponics Azamax Botanical Insecticide/Miticide/Nematicide Pro” have been added to or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Page 2 of 2

EPA Reg. No. 91865-4

OPP Decision No. 551122

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the U.S. Environmental Protection Agency (EPA). If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Menyon Adams by phone at (703) 347-8496 or via email at adams.menyon@epa.gov.


Sincerely,



Gina Burnett, Senior Regulatory Advisor
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)
Office of Pesticide Programs

Enclosure

Bold, italicized text is information for the reader and is not part of the label.
[Bracketed information is optional.] Text separated by / denotes and/or options.



GH NAMT

ACTIVE INGREDIENT:	% by wt.
Azadirachtin.....	1.2%
OTHER INGREDIENTS:	<u>98.8%</u>
Total	100.0%

Contains 0.0987 lb. azadirachtin per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back/side/top/bottom/booklet [panel/label] for [additional] directions for use/precautions/precautionary statements/first aid [statements].

NET CONTENTS XX.XX fl oz (X.XX L)

Bold, italicized text is information for the reader and is not part of the label.
[Bracketed information is optional.] Text separated by / denotes and/or options.

Alternate Brand Names Sublabels 1 through 4 COMMERCIAL USE:

- GENERAL HYDROPONICS AZAMAX
- GENERAL HYDROPONICS AZAMAX BOTANICAL INSECTICIDE / MITICIDE / NEMATICIDE
- GENERAL HYDROPONICS AZAMAX BOTANICAL INSECTICIDE
- GENERAL HYDROPONICS AZAMAX₁
- GENERAL HYDROPONICS AZAMAX⁴²⁰
- GENERAL HYDROPONICS AZAMAX PRO BOTANICAL INSECTICIDE / MITICIDE / NEMATICIDE
- GENERAL HYDROPONICS AZAMAX BOTANICAL INSECTICIDE / MITICIDE / NEMATICIDE PRO

Alternate Brand Names Sublabel 5 HOMEOWNER USE:

- AZAMAX BOTANICAL INSECTICIDE / MITICIDE / NEMATICIDE₁

EPA Reg. No. 91865-4
EPA Est. No. XXX-XX-X (*insert EPA Registered Establishment Number(s)*)
[Superscript used is first letter of lot code]

[Sold by:] General Hydroponics, 2877 Giffen Ave., Santa Rosa, CA 95407

Bold, italicized text is information for the reader and is not part of the label.
 [Bracketed information is optional.] Text separated by / denotes and/or options.

SUBLABEL 1

GH NAMT

AZADIRACHTIN-BASED BOTANICAL, ANTIFEEDANT, REPELLANT AND INSECT GROWTH REGULATOR

FOR USE ON GREENHOUSE AND OUTDOOR FOOD CROPS, ORNAMENTAL FLOWERS, HEMP, TREES, SHRUBS AND PLANTS

ACTIVE INGREDIENT:	% by wt.
Azadirachtin	1.2%
OTHER INGREDIENTS:	98.8%
Total	100.0%

Contains 0.0987 lb. (44.8g) azadirachtin per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back/side/top/bottom/booklet [panel/label] for [additional] directions for use/precautions/precautionary statements/first aid [statements].

FIRST AID	
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call poison control center or doctor for treatment advice.
If on skin or clothing	Take off contaminate clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. CALL THE POISON CONTROL HOTLINE 24 HOURS A DAY AT 1-877-465-5161	

[Sold by:] General Hydroponics, 2877 Giffen Ave., Santa Rosa, CA 95407

EPA Reg. No. 91865-4
 EPA Est. No. XXX-XX-X (*insert EPA Registered Establishment Number(s)*)
 [Superscript used is first letter of lot code]

NET CONTENTS XX.XX fl oz (X.XX L)

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

READ ALL DIRECTIONS BEFORE USING THIS PRODUCT

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, absorbed through skin or if inhaled. Causes moderate eye irritation. Avoid breathing vapor. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse. Wear chemical resistant gloves.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates.

For Terrestrial Uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate.

USER SAFETY RECOMMENDATIONS

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is:

- Coveralls
- Chemical resistant gloves
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in the box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

PRODUCT MODE OF ACTION

GH NAMT CONTROLS target pests on contact or by ingestion. The product acts on pests by way of repellence, anti-feeding and interference with the molting process. The buyer or user is reminded that the degree of efficacy of the product is largely dependent on weather conditions, intensity of pest population, area of application, type of pest, and physical stages of pests and crops.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

GENERAL INFORMATION

Read all directions before using this product.

Apply GH NAMT as directed to any food or non-food crop up to and including the day of harvest, at a maximum rate of 3 1/2 pints/Acre. Refer to the Use Site Section for a complete listing of crops.

MIXING

Shake well before using. Add required amount of GH NAMT to a clean spray tank with at least one-half of the water to be sprayed. Constant agitation is required, particularly with tank mixes. Agitate the mixture thoroughly and then fill the tank with remaining water and continue agitation. Thorough mixing is necessary for uniform coverage. Non-uniform mixing can cause crop injury or can result in lowered effectiveness. For tank mixes, add other components to the tank containing the GH NAMT spray mixture and agitate thoroughly. If tank mixture is allowed to sit, agitation is necessary prior to application. Adjusting the spray mixture pH between 5.5 and 7 will provide optimal performance. Always use this product promptly after mixing with water and do not let tank mix sit for any extended period.

COMPATIBILITY: GH NAMT has been found to be compatible with most commonly used pesticides and fertilizers. To avoid problems, conduct a compatibility test before using this product in a tank mix with other pesticides or with fertilizers. To test for compatibility, mix a small amount of each product, in the appropriate proportions, in a small jar.

PHYTOTOXICITY: GH NAMT has been evaluated for phytotoxicity on a wide range of crops and ornamentals. However, since testing on all varieties of all crops and ornamentals is not feasible, testing a small portion of the area to be treated for phytotoxicity is recommended before treating the entire area. Further, all possible combinations or sequences of pesticide sprays, including other fertilizers, surfactants, adjuvants and other pesticides, have not been tested, thus testing for phytotoxicity of spray mixtures is recommended. It is further recommended that spray equipment used to apply GH NAMT be thoroughly cleaned before use. The addition of spray adjuvants enhances control in some crops under ideal conditions. Addition of certain adjuvants may cause phytotoxicity therefore, the addition of crop oils and other adjuvants should be thoroughly tested before using. The addition of crop oils to spray mixtures is NOT recommended on ornamental crops. Captan, Bordeaux mixtures, and highly alkaline products cause unacceptable phytotoxicity and/or reduced effectiveness on target pests. Avoid tank mix combinations of GH NAMT plus compounds known to be incompatible with oil-based formulations to prevent phytotoxicity, "Waxy bloom" on certain crops and ornamental plants is reduced after a GH NAMT application.

APPLICATION INSTRUCTIONS

For optimal performance spray product as soon as possible when pests are expected or when pests first appear. For foliar applications, apply GH NAMT in sufficient spray volume and with adequate spray pressure to ensure complete and thorough coverage of all plant surfaces including both the top and bottom of leaves. Avoid excessive runoff. Best results are obtained following 2-3 applications made at 7-10 day intervals. When pest pressure is heavy or plant canopy is dense, use higher rates and increase spray frequency. Spraying in the morning or evening hours will provide the best results. Repeat application if rain occurs within two to three hours of spraying.

SPRAY DIRECTIONS

Apply GH NAMT as a foliar spray or a drench to soil or non-soil media to control insects. When needed, soil drenches can also be used to control soil-borne pests, including soil-borne larvae of foliar insect pests. When applying as a drench, avoid excessive leaching. GH NAMT can also be applied through sub-surface soil treatment equipment. Always follow equipment manufacturers use directions. GH NAMT may be applied using any powered or manual pesticide application equipment which includes, but is not restricted to, high volume, low volume, ultra-low volume, electrostatic, fogging and chemigation. Follow the original manufacturer's instructions when using these types of equipment.

DRENCH AND ORNAMENTAL SPRAY DIRECTIONS FOR LABELED PLANTS GROWN IN GREENHOUSES, SHADECLOTHS AND NURSERIES

When used as a soil drench, apply one pint of finished spray for each gallon of soil in the pot. For most pests apply 18-21 oz. GH NAMT per 100 gallons of water. For treatment of harder to control pests, such as Dipteran leafminers, use up to 27 ounces per 100 gallons of water. Do not exceed 57 oz. of GH NAMT per acre per application.

RATES

Use GH NAMT at 1-2 pints/acre for most pest and crop conditions. Under extremely heavy pest pressure up to 3 1/2 pints may be used. Do not use less than 5 oz. per acre of GH NAMT alone. When tank mixed with other insecticidal products, the rate of GH NAMT may be reduced by 1/2, but not less than 4 oz per acre. Use up to 2.6 oz/1000 square feet for manure and compost treatments.

GENERAL CHEMIGATION INSTRUCTIONS

Apply this product only through low pressure, drip (trickle) or sprinkler (center pivot, lateral move, end tow, side (wheel) roll, traveler, solid set or hand move) irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

Dilute GH NAMT with water before introduction into the system. Use the diluted mixture within 8 hours. Do not apply in irrigation water if the pH exceeds 7.0. The optimum pH range for application is 5.5 - 6.5. If needed, the pH of the irrigation water can be adjusted by use of a suitable buffering agent. Agitation is necessary. Apply at the rate indicated in APPLICATION INSTRUCTIONS using sufficient water to achieve an even distribution.

FOR CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection:
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

In addition, all directions and requirements specified for Sprinkler Irrigation Systems must be followed.

Sprinkler Irrigation Systems

The system must contain a functional check valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Center pivot, motorized lateral move, or traveling gun types of equipment: Inject into the system for one revolution or run. Shut off injection equipment after one revolution or run, but continue to operate irrigation system until GH NAMT has been cleared from the last sprinkler head. Do not use end guns. The system should be run at maximum speed for a foliar application.

Wheel move, side roll, end tow, solid set, or hand move types of equipment: Adjust equipment to inject GH NAMT over a 30-60 minute period. Shut off injection equipment. Continue to operate irrigation system until GH NAMT has been cleared from the last sprinkler head. GH NAMT can be injected at the end of the irrigation cycle or as a separate application. Do not use end guns. GH NAMT must be premixed in a supply tank with water and other appropriate tank-mix chemicals. Agitation is necessary at all times.

Attention must be exercised in irrigation waters with a pH greater than 7. If the irrigation cycle will last longer than 8 hours and the GH NAMT is premixed in the supply tank, the tank mix must be buffered to a pH of 6 or lower. Please contact your

Company sales representative should this situation apply. Application is to be made in sufficient water and of sufficient duration to apply the appropriate rate evenly over the entire treated area.

No field runoff can be permitted during chemigation.

USE SITES

AGRICULTURAL USE SITES - Use GH NAMT on agricultural use sites such as the following:

BERRIES Group, such as: Blackberry, Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Loganberry, Raspberry (black and red) Note: For Strawberries - see other crops.

BULB VEGETABLES, such as: Garlic, Leek, Onion (dry bulb, green and Welch), Shallot

CEREAL GRAINS AND GRAINS GROUP, such as: Barley, Buckwheat, Corn, Millet (pearl and Proso), Oats, Popcorn, Rice, Rye, Sorghum (milo), Teosinte, Triticale, Wheat, Wild rice

CITRUS FRUIT, such as: Calamondin, Citrus citron, Citrus hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (sour and sweet), Pummelo, Satsuma mandarin, White Sapote, Uniq Fruit

COTTON AND TOBACCO

CUCURBIT VEGETABLES, such as: Chayote, Chinese waxgourd, Citron melon, Cucumber, Gherkin, Gourd (edible), Muskmelon, Pumpkin, Squash (summer and winter), Watermelon

FORAGE CROPS, such as: Alfalfa, Clover, Trefoil or Vetch

FRUITING VEGETABLES, such as: Eggplant, Groundcherry, Pepino, Pepper (including bell pepper, chili pepper, cooking pepper,

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

pimiento, sweet pepper), Tomatillo, Tomato

HERBS AND SPICES GROUP, such as: Allspice, Angelica, Anise (anise seed and star), Annatto (seed), Balm (lemon balm), Basil, Borage, Burnet, Chamomile, Caper buds, Caraway. Caraway (black), Cardamom, Cassia bark, Cassia buds, Catnip, Celery seed, Chervil (dried), Chive, Chinese Chive, Cinnamon, Clary, Clove buds, Coriander (cilantro or Chinese parsley - leaf), Coriander (cilantro-seed), Costmary, Culantro (leaf and seed), Cumin, Curry (leaf), Dill (Dillweed and seed), Fennel (common, Florence), Fenugreek, Grains of paradise, Horehound, Hyssop, Juniper berry, Lavender, Lemongrass, Lovage (leaf and seed), Mace, Marigold, Marjoram, Mustard (seed), Nasturtium, Nutmeg, Parsley (dried), Pennyroyal, Pepper (black and white), Poppy (seed), Rosemary, Rue, Saffron, Sage, Savory (summer and winter), Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff, Wormwood

LEGUME VEGETABLES (Succulent or Dried), such as: Bean, Broad Bean, Chickpea, Guar, Jackbean, Lablab bean, Lentil, Pea, Pigeon Pea, Soybean, Sword bean

LEAFY AND BRASSICA(COLE) VEGETABLES, such as: Amaranth, Arugula, Broccoli, Broccoli raab (rapini), Brussels Sprouts, Cabbage, Cauliflower, Cardoon, Cavalo broccolo, Celery, Chinese Broccoli (gai ion), Chinese Cabbage (bok choy, Napa), Chinese mustard Cabbage (gai choy), Chinese Celery, Celtuce, Chervil, Chrysanthemum (edible-leaved, Garland), Collards, Corn salad, Cress (garden, upland), Dandelion, Dock (sorrel), Endive (escarole), Fennel (florence), Kale, Kohlrabi, Lettuce (head and leaf), Mizuna, Mustard Greens, Mustard Spinach, Orach, Parsley, Purslane (garden, winter), Radicchio (red chicory), Rape Greens, Rhubarb, Spinach, Spinach (New Zealand, vine), Swiss Chard, Turnip Greens

OTHER CROPS, such as: Asparagus, Avocado, Banana, Coffee, Cocoa, Cranberry, Fig, Globe artichoke. Grape, Hemp, Hops, Kiwifruit, Mango, Mushroom, Okra, Olives, Papaya, Pawpaw, Peanut, Persimmon, Pineapple, Pomegranate, Strawberry, Tea, Water chestnut, Watercress, and all other food crops

POME FRUITS GROUP, such as: Apple, Crabapple, Loquat, Mayhaw, Quince, Oriental Pear, or Pear: Comice Varieties: DO NOT apply more than 24 fl oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale usage.

ROOT AND TUBER VEGETABLE GROUP, such as: Arracacha, Arrowroot, Artichoke (Jerusalem, Chinese), Beet (garden, sugar), Burdock (edible), Canna (edible), Carrot, Cassava (bitter and sweet), Celeriac (celery root), Chayote (root), Chervil (turnip-rooted), Chicory, Chufa, Dasheen (taro), Ginger, Ginseng, Horseradish, Leren, Oriental Radish (daikon), Parsley (turnip-rooted), Parsnip, Potato, Radish, Rutabaga, Salsify (oyster plant, black, Spanish), Skirret, Sweet potato, Tanier, Turmeric, Turnip, Yam bean (jicama, manioc pea), Yam (true)

STONE FRUIT GROUP, such as: Apricot, Cherry (sweet and tart), Nectarine, Peach, Plum (Chickasaw, Damson, Japanese), Plumcot, Prune

TREE AND NUT GROUP, such as: Almond, Beech nut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory nut, Macadamia nut (bush nut), Pecan, Walnut (black and English), Pistachios

TROPICAL FRUITS, such as: Papaya, Black Sapote, Canistel, Mamey Sapote, Mango, Sapodilla, Star Apple, Guava, Feijoa, Jaboticaba, Wax Jambu, Star Fruit, Passion Fruit, Acerola, Lychee, Longan, Spanish Lime, Rambutan, Pulasan, Sugar Apple, Atemoya, Custard Apple, Cherimoya, Ilama, Soursop, and Biriba

ORNAMENTAL USE SITES – GH NAMT may be used on Ornamental Use sites, such as:

ORNAMENTAL SHRUBS AND PLANTS, such as: Amaranthus, Aster, Azalea, Ferns, Fuchsias, Caladium, Carnation, Chrysanthemum, Dahlia, Daisy, Lilies, Ivy, Ficus. Gardenia, Impatiens, Iris, Jasmine, Lilac, Marigold, Philodendron, Poinsettia, Rose, Zinnia

ORNAMENTAL TREES, such as: Ash, Birch, Cedar, Cyprus, Dogwood, Fir, Elm, Juniper, Maple, Oak, Pine, Spruce.

CHRISTMAS TREES AND CHRISTMAS TREE PLANTATIONS

NON-CROP USE SITES – Use GH NAMT on non-crop use sites, such as:

UNCULTIVATED AGRICULTURAL AREAS, such as: farm yards, fuel storage areas, fence rows, rights-of-way, fallow land; soil bank land, barrier strips.

GERNERAL SOIL TREATMENTS, such as: Manure, Composts, Cull piles, Mulches, soil application with no mention of crops to be grown (potting soil, tops soil)

PESTS

GH NAMT may be used against the following pests:

Aphids (such as pea aphid, Rosy Apple Aphid), Beetles (such as Japanese beetle), Borers, (such as peachtree borers, peach twig borers), True Bugs, (such as Lygus bugs, stink bugs), Caterpillars, (such as leafrollers, cutworms, loopers, armyworms), Flies (such as walnut husk fly, leafminers and fungus gnats), Leafhoppers, Leafminers, Whiteflies, Mealy Bugs, Mites, Psyllids (such as pear psylla), Weevils, Scales (such as San Jose scale), Thrips, (such as western flower thrips).

Bold, italicized text is information for the reader and is not part of the label.
[Bracketed information is optional.] Text separated by / denotes and/or options.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal

Pesticide Storage: Store in original container in a dry, cool, well ventilated area.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill container 1/4 full with water and recap. Shake 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times.

Bold, italicized text is information for the reader and is not part of the label.
 [Bracketed information is optional.] Text separated by / denotes and/or options.

OPTIONAL TEXT AND CLAIMS

GENERAL CLAIMS

1. For organic gardening
2. OMRI Logo
3. [Questions/Comments [Call]] ***insert company phone number***
4. [Visit [our website]] ***insert company website***

USE SITE CLAIMS

1. For use on outdoor and greenhouse food and non-food crops, hemp, ornamental flowers, trees, shrubs and plants, landscapes
2. For use on roses, vegetables, fruits, nuts, flowers, foliage, hemp, trees, and shrubs
3. Can be used on [the following plants:] vegetables; fruits; herbs; nuts; ornamental trees, hemp, shrubs and flowering plants

EFFICACY CLAIMS

1. Insect repellent/insecticide
2. Miticide
3. Kills listed insects such as mites, thrips, psyllids, leafhoppers, white flies, borers, mealybugs and true bugs

U.S. Patent No. X,XXX,XXX

© 20XX [Licensed by] ***insert company name***. [World rights reserved.]

Sold by: ***insert company name and address***

Made in/Product of _____

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK, OR FIRE), CALL CHEMTREC® (800) 424-9300.
 For other product information, contact General Hydroponics or see Material Safety Data Sheet.

NOTICE ON CONDITIONS OF SALE

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.
 Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of General Hydroponics. To the extent consistent with applicable law, all such risks are assumed by the Buyer and User.

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified. Country: United States of America.

Bold, italicized text is information for the reader and is not part of the label.
 [Bracketed information is optional.] Text separated by / denotes and/or options.

SUBLABEL 2

GH NAMT

BOTANICAL

- BOTANICAL INSECTICIDE, REPELLANT, ANTIFEEDANT AND INSECT GROWTH REGULATOR (IGR)
- BOTANICAL PRODUCT FOR CONTROL OF INSECTS ON INDOOR AND OUTDOOR TREES, SHRUBS, FLOWERS, FRUIT AND NUT TREES, HEMP, GARDEN VEGETABLES AND PLANTS.
- INDOOR AND OUTDOOR VEGETABLES, ORNAMENTAL FLOWERS, TREES, HEMP, TURFGRASS, SHRUBS AND PLANTS, INCLUDING PLANTS GROWN IN CONTAINERS, INTERIORSCAPES, HOME AND GARDEN USE.

ACTIVE INGREDIENT:	% by wt.
Azadirachtin.....	1.2%
OTHER INGREDIENTS:98.8%
Total	100.0%

Contains 0.0987 lb. (44.8g) azadirachtin per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back/side/top/bottom/booklet [panel/label] for [additional] directions for use/precautions/precautionary statements/first aid [statements].

FIRST AID	
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call poison control center or doctor for treatment advice.
If on skin or clothing	Take off contaminate clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. CALL THE POISON CONTROL HOTLINE 24 HOURS A DAY AT 1-877-465-5161	

[Sold by:] General Hydroponics, 2877 Giffen Ave., Santa Rosa, CA 95407

EPA Reg. No. 91865-4
 EPA Est. No. XXX-XX-X (*insert EPA Registered Establishment Number(s)*)
 [Superscript used is first letter of lot code]

NET CONTENTS XX.XX fl oz (X.XX L)

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin or if inhaled. Avoid breathing vapor. Causes moderate eye irritation. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing and wash before reuse. Wear chemical resistant gloves.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates.

For Terrestrial Uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170.

This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water is

- Coveralls
- Chemical resistant gloves
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in the box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

This is an end use product. General Hydroponics does not intend that this product be reformulated or repackaged except under a toll repackaging agreement.

PRODUCT DESCRIPTION:

GH NAMT is a botanical product for control of insects on indoor and outdoor plants including ornamental trees, shrubs, flowers, garden vegetables, turfgrass, hemp, fruit trees and nut trees.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

When used as a component of an Integrated Pest Management (IPM) program, GH NAMT provides an effective resistance management tool.

MODE OF ACTION

GH NAMT controls target pests on contact or by ingestion. The product acts on pests by way of repellence, anti-feedance, and interference with the molting process.

Azadirachtin, an insect growth regulator (IGR), mimics the pests' hormones and disrupts distinct stages of growth and development of insects and mites. The primary mode of action of azadirachtin is an interference with synthesis and metabolism of ecdysone and the juvenile hormone. Ecdysone is the molting hormone of insects, and azadirachtin can regulate growth leading to death before or during molting.

INDOOR AND OUTDOOR ORNAMENTAL TREES, SHRUBS; HEMP, FLOWERS, AND PLANTS ESTABLISHED IN RESIDENTIAL, LANDSCAPE PLANTINGS AROUND INSTITUTIONAL, PUBLIC, COMMERCIAL AND INDUSTRIAL BUILDINGS, PARKS, RECREATIONAL AREAS, GREENHOUSES, SHADECLOTHS, NURSERIES, AND ATHLETIC FIELDS.

GH NAMT has been evaluated for phytotoxicity on a wide range of ornamentals and crops. However, since testing on all plant varieties is not feasible, test a small portion of the area to be treated for phytotoxicity before treating the entire area. All possible combinations or sequences of pesticide sprays, including other fertilizers, surfactants, adjuvants and other pesticides have not been tested. Thus, testing for phytotoxicity of spray mixtures is recommended.

The professional user assumes the responsibility for determining the level of tolerance of treated plants to GH NAMT when applied alone or in tank-mix combinations under commercial growing conditions.

Waxy bloom on certain ornamental plants is reduced after a GH NAMT application.

Applications of GH NAMT will remove the glaucous 'blue' coloring from evergreens such as Colorado blue spruce and Koster spruce.

Use GH NAMT on the following plants:

Ornamental Plants and Flowers	Actinopteris, African violets*, ageratum, aglaonema, Algerian ivy, allamanda, alocaisia, amaranthus, anthurium, aphelandra, arborvitae, Artemisia, aster, aucuba ilex, azalea, baby's breath, begonia, Boston fern, bougainvillea, boxwood, brachycome, cacti, calabrese, caladium, calathea, calendula, calla, camellia, carnation, ceanothus, chrysanthemum, cineraria, coleus, columbine, cotoneaster, cyclamen, daffodil, dahlia, daisy, daylily, delphinium, dianthus, dieffenbachia, dogwood, dusty miller, Easter lily, English ivy, euphorbia, fern, ficus, foliage plants, foxglove, freesia, fuschia, gaillardia, gardenia, geranium, gerbera, gladiola, gloxinia, gypsophilla, hederia, hibiscus, hyacinth, hydrangea, ilex, impatiens, iris, ivy, . jasmine, lilac, lily, maidenhair fern, mandevilla, marigold, narcissus, nasturtium, orchid*, pansy, pelargonium, peony, peperomia, petunia, philodendron, phlox, photinia, pinks, pittosporum, poinsettia*, pothos, portulaca, primrose, pyracantha, rhododendron, rose*, rosemary, rubber plant, salvia, schefflera, sedum, sempervivum, snapdragon, spathiphyllum, stock, syngonium, tulip, verbena, vinca, wandering jew, yucca, zinnia *Please note that when making applications to these species, spotting of plant foliage is possible.
Ornamental Trees and Shrubs	Andromeda, arborvitae, ash, Austrian pine, azalea, beech, birch, birdsnest spruce, blue spruce, bougainvillea, boxwood, butternut, cedar, charmaecyparis, cherry, cotoneaster, crabapple, cyprus, dogwood, Douglas fir, elm, euonymus, firethorn, forsythia, hackberry, hawthorn, hemlock, hickory, holly, honey locust, horse chestnut, juniper, larch, laurel, lilac, linden, London planetree, magnolia, mandevilla, maple, mimosa, mountain ash, myrtle, oak, pachysandra, peach, photinia, pine, planetree, poplar, privet, purpleleaf wintercreeper, quince, sage, spruce, sycamore, white cedar, white pine, yew

PESTS CONTROLLED OR SUPPRESSED

Use GH NAMT against the following pests presented in Table 1.

TABLE 1. TARGET PEST SPECIES OF GH NAMT.

HEMIPTERA AND HOMOPTERA	true bugs including boxelder bugs, chinch bugs, lygus bugs and stink bug; lacebugs; leafhoppers including grape leafhopper, spittlebug, potato leafhopper and variegated leafhopper; mealy bugs including apple mealy bugs, citrus mealy bugs, grape mealy bugs; whiteflies including greenhouse whitefly, silverleaf whitefly and sweet potato whitefly and woolly whitefly; aphids including apple aphid, green peach aphid, melon aphid, pea aphid, potato aphid and rose aphid; psyllids including pear psyllids and scales including black scale, brown soft scale, California red scale, coffee scale, olive scale, San Jose scale, and cottony cushion scale.
-------------------------	---

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

LEPIDOPTERA	<p>Moths including European pine shoot moth, pine tip moth and Tussock moth; leafrollers including blueberry leafroller, filbert leafroller, fruitree leafroller, citrus leafminers, grape leafroller, oblique banded leafroller, omnivorous leafroller; Cutworms including black cutworm and citrus cutworm; Caterpillars and loopers including bagworms, budworms, cabbage looper, canker worms, case bearers, caseworms, corn earworm, diamondback moth, fruit worms, grapeleaf skeletonizer, gypsy moth, hornworms, imported cabbageworm, navel orangeworm, soybean looper, spruce budworm, tent caterpillar, tip moths, tent caterpillars, tobacco budworm, tobacco hornworm, tomato pinworm and tussock moth; Armyworms including beet armyworm, fall armyworm, lawn armyworm, southern armyworm and yellow striped armyworm; webworms and leaf perforators.</p>
COLEOPTERA	<p>beetles, grubs and weevils including Asian long-horned beetle, bark beetles, black vine weevil, Colorado potato beetle, elm bark beetle, European chafer, flea beetles, Japanese beetle, June beetle, leaf beetles, Mexican bean beetle, Northern masked chafer, rose chafer and Southern masked chafer and twig girders.</p>
DIPTERA*	<p>Flies including Caribbean fruit fly, cherry maggots, crane fly, fungus gnat, Hessian fly, oriental fruit fly, Mediterranean fruit fly, marsh crane flies, melon fly, shore fly and walnut husk fly; leafminers including citrus leafminers and serpentine leafminers.</p> <p>*Not intended for use on public health pests</p>
THYSANOPTERA	<p>thrips including citrus thrips, flower thrips, gladiolus thrips, onion thrips, thrips palmi and Western flower thrips.</p>
ACARINA*	<p>mites including, red spider mites, brown mite, clover mite, conifer spider mite, European red mite, spruce spider mite, and two-spotted spider mite.</p> <p>*Not intended for use on public health pests</p>
ORTHOPTERA	<p>crickets; grasshoppers; locusts</p>
HYMENOPTERA*	<p>sawflies including European sawflies, pear sawflies, red-headed pine sawflies, yellow-headed pin sawflies.</p> <p>*Not intended for use on public health pests</p>
NEMATODA	<p>nematodes (suppression)</p>

SPRAY PREPARATION

GH NAMT is an emulsifiable concentrate to be diluted with water.

Water as diluent:

Add one-half the required amount of water to the spray tank, then add GH NAMT slowly with agitation, and complete filling the tank with water. To prevent separation of the emulsion, mix thoroughly and continue agitation while spraying.

This product forms an emulsion and can separate upon extended or prolonged standing. Re-agitate to assure uniformity of the spray mixture.

Adjusting the mixture pH to between 5 and 7 will provide optimal performance. Do not use tank additives that alter the pH of the spray solution above pH 7. Buffer the spray solution to alter the pH range as appropriate.

Prepare only the volume needed for the intended application, and use the spray mixture within 24 hours of preparation.

TANK MIXTURES

GH NAMT is an emulsifiable concentrate and is compatible with commonly used pesticides and fertilizers. Always check the physical compatibility using a jar test in the correct proportions if needed.

If a broader spectrum of control is required tank-mix GH NAMT with insecticides or miticides. If a rapid knockdown of heavy populations is necessary, then include an effective contact insecticide/miticide in combination with GH NAMT.

Tank mixtures are for use only in states where the companion product(s) and the application site are registered.

Always read and follow the directions for use, precautions and limitations for use on all product labels used in combination. Applications must follow the precautions and limitations of the most restrictive product label in the mixture. Do not exceed the dosage rates of any product.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

Select the right companion products:

IPM uses a variety of control options including biological, chemical, and cultural practices. Azadirachtin is botanical with growth regulator effect on insects and mites. Companion products include pyrethroids, spinosyns, microbial toxins, and chloronicotinyls that can complement azadirachtin. Formulations of bifenthrin, spinosad, abamectins, and imidacloprid are effective for different pests. Select the product that has been proven to provide adequate performance for the pests you are trying to control.

Physical Incompatibility

Do not use GH NAMT with Captan, Bordeaux mixture, triphenyltin hydroxide, lime sulfur, Rayplex iron or other highly alkaline materials as they can cause phytotoxicity and/or reduced efficacy on some target pests. Tank-mix combinations with compounds known to be incompatible with oil-based formulations are not to be used or phytotoxicity will occur.

ADJUVANTS

The addition of adjuvants may enhance control under certain conditions; the use of adjuvants or oils will cause phytotoxicity and should be thoroughly tested prior to use. Do not add crop oils to spray mixtures intended for use on ornamental plants, flowers, trees, and shrubs.

APPLICATION EQUIPMENT

Ground Equipment

Apply GH NAMT with hand-operated (manual) or power spray equipment suitable for low volume and/or high volume applications. Follow the recommendations of the equipment manufacturer when using backpack sprayers; hose-end sprayers, compression (pump-up) sprayers, and other sprayers suitable for foliar applications of insecticides.

Chemigation and Subsurface Equipment

GH NAMT may also be applied through chemigation systems and sub-soil treatment equipment; always follow equipment manufacturer's directions.

APPLICATION SCHEDULE

For the most effective control, apply the product when pests are expected to appear or as soon as possible after pests appear and are in immature stages. Spray at an interval of seven (7) to ten (10) days or as the situation warrants.

During high pest infestation levels or when canopy is dense use higher dosage (use) rates and increase the spray frequency. Spraying in the morning or evening hours is recommended. Repeat spraying if rain occurs within two to three hours of spraying. For additional guidance, consult with the state agricultural experiment station or local extension horticulturalist/arborist for information on tactics and windows of application.

APPLICATION RATES

Use GH NAMT on ornamental pests as a spray concentration of 0.25 - 1.70% vol/vol per treatment with high volume applications in Table 2.

The application rates are specified as rate ranges depending upon the pest infestations:

Lower rate ranges with a spray concentration of 0.25 - 0.75% vol/vol: Use lower rate ranges for light infestations of lepidopterous insects, at the first sign or at the first observation of the early and uniform growth stages of the pest(s), and/or tank mixtures with contact insecticides.

Medium rate ranges with a spray concentration of 0.75 - 1.25% vol/vol: Use medium rate ranges for moderate infestations, when multiple growth stages of the pests are present, and/or heterogeneous pesticide populations are present.

Upper rate ranges with a spray concentration of 1.25 - 1.70% vol/vol: Use upper rate ranges for moderate to heavy pest populations of difficult to control pest species, for the late stages of larva/worms, for dense foliage, and/or when re-infestations occur.

High Volume Applications:

Apply GH NAMT at spray concentration of 0.25 - 1.70% v/v in sufficient amounts of water to achieve complete coverage. Use an adequate spray volume to wet the leaves (foliage) and stems. Spray volumes will vary with the plant size. Attempt to penetrate dense foliage. Thorough coverage of the upper and lower leaf surfaces is critical for effective levels of control.

Refer to Table 3 for the amounts of GH NAMT required to prepare spray concentrations of 0.25% to 1.70% for spray volumes of 1 gallon to 200 gallons.

Specialized Low Volume Applications:

Select a spray volume to achieve sufficient coverage. Uniform coverage of both upper and lower leaf surfaces is critical for effective insect control.

Apply GH NAMT in a minimum spray volume of 5 gallons per acre. Larger plants will require the higher spray volumes (20 - 25 gallons per acre) to obtain sufficient coverage. Do not exceed 20 grams active ingredient per acre per application or 57 fl. oz. of product per acre per application. Refer to Table 4 for the amounts of GH NAMT required to prepare spray concentrations of 0.25% to 1.70% for spray volumes of 5 - 25 gallons per acre.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

TABLE 2. APPLICATION RATES FOR ORNAMENTALS ESTABLISHED IN RESIDENTIAL, LANDSCAPE PLANTINGS AROUND INSTITUTIONAL, PUBLIC, COMMERCIAL AND INDUSTRIAL BUILDINGS, PARKS, RECREATIONAL AREAS, GREENHOUSES, SHADECLOTHS, NURSERIES, AND ATHLETIC FIELDS

USE	PESTS	SPRAY CONCENTRATION %	Amounts of GH NAMT		
			Fluid ounces per gallon	Fluid ounces per 100 gallons	Quarts per 100 gallons
Including trees, shrubs, flowers, conifers, evergreens, herbaceous ornamentals, foliage plants, container-grown ornamentals, plants and groundcovers	Armyworms	Lower rate ranges of 0.25 - 0.75% vol/vol:	0.32 – 1.0 fl. oz.	32 – 96 fl. oz.	1.0 – 3.0 qts.
	Azalea caterpillars				
	Aphids				
	Bagworms	Medium rate ranges of 0.75 – 1.25% vol/vol:	1.00– 1.60 fl. oz.	96 – 160 fl. oz.	3.0 – 5.0 qts.
	Black vine weevils				
	Boxelder bugs				
	Budworms	Upper rate ranges of 1.25 – 1.70% vol/vol:	1.60 – 2.18 fl. oz.	160 – 218 fl. oz.	5.0 – 6.8 qts.
	Cankerworms				
	Cutworms				
	Eastern tent caterpillars				
	Elm Leaf beetles				
	European sawflies				
	Fall webworms				
	Flea beetles				
	Forest tent caterpillars				
	Gypsy moth larvae				
	Japanese beetles				
	June beetles				
	Lace bugs				
	Leaf-feeding caterpillars				
	Leafhoppers				
	Leafminers				
	Leaf rollers				
	Leaf skeletonizer				
	Oleander moth larvae				
	Pine sawflies				
	Pine shoot beetles				
	Pine tip moths				
	Plant bugs				
	Sawflies (larva)				
	Scale insects (crawlers)				
	Spruce bud worm				
Striped beetles					
Striped oakworms					
Thrips					
Tussock moth larvae					
Brown softscale					
California redscale (crawler)					
Clover mites					
Mealybugs					
Pineneedlescale (crawler)					
Spider mites					
Whiteflies and other species identified in Table 1.					

TABLE 3. SPRAY PREPARATION FOR HIGH VOLUME APPLICATIONS FOR SPRAY CONCENTRATIONS OF 0.25% to 1.70%.

Gallons of Water	Amounts of GH NAMT for:						
	0.25%	0.50%	0.75%	1.00%	1.25%	1.50%	1.70%
1 gallon	0.32 fl. oz.	0.64 fl. oz.	0.96 fl. oz.	1.28 fl. oz.	1.60 fl. oz.	1.94 fl. oz.	2.18 fl. oz.
5 gallons	1.60 fl. oz.	3.2 fl. oz.	4.8 fl. oz.	6.4 fl. oz.	8.0 fl. oz.	9.7 fl. oz.	10.9 fl. oz.
10 gallons	3.2 fl. oz.	6.4 fl. oz.	9.6 fl. oz.	12.8 fl. oz.	16.0 fl. oz.	19.4 fl. oz.	21.8 fl. oz.
25 gallons	8.0 fl. oz.	16.0 fl. oz.	24.0 fl. oz.	32 fl. oz.	1.25 qts.	1.50 qts.	1.70 qts.
50 gallons	16.0 fl. oz.	32.0 fl. oz.	1.50 qts.	2.0 qts.	2.5 qts.	3.0 qts.	3.4 qts.
100 gallons	1.0 qt.	2.0 qts.	3.0 qts.	4.0 qts.	5.0 qts.	6.0 qts.	6.8 qts.
150 gallons	1.5 qts.	3.0 qts.	4.5 qts.	6.0 qts.	7.5 qts.	9.0 qts.	10.2 qts.
200 gallons	2.0 qts.	4.0 qts.	6.0 qts.	8.0 qts.	10.0 qts.	12.0 qts.	13.6 qts.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

TABLE 4. SPECIALIZED SPRAY PREPARATION FOR LOW VOLUME APPLICATIONS OF 5 - 25 GALLONS PER ACRE WITH SPRAY CONCENTRATIONS OF 0.25% to 1.70%.

Spray Concentration Desired. % vol/vol	Spray Volume, Gallons Per Acre				
	5 gpa	10 gpa	15 gpa	20 gpa	25 gpa
0.25% v/v	1.6 fl. oz./acre	3.2 fl. oz./acre	4.9 fl. oz./acre	6.5 fl. oz./acre	8.0 fl. oz./acre
0.50% v/v	3.2 fl. oz./acre	6.4 fl. oz./acre	9.6 fl. oz./acre	12.8 fl. oz./acre	16.0 fl. oz./acre
0.75% v/v	4.8 fl. oz./acre	9.6 fl. oz./acre	14.4 fl. oz./acre	19.2 fl. oz./acre	24.0 fl. oz./acre
1.00% v/v	6.4 fl. oz./acre	12.8 fl. oz./acre	19.2 fl. oz./acre	25.5 fl. oz./acre	32.0 fl. oz./acre
1.25% v/v	8.0 fl. oz./acre	16.0 fl. oz./acre	24.0 fl. oz./acre	32.0 fl. oz./acre	40.0 fl. oz./acre
1.50% v/v	9.6 fl. oz./acre	19.2 fl. oz./acre	28.9 fl. oz./acre	38.5 fl. oz./acre	48.0 fl. oz./acre
1.70% v/v	10.8 fl. oz./acre	21.6 fl. oz./acre	32.5 fl. oz./acre	43.3 fl. oz./acre	54.0 fl. oz./acre

SPECIFIC USE INSTRUCTIONS:

Decision-making for IPM:

Scouting, monitoring, sampling, record-keeping, and predictive models are techniques to determine if and when insecticide/miticide applications are needed. The application schedule should coincide with the most vulnerable stage of the pest. For azadirachtin, target the most vulnerable stages of young larvae and young nymphs. The early larval stages and the early instar stages are more susceptible to this IGR than the later stages of the same pests.

For Lepidoptera:

- Armyworms: Apply when larvae are small.
- Bagworms: Apply when bags are small and larvae are actively feeding.
- Gypsy moth larvae: Apply when larvae are small and all eggs have hatched.
- Spruce budworms: Apply when larvae are exposed and actively feeding.

For Acarina:

- Spider mites: Apply when nymphs are first observed and before mite populations have become severe. Use multiple applications with 7-10 day intervals until infestation is controlled. Thorough coverage of both upper and lower leaf surfaces is needed.

For Thysanoptera:

- Thrips: Apply early at first signs of infestation and repeat until infestation is controlled.

For Hymenoptera:

- Sawfly: Apply when larvae are small. Refer to tree injection method of this label.

For Hemiptera and Homoptera:

- Leafhoppers: .Apply when first observed and repeat applications at 5 - 7 day intervals.
- Mealybugs: Obtain thorough coverage of leaves and twigs.
- Scale: Obtain thorough coverage of leaves and twigs

For Coleoptera:

- Beetles: Apply early at first signs of infestation and repeat applications at 7 - 10 day intervals.
- Japanese beetle (adults): Use foliar applications to repel adult feeding and treat at 5 - 7day intervals.

For Diptera:

- Leafminers: Apply early to larvae when stippling or mining of leaves is first observed. Repeat applications at 7 - 10 day intervals until infestation is controlled.

TURFGRASS ESTABLISHED IN RESIDENTIAL (LAWNS), INSTITUTIONAL, PUBLIC, COMMERCIAL AND INDUSTRIAL SITES, PARKS, RECREATIONAL AREAS, GOLF COURSES, SOD FARMS, AND ATHLETIC FIELDS

Use GH NAMT to control the pests presented in Table 5. Dilute GH NAMT in water.

The most vulnerable stage to this product is young larvae and nymphs. Schedule treatments for the early larval stages and early instars when populations are established, but before turf damage becomes noticeable.

The maximum rate on turfgrass of GH NAMT is 57.0 fl. oz of product per acre per application or 1.3 fl. oz product per 1,000 sq. ft. per application. Apply at a rate up to 57 fl. oz of product per acre. Use the higher rate specified on this label for moderate to heavy infestations.

Irrigation:

Avoid (delay or postpone) irrigation for 12 - 24 hours after application of this product.

Bold, italicized text is information for the reader and is not part of the label.
 [Bracketed information is optional.] Text separated by / denotes and/or options.

Mowing:

Avoid (delay or postpone) mowing of the treated area for 12 - 24 hours after treatment.

Degree day and plant phenology models can assist in developing the appropriate application schedule for the target pests. Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area.

GH NAMT can be tank mixed with other insecticide/miticides if a broader spectrum of pest control is required. Observe all precautionary statements and follow all label directions of companion product(s).

Specific Use Instructions:

1. Armyworms: Apply during the early morning or late afternoon to maximize control.
2. Sod webworm larvae: Applications in the late afternoon or early evening can maximize control.

TABLE 5. APPLICATION RATES FOR TURFGRASS ESTABLISHED IN RESIDENTIAL (LAWNS), INSTITUTIONAL, PUBLIC, COMMERCIAL AND INDUSTRIAL SITES, PARKS, RECREATIONAL AREAS, GOLF COURSES, SOD FARMS, AND ATHLETIC FIELDS.

USE	PESTS	Amount of GH NAMT		Spray Volumes		Number of Applications & Interval Days
		fl. oz./acre	fl. oz./1000 sq. ft.	gals/acre	gals./1000sq. ft.	
Cool-season and warm-season Turfgrass	Larvae and nymphs of these pests: Armyworms Bermudagrass mite Cutworms Grasshopper Sod webworm	Up to 57.0 fl. oz.	Up to 1.3 fl. oz.	40 - 100 gpa	1-2 gal/1000 sq. ft.	As needed, 7 days

DRENCH APPLICATION FOR GREENHOUSES, NURSERIES, INTERIORSCAPES AND FOR PLANTS GROWN IN CONTAINERS:

Use GH NAMT as a soil drench for effective control of soil-borne insect larvae, including soil-borne larvae of foliar pests, such as fungus gnats, nematodes, or soil borne thrips. When applying as a drench, avoid excessive leaching.

Preventive applications as a soil drench may be warranted for certain pests. Soil drench applications of azadirachtin will have a slower rate of activity because of soil absorption when compared to foliar applications of azadirachtin. Target the initial application of a soil drench treatment to coincide with the early stages of young larvae and young nymphs.

Dilute GH NAMT with water for concentrations of 0.4 to 0.8% vol/vol. Drench the soil in the pot with one (1) pint of finished spray per 1.0 gallon of soil. For fungus gnats, use the 0.4% spray concentration. For mushroom fly maggot control, use the 0.6% vol/vol spray concentration. For leafminers and other difficult to control pests, use the 0.8% vol/vol spray concentration. Two to three (2-3) applications should be scheduled at 10-14 day intervals until the pest pressure has ended.

DILUTION TABLE FOR DRENCH APPLICATIONS

Gallons of Water	Amount of GH NAMT			Application Interval	Number of Applications
	0.4%	0.6%	0.8%		
1 gallon	1 Tbs.	1.5 Tbs.	2.0 Tbs.	10 – 14 days	2 - 3
5 gallons	2.7 fl. oz	4.0 fl. oz	5.5 fl. oz	10 – 14 days	2 – 3
10 gallons	5.4 fl. oz	8.0 fl. oz	11.0 fl. oz	10 – 14 days	2 – 3
100 gallons	1.7 qts.	2.5 qts.	3.4 qts.	10 – 14 days	2 - 3

GH NAMT can also be applied through sub-surface treatment equipment. Always follow manufacturer's use directions.

TREE INJECTION

Inject GH NAMT into mature trees established in landscapes, residential settings, nurseries, and forestry sites.

Use appropriate tree injection equipment and follow the instructions provided by the equipment manufacturer.

Application Schedule For Tree Injections

Consult with your state agricultural experiment station, extension specialist, or your local U.S. Forest Service authority for information on the application schedule for specific pests in your area.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

Pests Controlled and Hosts:

PESTS	HOSTS
Spruce budworm larva	White Spruce Black Spruce Balsam Fir
Pine false webworm	Eastern White Pine Red Pine
Pine sawfly larvae	White Pine
Cedar leafminer	White Cedar

Dosage Rate For Tree Injections

Use appropriate injection equipment. Inject at the rate of 0.37 -0.74 fl. oz. (11 - 22 ml) of product per inch tree trunk diameter at breast height or inject at the rate of 0.127 - 0.25 grams azadirachtin per inch tree trunk diameter at breast height.

INSTRUCTIONS FOR GARDEN CROPS, VEGETABLES, HERBS AND SPICES, HEMP, FRUITS, AND BERRIES

- For the most effective control, spray the product as soon as possible after pests appear and are in immature stages.
- Spray at an interval of seven to ten days or as the situation warrants. During high pest infestation levels use higher label rates and increase the spray frequency.
- For best results spray in the morning or evening hours.
- Repeat spraying if rain occurs within two to three hours of spraying.

SPRAY EQUIPMENT

Use any suitable application equipment to ensure uniform coverage.

USE RATES

Apply GH NAMT as directed to any" food or non-food crop up to and including the day of harvest, at a maximum rate of 57 fl ozs (20 grams active ingredient) per acre (1.33 fl. ozs. per 1,000 sq. ft.) per application. Rates in Table 6 pertain to typical pest infestations.

Apply GH NAMT alone to food/garden crops on the day of harvest.

Dilute GH NAMT with water at 0.5 - 4.0 tablespoons (Tbs) per gallon of water. For hose end sprayers, set the RATE PER GALLON at the dial setting of 1 to 4 Tbs. depending on the crop and pests. Use the lower RATE PER GALLON for low to moderate infestations and use the higher specified RATE PER GALLON for severe infestations.

TABLE 6. USE RATES FOR GARDEN CROPS, VEGETABLES, HERBS AND SPICES, HEMP, BERRIES AND FRUIT.

CROP	PESTS such as:	Dilution Rate for Sprayers	
		fl. ozs of product per 1000 sq. ft.	Tbs. of product per 1.0 gallon of water
Leafy vegetables: Broccoli, Brussel Sprouts, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce, Spinach	Leafrollers, Cutworms, Loopers, Armyworms	0.19 – 0.96 fl. oz	¾ Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Beetles, Weevils, Flies, Thrips, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Root Vegetables: Beet, Carrot, Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yams	Beetles, Weevils	0.11 – 0.25 fl. oz	1/2 Tbs. – 1 1/2 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Thrips, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Fruiting Vegetables: Eggplant, Pepper, Tomatillo, Tomato	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Cucurbit Vegetables: Cucumber, Gourd (edible), Muskmelon, Pumpkin, Squash, Watermelon,	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

Including Cantaloupe, Casaba, Gherkins, Melons (including hybrids), Zucchini	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Legume Vegetables: Bean, Chickpea, Lentil, Pea	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Bulb Vegetables: Garlic, Onion, Shallot	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Berries: Blackberry, Blueberry, Raspberry, Strawberry, others include: Boysenberry, Currants, Dewberry, Elderberry, Gooseberry, Loganberry	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Herbs and Spices: Chive, Dill, Fennel, Mustard, Sage, Sweet bay, others include: Anise, Balm, Basil, Black pepper, Borage, Caraway, Catnip, Chamomile, Coriander, Cumin, Curry leaf, Dandelion, Fenugreek, Horehound, Hyssop, Marjoram, Marigold, Mint, Nasturtium, Pennyroyal, Peppermint, Rosemary, Savory, Spearmint, Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Other crops: Hemp	True Bugs, Leafhoppers, Whiteflies, Aphids, Mites, Beetles, Weevils, Flies, Thrips	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
	Leafrollers, Cutworms, Loopers, Armyworms	0.19 – 0.96 fl. oz	¾ Tbs. – 4 Tbs./gal
Nut Trees: Almond, Brazil nut, Filbert, Hickory nut, Pecan, Pistachios, Walnut	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Pome Fruits: Apple, Quince, or Pear (Comice varieties: DO NOT apply more than 24 fl. oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale usage.)	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Stone Fruits: Apricot, Cherry, Nectarine,	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

Peach, Plum	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Citrus Fruits: Grapefruit, Lemon, Lime, Orange, others include: Citrus Citron, Mandarin (tangerine), Nectarine, Satsuma (orange mandarin), tangerine	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal

CHEMIGATION GENERAL INFORMATION

Apply this product only through drip (trickle) or sprinkler irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues can result from non-uniform distribution of treated water. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place. Direct your questions concerning calibration to your State Extension Service Specialist, the equipment manufacturer, or other expert. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of a responsible person, shall shut the system down and make necessary adjustments should the need arise.

Dilute GH NAMT with water before introduction into the system. Use the diluted solution within 8 hours. Do not apply in irrigation water if the pH exceeds 7.0. The optimum pH range for application is 5.5 to 6.5. The pH of the irrigation water can be adjusted by use of a suitable buffering agent. Agitation is necessary. Apply at the specified rate using sufficient water to achieve an even distribution within an 8-hour period. Do not apply GH NAMT at a rate that exceeds 3.5 pints active ingredient per acre (57 fl. oz).

If the irrigation cycle will last longer than 8 hours and the GH NAMT is premixed in the supply tank, the tank mix must be buffered to a pH of 8 or lower.

For Chemigation Systems Connected to A Public Water System: Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of a year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), backflow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction.

There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top of overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speeds favor drift beyond the area intended for treatment.

Operation Of Sprinkler Chemigation Or Drip (Trickle) Utilizing A Pressurized Water And Pesticide Injection System: The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering .pump,

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

STORAGE AND DISPOSAL
Do not contaminate water, food, or feed by storage or disposal
Pesticide Storage: Store in original container in a dry, cool, well ventilated area.
Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.
Container Disposal: Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill container 1/4 full with water and recap. Shake 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times.

Bold, italicized text is information for the reader and is not part of the label.
 [Bracketed information is optional.] Text separated by / denotes and/or options.

OPTIONAL TEXT AND CLAIMS

GENERAL CLAIMS

1. For organic gardening
2. OMRI Logo
3. Organic Materials Review Institute (OMRI) Certified.
4. Low-odor formulation
5. Low (Mild) odor
6. [Questions/Comments [Call]] ***insert company phone number***
7. [Visit [our website]] ***insert company website***

USE SITE CLAIMS

1. Can be applied the day of harvest
2. For use on a wide variety of trees, shrubs, flowers, fruit and nut trees, hemp, garden vegetables and plants
3. For use on outdoor and greenhouse food and non-food crops, ornamental flowers, trees, hemp, shrubs and plants, landscapes
4. For use on roses, vegetables, fruits, nuts, flowers, houseplants, foliage, hemp, trees, and shrubs
5. Can be used on [the following plants:] vegetables; fruits; herbs; nuts; ornamental trees, shrubs, hemp, flowering plants; houseplants, and tropical plants [grown in and around home gardens or home greenhouses]

EFFICACY CLAIMS

1. Controls mites, caterpillars, whiteflies, thrips, aphids, and other insects as listed on this label.
2. Controls chewing and sucking insects
3. Broad-spectrum control
4. Insect repellent/insecticide
5. Miticide
6. Kills listed insects such as mites, thrips, psyllids, leafhoppers, lepidoptera larvae, white flies, borers, mealybugs and true bugs

U.S. Patent No. X,XXX,XXX

© 20XX [Licensed by] ***insert company name***. [World rights reserved.]

Sold by: ***insert company name and address***

Made in/Product of _____

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK, OR FIRE), CALL CHEMTREC® (800) 424-9300.
 For other product information, contact General Hydroponics or see Material Safety Data Sheet.

NOTICE ON CONDITIONS OF SALE

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of General Hydroponics. To the extent consistent with applicable law, all such risks are assumed by the Buyer and User.

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified. Country: United States of America.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

SUBLABEL 3

GH NAMT

BOTANICAL INSECTICIDE, MITICIDE, AND NEMATICIDE

- REPELLANT, ANTIFEEDANT AND INSECT GROWTH REGULATOR (IGR)
- BOTANICAL PRODUCT FOR CONTROL OF INSECTS ON FRUIT AND NUT TREES.
- INDOOR AND OUTDOOR VEGETABLES, ORNAMENTAL FLOWERS, TREES, SHRUBS, HEMP AND PLANTS, INCLUDING PLANTS GROWN IN CONTAINERS, INTERIORSCAPES, HOME AND GARDEN USE.

See the Directions for Use for a Complete List of Insects Controlled.

ACTIVE INGREDIENT:	%by wt.
Azadirachtin.....	1.2%
OTHER INGREDIENTS:	98.8%
Total	100.0%

Contains 0.35 grams azadirachtin per fluid ounce.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back/side/top/bottom/booklet [panel/label] for [additional] directions for use/precautions/precautionary statements/first aid [statements].

FIRST AID	
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call poison control center or doctor for further treatment advice.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. CALL THE POISON CONTROL HOTLINE 24 HOURS A DAY AT 1-877-465-5161	

[Sold by:] General Hydroponics, 2877 Giffen Ave., Santa Rosa, CA 95407

EPA Reg. No. 91865-4

EPA Est. No. XXX-XX-X (*insert EPA Registered Establishment Number(s)*)

[Superscript used is first letter of lot code]

NET CONTENTS XX.XX fl oz (X.XX L)

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, absorbed through skin or if inhaled. Avoid breathing vapor. Causes moderate eye irritation. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse. Wear chemical resistant gloves.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt
- Long pants
- Socks and shoes
- Chemical resistant gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates.

For Terrestrial Uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate.

USER SAFETY RECOMMENDATIONS

User should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

User should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the Agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural products. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box apply to the uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow any worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, water wear coveralls, chemical resistant gloves, shoes plus socks and protective eyewear.

MODE OF ACTION

GH NAMT controls target pests on contact or by ingestion. The product acts on pests by way of repellence, anti-feedance, and interference with the molting process.

Azadirachtin, an insect growth regulator (IGR), mimics the pests' hormones and disrupts distinct stages of growth and development of insects and mites. The primary mode of action of azadirachtin is an interference with synthesis and metabolism of ecdysone and the juvenile hormone. Ecdysone is the molting hormone of insects, and azadirachtin can regulate growth leading to death before or during molting.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

I. USES FOR ORNAMENTALS AND LANDSCAPE PLANTINGS

Ornamental Plants and Flowers	Actinopteris, African violets*, ageratum, aglaonema, Algerian ivy, allamanda, alocaasia, amaranthus, anthurium, aphelandra, arborvitae, Artemisia, aster, aucuba ilex, azalea, baby's breath, begonia, Boston fern, bougainvillea, boxwood, brachycome, cacti, calabrese, caladium, calathea, calendula, calla, camellia, carnation, ceanothus, chrysanthemum, cineraria, coleus, columbine, cotoneaster, cyclamen, daffodil, dahlia, daisy, daylily, delphinium, dianthus, dieffenbachia, dogwood, dusty miller, Easter lily, English ivy, euphorbia, fern, ficus, foliage plants, foxglove, freesia, fuschia, gaillardia, gardenia, geranium, gerbera, gladiola, gloxinia, gypsophilla, heder, hibiscus, hyacinth, hydrangea, ilex, impatiens, iris, ivy, jasmine, lilac, lily, maidenhair fern, mandevilla, marigold, narcissus, nasturtium, orchid*, pansy, pelargonium, peony, peperomia, petunia, philodendron, phlox, photinia, pinks, pittosporum, poinsettia*, pothos, portulaca, primrose, pyracantha, rhododendron, rose*, rosemary, rubber plant, salvia, schefflera, sedum, sempervivum, snapdragon, spathiphyllum, stock, syngonium, tulip, verbena, vinca, wandering jew, yucca, zinnia *Please note that when making applications to these species, spotting of plant foliage and blossoms is possible.
Ornamental Trees and Shrubs	Andromeda, arborvitae, ash, Austrian pine, azalea, beech, birch, birdsnest spruce, blue spruce, bougainvillea, boxwood, butternut, cedar, charmaecyparis, cherry, cotoneaster, crabapple, cyprus, dogwood, Douglas fir, elm, euonymus, firethorn, forsythia, hackberry, hawthorn, hemlock, hickory, holly, honey locust, horse chestnut, juniper, larch, laurel, lilac, linden, London planetree, magnolia, mandevilla, maple, mimosa, mountain ash, myrtle, oak, pachysandra, peach, photinia, pine, planetree, poplar, privet, purpleleaf wintercreeper, quince, sage, spruce, sycamore, white cedar, white pine, yew

Waxy bloom on certain ornamental plants will be reduced after a GH NAMT application.

Applications of GH NAMT will remove the glaucous 'blue' coloring from evergreens such as Colorado blue spruce and Koster spruce.

PESTS CONTROLLED OR SUPPRESSED

Use GH NAMT against the following pests.

TARGET PEST SPECIES OF GH NAMT.

HEMIPTERA AND HOMOPTERA	true bugs including boxelder bugs, chinch bugs, lygus bugs and stink bug; lacebugs; leafhoppers including grape leafhopper, spittlebug, potato leafhopper and variegated leafhopper; mealy bugs including apple mealy bugs, citrus mealy bugs, grape mealybugs; whiteflies including whitefly, silverleaf whitefly and sweet potato whitefly and woolly whitefly; aphids including apple aphid, green peach aphid, melon aphid, pea aphid, potato aphid and rose aphid; psyllids including pear psyllids and scales including black scale, brown soft scale, California red scale, coffee scale, olive scale, San Jose scale, and cottony cushion scale.
LEPIDOPTERA	Moths including European pine shoot moth, pine tip moth and Tussock moth; leafrollers including blueberry leafroller, filbert leafroller, fruitree leafroller, citrus leafminers, grape leafroller, oblique banded leafroller, omnivorous leafroller; Cutworms including black cutworm and citrus cutworm; Caterpillars and loopers including bagworms, budworms, cabbage looper, canker worms, case bearers, caseworms, corn earworm, diamondback moth, fruit worms, grapeleaf skeletonizer, gypsy moth, hornworms, imported cabbageworm, navel orangeworm, soybean looper, spruce budworm, tent caterpillar, tip moths, tent caterpillars, tobacco budworm, tobacco hornworm, tomato pinworm and tussock moth; Amyworms including beet armyworm, fall armyworm, lawn armyworm, southern armyworm and yellow striped armyworm; webworms and leaf perforators.
COLEOPTERA	beetles, grubs and weevils including Asian long-horned beetle, bark beetles, black vine weevil, Colorado potato beetle, elm bark beetle, European chafer, flea beetles, Japanese beetle, June beetle, leaf beetles, Mexican bean beetle, Northern masked chafer, rose chafer and Southern masked chafer and twig girders.
DIPTERA*	Flies including Caribbean fruit fly, cherry maggots, crane fly, fungus gnat , Hessian fly, oriental fruit fly, Mediterranean fruit fly, marsh crane flies, melon fly, shore fly and walnut husk fly; leafminers including citrus leafminers and serpentine leafminers. *Not intended for use on public health pests

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

THYSANOPTERA	thrips including citrus thrips, flower thrips, gladiolus thrips, onion thrips, thrips palmi and Western flower thrips.
ACARINA*	mites including, red spider mites, brown mite, clover mite, conifer spider mite: European red mite, spruce spider mite, and two-spotted spider mite. *Not intended for use on public health pests
ORTHOPTERA	crickets; grasshoppers; locusts
HYMENOPTERA*	sawflies including European sawflies, pear sawflies, red-headed pine sawflies, yellow-headed pin sawflies. *Not intended for use on public health pests
NEMATODA	nematodes (suppression)

SPRAY PREPARATION

GH NAMT is an emulsifiable concentrate to be diluted with water.

This product forms an emulsion which separates upon extended or prolonged standing. Re-agitate to assure uniformity of the spray mixture.

Prepare only the volume needed for the intended application, and use the spray mixture within 24 hours of preparation.

TANK MIXTURES

GH NAMT is an emulsifiable concentrate and is compatible with commonly used pesticides and fertilizers. Always check the physical compatibility using a jar test in the correct proportions if needed.

If a broader spectrum of control is required tank mix GH NAMT with insecticides or miticides. If a rapid knockdown of heavy populations is necessary, then include an effective contact insecticide/miticide in combination with GH NAMT.

Always read and follow the directions for use, precautions and limitations for use on all product labels used in combination. Applications must follow the precautions and limitations of the most restrictive product label in the mixture. Do not exceed the dosage rates of any product.

Select the right companion products:

IPM uses a variety of control options including biological, chemical, and cultural practices. GH NAMT is botanical with growth regulator effect on insects and mites. Companion products include pyrethroids, spinosyns, microbial toxins, and chloronicotinyls that complement azadirachtin. Formulations of bifenthrin, spinosad, abamectins, and imidacloprid are effective for different pests. Select the product that has been proven to provide adequate performance for the pests you are trying to control.

Physical Incompatibility

Do not use GH NAMT with Captan, Bordeaux mixture, triphenyltin hydroxide, lime sulfur, Rayplex iron or other highly alkaline materials as they can cause phytotoxicity and/or reduced efficacy on some target pests. Phytotoxicity will occur if tank-mix combinations with compounds known to be incompatible with oil-based formulations are used.

APPLICATION EQUIPMENT

Ground Equipment

Apply GH NAMT with hand-operated (manual) or power spray equipment suitable for low volume and/or high volume applications. Follow the recommendations of the equipment manufacturer when using backpack sprayers; hose-end sprayers, compression (pump-up) sprayers, and other sprayers suitable for foliar applications of insecticides.

APPLICATION SCHEDULE

For the most effective control, apply the product when pests are expected to appear or as soon as possible after pests appear and are in immature stages. Spray at an interval of seven (7) to ten (10) days or as the situation warrants.

During high pest infestation levels or when canopy is dense use higher dosage (use) rates and increase the spray frequency. For best results spray in the morning or evening hours. Repeat spraying if rain occurs within two to three hours of spraying. For additional guidance, consult with your state agricultural experiment station or local extension horticulturalist/arborist for information on tactics and windows of application.

FOLIAR APPLICATION RATES FOR ORNAMENTALS AND LANDSCAPE PLANTINGS

USE	SPRAY CONCENTRATION %	Amounts of GH NAMT
		Fluid ounces per gallon
Including trees, shrubs, flowers, conifers, evergreens, herbaceous ornamentals, foliage plants, container-grown ornamentals, plants and groundcovers	Lower rate ranges of 0.25 - 0.75% vol/vol:	0.32 – 1.0 fl. oz.
	Medium rate ranges of 0.75 – 1.25% vol/vol:	1.00– 1.60 fl. oz.
	Upper rate ranges of 1.25 – 1.70% vol/vol:	1.60 – 2.18 fl. oz.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

DRENCH APPLICATION FOR INTERIORSCAPES AND FOR PLANTS GROWN IN CONTAINERS:

Use GH NAMT as a soil drench for effective control of soil-borne insect larvae, including soil-borne larvae of foliar pests, such as fungus gnats, nematodes, or soil borne thrips. When applying as a drench, avoid excessive leaching.

Preventive applications as a soil drench may be warranted for certain pests. Soil drench applications of azadirachtin will have a slower rate of activity because of soil absorption when compared to foliar applications of azadirachtin. Target the initial application of a soil drench treatment to coincide with the early stages of young larvae and young nymphs.

Dilute GH NAMT with water for concentrations of 0.4 to 0.8% vol/vol. Drench the soil in the pot with one (1) pint of finished spray per 1.0 gallon of soil. For fungus gnats, use the 0.4% spray concentration. For mushroom fly maggot control, use the 0.6% vol/vol spray concentration. For leafminers and other difficult to control pests, use the 0.8% vol/vol spray concentration make two to three (2-3) applications at 10-14 day intervals until the pest pressure has ended.

DILUTION TABLE FOR DRENCH APPLICATIONS

Gallons of Water	Amount of GH NAMT			Application Interval	Number of Applications
	0.4%	0.6%	0.8%		
1 gallon	1 Tbs.	1.5 Tbs.	2.0 Tbs.	10 – 14 days	2 - 3
5 gallons	2.7 fl. oz	4.0 fl. oz	5.5 fl. oz	10 – 14 days	2 – 3
10 gallons	5.4 fl. oz	8.0 fl. oz	11.0 fl. oz	10 – 14 days	2 – 3

GH NAMT can also be applied through sub-surface treatment equipment. Always follow manufacturer's use directions.

II. USES FOR GARDEN CROPS, VEGETABLES, HERBS, AND SPICES, HEMP, FRUITS AND BERRIES

INSTRUCTIONS FOR GARDEN CROPS, VEGETABLES, HERBS AND SPICES, HEMP, FRUITS, AND BERRIES

- For the most effective control, spray the product as soon as possible after pests appear and are in immature stages.
- Spray at an interval of seven to ten days or as the situation warrants. During high pest infestation levels use higher label rates and increase the spray frequency.
- For best results spray in the morning or evening hours.
- Repeat spraying if rain occurs within two to three hours of spraying.

USE RATES

Apply GH NAMT as directed to any food or non-food crop up to and including the day of harvest, at a maximum rate of 1.33 fl. ozs. per 1000 sq. ft. per application.

Apply GH NAMT alone to food/garden crops on the day of harvest.

Dilute GH NAMT with water at 0.5 - 4.0 tablespoons (Tbs) per gallon of water. For hose end sprayers, set the RATE PER GALLON at the dial setting of 1 to 4 Tbs. depending on the crop and pests. Use the lower RATE PER GALLON for low to moderate infestations and use the higher specified RATE PER GALLON for severe infestations.

USE RATES FOR GARDEN CROPS, VEGETABLES, HERBS AND SPICES, HEMP, BERRIES AND FRUIT.

CROP	PESTS such as:	Dilution Rate for Sprayers	
		fl. ozs of product per 1000 sq. ft.	Tbs. of product per 1.0 gallon of water
Leafy vegetables: Broccoli, Brussel Sprouts, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce, Spinach	Leafrollers, Cutworms, Loopers, Armyworms	0.19 – 0.96 fl. oz.	¾ Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Beetles, Weevils, Flies, Thrips, Mites	0.24 – 0.96 fl. oz.	1 Tbs. – 4 Tbs./gal
Root Vegetables: Beet, Carrot, Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yams	Beetles, Weevils	0.11 – 0.25 fl. oz.	1/2 Tbs. – 1 1/2 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Thrips, Mites	0.24 – 0.96 fl. oz.	1 Tbs. – 4 Tbs./gal
Fruiting Vegetables: Eggplant, Pepper, Tomatillo, Tomato	Beetles, Weevils	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz.	1 Tbs. – 4 Tbs./gal
Cucurbit Vegetables: Cucumber, Gourd (edible),	Beetles, Weevils	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

Muskmelon, Pumpkin, Squash, Watermelon, Including Cantaloupe, Casaba, Gherkins, Melons (including hybrids), Zucchini	Thrips,	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz.	1 Tbs. – 4 Tbs./gal
Legume Vegetables: Bean, Chickpea, Lentil, Pea	Beetles, Weevils	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz.	1 Tbs. – 4 Tbs./gal
Bulb Vegetables: Garlic, Onion, Shallot	Beetles, Weevils	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz.	1 Tbs. – 4 Tbs./gal
Berries: Blackberry, Blueberry, Raspberry, Strawberry, others include: Boysenberry, Currants, Dewberry, Elderberry, Gooseberry, Loganberry	Beetles, Weevils	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz.	1 Tbs. – 4 Tbs./gal
Herbs and Spices: Chive, Dill, Fennel, Mustard, Sage, Sweet bay, others include: Anise, Balm, Basil, Black pepper, Borage, Caraway, Catnip, Chamomile, Coriander, Cumin, Curry leaf, Dandelion, Fenugreek, Horehound, Hyssop, Marjoram, Marigold, Mint, Nasturtium, Pennyroyal, Peppermint, Rosemary, Savory, Spearmint, Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood	Beetles, Weevils	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz.	1 Tbs. – 4 Tbs./gal
Other crops: Hemp	True Bugs, Leafhoppers, Whiteflies, Aphids, Mites, Beetles, Weevils, Flies, Thrips	0.24 – 0.96 fl. oz.	1 Tbs. – 4 Tbs./gal
	Leafrollers, Cutworms, Loopers, Armyworms	0.19 – 0.96 fl. oz.	¾ Tbs. – 4 Tbs./gal
Nut Trees: Almond, Brazil nut, Filbert, Hickory nut, Pecan, Pistachios, Walnut	Beetles, Weevils	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz.	1 Tbs. – 4 Tbs./gal
Pome Fruits: Apple, Quince, or Pear (Comice varieties: DO NOT apply more than 24 fl. oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale usage.)	Beetles, Weevils	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz.	1 Tbs. – 4 Tbs./gal

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

Stone Fruits: Apricot, Cherry, Nectarine, Peach, Plum	Beetles, Weevils	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz.	1 Tbs. – 4 Tbs./gal
Citrus Fruits: Grapefruit, Lemon, Lime, Orange, others include: Citrus Citron, Mandarin (tangerine), Nectarine, Satsuma (orange mandarin), tangerine	Beetles, Weevils	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz.	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz.	1 Tbs. – 4 Tbs./gal

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal

Pesticide Storage: Store in original container in a dry, cool, well ventilated area.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill container 1/4 full with water and recap. Shake 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

OPTIONAL TEXT AND CLAIMS

GENERAL CLAIMS

1. For organic gardening
2. OMRI Logo
3. Organic certified
4. Organic Materials Review Institute (OMRI) Certified
5. Can be used in Organic Gardening FOR ORGANIC PRODUCTION
6. Low-odor formulation
7. Low (Mild) odor
8. Three in one product - Insecticide, Miticide, and Nematicide
9. Recommended by General Hydroponics Bringing Nature and Technology Together
10. Brought to you Exclusively by General Hydroponics - Bringing Nature and Technology Together
11. [Questions/Comments [Call]] ***insert company phone number***
12. [Visit [our website]] ***insert company website***

USE SITE CLAIMS

1. Can be applied the day of harvest
2. For use on a wide variety of trees, shrubs, flowers, fruit and nut trees, hemp, garden vegetables and plants
3. For use on outdoor and greenhouse food and non-food crops, ornamental flowers, hemp, trees, shrubs and plants, landscapes
4. Spray and Drench
5. Rose Spray
6. For uses on Roses and Flowers
7. Vegetable Garden Spray
8. For use on Tomatoes and Garden Vegetables
9. For use on roses, vegetables, fruits, nuts, flowers, houseplants, hemp, foliage, trees, and shrubs
10. Taken up through the roots, controls insects on the leaves
11. For foliar and systemic insect control
12. For Use on Flowers, Ornamentals, and Landscape Plantings
13. For Use on Garden Crops, Vegetables, Herbs, Spices, hemp, Fruits, and Berries
14. Can be used on [the following plants:] vegetables; fruits; herbs; nuts; ornamental trees, shrubs, hemp, flowering plants; houseplants, and tropical plants [grown in and around home gardens or home greenhouses]

EFFICACY CLAIMS

1. Controls mites, caterpillars, whiteflies, thrips, aphids, and other insects as listed on this label.
2. Controls chewing and sucking insects
3. Broad spectrum insect and mite control
4. Broad-spectrum control
5. Controls Japanese Beetles on Roses
6. Insect and Mite Control for Home Garden Vegetables
7. Complete Insect and Mite Control for Flowers
8. Three applications in 30 days provides complete plant protection
9. Quadruple action insect control – Antifeedant, Insect Growth Regulator, Anti-ovipository, Repellant
10. The Pure Power of Purer azadirachtin
11. Single product; multiple action
12. A proven botanical pesticide born out of a decade of intense global research
13. Works effectively against a broad spectrum of pests on a stand alone basis on a variety of vegetable crops
14. Controls insect pest during various stages of growth
15. The benefits to your insect control program are multiple
16. Controls Insects Systemically
17. Insects stop feeding when they feed on treated plants
18. Insect repellant/insecticide
19. Miticide
20. Kills listed insects such as mites, thrips, psyllids, leafhoppers, lepidoptera larvae, white flies, borers, mealybugs and true bugs

U.S. Patent No. X,XXX,XXX

© 20XX [Licensed by] ***insert company name***. [World rights reserved.]

Sold by: ***insert company name and address***

Made in/Product of _____

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK, OR FIRE), CALL CHEMTREC® (800) 424-9300.

For other product information, contact General Hydroponics or see Material Safety Data Sheet.

WARRANTY INFORMATION

General Hydroponics warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purpose referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the disease problem, condition of the plant, incompatibility with other influencing factors in the use of this product are beyond the control of the seller. To the extent consistent with applicable law, buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herein, with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OR MERCHANTABILITY IS MADE.

NOTICE ON CONDITIONS OF SALE

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of General Hydroponics. To the extent consistent with applicable law, all such risks are assumed by the Buyer and User.

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified. Country: United States of America.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

SUBLABEL 4

GH NAMT

BOTANICAL INSECTICIDE, MITICIDE, AND NEMATICIDE

REPELLANT, ANTIFEEDANT AND INSECT GROWTH REGULATOR (IGR)

BOTANICAL PRODUCT FOR CONTROL OF INSECTS ON INDOOR AND OUTDOOR ORNAMENTAL FLOWERS, TREES, SHRUBS, VEGETABLES, FRUIT AND NUT TREES, HEMP, PLANTS, INCLUDING PLANTS GROWN IN CONTAINERS, RECIRCULATORY, AEROPONIC, AND HYDROPONIC SYSTEMS, INTERIORSCAPES, GREENHOUSES, HOME AND GARDEN

See the Directions for Use for a Complete List of Insects Controlled

ACTIVE INGREDIENT:	% by wt.
Azadirachtin.....	1.2%
OTHER INGREDIENTS:	98.8%
Total	100.0%

Contains 0.35 grams azadirachtin per fluid ounce.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back/side/top/bottom/booklet [panel/label] for [additional] directions for use/precautions/precautionary statements/first aid [statements].

FIRST AID	
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
If on skin or clothing	Take off contaminate clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. CALL THE POISON CONTROL HOTLINE 24 HOURS A DAY AT 1-877-465-5161	

[Sold by:] General Hydroponics, 2877 Giffen Ave., Santa Rosa, CA 95407

EPA Reg. No. 91865-4

EPA Est. No. XXX-XX-X (*insert EPA Registered Establishment Number(s)*)

[Superscript used is first letter of lot code]

NET CONTENTS XX.XX fl oz (X.XX L)

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, absorbed through skin or if inhaled. Avoid breathing vapor. Causes moderate eye irritation. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash contaminated clothing before reuse. Wear chemical resistant gloves.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt
- Long pants
- Socks and shoes
- Chemical resistant gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates.

For Terrestrial Uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural products. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, water, wear coveralls, chemical resistant gloves, shoes plus socks and protective eyewear.

MODE OF ACTION

GH NAMT controls target pests on contact or by ingestion. The product acts on pests by way of repellence, anti-feedance, and interference with the molting process.

Azadirachtin, an insect growth regulator (IGR), mimics the pests' hormones and disrupts distinct stages of growth and development of insects and mites. The primary mode of action of azadirachtin is an interference with synthesis and metabolism of ecdysone and the juvenile hormone. Ecdysone is the molting hormone of insects, and azadirachtin can regulate growth leading to death before or during molting.

GH NAMT will provide control results comparable to the synthetic insecticide standards. GH NAMT provides broad spectrum control with very low environmental impact. GH NAMT provides all the benefits of azadirachtin, a proven anti-feedant, insect growth regulator (IGR), anti-ovipository, and repellent, as well as a toxin to soft bodied insect larvae.

The active ingredient in GH NAMT - Azadirachtin - is a unique botanical insecticide, miticide and nematicide.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

Mode of Action: Control of different orders of insects or insects in different phases of their life cycle is due to the complexity of the azadirachtin molecule and the many modes of action inherent in azadirachtin	Anti-Feedant	Insects feed less or not at all on treated plants. Foliage is not damaged and insects ultimately starve to death.
	Insect Growth Regulator (IGR)	Insects fail to mature and reproduce, eliminating populations over time.
	Anti-ovipository	Insects do not lay eggs on treated plants. The likelihood of insect infestation is greatly decreased in treated plants. This adds a preventive aspect to your insect control.
	Repellant	Insects do not prefer treated plants.

PESTS CONTROLLED OR SUPPRESSED

Use GH NAMT against the following pests.

TARGET PEST SPECIES OF GH NAMT.

HEMIPTERA AND HOMOPTERA	<p>true bugs including boxelder bugs, chinch bugs, lygus bugs and stink bug; lacebugs; leafhoppers including grape leafhopper, spittlebug, potato leafhopper and variegated leafhopper; mealy bugs including apple mealy bugs, citrus mealy bugs, grape mealybugs; whiteflies including greenhouse whitefly, silverleaf whitefly and sweet potato whitefly and woolly whitefly; aphids including apple aphid, green peach aphid, melon aphid, pea aphid, potato aphid and rose aphid; psyllids including pear psyllids and scales including black scale, brown soft scale, California red scale, coffee scale, olive scale, San Jose scale, and cottony cushion scale.</p>
LEPIDOPTERA	<p>Moths including European pine shoot moth, pine tip moth and Tussock moth; leafrollers including blueberry leafroller, filbert leafroller, fruitree leafroller, citrus leafminers, grape leafroller, oblique banded leafroller, omnivorous leafroller; Cutworms including black cutworm and citrus cutworm; Caterpillars and loopers including bagworms, budworms, cabbage looper, canker worms, case bearers, caseworms, corn earworm, diamondback moth, fruit worms, grapeleaf skeletonizer, gypsy moth, hornworms, imported cabbageworm, navel orangeworm, soybean looper, spruce budworm, tent caterpillar, tip moths, tent caterpillars, tobacco budworm, tobacco hornworm, tomato pinworm and tussock moth; Amyworms including beet armyworm, fall armyworm, lawn armyworm, southern armyworm and yellow striped armyworm; webworms and leaf perforators.</p>
COLEOPTERA	<p>beetles, grubs and weevils including Asian long-horned beetle, bark beetles, black vine weevil, Colorado potato beetle, elm bark beetle, European chafer, flea beetles, Japanese beetle, June beetle, leaf beetles, Mexican bean beetle, Northern masked chafer, rose chafer and Southern masked chafer and twig girders.</p>
DIPTERA*	<p>Flies including Caribbean fruit fly, cherry maggots, crane fly, fungus gnat, Hessian fly, oriental fruit fly, Mediterranean fruit fly, marsh crane flies, melon fly, shore fly and walnut husk fly; leafminers including citrus leafminers and serpentine leafminers.</p> <p>*Not intended for use on public health pests</p>
THYSANOPTERA	<p>thrips including citrus thrips, flower thrips, gladiolus thrips, onion thrips, thrips palmi and Western flower thrips.</p>
ACARINA*	<p>mites including, red spider mites, brown mite, clover mite, conifer spider mite: European red mite, spruce spider mite, and two-spotted spider mite.</p> <p>*Not intended for use on public health pests</p>
ORTHOPTERA	<p>crickets; grasshoppers; locusts</p>
HYMENOPTERA*	<p>sawflies including European sawflies, pear sawflies, red-headed pine sawflies, yellow-headed pin sawflies.</p> <p>*Not intended for use on public health pests</p>
NEMATODA	<p>nematodes (suppression)</p>

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

FOR USE ON ORNAMENTALS AND LANDSCAPE PLANTINGS

Ornamental Plants and Flowers:	Actinopterus, African violets*, ageratum, aglaonema, Algerian ivy, allamanda, alocasia, amaranthus, anthurium, aphelandra, arborvitae, Artemisia, aster, aucuba ilex, azalea, baby's breath, begonia, Boston fern, bougainvillea, boxwood, brachycome, cacti, calabrese, caladium, calathea, calendula, calla, camellia, carnation, ceanothus, chrysanthemum, cineraria, coleus, columbine, cotoneaster, cyclamen, daffodil, dahlia, daisy, daylily, delphinium, dianthus, dieffenbachia, dogwood, dusty miller, Easter lily, English ivy, euphorbia, fern, ficus, foliage plants, foxglove, freesia, fuschia, gaillardia, gardenia, geranium, gerbera, gladiola, gloxinia, gypsophilla, hederia, hibiscus, hyacinth, hydrangea, ilex, impatiens, iris, ivy, jasmine, lilac, lily, maidenhair fern, mandevilla, marigold, narcissus, nasturtium, orchid*, pansy, pelargonium, peony, peperomia, petunia, philodendron, phlox, photinia, pinks, pittosporum, poinsettia*, pothos, portulaca, primrose, pyracantha, rhododendron, rose*, rosemary, rubber plant, salvia, schefflera, sedum, sempervivum, snapdragon, spathiphyllum, stock, syngonium, tulip, verbena, vinca, wandering jew, yucca, zinnia *Please note that when making applications to these species, spotting of plant foliage and blooms is possible.
Ornamental Trees and Shrubs:	Andromeda, arborvitae, ash, Austrian pine, azalea, beech, birch, birdsnest spruce, blue spruce, bougainvillea, boxwood, butternut, cedar, charmaecyparis, cherry, cotoneaster, crabapple, cyprus, dogwood, Douglas fir, elm, euonymus, firethorn, forsythia, hackberry, hawthorn, hemlock, hickory, holly, honey locust, horse chestnut, juniper, larch, laurel, lilac, linden, London planetree, magnolia, mandevilla, maple, mimosa, mountain ash, myrtle, oak, pachysandra, peach, photinia, pine, planetree, poplar, privet, purpleleaf wintercreeper, quince, sage, spruce, sycamore, white cedar, white pine, yew

Waxy bloom on certain ornamental plants will be reduced after an application.

Applications of GH NAMT will remove the glaucous 'blue' coloring from evergreens such as Colorado blue spruce and Koster spruce.

FOR USE ON GARDEN CROPS, VEGETABLES, HERBS, SPICES, HEMP, FRUITS, AND BERRIES

Leafy Vegetables:	Broccoli, Brussel Sprouts, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce, Spinach
Root Vegetables:	Beet, Carrot, Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yams
Fruiting Vegetables:	Eggplant, Pepper, Tomatillo, Tomato
Cucurbit Vegetables:	Cucumber, Gourds (edible), Muskmelon, Pumpkin, Squash, Watermelon, including Cantaloupe, Casaba, Gherkins, Melons (including hybrids), Zucchini
Legume Vegetables:	Bean, Chickpea, Lentil, Pea
Bulb Vegetables:	Garlic, Onion, Shallot
Berries:	Blackberry, Blueberry, Raspberry, Strawberry, others include: Boysenberry, Currants, Dewberry, Elderberry, Gooseberry, Loganberry
Herb and Spices:	Chive, Dill, Fennel, Mustard, Sage, Sweet bay, others include: Anise, Balm, Basil, Black pepper, Borage, Caraway, Catnip, Chamomile, Coriander, Cumin, Curry Leaf, Dandelion, Fenugreek, Horehound, Hyssop, Marjoram, Marigold, Mint, Nasturtium, Pennyroyal, Peppermint, Rosemary, Savory, Spearmint, Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood.
Nut Trees:	Almond, Brazil nut, Filbert, Hickory nut, Pecan, Pistachios, Walnut
Other crops:	Hemp
Pome Fruits:	Apple, Quince, or Pear (Cornice varieties: DO NOT apply more than 24 fl oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale usage.)
Stone Fruits:	Apricot, Cherry, Nectarine, Peach, Plum
Citrus Fruits:	Grapefruit, Lemon, Lime, Orange others include: Citrus Citron, Mandarin (tangerine), Nectarine, Satsuma (orange mandarin), Tangerine

GH NAMT has been evaluated for phytotoxicity on a wide range of ornamentals and garden plants. However, since testing on all plant varieties is not feasible, test a small portion of the area to be treated for phytotoxicity before treating the entire area.

There are no restrictions on applying GH NAMT up to the time of harvest.

SPRAY PREPARATION

GH NAMT is an emulsifiable concentrate to be diluted with water.

This product forms an emulsion and can separate upon extended or prolonged standing. Re-agitate to assure uniformity of the spray mixture.

Prepare only the volume needed for the intended application, and use the spray mixture within 24 hours of preparation.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

TANK MIXTURES

GH NAMT is an emulsifiable concentrate and is compatible with commonly used pesticides and fertilizers. Always check the physical compatibility using a jar test in the correct proportions if needed.

If a broader spectrum of control is required tank-mix GH NAMT with insecticides or miticides. If a rapid knockdown of heavy populations is necessary, then include an effective contact insecticide/miticide in combination with GH NAMT.

Always read and follow the directions for use, precautions and limitations for use on all product labels used in combination. Applications must follow the precautions and limitations of the most restrictive product label in the mixture. Do not exceed the dosage rates of any product.

Select the right companion products:

IPM uses a variety of control options including biological, chemical, and cultural practices. GH NAMT is botanical with growth regulator effect on insects and mites. Companion products include pyrethroids, spinosyns, microbial toxins, and chloronicotinyls that complement azadirachtin. Formulations of bifenthrin, spinosad, abamectins, and imidacloprid are effective for different pests. Select the product that has been proven to provide adequate performance for the pests you are trying to control.

Physical Incompatibility

Do not use GH NAMT with Captan, Bordeaux mixture, triphenyltin hydroxide, lime sulfur, Rayplex iron or other highly alkaline materials as they can cause phytotoxicity and/or reduced efficacy on some target pests. Phytotoxicity will occur if tank-mix combinations with-compounds known to be incompatible with oil-based formulations are used.

APPLICATION EQUIPMENT

Apply GH NAMT with hand-operated (manual) or power spray equipment suitable for low volume and/or high volume applications. Follow the recommendations of the equipment manufacturer when using backpack sprayers; hose-end sprayers, compression (pump-up) sprayers, and other sprayers suitable for foliar applications of insecticides.

APPLICATION SCHEDULE

For the most effective control, apply GH NAMT when pests are expected to appear or as soon as possible after pests appear and are in immature stages. Spray at an interval of seven (7) to ten (10) days or as the situation warrants.

During high pest infestation levels or when canopy is dense use higher dosage (use) rates and increase the spray frequency. Spraying in the morning or evening hours. Repeat spraying if rain occurs within two to three hours of spraying. For additional guidance, consult with the state agricultural experiment station or local extension horticulturalist/arborist for information on tactics and windows of application.

APPLICATION METHODS

Apply GH NAMT as directed to any food or non-food crop up to and including the day of harvest, at a maximum rate of 1.33 fl. oz. per 1,000 sq. ft. per application.

Dilute GH NAMT with water at a rate of 0.5 - 4.0 tablespoons (Tbs) per gallon of water. For hose end sprayers, set the RATE PER GALLON at the dial setting of 1 to 4 Tbs. depending on the crop and pests. Use the lower RATE PER GALLON for low to moderate infestations and use the higher specified RATE PER GALLON for severe infestations.

FOLIAR APPLICATION

USE	SPRAY CONCENTRATION %	Amounts of GH NAMT	
		Fluid ounces (Tbs.) per quart	Fluid ounces (Tbs.) per gallon
Including trees, shrubs, flowers, conifers, evergreens, herbaceous, ornamentals, hemp, foliage plants, container-grown ornamentals & garden plants and groundcovers	Lower rate ranges of 0.25 - 0.75% vol/vol:	0.08 – 0.25 fl. oz (1/6 – 1/2 Tbs.)	0.32 – 1 fl. oz (2/3 – 2.0 Tbs.)
	Medium rate ranges of 0.75 – 1.25% vol/vol:	0.25 – 0.40 fl. oz (1/2 – 5/6 Tbs.)	1.0 – 1.6 fl. oz (2.0 – 3 1/3 Tbs.)
	Upper rate ranges of 1.25 – 1.70% vol/vol:	0.40 – 0.50 fl. oz (5/6 -1.0 Tbs.)	1.6 – 2.0 fl. oz (3 1/3 – 4 Tbs.)

DRENCH APPLICATION

Use GH NAMT as a soil drench for effective control of soil-borne insect larvae, including soil-borne larvae of foliar pests, such as fungus gnats, nematodes, or soil borne thrips. When applying as a drench, avoid excessive leaching.

Preventive applications as a soil drench may be warranted for certain pests. Soil drench applications of azadirachtin will have a slower rate of activity because of soil absorption when compared to foliar applications of azadirachtin. Target the initial application of a soil drench treatment to coincide with the early stages of young larvae and young nymphs.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

Dilute GH NAMT with water for concentrations of 0.4 to 0.8% volume/volume. See use rate table below. Add the required amount of GH NAMT to a clean bucket with at least one-half of the water to be drenched. Agitate the mixture thoroughly and then fill with the remaining water and continue agitation until the product is thoroughly dispersed.

Drench the soil in the pot with one (1) pint of finished product dilution per 1.0 gallon of soil. For fungus gnats, use the 0.4% spray concentration. For mushroom fly maggot control, use the 0.6% volume/volume spray concentration. For leafminers and other difficult to control pests, use the 0.8% volume/volume spray concentration. Make two to three (2 - 3) applications at 10 -14 day intervals until pest pressure has ended. With high insect pressure make applications every 5 to 6 days. Additional applications of GH NAMT may be required with increased and prolonged pest infestation.

DILUTION TABLE FOR DRENCH APPLICATIONS

Gallons of Water	Amount of GH NAMT			Application Interval
	0.4%	0.6%	0.8%	
1 gallon	1 Tbs.	1.5 Tbs.	2.0 Tbs.	10 – 14 days
1 gallon	0.5 fl. oz.	0.8 fl. oz.	1.0 fl. oz.	10 – 14 days
5 gallons	2.5 fl. oz.	4.0 fl. oz.	5.0 fl. oz.	10 – 14 days
10 gallons	5.0 fl. oz.	8.0 fl. oz.	10.0 fl. oz.	10 – 14 days

RECIRCULATORY, AEROPONIC, AND HYDROPONIC APPLICATION

Use GH NAMT in recirculatory, aeroponic, or hydroponic systems for the control of foliar pests, soil borne insect larvae, including soil borne larvae of foliar pests such as fungus gnats, nematodes or soil borne thrips for interiorscapes, hydroponic, aeroponic and container plants.

Dilute GH NAMT with water for concentrations of 0.1 % to 0.8% volume/volume in a recirculatory or in a hydroponic liquid system. See use rate table below. Agitate the mixture thoroughly until the product is thoroughly dispersed.

For fungus gnats, use the 0.6% volume/volume concentration. For mushroom fly maggot control, use the 0.6% volume/volume concentration. For leafminers and other difficult to control pests, use the 0.8% volume/volume concentration. Make two to three (2-3) applications at 10-14 day intervals until the pest pressure has ended. With high insect pressure applications make applications every 5 to 7 days. Additional applications of GH NAMT may be required with increased and prolonged pest infestation.

DILUTION TABLE FOR RECIRCULATORY, AEROPONIC, AND HYDROPONIC APPLICATIONS

Gallons of Water	Amount of GH NAMT					Application Interval
	0.1%	0.2%	0.4%	0.6%	0.8%	
1 gallon	1/4 Tbs.	1/2 Tbs.	1 Tbs.	1.5 Tbs.	2 Tbs.	7 – 14 days
1 gallon	0.14 fl. oz	0.25 fl. oz	0.5 fl. oz	0.8 fl. oz	1.0 fl. oz	7 – 14 days
5 gallons	0.7 fl. oz	1.3 fl. oz	2.5 fl. oz	4.0 fl. oz	5.0 fl. oz	7 – 14 days
10 gallons	1.4 fl. oz	2.6 fl. oz	5.0 fl. oz	8.0 fl. oz	10.0 fl. oz	7 – 14 days

Preventive applications as a recirculatory system application may be warranted for certain pests.

GH NAMT can also be applied through sub-surface treatment equipment. Always follow manufacturer's use directions.

FOLIAR APPLICATION – SPECIFIC PESTS OF GARDEN CROPS, VEGETABLES, HERBS AND SPICES, HEMP, BERRIES AND FRUIT.

CROP	PESTS such as:	Dilution Rate for Sprayers	
		fl. ozs of product per 1000 sq. ft.	Tbs. of product per 1.0 gallon of water
Leafy vegetables: Broccoli, Brussel Sprouts, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce, Spinach	Leafrollers, Cutworms, Loopers, Armyworms	0.19 – 0.96 fl. oz	¾ Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Beetles, Weevils, Flies, Thrips, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Root Vegetables: Beet, Carrot, Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yams	Beetles, Weevils	0.11 – 0.25 fl. oz	1/2 Tbs. – 1 1/2 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Thrips, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

Fruiting Vegetables: Eggplant, Pepper, Tomatillo, Tomato	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Cucurbit Vegetables: Cucumber, Gourd (edible), Muskmelon, Pumpkin, Squash, Watermelon, Including Cantaloupe, Casaba, Gherkins, Melons (including hybrids), Zucchini	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Legume Vegetables: Bean, Chickpea, Lentil, Pea	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Bulb Vegetables: Garlic, Onion, Shallot	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Berries including: Blackberry, Blueberry, Raspberry, Strawberry, others include: Boysenberry, Currants, Dewberry, Elderberry, Gooseberry, Loganberry	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Herbs and Spices: Chive, Dill, Fennel, Mustard, Sage, Sweet bay, others include: Anise, Balm, Basil, Black pepper, Borage, Caraway, Catnip, Chamomile, Coriander, Cumin, Curry leaf, Dandelion, Fenugreek, Horehound, Hyssop, Marjoram, Marigold, Mint, Nasturtium, Pennyroyal, Peppermint, Rosemary, Savory, Spearmint, Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Other crops: Hemp	True Bugs, Leafhoppers, Whiteflies, Aphids, Mites, Beetles, Weevils, Flies, Thrips	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
	Leafrollers, Cutworms, Loopers, Armyworms	0.19 – 0.96 fl. oz	¼ Tbs. – 4 Tbs./gal

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

Nut Trees: Almond, Brazil nut, Filbert, Hickory nut, Pecan, Pistachios, Walnut	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Pome Fruits: Apple, Quince, or Pear (Comice varieties: DO NOT apply more than 24 fl. oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale usage.)	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Stone Fruits: Apricot, Cherry, Nectarine, Peach, Plum	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Citrus Fruits: Grapefruit, Lemon, Lime, Orange, others include: Citrus Citron, Mandarin (tangerine), Nectarine, Satsuma (orange mandarin), tangerine	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal

Pesticide Storage: Store in original container in a dry, cool, well ventilated area.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill container 1/4 full with water and recap. Shake 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

OPTIONAL TEXT AND CLAIMS

GENERAL CLAIMS

1. For organic gardening
2. OMRI Logo
3. Organic Materials Review Institute (OMRI) Certified.
4. Low-odor formulation
5. Low (Mild) odor
6. [Questions/Comments [Call]] ***insert company phone number***
7. [Visit [our website]] ***insert company website***

USE SITE CLAIMS

1. Can be applied the day of harvest
2. For use on a wide variety of trees, shrubs, flowers, fruit and nut trees, hemp, garden vegetables and plants
3. For use on outdoor and greenhouse food and non-food crops, ornamental flowers, trees, shrubs and plants, hemp and landscapes
4. For use on roses, vegetables, fruits, nuts, flowers, houseplants, foliage, hemp, trees, and shrubs
5. Can be used on [the following plants:] vegetables; fruits; herbs; nuts; ornamental trees, shrubs, hemp, flowering plants; houseplants, and tropical plants [grown in and around home gardens or home greenhouses]

EFFICACY CLAIMS

1. Controls mites, caterpillars, whiteflies, thrips, aphids, and other insects as listed on this label.
2. Controls chewing and sucking insects
3. Broad-spectrum control
4. Insect repellent/insecticide
5. Miticide
6. Kills listed insects such as mites, thrips, psyllids, leafhoppers, lepidoptera larvae, white flies, borers, mealybugs and true bugs

U.S. Patent No. X,XXX,XXX

© 20XX [Licensed by] ***insert company name***. [World rights reserved.]

Sold by: ***insert company name and address***

Made in/Product of _____

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK, OR FIRE), CALL CHEMTREC® (800) 424-9300.

For other product information, contact General Hydroponics or see Material Safety Data Sheet.

NOTICE ON CONDITIONS OF SALE

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of General Hydroponics. To the extent consistent with applicable law, all such risks are assumed by the Buyer and User.

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified. Country: United States of America.

Bold, italicized text is information for the reader and is not part of the label.
 [Bracketed information is optional.] Text separated by / denotes and/or options.

SUBLABEL 5 HOMEOWNER USE

GH NAMT

BOTANICAL INSECTICIDE, MITICIDE, AND NEMATOCIDE

REPELLANT, ANTIFEEDANT AND INSECT GROWTH REGULATOR (IGR)

BOTANICAL PRODUCT FOR CONTROL OF INSECTS ON INDOOR AND OUTDOOR ORNAMENTAL FLOWERS, TREES, SHRUBS, VEGETABLES, FRUIT AND NUT TREES, HEMP, PLANTS, INCLUDING PLANTS GROWN IN CONTAINERS, RECIRCULATORY, AEROPONIC, AND HYDROPONIC SYSTEMS, INTERIORSCAPES, GREENHOUSES, HOME AND GARDEN

ACTIVE INGREDIENT:	% by wt.
Azadirachtin.....	1.2%
OTHER INGREDIENTS:	98.8%
Total	100.0%

Contains 0.35 grams azadirachtin per fluid ounce.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back/side/top/bottom/booklet [panel/label] for [additional] directions for use/precautions/precautionary statements/first aid [statements].

FIRST AID	
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
If on skin or clothing	Take off contaminate clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. CALL THE POISON CONTROL HOTLINE 24 HOURS A DAY AT 1-877-465-5161	

[Sold by:] General Hydroponics, 2877 Giffen Ave., Santa Rosa, CA 95407

EPA Reg. No. 91865-4

EPA Est. No. XXX-XX-X (*insert EPA Registered Establishment Number(s)*)

[Superscript used is first letter of lot code]

NET CONTENTS XX.XX fl oz (X.XX L)

Bold, italicized text is information for the reader and is not part of the label.
 [Bracketed information is optional.] Text separated by / denotes and/or options.

**PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed, absorbed through skin or if inhaled. Avoid breathing vapor. Causes moderate eye irritation. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash contaminated clothing before reuse. Wear chemical resistant gloves.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product through any type of irrigation system.

MODE OF ACTION

GH NAMT controls target pests on contact or by ingestion. The product acts on pests by way of repellence, anti-feedance, and interference with the molting process.

Azadirachtin, an insect growth regulator (IGR), mimics the pests' hormones and disrupts distinct stages of growth and development of insects and mites. The primary mode of action of azadirachtin is an interference with synthesis and metabolism of ecdysone and the juvenile hormone. Ecdysone is the molting hormone of insects, and azadirachtin can regulate growth leading to death before or during molting.

GH NAMT will provide control results comparable to the synthetic insecticide standards. GH NAMT provides broad spectrum control with very low environmental impact. GH NAMT provides all the benefits of azadirachtin, a proven anti-feedant, insect growth regulator (IGR), anti-ovipository, and repellent, as well as a toxin to soft bodied insect larvae.

The active ingredient in GH NAMT - Azadirachtin - is a unique botanical insecticide, miticide and nematicide.

Mode of Action: Control of different orders of insects or insects in different phases of their life cycle is due to the complexity of the azadirachtin molecule and the many modes of action inherent in azadirachtin	Anti-Feedant	Insects feed less or not at all on treated plants. Foliage is not damaged and insects ultimately starve to death.
	Insect Growth Regulator (IGR)	Insects fail to mature and reproduce, eliminating populations over time.
	Anti-ovipository	Insects do not lay eggs on treated plants. The likelihood of insect infestation is greatly decreased in treated plants. This adds a preventive aspect to your insect control.
	Repellent	Insects do not prefer treated plants.

PESTS CONTROLLED OR SUPPRESSED

Use GH NAMT against the following pests.

TARGET PEST SPECIES OF GH NAMT.

HEMIPTERA AND HOMOPTERA	<p>true bugs including boxelder bugs, chinch bugs, lygus bugs and stink bug; lacebugs; leafhoppers including grape leafhopper, spittlebug, potato leafhopper and variegated leafhopper; mealy bugs including apple mealy bugs, citrus mealy bugs, grape mealybugs; whiteflies including greenhouse whitefly, silverleaf whitefly and sweet potato whitefly and woolly whitefly; aphids including apple aphid, green peach aphid, melon aphid, pea aphid, potato aphid and rose aphid; psyllids including pear psyllids and scales including black scale, brown soft scale, California red scale, coffee scale, olive scale,</p>
-------------------------	--

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

	San Jose scale, and cottony cushion scale.
LEPIDOPTERA	Moths including European pine shoot moth, pine tip moth and Tussock moth; leafrollers including blueberry leafroller, filbert leafroller, fruitree leafroller, citrus leafminers, grape leafroller, oblique banded leafroller, omnivorous leafroller; Cutworms including black cutworm and citrus cutworm; Caterpillars and loopers including bagworms, budworms, cabbage looper, canker worms, case bearers, caseworms, corn earworm, diamondback moth, fruit worms, grapeleaf skeletonizer, gypsy moth, hornworms, imported cabbageworm, navel orangeworm, soybean looper, spruce budworm, tent caterpillar, tip moths, tent caterpillars, tobacco budworm, tobacco hornworm, tomato pinworm and tussock moth; Armyworms including beet armyworm, fall armyworm, lawn armyworm, southern armyworm and yellow striped armyworm; webworms and leaf perforators.
COLEOPTERA	beetles, grubs and weevils including Asian long-horned beetle, bark beetles, black vine weevil, Colorado potato beetle, elm bark beetle, European chafer, flea beetles, Japanese beetle, June beetle, leaf beetles, Mexican bean beetle, Northern masked chafer, rose chafer and Southern masked chafer and twig girders.
DIPTERA*	Flies including Caribbean fruit fly, cherry maggots, crane fly, fungus gnat , Hessian fly, oriental fruit fly, Mediterranean fruit fly, marsh crane flies, melon fly, shore fly and walnut husk fly; leafminers including citrus leafminers and serpentine leafminers.
	*Not intended for use on public health pests
THYSANOPTERA	thrips including citrus thrips, flower thrips, gladiolus thrips, onion thrips, thrips palmi and Western flower thrips.
ACARINA*	mites including, red spider mites, brown mite, clover mite, conifer spider mite: European red mite, spruce spider mite, and two-spotted spider mite.
	*Not intended for use on public health pests
ORTHOPTERA	crickets; grasshoppers; locusts
HYMENOPTERA*	sawflies including European sawflies, pear sawflies, red-headed pine sawflies, yellow-headed pin sawflies.
	*Not intended for use on public health pests
NEMATODA	nematodes (suppression)

FOR USE ON ORNAMENTALS AND LANDSCAPE PLANTINGS

Ornamental Plants and Flowers:	Actinopteris, African violets*, ageratum, aglaonema, Algerian ivy, allamanda, alocaasia, amaranthus, anthurium, aphelandra, arborvitae, Artemisia, aster, aucuba ilex, azalea, baby's breath, begonia, Boston fern, bougainvillea, boxwood, brachycome, cacti, calabrese, caladium, calathea, calendula, calla, camellia, carnation, ceanothus, chrysanthemum, cineraria, coleus, columbine, cotoneaster, cyclamen, daffodil, dahlia, daisy, daylily, delphinium, dianthus, dieffenbachia, dogwood, dusty miller, Easter lily, English ivy, euphorbia, fern, ficus, foliage plants, foxglove, freesia, fuschia, gaillardia, gardenia, geranium, gerbera, gladiola, gloxinia, gypsophilla, hederia, hibiscus, hyacinth, hydrangea, ilex, impatiens, iris, ivy, jasmine, lilac, lily, maidenhair fern, mandevilla, marigold, narcissus, nasturtium, orchid*, pansy, pelargonium, peony, peperomia, petunia, philodendron, phlox, photinia, pinks, pittosporum, poinsettia*, pothos, portulaca, primrose, pyracantha, rhododendron, rose*, rosemary, rubber plant, salvia, schefflera, sedum, sempervivum, snapdragon, spathiphyllum, stock, syngonium, tulip, verbena, vinca, wandering jew, yucca, zinnia *Please note that when making applications to these species, spotting of plant foliage and blooms is possible.
Ornamental Trees and Shrubs:	Andromeda, arborvitae, ash, Austrian pine, azalea, beech, birch, birdsnest spruce, blue spruce, bougainvillea, boxwood, butternut, cedar, charmaecyparis, cherry, cotoneaster, crabapple, cyprus, dogwood, Douglas fir, elm, euonymus, firethorn, forsythia, hackberry, hawthorn, hemlock, hickory, holly, honey locust, horse chestnut, juniper, larch, laurel, lilac, linden, London planetree, magnolia, mandevilla, maple, mimosa, mountain ash, myrtle, oak, pachysandra, peach, photinia, pine, planetree, poplar, privet, purpleleaf wintercreeper, quince, sage, spruce, sycamore, white cedar, white pine, yew

Waxy bloom on certain ornamental plants will be reduced after an application.

Applications of GH NAMT will remove the glaucous 'blue' coloring from evergreens such as Colorado blue spruce and Koster spruce.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

FOR USE ON GARDEN CROPS, VEGETABLES, HERBS, SPICES, HEMP, FRUITS, AND BERRIES

Leafy Vegetables:	Broccoli, Brussel Sprouts, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce, Spinach
Root Vegetables:	Beet, Carrot, Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yams
Fruiting Vegetables:	Eggplant, Pepper, Tomatillo, Tomato
Cucurbit Vegetables:	Cucumber, Gourds (edible), Muskmelon, Pumpkin, Squash, Watermelon, including Cantaloupe, Casaba, Gherkins, Melons (including hybrids), Zucchini
Legume Vegetables:	Bean, Chickpea, Lentil, Pea
Bulb Vegetables:	Garlic, Onion, Shallot
Berries:	Blackberry, Blueberry, Raspberry, Strawberry, others include: Boysenberry, Currants, Dewberry, Elderberry, Gooseberry, Loganberry
Herb and Spices:	Chive, Dill, Fennel, Mustard, Sage, Sweet bay, others include: Anise, Balm, Basil, Black pepper, Borage, Caraway, Catnip, Chamomile, Coriander, Cumin, Curry Leaf, Dandelion, Fenugreek, Horehound, Hyssop, Marjoram, Marigold, Mint, Nasturtium, Pennyroyal, Peppermint, Rosemary, Savory, Spearmint, Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood.
Nut Trees:	Almond, Brazil nut, Filbert, Hickory nut, Pecan, Pistachios, Walnut
Other crops:	Hemp
Pome Fruits:	Apple, Quince, or Pear (Cornice varieties: DO NOT apply more than 24 fl oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale usage.)
Stone Fruits:	Apricot, Cherry, Nectarine, Peach, Plum
Citrus Fruits:	Grapefruit, Lemon, Lime, Orange others include: Citrus Citron, Mandarin (tangerine), Nectarine, Satsuma (orange mandarin), Tangerine

GH NAMT has been evaluated for phytotoxicity on a wide range of ornamentals and garden plants. However, since testing on all plant varieties is not feasible, test a small portion of the area to be treated for phytotoxicity before treating the entire area.

There are no restrictions on applying GH NAMT up to the time of harvest.

SPRAY PREPARATION

GH NAMT is an emulsifiable concentrate to be diluted with water.

This product forms an emulsion and can separate upon extended or prolonged standing. Re-agitate to assure uniformity of the spray mixture.

Prepare only the volume needed for the intended application, and use the spray mixture within 24 hours of preparation.

TANK MIXTURES

GH NAMT is an emulsifiable concentrate and is compatible with commonly used pesticides and fertilizers. Always check the physical compatibility using a jar test in the correct proportions if needed.

If a broader spectrum of control is required tank-mix GH NAMT with insecticides or miticides. If a rapid knockdown of heavy populations is necessary, then include an effective contact insecticide/miticide in combination with GH NAMT.

Always read and follow the directions for use, precautions and limitations for use on all product labels used in combination. Applications must follow the precautions and limitations of the most restrictive product label in the mixture. Do not exceed the dosage rates of any product.

Select the right companion products:

IPM uses a variety of control options including biological, chemical, and cultural practices. GH NAMT is botanical with growth regulator effect on insects and mites. Companion products include pyrethroids, spinosyns, microbial toxins, and chloronicotinyls that complement azadirachtin. Formulations of bifenthrin, spinosad, abamectins, and imidacloprid are effective for different pests. Select the product that has been proven to provide adequate performance for the pests you are trying to control.

Physical Incompatibility

Do not use GH NAMT with Captan, Bordeaux mixture, triphenyltin hydroxide, lime sulfur, Rayplex iron or other highly alkaline materials as they can cause phytotoxicity and/or reduced efficacy on some target pests. Phytotoxicity will occur if tank-mix combinations with-compounds known to be incompatible with oil-based formulations are used.

APPLICATION EQUIPMENT

Apply GH NAMT with hand-operated (manual) or power spray equipment suitable for low volume and/or high volume applications. Follow the recommendations of the equipment manufacturer when using backpack sprayers; hose-end sprayers, compression (pump-up) sprayers, and other sprayers suitable for foliar applications of insecticides.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

APPLICATION SCHEDULE

For the most effective control, apply GH NAMT when pests are expected to appear or as soon as possible after pests appear and are in immature stages. Spray at an interval of seven (7) to ten (10) days or as the situation warrants.

During high pest infestation levels or when canopy is dense use higher dosage (use) rates and increase the spray frequency. Spraying in the morning or evening hours. Repeat spraying if rain occurs within two to three hours of spraying. For additional guidance, consult with the state agricultural experiment station or local extension horticulturalist/arborist for information on tactics and windows of application.

APPLICATION METHODS

Apply GH NAMT as directed to any food or non-food crop up to and including the day of harvest, at a maximum rate of 1.33 fl. oz. per 1,000 sq. ft. per application.

Dilute GH NAMT with water at a rate of 0.5 - 4.0 tablespoons (Tbs) per gallon of water. For hose end sprayers, set the RATE PER GALLON at the dial setting of 1 to 4 Tbs. depending on the crop and pests. Use the lower RATE PER GALLON for low to moderate infestations and use the higher specified RATE PER GALLON for severe infestations.

FOLIAR APPLICATION

USE	SPRAY CONCENTRATION %	Amounts of GH NAMT	
		Fluid ounces (Tbs.) per quart	Fluid ounces (Tbs.) per gallon
Including trees, shrubs, flowers, conifers, evergreens, herbaceous, ornamentals, hemp, foliage plants, container-grown ornamentals & garden plants and groundcovers	Lower rate ranges of 0.25 - 0.75% vol/vol:	0.08 – 0.25 fl. oz (1/6 – 1/2 Tbs.)	0.32 – 1 fl. oz (2/3 – 2.0 Tbs.)
	Medium rate ranges of 0.75 – 1.25% vol/vol:	0.26 – 0.40 fl. oz (1/2 – 5/6 Tbs.)	2.0 – 1.6 fl. oz (2.0 – 3 1/3 Tbs.)
	Upper rate ranges of 1.25 – 1.70% vol/vol:	0.40 – 0.50 fl. oz (5/6 -1.0 Tbs.)	1.6 – 2.0 fl. oz (3 1/3 – 4 Tbs.)

DRENCH APPLICATION

Use GH NAMT as a soil drench for effective control of soil-borne insect larvae, including soil-borne larvae of foliar pests, such as fungus gnats, nematodes, or soil borne thrips. When applying as a drench, avoid excessive leaching.

Preventive applications as a soil drench may be warranted for certain pests. Soil drench applications of azadirachtin will have a slower rate of activity because of soil absorption when compared to foliar applications of azadirachtin. Target the initial application of a soil drench treatment to coincide with the early stages of young larvae and young nymphs.

Dilute GH NAMT with water for concentrations of 0.4 to 0.8% volume/volume. See use rate table below. Add the required amount of GH NAMT to a clean bucket with at least one-half of the water to be drenched. Agitate the mixture thoroughly and then fill with the remaining water and continue agitation until the product is thoroughly dispersed.

Drench the soil in the pot with one (1) pint of finished product dilution per 1.0 gallon of soil. For fungus gnats, use the 0.4% spray concentration. For mushroom fly maggot control, use the 0.6% volume/volume spray concentration. For leafminers and other difficult to control pests, use the 0.8% volume/volume spray concentration. Make two to three (2 - 3) applications at 10 -14 day intervals until pest pressure has ended. With high insect pressure make applications every 5 to 6 days. Additional applications of GH NAMT may be required with increased and prolonged pest infestation.

DILUTION TABLE FOR DRENCH APPLICATIONS

Gallons of Water	Amount of GH NAMT			Application Interval
	0.4%	0.6%	0.8%	
1 gallon	1 Tbs.	1.5 Tbs.	2.0 Tbs.	10 – 14 days
1 gallon	0.5 fl. oz.	0.8 fl. oz.	1.0 fl. oz.	10 – 14 days
5 gallons	2.5 fl. oz.	4.0 fl. oz.	5.0 fl. oz.	10 – 14 days
10 gallons	5.0 fl. oz.	8.0 fl. oz.	10.0 fl. oz.	10 – 14 days

RECIRCULATORY, AEROPONIC, AND HYDROPONIC APPLICATION

Use GH NAMT in recirculatory, aeroponic, or hydroponic systems for the control of foliar pests, soil borne insect larvae, including soil borne larvae of foliar pests such as fungus gnats, nematodes or soil borne thrips for interiorscapes, hydroponic, aeroponic and container plants.

Dilute GH NAMT with water for concentrations of 0.1 % to 0.8% volume/volume in a recirculatory or in a hydroponic liquid system. See use rate table below. Agitate the mixture thoroughly until the product is thoroughly dispersed.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

For fungus gnats, use the 0.6% volume/volume concentration. For mushroom fly maggot control, use the 0.6% volume/volume concentration. For leafminers and other difficult to control pests, use the 0.8% volume/volume concentration. Make two to three (2-3) applications at 10-14 day intervals until the pest pressure has ended. With high insect pressure applications make applications every 5 to 7 days. Additional applications of GH NAMT may be required with increased and prolonged pest infestation.

DILUTION TABLE FOR RECIRCULATORY, AEROPONIC, AND HYDROPONIC APPLICATIONS

Gallons of Water	Amount of GH NAMT					Application Interval
	0.1%	0.2%	0.4%	0.6%	0.8%	
1 gallon	1/4 Tbs.	1/2 Tbs.	1 Tbs.	1.5 Tbs.	2 Tbs.	7 – 14 days
1 gallon	0.14 fl. oz	0.25 fl. oz	0.5 fl. oz	0.8 fl. oz	1.0 fl. oz	7 – 14 days
5 gallons	0.7 fl. oz	1.3 fl. oz	2.5 fl. oz	4.0 fl. oz	5.0 fl. oz	7 – 14 days
10 gallons	1.4 fl. oz	2.6 fl. oz	5.0 fl. oz	8.0 fl. oz	10.0 fl. oz	7 – 14 days

Preventive applications as a recirculatory system application may be warranted for certain pests.

GH NAMT can also be applied through sub-surface treatment equipment. Always follow manufacturer's use directions.

FOLIAR APPLICATION – SPECIFIC PESTS OF GARDEN CROPS, VEGETABLES, HERBS AND SPICES, HEMP, BERRIES AND FRUIT.

CROP	PESTS such as:	Dilution Rate for Sprayers	
		fl. ozs of product per 1000 sq. ft.	Tbs. of product per 1.0 gallon of water
Leafy vegetables: Broccoli, Brussel Sprouts, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce, Spinach	Leafrollers, Cutworms, Loopers, Armyworms	0.19 – 0.96 fl. oz	¾ Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Beetles, Weevils, Flies, Thrips, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Root Vegetables: Beet, Carrot, Horseradish, Parsnip, Potato, Radish, Sweet potato, Turnip, Yams	Beetles, Weevils	0.11 – 0.25 fl. oz	1/2 Tbs. – 1 1/2 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Thrips, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Fruiting Vegetables: Eggplant, Pepper, Tomatillo, Tomato	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Cucurbit Vegetables: Cucumber, Gourd (edible), Muskmelon, Pumpkin, Squash, Watermelon, Including Cantaloupe, Casaba, Gherkins, Melons (including hybrids), Zucchini	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Legume Vegetables: Bean, Chickpea, Lentil, Pea	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Bulb Vegetables: Garlic, Onion, Shallot	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Berries including: Blackberry, Blueberry,	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional.] Text separated by / denotes and/or options.

Raspberry, Strawberry, others include: Boysenberry, Currants, Dewberry, Elderberry, Gooseberry, Loganberry	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Herbs and Spices: Chive, Dill, Fennel, Mustard, Sage, Sweet bay, others include: Anise, Balm, Basil, Black pepper, Borage, Caraway, Catnip, Chamomile, Coriander, Cumin, Curry leaf, Dandelion, Fenugreek, Horehound, Hyssop, Marjoram, Marigold, Mint, Nasturtium, Pennyroyal, Peppermint, Rosemary, Savory, Spearmint, Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Other crops: Hemp	True Bugs, Leafhoppers, Whiteflies, Aphids, Mites, Beetles, Weevils, Flies, Thrips	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
	Leafrollers, Cutworms, Loopers, Armyworms	0.19 – 0.96 fl. oz	¾ Tbs. – 4 Tbs./gal
Nut Trees: Almond, Brazil nut, Filbert, Hickory nut, Pecan, Pistachios, Walnut	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Pome Fruits: Apple, Quince, or Pear (Comice varieties: DO NOT apply more than 24 fl. oz/A. DO NOT apply after pink stage of flowering; test small areas of other varieties of pears for plant safety prior to full scale usage.)	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Stone Fruits: Apricot, Cherry, Nectarine, Peach, Plum	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal
Citrus Fruits: Grapefruit, Lemon, Lime, Orange, others include: Citrus Citron, Mandarin (tangerine), Nectarine, Satsuma (orange mandarin), tangerine	Beetles, Weevils	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	Thrips,	0.29 – 0.96 fl. oz	2 Tbs. – 4 Tbs./gal
	True Bugs, Leafhoppers, Whiteflies, Aphids, Leafrollers, Cutworms, Loopers, Armyworms, Flies, Mites	0.24 – 0.96 fl. oz	1 Tbs. – 4 Tbs./gal

Bold, italicized text is information for the reader and is not part of the label.
[Bracketed information is optional.] Text separated by / denotes and/or options.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Keep from freezing. To be stored in original container and placed in areas inaccessible to children.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: If empty: Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

Bold, italicized text is information for the reader and is not part of the label.
 [Bracketed information is optional.] Text separated by / denotes and/or options.

OPTIONAL TEXT AND CLAIMS

GENERAL CLAIMS

8. For organic gardening
9. OMRI Logo
10. Organic Materials Review Institute (OMRI) Certified.
11. Low-odor formulation
12. Low (Mild) odor
13. [Questions/Comments [Call]] ***insert company phone number***
14. [Visit [our website]] ***insert company website***

USE SITE CLAIMS

6. Can be applied the day of harvest
7. For use on a wide variety of trees, shrubs, flowers, fruit and nut trees, hemp, garden vegetables and plants
8. For use on outdoor and greenhouse food and non-food crops, ornamental flowers, trees, shrubs and plants, hemp and landscapes
9. For use on roses, vegetables, fruits, nuts, flowers, houseplants, foliage, hemp, trees, and shrubs
10. Can be used on [the following plants:] vegetables; fruits; herbs; nuts; ornamental trees, shrubs, hemp, flowering plants; houseplants, and tropical plants [grown in and around home gardens or home greenhouses]

EFFICACY CLAIMS

7. Controls mites, caterpillars, whiteflies, thrips, aphids, and other insects as listed on this label.
8. Controls chewing and sucking insects
9. Broad-spectrum control
10. Insect repellent/insecticide
11. Miticide
12. Kills listed insects such as mites, thrips, psyllids, leafhoppers, lepidoptera larvae, white flies, borers, mealybugs and true bugs

U.S. Patent No. X,XXX,XXX

© 20XX [Licensed by] ***insert company name***. [World rights reserved.]

Sold by: ***insert company name and address***

Made in/Product of _____

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK, OR FIRE), CALL CHEMTREC® (800) 424-9300.
 For other product information, contact General Hydroponics or see Material Safety Data Sheet.

NOTICE ON CONDITIONS OF SALE

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.
 Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of General Hydroponics. To the extent consistent with applicable law, all such risks are assumed by the Buyer and User.

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified. Country: United States of America.