

## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

August 10, 2018

Ashwini Kulkarni Regulatory Specialist SBM Life Science Corporation 1001 Winstead Drive, Suite 500, Cary, North Carolina, 27513

Subject: Notification per PRN 98-10 – minor label revisions to correct misspellings, replace brackets with parentheses, add trademark logos, add a warranty statement, add back erroneously omitted marketing statements Product Name: Lawn Herbicide JD Concentrate EPA Registration Number: 92564-4 Application Date: 12/19/17 Decision Number: 543348

Dear Ashwini:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you have any questions, you may contact Beth Benbow via email at Benbow.bethany@epa.gov.

Sincerely,

astryn V. Wontague

Kathryn Montague, Product Manager 23 Herbicide Branch Registration Division (7505P) Office of Pesticide Programs (FRONT PANEL)

NOTIFICATION 92564-4 The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

# Lawn Herbicide JD Concentrate

(ABN) Bayer Advanced Weed Killer for Lawns Concentrate; Bayer Advanced Lawn Weed Killer Concentrate; Bayer Advanced Southern Weed Killer for Lawns Concentrate; Bianova Advanced<sup>™</sup> Weed Killer for Lawns Concentrate; BioAdvanced<sup>™</sup> Weed Killer for Lawns Concentrate; BioAdvanced<sup>™</sup> Science-Based Solutions Weed Killer for Lawns Concentrate; Fortifend Advanced<sup>™</sup> Weed Killer for Lawns Concentrate; Bianova Advanced<sup>™</sup> Lawn Weed Killer Concentrate; BioAdvanced<sup>™</sup> Lawn Weed Killer Concentrate; BioAdvanced<sup>™</sup> Science-Based Solutions Lawn Weed Killer Concentrate; Fortifend Advanced<sup>™</sup> Lawn Weed Killer Concentrate; Bianova Advanced<sup>™</sup> Southern Weed Killer for Lawn Weed Killer Concentrate; Bianova Advanced<sup>™</sup> Southern Weed Killer for Lawns Concentrate; BioAdvanced<sup>™</sup> Southern Weed Killer for Lawns Concentrate; BioAdvanced<sup>™</sup> Science-Based Solutions Southern Weed Killer for Lawns Concentrate; Fortifend Advanced<sup>™</sup> Southern Weed Killer for Lawns Concentrate; BioAdvanced<sup>™</sup> Science-Based Solutions Southern Weed Killer for Lawns Concentrate; Fortifend Advanced<sup>™</sup> Southern Weed Killer for Lawns Concentrate

This label contains separate label statements for **Outdoor Residential Pump Sprayer/ Hose-end Sprayer Concentrate** use product and **Outdoor Residential Ready-To-Spray** use product.

- Effective -Listed [\*] Weed Protection
- Effective Listed [\*] Broadleaf Weed Protection
- Effectively Protects Lawns from Listed [\*] Broadleaf Weeds
- Kills Major Listed [\*] Broadleaf Weeds
- Kills Listed [\*] Weeds Won't Harm Lawns<sup>†</sup>
- Kills to the Root
- Protects Against Major Listed [\*] Broadleaf Weeds
- Tough Listed [\*] Weed Protection
- Listed [\*] Weed Protection System
- Systemic
- 1 Quart treats up to 16,000 square feet
- Convenient MEASURE-UP™ Cup
- Kills Listed [\*] Broadleaf weeds in St. Augustine & other Southern Lawns
- For St. Augustine & other Southern Lawns

Marketing Claim for Brand Name Change: New Name. Same Great Results. New Name. Same Familiar Formula. <u>New Name. Same Performance.</u>

- Kills Over 200 Listed [\*] Broadleaf Weeds (as listed on this label)
- Won't Harm Your Lawn<sup>†</sup>
- Rainproof in hours
- 100% Satisfaction Guaranteed\* (or your money back)
- Rainproof in 1 hour!
- Attach to Hose & Spray
- Kills 200+ Listed [\*] Weeds (as listed on this label)
- [optional advertising claims when product is sold in "low-water" hose-end sprayers, as determined by testing]
- Water [Saver][Saving] Formula
- Water [Saver][Saving] Applicator
- Water [Saver][Saving] Sprayer
- Uses Up To [80][XX]% Less Water\*
- [with footnote disclaimer] \*vs leading Southern weed hose end sprayers
- won't harm lawns†
- †when used as directed

- Bonus [X] oz size [X]% more than the [X] oz size
- Now covers [XXX] [square] [sq] feet
- Bonus [X] oz size

#### Kills the Root

- [For] [Use] [On] [Most] [Many] [Majority of] [Northern] [and] [Southern] [Turfs] [Lawns] [Types] [Varieties]\*
- Use on Northern Turfs: Fescue, Kentucky Bluegrass & [and] Perennial Rye [grass]
- Use on Southern Turfs: Bermuda, Zoysia, Centipede & [and] St. Augustine [grass]

<u>\*do not use on Carpetgrass,</u> Dichondra or Floratam variety of St. Augustinegrass

\*listed (Note to PM: The word "listed" can appear as a footnote with the appropriate phrase, such as \*listed plants/pests/weeds)

| ACTIVE INGREDIENTS:            |         |
|--------------------------------|---------|
| 2,4-D, dimethylamine salt      |         |
| Mecoprop-p, dimethylamine salt | 1.83 %  |
| Dicamba, dimethylamine salt    |         |
| OTHER INGREDIENTS:             |         |
| TOTAL                          | 100.00% |

Net Contents: 2 fl oz (59 mL) or 4 fl oz (118 mL) or 16 fl oz (473 mL) or 24 fl oz (709 mL) or 32 fl oz (946 mL), 40 oz (1.18L), or 1 Gal (3.78 L)

# KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel Booklet for Additional Ingredients and Precautionary Statements

Editorial Note - [Bracketed text] is optional language

# Lawn Herbicide JD Concentrate

(ABN) Bayer Advanced Weed Killer for Lawns Concentrate; Bayer Advanced Lawn Weed Killer Concentrate; Bayer Advanced Southern Weed Killer for Lawns Concentrate; Bianova Advanced<sup>™</sup> Weed Killer for Lawns Concentrate; BioAdvanced<sup>™</sup> Weed Killer for Lawns Concentrate; BioAdvanced<sup>™</sup> Science-Based Solutions Weed Killer for Lawns Concentrate; Fortifend Advanced<sup>™</sup> Weed Killer for Lawns Concentrate; Bianova Advanced<sup>™</sup> Lawn Weed Killer Concentrate; BioAdvanced<sup>™</sup> Lawn Weed Killer Concentrate; BioAdvanced<sup>™</sup> Science-Based Solutions Lawn Weed Killer Concentrate; Fortifend Advanced<sup>™</sup> Lawn Weed Killer Concentrate; Bianova Advanced<sup>™</sup> Southern Weed Killer for Lawns Concentrate; BioAdvanced<sup>™</sup> Southern Weed Killer for Lawns Concentrate; BioAdvanced<sup>™</sup> Science-Based Solutions Southern Weed Killer for Lawns Concentrate; Fortifend Advanced<sup>™</sup> Southern Weed Killer for Lawns Concentrate; BioAdvanced<sup>™</sup> Science-Based Solutions Southern Weed Killer for Lawns Concentrate; Fortifend Advanced<sup>™</sup> Southern Weed Killer for Lawns Concentrate

- Effective Listed [\*] Weed Protection
- Effective Listed [\*] Broadleaf Weed Protection
- Effectively Protects Lawns from Listed [\*] Broadleaf Weeds
- Kills Major Listed [\*] Broadleaf Weeds
- Kills Listed [\*] Weeds Won't Harm Lawns<sup>†</sup>
- Kills to the Root
- Protects Against Major Listed
   [\*] Broadleaf Weeds
- Tough Listed [\*] Weed
   Protection
- Listed [\*] Weed Protection
   System
- Systemic
- 1 Quart treats up to 16,000 square feet
- Convenient MEASURE-UP™ Cup
   Kills Listed [\*] Broadleaf weeds in St. Augustine &
- other Southern Lawns For St. Augustine & other Southern Lawns
- Kills Over 200 Listed [\*] Broadleaf Weeds (as listed on this label)
- Won't Harm Your Lawn<sup>†</sup>
- Rainproof in hours

- 100% Satisfaction Guaranteed\* (or your money back)
- Rainproof in 1 hour!
- Kills 200+ Listed [\*] Weeds (as listed on this label)
- [optional advertising claims when product is sold in "lowwater" hose-end sprayers, as determined by testing]
- Water [Saver][Saving] Formula
- Water [Saver][Saving]
   Applicator
- Water [Saver][Saving] Sprayer
- Uses Up To [80][XX]% Less
  Water\*
- [with footnote disclaimer] \*vs leading Southern weed hose end sprayers
- won't harm lawns†
- †when used as directed
- Bonus 40 oz size 25% more than the 32 oz size
- Now covers [XXX] sq ft
- Bonus [xx] oz. size [x]% more than [X] oz size
- Bonus [x] oz size
- [Free] refill [ for 1.3 gallon Ergo-Grip<sup>™</sup> Weed Killer for [Southern] Lawns Ready-To-Use]
- [Free] bonus refill
- [xxx] oz makes [xxx] gallon[s] of spray

- Refills 1.3 gallon Ergo–Grip<sup>™</sup> Weed Killer for [Southern] Lawns Ready-To-Use
- Easy hold
- EZ Spray
- One-touch sprayer
- ●\_\_\_EZ<mark>-G-g</mark>rip<mark>™</mark>
- <u>EZ-Grip One-Touch</u> Sprayer™
- Ergo<u>-<mark>G-g</mark>rip</mark>™</u>
- Simple spray
- One-touch
- Spray n go
- Thumb switch
- Handy spray
- EZ touch
- One-touch application
- [For] [Use] [On] [Most] [Many] [Majority of] [Northern] [and] [Southern] [Turfs] [Lawns] [Types] [Varieties]\*
- Use on Northern Turfs: Fescue, Kentucky Bluegrass & [and] Perennial Rye [grass]
- Use on Southern Turfs: Bermuda, Zoysia, Centipede & [and] St. Augustine [grass]

\*<u>do not use on Carpetgrass.</u> Dichondra or Floratam variety of St. Augustinegrass Marketing Claim for Brand Name Change: New Name. Same Great Results. New Name. Same Familiar Formula. New Name. Same Performance.

\*listed (Note to PM: The word "listed" can appear as a footnote with the appropriate phrase, such as \*listed plants/pests/weeds)

| ACTIVE INGREDIENTS:            |  |
|--------------------------------|--|
| 2,4-D, dimethylamine salt      |  |
| Mecoprop-p, dimethylamine salt |  |
| Dicamba, dimethylamine salt    |  |
| OTHER INGREDIENTS:             |  |
| TOTAL                          |  |

Net Contents: 2 fl. oz. (59 mL) or 4 fl oz (118 mL) or 16 fl oz (473 mL) or 24 fl oz (709 mL) or 32 fl oz (946 mL), 40 oz (1.18L), or 1 Gal (3.78 L)

# KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel Booklet for Additional Ingredients and Precautionary Statements Editorial Note – [Bracketed text] is optional language

#### (BACK PANEL)

**OPEN** Resealable Label for Directions & Precautions **PRESS TO RESEAL** 

This product kills over 200 types of [listed] [\*] broadleaf weeds in St. Augustine and other Southern lawns.

This formula kills Dollarweed, Spurge, Oxalis and Dandelion without harming your lawn\*. Controlling [listed] [\*] weeds in your lawn has never been easier. (\*when used as directed.)

[modify representative weeds or add approved marketing claims as needed]

#### [READ THE LABEL FIRST!]

#### QUICK FACTS

- Kills over 200 listed [\*] broadleaf weeds (as listed on this label)
- Won't harm your lawn<sup>†</sup>
- Treats up to 16,000 sq ft

| KILLS           | Over 200 listed [*] broadleaf weeds such as Dollarweed, Spurge and Oxalis, Dandelion, Clover, Chickweed and other listed weeds [or choose from marketing claims]            |
|-----------------|---|
| WHERE TO USE    | On Lawns  |
|                 | ំ (Do not use on Carpetgrass, Dichondra or Floratam variety of St. Augustinegrass.)   |
| WHEN TO USE     | When weeds are actively growing   |
| HOW MUCH TO USE | Northern Lawns: Fescue, Kentucky Bluegrass, and Perennial Ryegrasses  |
|                 | and   |
|                 | Hardy Southern Lawns: Bermuda, Buffalo and Zoysia grasses<br>2 fl oz per 500 sq ft  |
|                 | Southern lawns: Bahia, Centipede and St. Augustinegrass grasses<br>1 fl oz per 500 sq ft  |
| <b>7</b> -      | For questions or comments, call toll-free 1-877-229-3724 or visit us on-line at<br>[www.bianovaadvanced.com][www.BioAdvanced.com][www.BioAdvancedScienceBasedSolutions.com] |
|                 | [www.fortifendadvanced.com]   |

#### FOR OUTDOOR RESIDENTIAL USE ONLY

#### **ACTIVE INGREDIENTS**

| 2,4-D, dimethylamine salt      | 7.59 % |
|--------------------------------|--------|
| Mecoprop-p, dimethylamine salt | 1.83 % |
| Dicamba, dimethylamine salt    | 0.84 % |
| OTHER INGREDIENTS*             | 9.74 % |
| TOTAL                          | 0.00 % |

[\* Other ingredients include: Water and surfactants (to keep the active ingredients dispersed in water).]

#### THIS PRODUCT CONTAINS:

0.00427 lb 2,4-D acid equivalent per fl oz or 6.30% 0.00102 lb Mecoprop-p acid equivalent per fl oz or 1.51% 0.00047 lb Dicamba acid equivalent per fl oz or 0.69% Isomer specific by AOAC Method

EPA Reg. No. 92564-4

EPA Est. No. \_-AA-\_

| FIRST AID              |   |  |
|------------------------|---|--|
| If swallowed           | <ul> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything to an unconscious person.</li> </ul> |  |
| If in eyes             | <ul> <li>Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>  |  |
| If on skin or clothing | <ul> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>  |  |
| If inhaled             | <ul> <li>Move person to fresh air.</li> <li>If not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth to-mouth, if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>   |  |

**[Active Ingredients:** 7.59% 2,4-D, dimethylamine salt **[**CAS # 2008-39-1**]]**, 1.83% Mecoprop-p, dimethylamine salt **[**CAS # 66423-09-4**]**], and 0.84% Dicamba, dimethylamine salt **[**CAS # 2300-66-5**]**].

# PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

- Harmful if absorbed through skin or if swallowed.
- · Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

After using this product:

- Rinse gloves before removing.
- · Remove clothing and launder separately before reuse.
- · Wash hands thoroughly with soap and water.
- If product gets on clothing, remove clothing as soon as possible and shower.

## **ENVIRONMENTAL HAZARDS**

- This product is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.
- Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.
- This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

# **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read and follow all label directions.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes. Do not apply through any type of irrigation system.

|   | · · · · · · · · · · · · · · · · · · ·  |  |  |  |
|---|--|--|--|--|
| BEFORE YOU USE<br>Read and follow<br>directions when using. | <ul> <li>Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn or garden area only.</li> <li>Spray when most of the day is expected to remain cooler than 90° F.</li> <li>Do not apply as a fine mist because of potential injury to desirable plants.</li> </ul> |  |  |  |
|   | The maximum number of broadcast applications per treatment site is two per year.   |  |  |  |
| HOW TO USE  | Determine amount of product to use based on grass type and size of area to be treated.   |  |  |  |
| Shake Well  | Prepare product for use according to How to Apply.   |  |  |  |
| Before Use  | Apply to lawn.   |  |  |  |
| HOW MUCH TO APPLY   | Northern Lawns:  |  |  |  |
|   | <ul> <li>Fescue, Kentucky Bluegrass, and Perennial Rye</li> </ul>  |  |  |  |
|   | 2 fl oz (4 Tbs) per gallon of water  |  |  |  |
|   | Hardy Southern Lawns:  |  |  |  |
|   | Bermuda, Buffalo and Zoysia grasses  |  |  |  |
|   | 2 fl oz (4 Tbs) per gallon of water  |  |  |  |
|   | Southern Lawns:  |  |  |  |
|   | Bahia, Centipede, and<br>St. Augustinegrass  |  |  |  |
|   | 1 fl oz (2 Tbs) per gallon of water  |  |  |  |
|   | For lawns with mixed grass types, use the lowest rate.   |  |  |  |
|   | Each gallon of solution treats 500 sq ft   |  |  |  |
|   | <b>TIP</b> : Measure length and width of area to be treated in feet. Multiply these 2 numbers together. This is the area to be treated in square feet (sq ft).   |  |  |  |
|   | Application Restrictions:  |  |  |  |
|   | • Limited to 2 applications per year with a 30 day interval between applications   |  |  |  |
|   | • Maximum single application rate of 8.0 fl oz/1,000 sq ft (1.47 lbs 2,4-D ae, 0.35 lb MCPP-p ae and 0.16 lb dicamba ae per acre per application)  |  |  |  |
|   | • Maximum seasonal rate of 16.0 fl oz/1,000 sq ft (2.94 lbs 2,4-D ae, 0.71 lb MCPP-p ae and 0.33 lb dicamba ae per acre per application)   |  |  |  |
| HOW TO APPLY  | Pump Style Sprayers  |  |  |  |
|   | 1. Add the appropriate amounts of water and concentrate to the sprayer tank.   |  |  |  |
|   | [For 2 or 4 fl oz container: Pour entire contents into spray tank and fill with water to [1] [2] gallon mark.]   |  |  |  |
|   | 2. Close sprayer, shake well and pressurize.   |  |  |  |
|   | 3. Adjust nozzle to a coarse spray pattern.  |  |  |  |
|   | 4. Large Areas – Apply evenly over area.   |  |  |  |
|   | Individual Weeds – Hold tip of wand about 12 inches from weed, aim at center of weed and spray.  |  |  |  |
|   | 5. Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.  |  |  |  |
|   | Dial Style Hose-End Sprayers   |  |  |  |
|   |  |  |  |  |

|                           | [Best for spraying large areas such as the entire lawn.  |
|---------------------------|--|
|                           | 1. Fill sprayer jar with enough product for the area to be treated. DO NOT ADD WATER.  |
|                           | 2. Close sprayer and set dial as recommended below:  |
|                           | TIP: The numbers on dial indicate amount of concentrate per gallon of spray.   |
|                           | - Ortho Dial 'N Spray Settings:  |
|                           | 1.5 OZ for Bermudagrass, Buffalograss, Fescue, Kentucky Bluegrass, Perennial Ryegrass, Zoysiagrass   |
|                           | 4 TSP for Bahiagrass, Centipedegrass, St. Augustinegrass (not for use on Carpetgrass, Dichondra or<br>Floratam variety of St. Augustinegrass)  |
|                           | - Chameleon Hose-End Sprayer Settings  |
|                           | 4 TSP for Bermudagrass, Buffalograss, Fescue, Kentucky Bluegrass, Perennial Ryegrass, Zoysiagrass  |
|                           | 2 TSP Bahiagrass, Centipedegrass, St. Augustinegrass (not for use on Carpetgrass, Dichondra or Floratam variety of St. Augustinegrass)   |
|                           | 3. Hook to garden hose and apply. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.  |
|                           | 4. Pour any unused product back into this original container.]   |
|                           |  |
|                           | [This concentrate can also be used to refill Bayer Advanced Weed Killer for Southern Lawns [1.3] Gallon<br>Ready-To-Use [Ergo <mark>–</mark> Grip <mark>™</mark> ] Sprayer]  |
|                           | Concentrate Plus Water Hose-End Sprayer  |
|                           | 1. Pour concentrate into sprayer jar to the required fluid ounce level.  |
|                           | 2. Add the appropriate amount of water to the gallons level.   |
|                           | 3. Replace sprayer top and shake well before applying.   |
|                           | 4. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.   |
| FOR BEST RESULTS          | Apply when weeds are young and actively growing.   |
|                           | If the lawn is dry, water 24 hours before using this product.  |
|                           | Rain or watering one hour after application will not wash away effectiveness.  |
|                           | Apply 1 to 3 days after mowing.  |
|                           | Certain hard-to-kill weeds may need to be spot-treated in 30 days.   |
|                           | Do not apply to new lawns until the grass has reached a height of at least 2 inches.   |
|                           | Bare spots may be seeded 3 to 4 weeks after application.   |
| HOW THIS PRODUCT<br>WORKS | This product is a selective herbicide - it selectively kills weeds in the lawn but won't harm the lawn itself. It is a blend of proven weed killers designed to get broadleaf weeds such as Dollarweed, Spurge and Oxalis. |
|                           | [modify representative weeds or add approved marketing claims as needed]   |

| Do not allow people (other than applicator) or pets on treated area during application.<br>Do not allow people or pets to re-enter the treated area until spray has dried. |
|--|
|  |

| CONTROLS | Annual yellow sweetclover | Cockle            | Mugwort                     | Spiny amaranth         |
|----------|---------------------------|-------------------|-----------------------------|------------------------|
| KILLS]   | Aster                     | Cocklebur         | Musk thistle                | Spiny cocklebur        |
|          | Austrian fieldcress       | Common mullein    | Mustard                     | Spiny sowthistle       |
|          | Bedstraw                  | Creeping Charlie  | Narrowleaf plantain         | Spotted catsear        |
|          | Beggarticks               | Creeping Jenny    | Narrowleaf vetch            | Spotted spurge         |
|          | Betony, Florida           | Cudweed           | Nettle                      | Spurweed               |
|          | Bindweed, field           | Curly dock        | Orange hawkweed             | Stinging nettle        |
|          | Bird vetch                | Daisy, English    | Oriental cocklebur          | Strawberry, India mock |
|          | Bitter wintercress        | Daisy, fleabane   | Oxalis                      | Tall nettle            |
|          | Bittercress, hairy        | Daisy, oxeye      | Parsley-piert               | Tall vervain           |
|          | Bitterweed                | Dandelion         | Parsnip                     | Tansy mustard          |
|          | Black-eyed Susan          | Dichondra         | Pearlwort                   | Tanweed                |
|          | Black medic               | Dogbane           | Pennycress                  | Thistle                |
|          | Black mustard             | Dogfennel         | Pennywort                   | Trailing crownvetch    |
|          | Blackseed plantain        | Dollarweed        | Pepperweed                  | Tumble mustard         |
|          | Blessed thistle           | Elderberry        | Pigweed                     | Tumble pigweed         |
|          | Bloodflower milkweed      | False dandelion   | Pineywoods bedstraw         | Velvetleaf             |
|          | Blue lettuce              | False flax        | Plains coreopsis (tickseed) | Venice mallow          |
|          | Blue vervain              | False sunflower   | Plantain                    | Virginia buttonweed    |
|          | Bracted plantain          | Fiddleneck        | Poison ivy                  | Virginia creeper       |
|          | Brassbuttons              | Florida pusley    | Poison oak                  | Virginia pepperweed    |
|          | Bristly oxtongue          | Frenchweed        | Pokeweed                    | Water pennywort        |
|          | Broadleaf dock            | Galinsoga         | Poorjoe                     | Wavyleaf bullthistle   |
|          | Broadleaf plantain        | Goathead          | Prairie sunflower           | Western clematis       |
|          | Broomweed                 | Goldenrod         | Prickly lettuce             | Western salsify        |
|          | Buckhorn                  | Ground ivy        | Prickly sida                | White mustard          |
|          | Buckhorn plantain         | Gumweed           | Prostrate knotweed          | Wild aster             |
|          | Bulbous buttercup         | Hairy fleabane    | Prostrate pigweed           | Wild buckwheat         |
|          | Bull thistle              | Hawkweed          | Prostrate spurge            | Wild carrot            |
|          | Bullnettle                | Healall           | Prostrate vervain           | Wild four-o'clock      |
|          | Burclover                 | Heartleaf drymary | Puncturevine                | Wild garlic            |
|          | Burdock                   | Heath aster       | Purslane, common            | Wild geranium          |
|          | Burning nettle            | Hedge bindweed    | Ragweed                     | Wild lettuce           |
|          | Burweed                   | Hedge mustard     | Red sorrel                  | Wild marigold          |
|          | Buttercup                 | Hemp              | Redroot pigweed             | Wild mustard           |
|          | Buttonweed                | Henbit            | Redstem filaree             | Wild onion             |
|          | Canada thistle            | Hoary cress       | Rough cinquefoil            | Wild parsnip           |
|          | Carolina geranium         | Hoary plantain    | Rough fleabane              | Wild radish            |
|          | Carpetweed                | Hoary vervain     | Russian pigweed             | Wild rape              |
|          | Catchweed bedstraw        | Horsenettle       | Russian thistle             | Wild strawberry        |
|          | Catnip                    | Jimsonweed        | Scarlet pimpernel           | Wild sweet potato      |
|          | Catsear                   | Knawel            | Scotch thistle              | Wild vetch             |
|          | Chickweed, common         | Knotweed          | Sheep sorrel                | Woodsorrel             |
|          | Chickweed, mouseear       | Kochia            | Shepherdspurse              | Woolly croton          |
|          | Chicory                   | Lambsquarters     | Slender plantain            | Woolly morningglory    |
|          | Cinquefoil                | Lespedeza         | Smallflower galinsoga       | Woolly plantain        |
|          | Clover, crimson           | Mallow            | Smooth dock                 | Wormseed               |
|          | Clover, hop               | Matchweed         | Smooth pigweed              | Yarrow                 |
|          | Clover, red               | Mexicanweed       | Sorrel                      | Yellow rocket          |
|          | Clover, strawberry        | Milk vetch        | Sowthistle                  | Yellowflower pepperwee |
|          | Clover, sweet             | Morningglory      | Spanishneedles              |                        |
|          | Clover, white             | Mouseear hawkweed | Speedwell                   |                        |

### STORAGE AND DISPOSAL

| Do not contaminate food, or feed by sto       | rage and disposal.   |
|---|--|
| Pesticide Storage:                            | Store in original container in a cool dry place, out of the reach of children, preferably in a locked storage cabinet. Protect from freezing.  |
| Pesticide Disposal and<br>Container Handling: | <ul> <li>If Empty:</li> <li>Non-refillable container. Do not refill or reuse empty container.</li> <li>Place empty container in trash or offer for recycling if available.</li> <li>If Partly Filled:</li> <li>If it is necessary to dispose of unused product, please call your local solid waste agency or you may call toll-free 1-877-229-3724 for disposal instructions.</li> <li>Never place unused product down any indoor or outdoor drain.</li> </ul> |

**NOTICE:** Research and testing have determined that the "Directions For Use" are appropriate for the proper use of this product under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

\*Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

BioAdvanced<sup>™</sup> [Fortifend Advanced<sup>™</sup>] [Bianova Advanced<sup>™</sup>] is a trademark of SBM Life Science Corp.

EZ-Grip<sup>™</sup> and Ergo-Grip<sup>™</sup> are trademarks of SBM Life Science Corp.

PRODUCED FOR [DISTRIBUTED BY]

SBM Life Science Corp. 1001 Winstead Drive, Suite 500 Cary, North Carolina 27513

## (Ready - To - Spray label) (FRONT PANEL)

# Lawn Herbicide JD Concentrate

(ABN) Bayer Advanced Weed Killer for Lawns Ready-to-Spray; Bayer Advanced Lawn Weed Killer Ready-to-Spray; Bayer Advanced Southern Weed Killer for Lawns Ready-to-Spray; Bianova Advanced<sup>™</sup> Weed Killer for Lawns Ready-to-Spray; BioAdvanced<sup>™</sup> Weed Killer for Lawns Ready-to-Spray; BioAdvanced<sup>™</sup> Science-Based Solutions Weed Killer for Lawns Ready-to-Spray; Fortifend Advanced<sup>™</sup> Weed Killer for Lawns Ready-to-Spray; Bianova Advanced<sup>™</sup> Lawn Weed Killer Ready-to-Spray; BioAdvanced<sup>™</sup> Lawn Weed Killer Ready-to-Spray; BioAdvanced<sup>™</sup> Science-Based Solutions Lawn Weed Killer Ready-to-Spray; Fortifend Advanced<sup>™</sup> Lawn Weed Killer Ready-to-Spray; Bianova Advanced<sup>™</sup> Southern Weed Killer for Lawns Ready-to-Spray; BioAdvanced<sup>™</sup> Southern Weed Killer for Lawns Ready-to-Spray;

- Effective Listed [\*] Weed Protection
- Effective Listed [\*] Broadleaf Weed Protection
- Effectively Protects Lawns from Listed [\*] Broadleaf Weeds
- Kills Major Listed [\*] Broadleaf Weeds
- Kills Listed [\*] Weeds Won't Harm Lawns<sup>†</sup>
- Kills to the Root
- Protects Against Major Listed [\*] Broadleaf Weeds
- Tough Listed [\*] Weed Protection
- Listed [\*] Weed Protection System
- Systemic
- 1 Quart treats [8,000] [up to 16,000] square feet
- Convenient MEASURE-UP™ Cup
- Kills Broadleaf weeds in St. Augustine & other Southern Lawns
- For St. Augustine & other Listed [\*] Southern Lawns
- Kills Over 200 Listed [\*] Broadleaf Weeds
- Won't Harm Your Lawn<sup>†</sup>
- Rainproof in hours

Marketing Claim for Brand Name Change: New Name. Same Great Results. New Name. Same Familiar Formula. New Name. Same Performance.

- 100% Satisfaction Guaranteed\* (or your money back)
- Rainproof in 1 hour!
- Attach to Hose & Spray
- Kills 200+ Listed [\*] Weeds
- [optional advertising claims when product is sold in "low-water" hose-end sprayers, as determined by testing]
- Water [Saver][Saving] Formula
- Water [Saver][Saving] Applicator
- Water [Saver][Saving] Sprayer
- Uses Up To [80][XX]% Less Water\*
- [with footnote disclaimer] \*vs leading Southern weed hose end sprayers
- won't harm lawns†
- twhen used as directed
- Bonus 40 oz size 25% more than the 32 oz size
- Now covers [XXX] sq ft
- Easy hold
- EZ Spray
- One-touch sprayer
- EZ<mark>-G-g</mark>rip<mark>™</mark>
- Ergo<mark>-G-g</mark>rip<mark>™</mark>

- Simple spray
- EZ<mark>--<u>G</u>g</mark>rip <u>One-Touch Sprayer™</u>
- One touch
- Spray n go
- Thumb switch
- Handy spray
- EZ touch
- One-touch application
- Bonus [xx] oz. size [x]% more than [X] oz size
- Bonus [x] oz size
- [For] [Use] [On] [Most] [Many]
   [Majority of] [Northern] [and]
   [Southern] [Turfs] [Lawns] [Types]
   [Varieties]\*
- Use on Northern Turfs: Fescue, Kentucky Bluegrass & [and] Perennial Rye [grass]
- Use on Southern Turfs: Bermuda, Zoysia, Centipede & [and] St. Augustine [grass]

\*do not use on Carpetgrass, Dichondra or Floratam variety of St. Augustinegrass

\*listed (Note to PM: The word "listed" can appear as a footnote with the appropriate phrase, such as \*listed plants/pests/weeds)

ers

Net Contents: 16 fl oz (473 mL) or 24 fl oz (709 mL) or 32 fl oz (946 mL), 40 oz (1.18L), or 1 Gal (3.78 L)

# KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel Booklet for Additional Ingredients and Precautionary Statements

#### (BACK PANEL)

#### **OPEN** Resealable Label for Directions & Precautions **PRESS TO RESEAL**

This product kills over 200 types of broadleaf weeds in St. Augustine and other Southern lawns.

This formula kills Dollarweed, Spurge, Oxalis and Dandelion without harming your lawn \*. Controlling weeds in your lawn has never been easier. (\*when used as directed.)

[modify representative weeds or add approved marketing claims as needed]

#### [READ THE LABEL FIRST!]

#### QUICK FACTS

- Kills over 200 listed [\*] broadleaf weeds (as listed on this label)
- Won't harm your lawn<sup>†</sup>
- Treats [8,000] [up to 16,000] sq ft

| KILLS        | Over 200 broadleaf weeds such as Dollarweed, Spurge and Oxalis, Dandelion, Clover, Chickweed and other listed weeds [ <i>or choose from marketing claims</i> ]              |  |  |  |
|--------------|---|--|--|--|
| WHERE TO USE | On Lawns  |  |  |  |
|              | $\dagger$ (Do not use on Carpetgrass, Dichondra or Floratam variety of St. Augustinegrass.)   |  |  |  |
| WHEN TO USE  | When weeds are actively growing   |  |  |  |
| HOW MUCH TO  | Northern Lawns: Fescue, Kentucky Bluegrass, and Perennial Ryegrasses  |  |  |  |
| USE          | and   |  |  |  |
|              | Hardy Southern Lawns: Bermuda, Buffalo and Zoysia grasses<br>One qt treats up to 8,000 sq ft  |  |  |  |
|              | Southern lawns: Bahia, Centipede and St. Augustinegrass grasses<br>One qt treats up to 16,000 sq ft   |  |  |  |
|              | No measuring needed.<br>Dilutes automatically.  |  |  |  |
| ☎ 🗖          | For questions or comments, call toll-free 1-877-229-3724 or visit us on-line at<br>[www.bianovaadvanced.com][www.BioAdvanced.com][www.BioAdvancedScienceBasedSolutions.com] |  |  |  |
|              | [www.fortifendadvanced.com]   |  |  |  |

#### FOR OUTDOOR RESIDENTIAL USE ONLY

#### **ACTIVE INGREDIENTS:**

| 2,4-D, dimethylamine salt      | % |
|--------------------------------|---|
| Mecoprop-p, dimethylamine salt | % |
| Dicamba, dimethylamine salt    | % |
| OTHER INGREDIENTS*:            | % |
| TOTAL                          | % |

\* Other ingredients include: Water and surfactants (to keep the active ingredients dispersed in water).

#### THIS PRODUCT CONTAINS:

0.00427 lb 2,4-D acid equivalent per fl oz or 6.30% 0.00102 lb Mecoprop-p acid equivalent per fl oz or 1.51% 0.00047 lb Dicamba acid equivalent per fl oz or 0.69% Isomer specific by AOAC Method

#### EPA Reg. No. 92564-4

EPA Est. No.\_-AA-\_

| FIRST AID   |   |  |
|---|---|--|
| If swallowed  | Call a poison control center or doctor immediately for treatment advice.  |  |
|   | Have person sip a glass of water if able to swallow.  |  |
|   | Do not induce vomiting unless told to do so by a poison control center or doctor.   |  |
|   | Do not give anything to an unconscious person.  |  |
| If in eyes  | Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.  |  |
|   | Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.  |  |
|   | Call a poison control center or doctor for treatment advice.  |  |
| If on skin or clothing  | Take off contaminated clothing.   |  |
|   | Rinse skin immediately with plenty of water for 15 to 20 minutes.   |  |
|   | Call a poison control center or doctor for treatment advice.  |  |
| If inhaled  | Move person to fresh air.   |  |
|   | <ul> <li>If not breathing, call 911 or an ambulance, then give artificial respiration, preferably by<br/>mouth-to-mouth, if possible.</li> </ul>  |  |
|   | Call a poison control center or doctor for further treatment advice.  |  |
| office, bring this product label to [Active Ingredients: 7.59% 2,4- | rol center, have this product label accessible. If seeking treatment at an emergency room or doctor's show medical personnel. You may call toll-free 877-229-3763 for medical emergency information. D, dimethylamine salt {CAS # 2008-39-1}, 1.83% Mecoprop-p, dimethylamine salt {CAS # 66423-09-/lamine salt {CAS # 2300-66-5]}. |  |

# PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

## CAUTION

- Harmful if absorbed through skin or if swallowed.
- Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

After using this product:

- Rinse gloves before removing.
- Remove clothing and launder separately before reuse.
- Wash hands thoroughly with soap and water.
- If product gets on clothing, remove clothing as soon as possible and shower.


### **ENVIRONMENTAL HAZARDS**


- This product is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.
- Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.
- This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

|  | DIRECTIONS FOR USE   |
|--|--|
| It is a violation of Federal La                                    | w to use this product in a manner inconsistent with its labeling.  |
| Read and follow all label dire                                     | ections.   |
| Do not apply this product in area during application.              | a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the   |
| Not for use on turf being gro                                      | wn for sale or other commercial use as sod, or for commercial seed production or for research purposes.  |
| Do not apply through any ty  | pe of irrigation system.   |
| <b>BEFORE YOU USE</b><br>Read and follow directions<br>when using. | <ul> <li>Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn or garden area only.</li> <li>Spray when most of the day is expected to remain cooler than 90° F.</li> </ul> |
|  | <ul><li>Do not apply as a fine mist because of potential injury to desirable plants.</li><li>The maximum number of broadcast applications per treatment site is two per year.</li></ul>  |

#### HOW TO USE [For Ready to spray hose end bottle]

#### Shake Well -Before Use





OFF CON

### [Optional Sprayer Directions 1]

- CONNECT
- 1. Shake well before use.
- 2. Connect sprayer to garden hose.
- 3. Turn on water.

#### SPRAY

- 1. Extend hose to farthest area to be treated, point sprayer nozzle in direction to be sprayed.
- 2. Begin spraying <*Insert sprayer-specific language here> Hayes HP-2 and Calmar RTS sprayers:* Press lock tab while using other hand to turn sprayer valve to ON position. *Green Garden RTS sprayer:* Turn sprayer valve to ON position.
- 3. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap the treated areas. Work back toward the faucet to avoid walking over wet, treated grass.

#### FINISH

1. To stop spraying, < *Insert sprayer-specific language here.*> *Hayes HP-2 RTS sprayer:* Press lock tab while using other hand to turn sprayer valve to OFF position. *Calmar & Green Garden RTS sprayers:* turn sprayer valve to OFF position.

- 2. Turn off water at faucet.
- 3. Relieve water pressure in hose by turning sprayer valve back to ON position until water slows to a drip then turn sprayer valve to OFF position.
- 4. Disconnect sprayer from hose.

## [Optional Sprayer Directions 2]

- [CONNECT
- 1. Shake well before use.
- 2. Securely Connect sprayer to garden hose.
- 3. Be sure that ON/OFF Switch is in OFF position, toward hose end of sprayer.
- 4. Turn on water at faucet.
- 5. For first time use, remove plastic shipping lock from beneath switch.

### SPRAY

- 1. Extend hose to farthest area to be treated, point sprayer nozzle in direction to be sprayed.
- 2. Begin spraying Push ON/OFF Switch forward into ON position.


3. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap treated areas. Work back toward faucet to avoid walking over wet, treated grass.

#### FINISH

- 1. To stop spraying, push ON/OFF Switch to OFF position.
- 2. Turn off water at faucet.
- 3. Relieve water pressure in hose by turning ON/OFF Switch back to ON position until water slows to a drip, then return ON/OFF switch to OFF position.
- 4. Disconnect sprayer from hose.]

#### **Application Restrictions:**

- · Limited to 2 applications per year with a 30 day interval between applications
- **Maximum single application rate** of 8.0 fl oz/1,000 sq ft (1.47 lbs 2,4-D ae, 0.35 lb MCPP-p ae and 0.16 lb dicamba ae per acre per application)
- Maximum seasonal rate of 16.0 fl oz/1,000 sq ft (2.94 lbs 2,4-D ae, 0.71 lb MCPP-p ae and 0.33 lb dicamba ae per acre per application)



| TIPS FOR APPLYING         |   |  |
|---------------------------|---|--|
| HOW TO APPLY              | 1. Extend hose to the farthest area to be treated and work back toward the faucet to avoid walking over wet, treated grass.   |  |
|                           | 2. Walk at a steady pace while spraying, using an even sweeping motion, slightly overlapping the treated areas.   |  |
| FOR BEST RESULTS          | Apply when weeds are young and actively growing.  |  |
|                           | If the lawn is dry, water 24 hours before using this product.   |  |
|                           | Rain or watering one hour after application will not wash away effectiveness.   |  |
|                           | Apply 1 to 3 days after mowing.   |  |
|                           | Certain hard-to-kill weeds may need to be spot-treated in 30 days.  |  |
|                           | Do not apply to new lawns until the grass has reached a height of at least 2 inches.  |  |
|                           | Bare spots may be seeded 3 to 4 weeks after application.  |  |
| HOW THIS PRODUCT<br>WORKS | This product is a selective herbicide - it selectively kills weeds in the lawn but won't harm the lawn itself.<br>is a blend of proven weed killers designed to get broadleaf weeds such as Dollarweed, Spurge and<br>Oxalis. |  |
|                           | [modify representative weeds or add approved marketing claims as needed]  |  |

| Do not allow people (other than applicator) or pets on treated area during application.<br>Do not allow people or pets to re-enter the treated area until spray has dried. |
|--|
|  |

| CONTROLS | Annual yellow sweetclover | Cockle              | Mugwort                     | Spiny amaranth         |
|----------|---------------------------|---------------------|-----------------------------|------------------------|
| [KILLS]  | Aster                     | Cocklebur           | Musk thistle                | Spiny cocklebur        |
|          | Austrian fieldcress       | Common mullein      | Mustard                     | Spiny sowthistle       |
|          | Bedstraw                  | Creeping Charlie    | Narrowleaf plantain         | Spotted catsear        |
|          | Beggarticks               | Creeping Jenny      | Narrowleaf vetch            | Spotted spurge         |
|          | Betony, Florida           | Cudweed             | Nettle                      | Spurweed               |
|          | Bindweed, field           | Curly dock          | Orange hawkweed             | Stinging nettle        |
|          | Bird vetch                | Daisy, English      | Oriental cocklebur          | Strawberry, India mock |
|          | Bitter wintercress        | Daisy, fleabane     | Oxalis                      | Tall nettle            |
|          | Bittercress, hairy        | Daisy, oxeye        | Parsley-piert               | Tall vervain           |
|          | Bitterweed                | Dandelion           | Parsnip                     | Tansy mustard          |
|          | Black-eyed Susan          | Dichondra           | Pearlwort                   | Tanweed                |
|          | Black medic               | Dogbane             | Pennycress                  | Thistle                |
|          | Black mustard             | Dogfennel           | Pennywort                   | Trailing crownvetch    |
|          | Blackseed plantain        | Dollarweed          | Pepperweed                  | Tumble mustard         |
|          | Blessed thistle           | Elderberry          | Pigweed                     | Tumble pigweed         |
|          | Bloodflower milkweed      | False dandelion     | Pineywoods bedstraw         | Velvetleaf             |
|          | Blue lettuce              | False flax          | Plains coreopsis (tickseed) | Venice mallow          |
|          | Blue vervain              | False sunflower     | Plantain                    | Virginia buttonweed    |
|          | Bracted plantain          | Fiddleneck          | Poison ivy                  | Virginia creeper       |
|          | Brassbuttons              | Florida pusley      | Poison oak                  | Virginia pepperweed    |
|          | Bristly oxtongue          | Frenchweed          | Pokeweed                    | Water pennywort        |
|          | Broadleaf dock            | Galinsoga           | Poorjoe                     | Wavyleaf bullthistle   |
|          | Broadleaf plantain        | Goathead            | Prairie sunflower           | Western clematis       |
|          | Broomweed                 | Goldenrod           | Prickly lettuce             | Western salsify        |
|          | Buckhorn                  | Ground ivy          | Prickly sida                | White mustard          |
|          | Buckhorn plantain         | Gumweed             | Prostrate knotweed          | Wild aster             |
|          | Bulbous buttercup         | Hairy fleabane      | Prostrate pigweed           | Wild buckwheat         |
|          | Bull thistle              | Hawkweed            | Prostrate spurge            | Wild carrot            |
|          | Bullnettle                | Healall             | Prostrate vervain           | Wild four-o'clock      |
|          | Burclover                 | Heartleaf drymary   | Puncturevine                | Wild garlic            |
|          | Burdock                   | Heath aster         | Purslane, common            | Wild geranium          |
|          | Burning nettle            | Hedge bindweed      | Ragweed                     | Wild lettuce           |
|          | Burweed                   | Hedge mustard       | Red sorrel                  | Wild marigold          |
|          | Buttercup                 | Hemp                | Redroot pigweed             | Wild mustard           |
|          | Buttonweed                | Henbit              | Redstem filaree             | Wild onion             |
|          | Canada thistle            | Hoary cress         | Rough cinquefoil            | Wild parsnip           |
|          | Carolina geranium         | Hoary plantain      | Rough fleabane              | Wild radish            |
|          | Carpetweed                | Hoary vervain       | Russian pigweed             | Wild rape              |
|          | Catchweed bedstraw        | Horsenettle         | Russian thistle             | Wild strawberry        |
|          | Catnip                    | Jimsonweed          | Scarlet pimpernel           | Wild sweet potato      |
|          | Catsear                   | Knawel              | Scotch thistle              | Wild vetch             |
|          | Chickweed, common         | Knotweed            | Sheep sorrel                | Woodsorrel             |
|          | Chickweed, mouseear       | Kochia              | Shepherdspurse              | Woolly croton          |
|          | Chicory                   | Lambsquarters       | Slender plantain            | Woolly morningglory    |
|          | Cinquefoil                | Lespedeza           | Smallflower galinsoga       | Woolly plantain        |
|          | Clover, crimson           | Mallow              | Smooth dock                 | Wormseed               |
|          |                           | Mailow<br>Matchweed |                             | Yarrow                 |
|          | Clover, hop               | Matchweed           | Smooth pigweed              |                        |
|          | Clover, red               |                     | Sorrel<br>Sowthistle        | Yellow rocket          |
|          | Clover, strawberry        | Milk vetch          |                             | Yellowflower pepperwee |
|          | Clover, sweet             | Morningglory        | Spanishneedles              |                        |
|          | Clover, white             | Mouseear hawkweed   | Speedwell                   |                        |

|  | STORAGE AND DISPOSAL   |
|--|--|
| Do not contaminate food or feed by st            | orage and disposal.  |
| Pesticide Storage:                               | Store in original container in a cool dry place, out of the reach of children, preferably in a locked storage cabinet. Protect from freezing.  |
| Pesticide Disposal<br>and Container<br>Handling: | <ul> <li>If Empty:</li> <li>Non-refillable container. Do not refill or reuse empty container.</li> <li>Place empty container in trash or offer for recycling if available.</li> <li>If Partly Filled:</li> <li>If it is necessary to dispose of unused product, please call your local solid waste agency or you may call toll-free 1-877-229-3724 for disposal instructions.</li> <li>Never place unused product down any indoor or outdoor drain.</li> </ul> |

**NOTICE:** Research and testing have determined that the "Directions For Use" are appropriate for the proper use of this product under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

\*Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

BioAdvanced<sup>™</sup> [Fortifend Advanced<sup>™</sup>] [Bianova Advanced<sup>™</sup>] is a trademark of SBM Life Science Corp.

EZ-Grip<sup>™</sup> and Ergo-Grip<sup>™</sup> are trademarks of SBM Life Science Corp.

PRODUCED FOR [DISTRIBUTED BY]

SBM Life Science Corp. 1001 Winstead Drive, Suite 500 Cary, North Carolina 27513

Lawn Herbicide JD Concentrate (PENDING) 04/07/201711/20/2017

## [ART]

| UR LABS                                     | READ   |
|---|--|
| LAB-DEVELOPED<br>SOLUTIONS                  | CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR<br>CONTRACTOR |
| PROTECTION ROB<br>POUR YARD<br>Togenou List |  |

# Lawn Herbicide JD Concentrate

(ABN) Bayer Advanced Weed Killer for Lawns Concentrate; Bayer Advanced Lawn Weed Killer Concentrate; Bayer Advanced Southern Weed Killer for Lawns Concentrate; Bianova Advanced<sup>™</sup> Weed Killer for Lawns Concentrate; BioAdvanced<sup>™</sup> Weed Killer for Lawns Concentrate; BioAdvanced<sup>™</sup> Science-Based Solutions Weed Killer for Lawns Concentrate; Fortifend Advanced<sup>™</sup> Weed Killer for Lawns Concentrate; Bianova Advanced<sup>™</sup> Lawn Weed Killer Concentrate; BioAdvanced<sup>™</sup> Lawn Weed Killer Concentrate; BioAdvanced<sup>™</sup> Science-Based Solutions Lawn Weed Killer Concentrate; Fortifend Advanced<sup>™</sup> Science-Based Solutions Lawn Weed Killer Concentrate; Fortifend Advanced<sup>™</sup> Lawn Weed Killer Concentrate; Bianova Advanced<sup>™</sup> Southern Weed Killer for Lawns Concentrate; BioAdvanced<sup>™</sup> Southern Weed Killer for Lawns Concentrate; BioAdvanced<sup>™</sup> Southern Weed Killer for Lawns Concentrate; Fortifend Advanced<sup>™</sup> Southern Weed Killer for Lawns

| ACTIVE INGREDIENTS:            |        |
|--------------------------------|--------|
| 2,4-D, dimethylamine salt      | 7.59 % |
| Mecoprop-p, dimethylamine salt |        |
| Dicamba, dimethylamine salt    | 0.84 % |
| OTHER INGREDIENTS:             |        |
| TOTAL:                         |        |

# KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel Booklet for Additional Ingredients and Precautionary Statements Editorial Note – [Bracketed text] is optional language

# PRECAUTIONARY STATEMENTS

# HAZARDS TO HUMANS AND DOMESTIC ANIMALS

## CAUTION

- Harmful if absorbed through skin or if swallowed.
- Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

After using this product:

- Rinse gloves before removing.
- Remove clothing and launder separately before reuse.
- Wash hands thoroughly with soap and water.
- If product gets on clothing, remove clothing as soon as possible and shower.

Read the label on the container in the box for full directions for use.

[Marketing claims] [For promotional use only. Not for resale.] [Not for individual sale]

BioAdvanced<sup>™</sup> [Fortifend Advanced<sup>™</sup>] [Bianova Advanced<sup>™</sup>] is a trademark of SBM Life Science Corp.

PRODUCED FOR [DISTRIBUTED BY]

SBM Life Science Corp<mark>.</mark> 1001 Winstead Drive, Suite 500 Cary, North Carolina 27513