



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

April 21, 2017

Norma C. Pangilinan, Ph.D.
Head of Regulatory Affairs
SBM Life Science Corp.
1001 Winstead Drive Suite 500
Cary, NC 27513

Subject: Notification per PRN 98-10 – ABNs and marketing claims
Product Name: TEMPO READY-TO-USE INSECTICIDE
EPA Registration Number: 92564-18
Application Date: 4/4/2017
Decision Number: 528006

Dear Dr. Pangilinan:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The alternate brand names "*Bianova Advanced Vegetable & Garden Insect Spray Ready-to-Use; BioAdvanced Vegetable & Garden Insect Spray Ready-to-Use; BioAdvanced Science-Based Solutions Vegetable & Garden Insect Spray Ready-to-Use; Fortifend Advanced Vegetable & Garden Insect Spray Ready-to-Use; Bianova Advanced Flower Insect Killer Ready-to-Use II; BioAdvanced Flower Insect Killer Ready-to-Use II; BioAdvanced Science-Based Solutions Flower Insect Killer Ready-to-Use II; Fortifend Advanced Flower Insect Killer Ready-to-Use II; Bianova Advanced Rose & Flower Insect Killer Ready-to-Use; BioAdvanced Rose & Flower Insect Killer Ready-to-Use; BioAdvanced Science-Based Solutions Rose & Flower Insect Killer Ready-to-Use; Fortifend Advanced Rose & Flower Insect Killer Ready-to-Use*" have been added to the product record.

The label submitted with the application has been stamped "Notification" and will be placed in our records.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false

Page 2 of 2
EPA Reg. No. 92564-18
Decision No. 528006

or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you have any questions, you may contact Angela Hollis at 703-347-0216 or via email at hollis.angela@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read 'Kable Bo Davis', enclosed within a hand-drawn oval.

Kable Bo Davis, Product Manager 3
Invertebrate and Vertebrate Branch 1
Registration Division (7505P)
Office of Pesticide Programs

NOTIFICATION

92564-18

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

04/21/2017

(FRONT PANEL)

Tempo[®] Ready-to-Use Insecticide

(ABN) Bayer Advanced Vegetable & Garden Insect Spray Ready-to-Use; Flower Insect Killer Ready-to-Use II; Bayer Advanced Rose & Flower Insect Killer

Ready to Use; **Bianova Advanced Vegetable & Garden Insect Spray Ready-to-Use; BioAdvanced Vegetable & Garden Insect Spray Ready-to-Use; BioAdvanced Science-Based Solutions Vegetable & Garden Insect Spray Ready-to-Use; Fortifend Advanced Vegetable & Garden Insect Spray Ready-to-Use; Bianova Advanced Flower Insect Killer Ready-to-Use II; BioAdvanced Flower Insect Killer Ready-to-Use II; BioAdvanced Science-Based Solutions Flower Insect Killer Ready-to-Use II; Fortifend Advanced Flower Insect Killer Ready-to-Use II; Bianova Advanced Rose & Flower Insect Killer Ready-to-Use; BioAdvanced Rose & Flower Insect Killer Ready-to-Use; BioAdvanced Science-Based Solutions Rose & Flower Insect Killer Ready-to-Use; Fortifend Advanced Rose & Flower Insect Killer Ready-to-Use**

Formatted: Highlight

For use on: Vegetables, Peanuts, Sunflowers, Grapes, Flowers, Shrubs, Groundcovers, Lawns and Around Homes

GUARANTEED SATISFACTION

READY TO USE

KILLS LISTED* INSECTS

MULTIPURPOSE INSECTICIDE

- Fast Acting...Kills Listed* Insects On Contact
- Long Lasting...Keeps On Killing For Up To 14 Days

FAST ACTING

LONG LASTING FOR UP TO 2 WEEKS

- Kills outdoor listed* pests on contact
- Lasts up to 14 days
- For use on flowers, shrubs, and listed* vegetables
- Just One Application Quickly Kills Listed* Insects
- Keeps On Killing For Up To 14 Days
- Use On Tomatoes Up To Day Of Harvest
- Contains Tempo®
- Low odor [formula]
- Won't harm plants
- Protects plants
- Water-based
- Kills Beetles (including Japanese Beetle adults), Caterpillars (including Tomato Hornworm), Ants, Armyworms, Ticks (including ticks carrying Lyme disease) and [&] other listed* garden and [&] nuisance pests [insects] [bugs]
- Kills Beetles, Caterpillars, Ants, Ticks and [&] other listed* insects
- Kills [Controls over] [more than] 100 listed* insects [bugs]

[pests] on vegetables, flowers, small trees and [&] shrubs

- Kills [Controls over] [more than] 100 listed* garden and nuisance pests without harming roses, flower, or shrubs
- Kill listed* insects on Contact
- Kills over 100 listed* insects [bugs] [pests] on contact
- On [For] [For use on] Roses, Flowers, Small trees, Shrubs, Fruits, Vegetables and [&] other Outdoor Surface [Areas]
- Can be used on [most popular] fruits and [&] vegetables (see back panel for details)
- User in (on) Vegetable Gardens
- Fast Acting Formula
- Reapply as directed for a more beautiful garden
- Guaranteed Kill or your money back
- Keeping your vegetable garden and lawn free of listed* pests has never been easier
- Kills aphids, Japanese beetles (adults), cutworms, tomato hornworms, and other listed* pests
- Also for use on lawns
- Kills listed* insects on contact & protects up to 14 days
- Protects up to 14 days
- This product provides up to 14 days protection with a single application. Now keeping your roses and flowers free of listed pests has never been easier.
- Kills 100+ listed* insects on contact
- Kills Aphids, Japanese Beetles (adult), Leafminers, Whiteflies, [Thrips], [Scale (crawler stages)] and other listed* pests [insects] [bugs]
- On [For] [For use on] Roses and Flowers

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Marketing Claim for Brand Name Change:

New Name. Same Great Results.

New Name. Same Familiar Formula.

New Name. Same Performance

* [The word "listed" can appear as a footnote with the appropriate phrase, such as *listed insects]

ACTIVE INGREDIENT:

Cyfluthrin 0.003 %

OTHER INGREDIENTS: 99.997 %

Net Contents: 24 fl oz, 1/2 gallon, 1 gallon

STOP - Read This Entire Label Before Use
KEEP OUT OF REACH OF CHILDREN
FOR OUTDOOR RESIDENTIAL USE ONLY

CAUTION

See back panel / booklet for additional ingredients and precautionary statements.

First Aid




| | |
|---------------------|---|
| If Swallowed | <ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person. |
| If In Eyes | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes.• Call a poison control center or doctor for treatment advice. |
| If Inhaled | <ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice. |

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free (877) 229-3763 for medical emergency information.

Active Ingredient: 0.003% Cyfluthrin [CAS # 68359-37-5]

Note to Physician: No specific antidote is available. Treat the patient symptomatically.

(BACK PANEL)

| [For 1 gallon] | | |
|---|--|--|
|  ATTACH SPRAYER <ul style="list-style-type: none">Remove sprayer from bottle and extend hose.Open flip-cap on bottle.Insert hose and plug into flip-cap opening until it snaps into place. |  SPRAY <ul style="list-style-type: none">Adjust nozzle to stream or spray position.Point sprayer downward and squeeze trigger repeatedly until liquid is released. |  AFTER USE <ul style="list-style-type: none">Turn sprayer nozzle to X (off) position.Do not unplug hose.Return sprayer to bottle. Sprayer will snap into place. |

QUICK FACTS

- Ready to Use...No Mixing, No Measuring
- Quickly Stops Harmful Listed* Insects Before They Destroy Vegetables
- May Be Used Up to Day of Harvest on Tomatoes
- Effective Against Broad Range of Listed* Pests
- Quickly Stops Harmful Listed* Insects Before They Do Damage
- Kills Household Nuisance Pest Outdoors
- Kills Listed* Insects on Contact
- Keeps on killing for up to 14 days
- Kills stink bugs

| KILLS | (ART: aphid, cabbage looper, tomato hornworm, whitefly) | | |
|---|---|---|--|
| Ants Aphids Armyworms Aster leafhoppers Bagworms Banded sunflower moths Boxelder bugs Budmoths Cabbage loopers Carrot weevils (adults) Caterpillars Celery leaf tiers Chinch bugs Clover mites Cockroaches Colorado potato beetles Corn borers Corn earworms Corn rootworms (adults) Crickets Cutworms Earwigs | Elm leaf beetles Flea beetles Flies Garden webworms Grasshoppers Gypsy moth larvae Head clipper weevils (adults) Japanese beetles (adults) Lace bugs Leaf beetles Leaf-feeding beetles Leaf-feeding caterpillars Leafhoppers (including Glassy-Winged Sharpshooter) Leafminers Mealybugs Midges Mosquitoes Pepper weevils (adults) Pillbugs Plant bugs | Potato aphids Potato leafhoppers Scale (crawler stages) Seed weevils (adults) Sod webworms Sowbugs Spiders Spittlebugs Stem weevil (adults) Stink bugs Sunflower beetles Sunflower moths Thrips Tomato fruitworms Tomato hornworms Tomato pinworms Wasps Weevils (adults) Western flower thrips Whiteflies | |
| WHERE TO USE | Roses, Flowers, Vegetable and flower gardens, shrubs, ground covers, lawns and around homes | | |
| WHEN TO USE | When insects or damage first appear For severe infestations, repeat every 7-14 days as specified. | | |
| WHEN TO HARVEST | See plant specific restrictions for pre-harvest interval. If no pre-harvest interval is listed, the product may be applied up to and including day of harvest. | | |



Questions or comments, call toll-free (877) 229-3724

www.BayerAdvanced.com [INSERT COMPANY WEBSITE HERE]

Formatted: Highlight

ACTIVE INGREDIENT:

Cyfluthrin.....0.003 %

OTHER INGREDIENTS:*.....99.997 %
100.000 %

* Other ingredients include: Water, organic solvent (less than 0.1%, to dissolve the active ingredient for mixing) and surfactants (to keep the active ingredient dispersed in water).

Cyfluthrin is the chemical name for Tempo®

EPA Reg. No. **72155-40 92564-18**

EPA Est. No. XXXX-AA-1

Formatted: Highlight

Formatted: Highlight

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

- Harmful if swallowed or inhaled.
- Causes eye irritation.
- Avoid contact with skin, eyes, or clothing.
- Avoid breathing spray mist.
- Wash thoroughly with soap and water after handling.

Environmental Hazards

- This pesticide is toxic to fish. Do not apply directly to water.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.
- This pesticide is highly toxic to bees. Do not apply TEMPO Ready-to-Use Insecticide when bees are active.
- Apply this product only as specified on this label.
-

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For best results, read and follow all label directions.

BEFORE YOU USE

Read and follow these directions when using:

- Apply the product to flowering plants during early morning or late evening when bees are not present.
- Do not spray the product into fish pools, ponds, streams, or lakes.
- Do not use in small, enclosed areas where spray can be inhaled.
- Do not water the treated area to the point of run-off.
- Do not make application during rain.
- All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:
 - (1) Treatment to soil or vegetation around structures;
 - (2) Applications to lawns, turf, and other vegetation;
 - (3) Applications to building foundations, up to a maximum height of 3 feet.
 Other than applications to building foundations, all outdoor applications to impervious surfaces such as

| | |
|---|---|
| | <p>sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are to spot and crack-and crevice applications, only.</p> <ul style="list-style-type: none"> Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application |
| WHEN TO USE | When insects or damage first appear. |
| HOW TO USE Shake well before using. | <p>To apply TEMPO Ready-to-Use Insecticide:</p> <ul style="list-style-type: none"> Adjust nozzle to desired spray pattern. Hold sprayer about 12 inches from foliage. Spray leaves, stems, and tender new shoots. To kill insects that may hide, spray the underside of the leaves. Apply until leaves are evenly coated with spray, but not dripping. <p>(ART: spraying undersides of leaves)</p> |
| HOW OFTEN TO USE | Kills insects on contact and keeps on killing for weeks. For severe infestations, repeat every 7 to 14 days as necessary. |

| | | | |
|---|---------------------------|---|---------------------------|
| FLOWERS, SHRUBS, GROUND COVERS | | | |
| PEST | | | |
| Aphids | Grasshoppers | Leaf-feeding beetles | Scale (crawler stages) |
| Bagworms | Gypsy moth larvae | Leafhoppers (including Glassy-winged Sharpshooter) | Spittlebugs |
| Boxelder bugs | Japanese beetle (adults) | Mealybugs | Thrips |
| Clover mites | Lace bugs | Pillbugs | Whiteflies |
| Elm leaf beetles | Leaf-feeding caterpillars | | |
| HOW TO APPLY | | | |
| <p>Spray foliage, tender new shoots and the underside of leaves thoroughly.</p> <p>To kill insects that hide, spray the underside of the leaves. Apply until leaves are evenly coated with spray droplets, but not dripping.</p> <p>(ART)</p> | | | |

| FIELD CROPS | | | |
|--|--|--|-----------------------------|
| Peanut | | | |
| After spraying, wait 14 days before harvesting. | | | |
| Do not spray more than 3 times per season. | | | |
| PEST | | | |
| Armyworm (1 st and 2 nd instar) | Fall armyworm (1 st and 2 nd instar) | Potato leafhopper | Thrips |
| Bean leaf beetle | Green cloverworm | Rednecked peanutworm | Vegetable weevil |
| Beet armyworm (1 st and 2 nd instar) | Grape colaspis (adult) | Stink bugs | Velvetbean caterpillar |
| Cutworms | Grasshoppers | Threecornered alfalfa hopper | Whitefringed beetle (adult) |
| Corn earworm | Japanese beetle (adult) | Southern armyworm (1 st and 2 nd instar) | Whitefly (adult) |
| Corn rootworms (adult) | June beetle (adult) | Soybean looper | |

| Sunflowers | | |
|---|-------------------------------|-------------------------------|
| After spraying, wait 30 days before harvesting. | | |
| Do not spray more than 3 times per season. | | |
| PEST | | |
| Banded sunflower moth | Sunflower beetle | Sunflower midge |
| Cutworms | Stink bugs | Sunflower moth |
| Grasshoppers | Sunflower bud moth | Sunflower seed weevil |
| Palestriped flea beetle | Sunflower headclipping weevil | Sunflower stem weevil (adult) |

| VEGETABLE CROPS | | | |
|---|--|--|---|
| Brassica (Cole) Leafy Vegetables: | | | |
| Includes all members of Crop Group 5: | | | |
| Broccoli, Broccoli raab (rapini), Chinese (gai lon) broccoli, Brussels sprouts, Cabbage, Chinese (bok choy) cabbage, Chinese (napa) cabbage, Chinese mustard (gai choy) cabbage, Cauliflower, Cavalo broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens, and Turnip greens. | | | |
| No waiting to harvest. | | | |
| Do not spray more than 4 times per season. | | | |
| PEST | | | |
| Alfalfa looper | Corn earworm | Japanese beetle (adult) | Southern cabbageworm |
| Armyworm (1 st and 2 nd instar) | Cutworms | Lygus bug | Tarnished plant bug |
| Beet armyworm (1 st and 2 nd instar) | Diamondback moth (larvae) | Meadow spittlebug | Thrips |
| Cabbage flea beetle | Fall armyworm (1 st and 2 nd instar) | Potato leafhopper | Vegetable weevil (adult) |
| Cabbage looper | Grasshoppers | Southern armyworm (1 st and 2 nd instar) | Whitefly (adult) |
| Cabbage webworm | Imported cabbageworm | Stink bugs | Yellowstriped armyworm (1 st and 2 nd instar) |

Cucurbits (except crops grown for seed):

Includes all members of Crop Group 9:

Balsam apple, Balsam pear, Bitter melon, Chayote, Chinese cucumber, Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Edible gourd (includes: hyotan, cucuzza, henchmia and Chinese okra), Muskmelon (includes: cantaloupe, true cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), Pumpkin, Summer squash (includes: crookneck squash, scallop squash, straightneck squash, vegetable marrow, and zucchini), Watermelon, Winter squash (includes: butternut squash, calabaza, hubbard squash, acorn squash and spaghetti squash)

No waiting to harvest.

Do not spray more than 4 times per season.

PEST

| | | | |
|---|-----------------|-------------------|--------------------------|
| Armyworm (1 st and 2 nd instar) | Cutworms | Pickleworm | Stripped cucumber beetle |
| Brown marmorated stink bug | Cucumber beetle | Potato leafhopper | Tarnished plant bug |
| Cabbage looper | Grasshoppers | Rindworm | Tobacco budworm |
| Colorado potato beetle | Lygus bug | Stink bugs | Whitefly (adult) |
| Corn earworm | Melonworm | | |

Fruiting Vegetables

Includes all members of Crop Group 8:

Eggplant, Groundcherry, Pepino, Pepper (includes: bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo, and Tomato

No waiting to harvest for tomato. After spraying, wait 7 days before harvesting for all other crops.

Do not spray more than 6 times per season.

PEST

| | | | |
|--|--------------------|--|---|
| Beet armyworm (1 st and 2 nd instar) | Flea beetles | Southern armyworm (1 st and 2 nd instar) | Tomato hornworm |
| Brown marmorated stink bug | Garden symphylan** | Stink bugs | Tomato pinworm |
| Cabbage looper | Garden webworm | Tarnished plant bug * | Variegated cutworm |
| Celery leaf-tier | Leafminers | Thrips (except <i>Thrips palmi</i>) | Whitefly (adult) |
| Colorado potato beetle | Pepper weevil | Tomato fruitworm (corn earworm) | Western yellowstriped armyworm (1 st and 2 nd instar) |
| European corn borer | Potato aphid | | |
| | Potato leafhopper | | |

Leafy Vegetables:

Includes all members of Crop Group 4:

Amaranth (Chinese spinach), Arugula (rouquette), Cardoon, Celery, Chinese celery, Celtuce, Chervil, Chrysanthemum (edible-leaved and garland), Corn salad, Cress (garden and upland), Dandelion, Dock (sorrel), Endive (escarole), Florence fennel, Lettuce (head and leaf), New Zealand spinach, Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Rhubarb, Spinach, Swiss chard, Vine spinach

No waiting to harvest. Do not spray more than 4 times per season.

PEST

| | | | |
|--|--|--|---|
| Alfalfa looper | Diamondback moth (larvae) | Japanese beetle (adult) | Stink bugs |
| Beet armyworm (1 st and 2 nd instar) | European corn borer | Leafhoppers | Tarnished plant bug |
| Brown marmorated stink bug | Fall armyworm (1 st and 2 nd instar) | Lygus bug | Thrips |
| Cabbage looper | Flea beetles | Meadow spittlebug | Vegetable weevil (adult) |
| Corn earworm | Grasshoppers | Saltmarsh caterpillar | Whitefly (adult) |
| Cutworms | Green cloverworm | Southern armyworm (1 st and 2 nd instar) | Yellowstriped armyworm (1 st and 2 nd instar) |
| Colorado potato beetle | Imported cabbageworm | | |

Potato and Other Tuberous and Corm Vegetables:

Includes all members of Crop Subgroup 1C:

Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Edible canna, Cassava (bitter and sweet), Chayote root, Chufa, Dasheen (taro), Ginger, Leren, Potato, Sweet potato, Tanier, True yam, Turmeric, Yam bean

No waiting to harvest.

Do not spray more than 6 times per season.

PEST

| | | | |
|----------------|------------------------|-------------------|-----------------------------|
| Aphids | European corn borer | Potato psyllid | Sweetpotato weevil (adults) |
| Cabbage looper | Flea beetles | Potato tuberworm | Tarnished plant bug |
| Cutworms | Colorado Potato Beetle | Potato Leafhopper | |

Root Vegetables (except sugarbeet):

Includes all members of Crop Subgroup 1B:

Garden beet, Edible burdock, Carrot, Celeriac, Turnip-rooted chervil, Chicory, Ginseng, Horseradish, Turnip-rooted parsley, Parsnip, Radish, Oriental radish, Rutabaga, Salsify (black, Spanish, and oyster plant), Skirret, Turnip

No waiting to harvest.

Do not spray more than 5 times per season.

PEST

| | | | |
|------------------|----------|--------------|-------------------|
| Aster leafhopper | Cutworms | Flea beetles | Potato leafhopper |
| Carrot weevil | | | |

Sweet Corn

No waiting to harvest.

Do not spray more than 10 times per season.

PEST

| | | | |
|---|--|--|---|
| Armyworm (1 st and 2 nd instar) | Corn silk fly | Japanese beetle(adult) | Southwestern corn borer |
| Bean leaf beetle | European corn borer | June beetle (adult) | Stalk borer |
| Black cutworm | Fall armyworm (1 st and 2 nd instar) | Leafhoppers | Stink bugs |
| Cereal leaf beetle | Flea beetle | Masked chafer (adult) | Webworm |
| Chinch bug | Granulate cutworm | Sand hill cutworm | Western bean cutworm |
| Click beetle (adult) | Grape colaspis (adult) | Southern armyworm (1 st and 2 nd instar) | Yellowstriped armyworm (1 st and 2 nd instar) |
| Corn earworm | Grasshoppers | Southern corn leaf beetle | |
| Corn rootworms (adult) | | | |

| TREE and VINE CROPS | | | |
|---|-------------------|---------------------------|---------------------------------|
| Grapes | | | |
| Includes: Table grape, Raisin, Wine and Muscadine grape | | | |
| After spraying, wait 3 days before harvesting | | | |
| Do not spray more than 4 times per season | | | |
| PEST | | | |
| Climbing cutworm | Grape flea beetle | Grape mealybug (crawlers) | Orange tortrix |
| Grape berry moth | Grape leaffolder | Glassywinged sharpshooter | Thrips |
| Grape bud beetle | Grape leafhopper | Grape leaf skeletonizer | Variegated leafhopper |
| Grape cane gallmaker (adult) | Grape leafroller | Omnivorous leafroller | Western grape leaf skeletonizer |

| FOR BEST RESULTS: |
|---|
| Do not water the foliage within one hour after applying. To avoid serious damage to plants, eliminate pests early in the season before they multiply. |

| DIRECTIONS FOR USE |
|---------------------------|
|---------------------------|

| HOUSEHOLD NUISANCE PESTS OUTDOORS | | | |
|---|--------------------------|------------|---------|
| PEST | | | |
| Ants | Crickets | Midges | Sowbugs |
| Boxelder bugs | Earwigs | Mosquitoes | Spiders |
| Clover mites | Flies | Pillbugs | Wasps |
| Cockroaches | Brown Mamoraed Stink Bug | | |
| HOW TO APPLY | | | |
| Spray around doors, window frames and other places where insects may enter house. Spray around foundations and porches. Spray on ant hills and ant trails. (ART) | | | |


| LAWN PESTS | | | |
|--|----------|--------------|---------------------------|
| PEST | | | |
| Ants | Crickets | Earwigs | Japanese beetles (adults) |
| Armyworms | Cutworms | Grasshoppers | Sod webworms (lawn moths) |
| Chinch bugs | | | |
| HOW TO APPLY | | | |
| For localized infestations, spray to penetrate the lawn and reach the soil. Do not mow for one day after application. For large areas, use Tempo® Concentrate Insecticide. | | | |



Do not allow entry into treated area until spray has dried.

Formatted: Highlight

Formatted Table

| Storage and Disposal | |
|--|---|
| Pesticide Storage Conditions: | Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing. |
| Pesticide Disposal: |  <ul style="list-style-type: none"> • Non-refillable container. Do not reuse or refill empty spray bottle. • If empty: Place in trash or offer for recycling if available. • If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain. |
| Important: It is best to use all of the product in accordance with label directions. If it is necessary to dispose of unused product, please follow any applicable state or local guidelines. | |

| NOTICE |
|--|
| Research and testing have determined that the "Directions for Use" are appropriate for the proper use of TEMPO Ready-to-Use Insecticide under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions. |

PRODUCED FOR

Formatted: Highlight

DISTRIBUTED BY

Formatted: Highlight



SBM Life Science Corp
1001 Winstead Drive, Suite 500
Carv, North Carolina 27513
A Business Unit of Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

Formatted: Highlight

Tempo RTU Insecticide (PENDING) 04/01/2017

Formatted: Underline