

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

July 16, 2021

Prachi Pandya Federal Regulatory Affairs Associate SBM Life Science Corp. 1001 Winstead Drive Suite 500 Cary, NC 27513

Subject: Label Amendment – Added formatting changes and graphics; revised marketing

claims and directions for use, registered ABN

Product Name: Tempo® Ready-To-Use Insecticide

EPA Registration Number: 92564-18 Application Date: November 11, 2020

Decision Number: 575941

Dear Ms. Pandya:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2 EPA Reg. No. 92564-18 Decision No. 575941

with FIFRA section 6. If you have any questions, please contact Elizabeth Andrews by phone at 703-347-8770, or via email at Andrews. Elizabeth@epa.gov.

Id Herrick

Sincerely,

Jacquelyn Herrick, Product Manager 03 Invertebrate & Vertebrate Branch 1 Registration Division (7505P) Office of Pesticide Programs

Enclosure

(FRONT PANEL)

Tempo® Ready-to-Use Insecticide

07/16/2021

ACCEPTED

Under the Federal Insecticide, Fungicide

and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 2000 4 4 9

92564-18

[ABN's]

Flower Insect Killer Ready-to-Use II

Bianova Advanced™ Vegetable & Garden Insect Spray Ready-to-Use

BioAdvanced® Vegetable & Garden Insect Spray Ready-to-Use

BioAdvanced® Science-Based Solutions Vegetable & Garden Insect Spray Ready-to-Use

Fortifend Advanced™ Vegetable & Garden Insect Spray Ready-to-Use

Bianova Advanced™ Flower Insect Killer Ready-to-Use II

BioAdvanced® Flower Insect Killer Ready-to-Use II

BioAdvanced® Science-Based Solutions Flower Insect Killer Ready-to-Use II

Fortifend Advanced™ Flower Insect Killer Ready-to-Use II

Bianova Advanced™ Rose & Flower Insect Killer Ready-to-Use

BioAdvanced® Rose & Flower Insect Killer Ready-to-Use

BioAdvanced® Science-Based Solutions Rose & Flower Insect Killer Ready-to-Use

Fortifend Advanced™ Rose & Flower Insect Killer Ready-to-Use

BioAdvanced® Science-Based Solutions Rose & Flower Insect Killer Spray

BioAdvanced® Science-Based Solutions Tomato & Vegetable Garden Insect Killer Ready-to-Use

BioAdvanced® Tomato & Vegetable Garden Insect Killer Ready-to-Use

BioAdvanced® Science-Based Solutions Tomato & Vegetable Insect Killer Ready-to-Use

BioAdvanced® Tomato & Vegetable Insect Killer Ready-to-Use

BioAdvanced® Science-Based Solutions Vegetable & Tomato Insect Killer Ready-to-Use

BioAdvanced® Vegetable & Tomato Insect Killer Ready-to-Use

For use on: [Vegetables,] [Peanuts,] [Sunflowers,] [Grapes,] Flowers, [Shrubs,] [Groundcovers,] [houseplants (treated outside),] [container plants (treated outside),] [Lawns] and [Around Homes]

ACTIVE INGREDIENT:

| Cyfluthrin | |
|--------------------|----------|
| OTHER INGREDIENTS: | 99.997 % |
| TOTAL: | |

Net Contents: 24 fl oz, 1/2 gallon, 1 gallon, 1.3 gallons

Editorial Note - [Bracketed text] is optional language. Any text separated by a backslash "/" and "or" are options

STOP - Read This Entire Label Before Use

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel / booklet for additional ingredients and precautionary statements.

FOR OUTDOOR RESIDENTIAL USE ONLY

| First Aid | | | |
|--------------|--|--|--|
| If Swallowed | Call a poison control center or doctor immediately for treatment advice. | | |
| | Have person sip a glass of water if able to swallow. | | |
| | Do not induce vomiting unless told to do so by a poison control center or doctor. | | |
| | Do not give anything by mouth to an unconscious person. | | |
| If In Eyes | Hold eye open and rinse slowly and gently with water for 15-20 minutes. | | |
| | Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. | | |
| | Call a poison control center or doctor for treatment advice. | | |
| If Inhaled | Move person to fresh air. | | |
| | If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. | | |
| | Call a poison control center or doctor for further treatment advice. | | |

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 1-(877) 229-3763 for medical emergency information.

[Active Ingredient: 0.003% Cyfluthrin (CAS # 68359-37-5)]

Note to Physician: No specific antidote is available. Treat the patient symptomatically.

(BACK PANEL)

{Note to PM: Any section from the below Quick [Product] Facts/Information will appear on the Marketplace label}

[Read the Label First!]

QUICK FACTS/ PRODUCT FACTS/ PRODUCT INFORMATION

- · Ready to Use...No Mixing, No Measuring
- Quickly Stops Harmful Listed* Insects Before They Destroy Vegetables
- May Be Used Up to Day of Harvest on Tomatoes
- Effective Against Broad Range of Listed* Pests
- Quickly Stops Harmful Listed* Insects Before They Do Damage
- Kills Household Nuisance Pest Outdoors
- Kills Listed* Insects on Contact
- · Keeps on killing for up to 14 days
- Kills stink bugs

{Note to PM: Marketing may choose any claims from the marketing claims list, Pg 15.}

| KILLS | Ants | Elm leaf beetles | Potato aphids |
|-----------------|--|--|------------------------|
| | Aphids | Flea beetles | Potato leafhoppers |
| | Armyworms | Flies | Scale (crawler stages) |
| | Aster leafhoppers | Garden webworms | Seed weevils (adults) |
| | Bagworms | Grasshoppers | Sod webworms |
| | Banded sunflower moths | Gypsy moth larvae | Sowbugs |
| | Boxelder bugs | Head clipper weevils (adults) | Spiders |
| | Budmoths | Japanese beetles (adults) | Spittlebugs |
| | Cabbage loopers | Lace bugs | Stem weevil (adults) |
| | Carrot weevils (adults) | Leaf beetles | Stink bugs |
| | Caterpillars | Leaf-feeding beetles | Sunflower beetles |
| | Celery leaf tiers | Leaf-feeding caterpillars | Sunflower moths |
| | Chinch bugs | Leafhoppers (including Glassy- | Thrips |
| | Clover mites | Winged Sharpshooter) | Tomato fruitworms |
| | Cockroaches | Leafminers | Tomato hornworms |
| | Colorado potato beetles | Mealybugs | Tomato pinworms |
| | Corn borers | Midges | Wasps |
| | Corn earworms | Mosquitoes | Weevils (adults) |
| | Corn rootworms (adults) | Pepper weevils (adults) | Western flower thrips |
| | Crickets | Pillbugs | Whiteflies |
| | Cutworms | Plant bugs | |
| | Earwigs | | |
| WHERE TO USE | [Roses,] [Flowers,] Vegetable and flower gardens, shrubs, ground covers, lawns and around homes | | |
| WHEN TO USE | When insects or damage first appear | | |
| | For severe infestations, repeat every 7-14 days as specified. | | |
| WHEN TO HARVEST | See plant specific restrictions for pre-harvest interval. If no pre-harvest interval is listed, the product may be applied up to and including day of harvest. | | |
| | For questions or comments, call tol | I-free 1-(877) 229-3724 or visit our w | rebsite at |
| | [www.BioAdvanced.com] | | |

ACTIVE INGREDIENT:

| Cyfluthrin | 0.003 % |
|--------------------|-----------|
| OTHER INGREDIENTS: | 99.997 % |
| TOTAL: | 100.000 % |

{Note to PM: Other ingredients include: Water, organic solvent (less than 0.1%, to dissolve the active ingredient for mixing) and surfactants (to keep the active ingredient dispersed in water).}

[Cyfluthrin is the chemical name for Tempo®]

EPA Reg. No. 92564-18

EPA Est. No. XXXX-AA-1

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

- Harmful if swallowed or inhaled.
- Causes eye irritation.
- Avoid contact with skin, eyes, or clothing.
- · Avoid breathing spray mist.
- Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

- This pesticide is toxic to fish. Do not apply directly to water.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.
- This pesticide is highly toxic to bees. Do not apply TEMPO Ready-to-Use Insecticide when bees are active.
- Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For best results, read and follow all label directions.

Open resealable label for directions & precautions.

BEFORE YOU USE


Read and follow these directions when using:


- · Apply the product to flowering plants during early morning or late evening when bees are not present.
- Do not spray the product into fish pools, ponds, streams, or lakes.
- For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds
- Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.
- Do not allow the product to enter any drain during or after application.
- Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios
 except as a spot or crack-and-crevice treatment.
- Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours)
- Rainfall within 24 hours after application may cause unintended runoff of pesticide application.
- Do not apply when the wind speed is greater than 15 mph.
- Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.)
- Treat surfaces to ensure thorough coverage but avoid runoff.
- Do not use in small, enclosed areas where spray can be inhaled.
- · Do not apply or irrigate to the point of run-off.
- Do not make application during rain.
- All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:
 - (1) Applications to pervious surfaces such as lawns, turf, and other vegetation;
 - (2) Applications to vertical surfaces (such as the side if a man-made structure) directly above impervious surfaces (e.g. driveways, sidewalks etc.) upto 2 feet above ground level.

(3) Applications to vertical surfaces directly above pervious surfaces, such as soil, lawn, turf, mulch or other vegetation) only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters. WHEN TO USE When insects or damage first appear. **HOW TO USE** Shake well before using. [Option 1] [Adjust nozzle to desired spray pattern.] Hold sprayer about 12 inches from foliage. Spray leaves, stems, and tender new shoots. To kill insects that may hide, spray the underside of the leaves. Apply until leaves are evenly coated with spray, but not dripping. [Option 2] [Nested Sprayer Directions for Use:] Attach Sprayer Remove sprayer from bottle and extend hose. ■ Adjust nozzle to stream ■ Turn sprayer nozzle or spray position. to X (off) position. Open flip-cap on bottle. ■ Point sprayer downward Do not unplug hose. and squeeze trigger repeatedly until liquid is ■ Insert hose and plug into Return sprayer to bottle. Sprayer will snap into placeflip-cap opening until it snaps into place. [Option 3] [Directions for Use for 1 Gallon Nested Sprayer] [Attach Sprayer •Remove sprayer from bottle. Remove twist tie. •Pull down and extend hose. Open flip-cap on bottle. •Insert hose and plug into flip-cap opening until it snaps into place. ■Move tab to open position. **Spray** Adjust nozzle to stream or spray position. Point sprayer downward and squeeze trigger repeatedly until liquid is released. After Use Turn sprayer nozzle to X (off) position. Do not unplug hose. Return sprayer to bottle. Sprayer will snap into place.] [Option 4] [1.3 Gallon ERGO-GRIP™ Sprayer Directions for Use:] [Remove ERGO- GRIP™ spray wand from pocket. Remove protective battery strip from sprayer handle.









CONNECT HOSE TO CAP

- Remove wrap from coiled hose.
- In down position, plug at end of hose into flip-spout opening on the cap until it snaps into place.
- Flip up spout until spraying.
 fully upright. Spout must remain upright while spraying.

OPEN SPRAY WAND

- Unfold spray wand until it snaps into place
- Point spray nozzle away from body.
- Adjust spray nozzle to desired setting.

TURN POWER ON

Slide OFF-ON switch to ON position.

SQUEEZE

- Squeeze trigger to dispense coiled product.
 Sprayer may require insert several seconds to prime before product is released.
- · Release trigger to stop

Storage

- Slide OFF-ON switch to OFF position.
- Return flip spout on cap to the down position. Do not unplug hose. Turn spray nozzle to the off position.
- Fold spray wand down against sprayer handle.
- Return ERGO-GRIP™ spray wand to pocket for storage.

4 AA Batteries Included

For important battery information - see booklet.

To Replace Batteries

- Open battery compartment [with a small screwdriver].
- Remove used batteries and replace with new batteries in correct position as indicated inside battery compartment.
- Securely close battery compartment door [and screw closed].

Proper use of Batteries

- Never mix old batteries with new.
- Do Not mix Alkaline with Standard (carbon zinc) or Rechargeable (nickel cadmium) batteries.
- When storing sprayer for long periods, remove batteries to prevent leakage. Dispose of used batteries according to manufacturer's instructions or in household trash.]

[Option 5]

[PROPOSED GENERIC WAND-STYLE SPRAYERS]

PREPARE TO SPRAY

- Remove spray wand from carrier. Extend hose to full length.
- Remove safety tab to activate batteries.
- [Connect hose to bottle by removing cap cover and inserting hose plug into cap, pushing firmly until

secure.]

•[Connect hose to bottle by removing existing cap and applying hose-end cap, turning until secure.]

SPRAY

- Unfold spray wand until it snaps into place.
- Point spray nozzle away from body.
- [Unlock trigger.]
- Squeeze to spray. Sprayer may require several seconds to prime before spraying begins. [If necessary, adjust nozzle to desired setting.]

AFTER USE

- [Move trigger lock to CLOSED position.]
- Fold wand down against handle.
- Return sprayer to carrier for storage.

[PROPOSED GENERIC GUN-STYLE SPRAYERS]

PREPARE TO SPRAY

- Remove sprayer from carrier. Extend hose to full length.
- Remove safety tab to activate batteries.
- •[Connect hose to bottle by removing cap cover and inserting hose plug into cap, pushing firmly until secure.]
- •[Connect hose to bottle by removing existing cap and applying hose-end cap, turning until secure.]

SPRAY

- Point spray nozzle away from body.
- [Unlock trigger.]
- Squeeze to spray. Sprayer may require several seconds to prime before spraying begins. [If necessary, adjust nozzle to desired setting.]

AFTER USE

- [Move trigger lock to CLOSED position.]
- Return sprayer to carrier for storage.

HOW OFTEN TO USE

Kills insects on contact and keeps on killing for up to two weeks. For severe infestations, repeat every 7 to 14 days as necessary.

FLOWERS, SHRUBS, GROUND COVERS and OUTDOOR CONTAINER PLANTS

Aphids Grasshoppers
Bagworms Gypsy moth larvae

Boxelder bugs Japanese beetle (adults)

Clover mites Lace bugs

Elm leaf beetles Leaf-feeding caterpillars

Leaf-feeding beetles Sca

Leafhoppers (including Glassy-winged Sharpshooter)

Mealybugs Pillbugs Scale

(crawler stages)
Spittlebugs

Thrips Whiteflies

HOW TO APPLY

- Spray foliage, tender new shoots and the underside of leaves thoroughly.
- To kill insects that hide, spray the underside of the leaves. Apply until leaves are evenly coated with spray droplets, but not dripping.

CONTAINER PLANTS [HOUSEPLANTS TREATED OUTSIDE]

HOW TO APPLY

- Move houseplants [indoor container plants] outside to treat. Spray foliage, tender new shoots and the underside of leaves thoroughly.
- To kill insects that hide, spray the underside of the leaves. Apply until the leaves are evenly coated with spray droplets, but not dripping. When the leaves are dry, move plant back inside.

HOME GARDENS

Peanut

After spraying, wait 14 days before harvesting.

Do not spray more than 3 times per season.

PEST[S]

Armyworm (1st and 2nd instar) Fall armyworm (1st and 2nd Potato leafhopper Thrips

Bean leaf beetle restary Rednecked peanutworm Vegetable weevil

Beet armyworm (1st and 2nd Green cloverworm Stink bugs Velvetbean caterpillar

instar) Grape colaspis (adult) Threecornered alfalfa hopper Whitefringed beetle (adult)

Cutworms Grasshoppers Southern armyworm (1st and 2nd Whitefly (adult)

Corn earworm Japanese beetle (adult) instar)

Corn earworm Japanese beetle (adult) instar)
Corn rootworms (adult) June beetle (adult) Soybean looper

Sunflowers

After spraying, wait 30 days before harvesting.

Do not spray more than 3 times per season.

Banded sunflower moth
Cutworms
Stink bugs
Sunflower moth
Grasshoppers
Palestripped flea beetle
Sunflower headclipping weevil
Sunflower stem weevil (adult)

VEGETABLE CROPS

Brassica (Cole) Leafy Vegetables:

Includes all members of Crop Group 5:

Broccoli, Broccoli raab (rapini), Chinese (gai lon) broccoli, Brussels sprouts, Cabbage, Chinese (bok choy) cabbage, Chinese (napa) cabbage, Chinese mustard (gai choy) cabbage, Cauliflower, Cavalo broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens, and Turnip greens.

No waiting to harvest.

Do not spray more than 4 times per season.

| Alfalfa looper | Corn earworm | Japanese beetle (adult) | Southern cabbageworm |
|-------------------------------|----------------------------|--------------------------------|-----------------------------|
| Armyworm (1st and 2nd instar) | Cutworms | Lygus bug | Tarnished plant bug |
| Beet armyworm (1st and 2nd | Diamondback moth (larvae) | Meadow spittlebug | Thrips |
| instar) | Fall armyworm (1st and 2nd | Potato leafhopper | Vegetable weevil (adult) |
| Cabbage flea beetle | instar) | Southern armyworm (1st and 2nd | Whitefly (adult) |
| Cabbage looper | Grasshoppers | instar) | Yellowstriped armyworm (1st |
| Cabbage webworm | Imported cabbageworm | Stink bugs | and 2 nd instar) |

Cucurbits (except crops grown for seed):

Includes all members of Crop Group 9:

Balsam apple, Balsam pear, Bitter melon, Chayote, Chinese cucumber, Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Edible gourd (includes: hyotan, cucuzza, henchmia and Chinese okra), Muskmelon (includes: cantaloupe, true cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), Pumpkin, Summer squash (includes: crookneck squash, scallop squash, straightneck squash, vegetable marrow, and zucchini), Watermelon, Winter squash (includes: butternut squash, calabaza, hubbard squash, acorn squash and spaghetti squash)

No waiting to harvest.

Do not spray more than 4 times per season.

PEST[S]

Armyworm (1st and 2nd instar) Cutworms Pickleworm Stripped cucumber beetle Brown marmorated stink bug Cucumber beetle Potato leafhopper Tarnished plant bug Tobacco budworm Cabbage looper Grasshoppers Rindworm Whitefly (adult) Colorado potato beetle Lygus bug Stink bugs Corn earworm Melonworm

Fruiting Vegetables

Includes all members of Crop Group 8:

Eggplant, Groundcherry, Pepino, Pepper (includes: bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo, and Tomato

No waiting to harvest for tomato. After spraying, wait 7 days before harvesting for all other crops.

Do not spray more than 6 times per season.

PEST[S]

Beet armyworm (1st and 2nd Flea beetles Southern armyworm (1st and 2nd Tomato hornworm instar) instar) Garden symphylan Tomato pinworm Brown marmorated stink bug Stink bugs Garden webworm Variegated cutworm Tarnished plant bug Cabbage looper Leafminers Whitefly (adult) Thrips (except Thrips palmi) Celery leaftier Western yellowstriped Pepper weevil Tomato fruitworm (corn armyworm (1st and 2nd instar) Colorado potato beetle Potato aphid earworm) European corn borer Potato leafhopper

Leafy Vegetables:

Includes all members of Crop Group 4:

Amaranth (Chinese spinach), Arugula (rouquette), Cardoon, Celery, Chinese celery, Celtuce, Chervil, Chrysanthemum (edible-leaved and garland), Corn salad, Cress (garden and upland), Dandelion, Dock (sorrel), Endive (escarole), Florence fennel, Lettuce (head and leaf), New Zealand spinach, Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Rhubarb, Spinach, Swiss chard, Vine spinach

No waiting to harvest. Do not spray more than 4 times per season.

Alfalfa looper Diamondback moth (larvae) Japanese beetle (adult) Stink bugs

Beet armyworm (1st and 2nd European corn borer Leafhoppers Tarnished plant bug

instar) Fall armyworm (1st and 2nd Lygus bug

Brown marmorated stink bug instar) Meadow spittlebug Vegetable weevil (adult)

Cabbage looper Flea beetles Saltmarsh caterpillar Whitefly (adult)

Corn earworm

Grasshoppers

Southern armyworm (1st and 2nd

Cutworms

Green cloverworm

Southern armyworm (1st and 2nd
instar)

Yellowstriped armyworm (1st and 2nd instar)

Thrips

Cutworms Green cloverworm instar) and 2nd In Colorado potato beetle Imported cabbageworm

Potato and Other Tuberous and Corm Vegetables:

Includes all members of Crop Subgroup 1C:

Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Edible canna, Cassava (bitter and sweet), Chayote root, Chufa, Dasheen (taro), Ginger, Leren, Potato, Sweet potato, Tanier, True yam, Turmeric, Yam bean

No waiting to harvest.

Do not spray more than 6 times per season.

PEST[S]

Aphids European corn borer Potato psyllid Sweetpotato weevil (adults)

Cabbage looper Flea beetles Potato tuberworm Tarnished plant bug

Cutworms Colorado Potato Beetle Potato Leafhopper

Root Vegetables (except sugarbeet):

Includes all members of Crop Subgroup 1B:

Garden beet, Edible burdock, Carrot, Celeriac, Turnip-rooted chervil, Chicory, Ginseng, Horseradish, Turnip-rooted parsley, Parsnip, Radish, Oriental radish, Rutabaga, Salsify (black, Spanish, and oyster plant), Skirret, Turnip

No waiting to harvest.

Do not spray more than 5 times per season.

PEST[S]

Aster leafhopper Cutworms Flea beetles Potato leafhopper

Carrot weevil

Sweet Corn

No waiting to harvest.

Do not spray more than 10 times per season.

PEST[S]

Armyworm (1st and 2nd instar) Corn silk fly Japanese beetle(adult) Southwestern corn borer

Bean leaf beetle European corn borer June beetle (adult) Stalk borer

Black cutworm Fall armyworm (1st and 2nd Leafhoppers Stink bugs

Cereal leaf beetle Masked chafer (adult) Webworm

Chinch bug Flea beetle Sand hill cutworm Western bean cutworm

Click beetle (adult)

Granulate cutworm

Southern armyworm (1st and 2nd Yellowstriped armyworm (1st an

Corn earworm Grape colaspis (adult) instar) and 2nd instar)

Corn rootworms (adult) Grasshoppers Southern corn leaf beetle

TREE and VINE CROPS

Grapes

Includes: Table grape, Raisin, Wine and Muscadine grape

After spraying, wait 3 days before harvesting Do not spray more than 4 times per season

PEST[S]

Climbing cutworm Grape flea beetle Grape mealybug (crawlers) Orange tortrix

Grape berry moth Grape leaffolder Glassywinged sharpshooter Thrips

Grape bud beetle Grape leafhopper Grape leaf skeletonizer Variegated leafhopper
Grape cane gallmaker (adult) Grape leafroller Omnivorous leafroller Western grape leaf skeletonizer

FOR BEST RESULTS:

Do not water the foliage within one hour after applying. To avoid serious damage to plants, eliminate pests early in the season before they multiply.

HOUSEHOLD NUISANCE PESTS OUTDOORS

PEST[S]

AntsCricketsMidgesSowbugsBoxelder bugsEarwigsMosquitoesSpidersClover mitesFliesPillbugsWasps

Cockroaches Brown Marmorated Stink Bug

HOW TO APPLY

Spray around doors, window frames and other places where insects may enter house. Spray around foundations and porches. Spray on ant hills and ant trails.

LAWN PESTS

PEST[S]

Ants Crickets Earwigs Japanese beetles (adults)
Armyworms Cutworms Grasshoppers Sod Webworms (lawn moths)

Chinch bugs

HOW TO APPLY

[For localized infestations, spray to penetrate the lawn and reach the soil. Do not mow for one day after application. For large areas, use Tempo® Concentrate Insecticide.]



Do not allow entry into treated area until spray has dried.

{Note to PM: Any or all of the following listed pests/insects may appear on the marketplace label}

| Storage and Disposal | | |
|-------------------------------|--|--|
| Pesticide Storage Conditions: | Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing. | |
| Pesticide Disposal: | Non-refillable container. Do not reuse or refill empty spray bottle. | |
| <u> </u> | If empty: Place in trash or offer for recycling if available. | |
| [[[]]] | If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain. | |

Important: It is best to use all of the product in accordance with label directions. If it is necessary to dispose of unused product, please follow any applicable state or local guidelines.

NOTICE

Research and testing have determined that the "Directions for Use" are appropriate for the proper use of TEMPO® Ready-to-Use Insecticide under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

+++ 100% satisfaction guaranteed or your money back

{Note to PM: Batch code to be added at time of production.}

{Note to PM: Product name Tempo® Ready-to-Use will be replaced with marketed Alternate Brand Name [it] [this product].} {Note to PM: For a list of pests/insects that [our/this product/insert product name] targets visit: www.bioadvanced.com

BioAdvanced® is a registered trademark of SBM Life Science Corp.

[Fortifend Advanced[™]] [Bianova Advanced[™]] [is/are] [a] trademark of SBM Life Science Corp.

[PRODUCED FOR] [PRODUCED BY] [DISTRIBUTED BY] [GUARANTEED BY]

SBM Life Science Corp. 1001 Winstead Drive, Suite 500 Cary, North Carolina 27513

Tempo RTU Insecticide (Master) 11/03/2020

Marketing Claims

Pesticide Claims

- 1. Guaranteed Satisfaction or your money back
- 2. Ready To Use
- 3. Kills Listed* Insects
- 4. Multipurpose Insecticide
- Fast Acting...Kills Listed* Insects On Contact [Insert Graphic +*]¹
- 6. Long Lasting...Keeps On Killing For Up To 14 Days1
- 7. Fast Acting¹
- 8. Long Lasting For Up To 2 Weeks1
- 9. Kills outdoor listed* pests on contact
- 10. Lasts up to 14 days
- 11. For use on flowers, shrubs, and listed* vegetables
- 12. Just One Application Quickly Kills Listed* Insects
- 13. Keeps On Killing For Up To 14 Days
- 14. Use On Tomatoes Up To Day Of Harvest [Insert Graphic ¥]
- 15. Low odor [formula]
- 16. Won't harm plants
- 17. Protects plants
- 18. Water-based
- 19. Kills Beetles (including Japanese Beetle adults), [Caterpillars (including Tomato Hornworm),] [Ants,] [Armyworms,] [Ticks (including ticks carrying Lyme disease)] and [&] other listed* garden and [&] nuisance pests [insects] [bugs]
- 20. Kills Beetles, Caterpillars, Ants, [Ticks] and [&] other listed* insects
- 21. Kills [Controls over] [more than] 100 listed* insects [bugs] [pests] on vegetables, flowers, small trees and [&] shrubs
- 22. Kills [Controls over] [more than 100 listed* garden and nuisance pests without harming roses, flower, or shrubs
- 23. Kill listed* insects on Contact
- 24. Kills On Contact
- 25. Fast Acting......Kills On Contact1
- 26. [For] use on Houseplants [& Container plants](treated outside)
- 27. Kills [insert any listed insect/pests]
- 28. Protect your houseplants [& container plants] (treated outside) from damaging [destructive] insects
- 29. For use on flowers, shrubs & container plants
- 30. Protect your flowers from destructive [damaging] insects
- 31. Kills over 100 listed* insects [bugs] [pests] on contact
- 32. On [For] [For use on] Roses, Flowers, Small trees, Shrubs, Fruits, Vegetables and [&] other Outdoor Surface [Areas]
- 33. Can be used on [most popular] fruits and [&] vegetables (see back panel for details)
- 34. User in (on) Vegetable Gardens
- 35. Kills Caterpillars, Beetles and [&] Aphids
- 36. Fast Acting Formula¹
- 37. Reapply as directed for a more beautiful garden
- 38. Guaranteed Kill or your money back+++
- 39. Keeping your vegetable garden and lawn free of listed[*] pests has never been easier
- 40. Kills aphids, Japanese beetles (adults), cutworms, tomato hornworms, and other listed* pests
- 41. Also for use on lawns
- 42. Kills listed* insects on contact [& protects up to 14 days]
- 43. Protects up to 14 days
- 44. This product provides up to 14 days protection with a single application. Now keeping your roses and flowers free of listed pests has never been easier
- 45. Kills 100+ listed* insects on contact
- 46. Kills Aphids, Japanese Beetles (adult), Leafminers, Whiteflies, [Thrips], [Scale (crawler stages)] and other listed* pests [insects] [bugs]
- 47. On [For] [For use on] Roses and Flowers
- 48. Kills damaging insects [that eat your tomatoes]
- 49. Save your tomatoes [& vegetables] from insects
- 50. Kills Tomato Hornworms
- 51. Kills Tomato Pinworms
- 52. Stop insect infestation
- 53. Stop insects from eating your tomatoes [& vegetables]
- 54. Don't let worms eat your tomatoes [& vegetables]
- 55. Stop worms from eating your tomatoes [& vegetables]
- 56. Protect your tomatoes [& vegetables] from destructive[damaging] pests

Marketing Claim for Brand Name Change: 1. New Name. Same Great Results.

- 2. New Name. Same Familiar Formula.
- 3. New Name. Same Performance.
- 4. New Look. Same Great Formula.
- 5. New Look. Same Great Results.

{Note to reviewer - "New" may be added anywhere to a claim and will only be used for the first 6 months of product launch}

¹ except Ants ,Flies, Mosquitoes, Spiders and Wasps

^{* [}The word "listed" can appear as a footnote with the appropriate phrase, such as listed[*] insects/pests]

[ART]

| READ | PROTECTION FOR YARD AND VALUE USE THE PROTECTION FOR YARD THE PROTEC |
|-----------------------------------|--|
| LAB-DEVELOPED SOLUTIONS OUR YARD | OUR LABS |
| +++ your money back | CUTWORMS |
| TOMATO HORNWORMS | LEAFY VEGETABLES |
| ADULT JAPANESE BEETLES | APHIDS (Control of the control of th |

| FRUITING VEGETABLES | ROOT VEGETABLES | CUCURBITS |
|---|-----------------|-----------|
| USE ON TOMATOES UP TO DAY OF HARVEST ¥ | +* | |