



## OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

January 17, 2024

Michael Barber  
Regulatory Affairs Specialist  
SBM Life Science Corp.  
1001 Winstead Drive Suite 500  
Cary, NC 27513

Subject: Label Amendment - Registration Review Mitigation for Fenoxaprop-p-ethyl and Mecoprop-p  
Product Name: LAWN HERBICIDE 3F CONCENTRATE  
EPA Registration Number: 92564-34  
Application Dates: July 5, 2022, and December 22, 2020  
Decision Numbers: 585602, 592193

Dear Michael Barber:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Fenoxaprop-p-ethyl and Mecoprop-p Interim Decisions, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling

before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact DeMariah Koger by phone at (202)-566-2288, or via email at [Koger.demariah@epa.gov](mailto:Koger.demariah@epa.gov).

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington", with a stylized flourish at the end.

Linda Arrington, Branch Chief  
Risk Management and Implementation Branch 4  
Pesticide Re-Evaluation Division  
Office of Pesticide Programs

ENCLOSURE: Stamped label

(Front Panel)

# Bayer Advanced Lawn Herbicide 3F Concentrate

This label contains two separate label statements for Outdoor Residential Pump sprayer/Hose-end Sprayer Concentrate product use and Outdoor Residential Ready-To-Spray product use.

- Kills Lawn Weeds plus Crabgrass [in One Step]
- Kills Over 200 Lawn Weeds Plus Crabgrass in One Easy Step
- Kills Crabgrass plus Broadleaf Weeds
- Kills Listed Weeds
- Kills Hard-to-Control Weeds
- Systemic [Herbicide] [Activity]
- Kills Both Broadleaf and Grassy Weeds
- Kills Dandelion, Crabgrass, Clover, plus 200 Other listed Weeds
- Rainproof in [1] hour[s]
- New!
- Two-in-One Product that Kills Crabgrass & Broadleaf Weeds
- Two Types of Weeds; One Solution
- Guaranteed\* Not to harm your lawn
- All-In-One Weed Killer [for lawns]
- Starts working on contact
- Kills even tough weeds – roots & all
- Kills 200+ Weeds
- Unique Formula
- Fast results
- Kills lawn weeds plus crabgrass
- Gets the root
- Won't Harm [your] lawn [s]
- Will not harm your lawn
- Kills tough weeds
- Kills the weeds, not your lawn
- Kills to the root
- Kills weeds, not lawns
- One [easy] step
- Two products in one
- Kills all major broadleaf weeds
- Weed protection system
- Visible results
- Kills weeds - Fast
- Kills both Crabgrass & Broadleaf weeds
- Proprietary Formula that Kills Both Crabgrass and Broadleaf Weeds
- Effectively Kills Both Broadleaf and Weed grasses
- Save time and energy
- Always ready to spray
- Easy to use
- Fast coverage
- Easy & Fast coverage
- Targets weeds
- Kills weeds the first time, every time
- Guaranteed\* not to harm lawns
- Guaranteed\* Results
- 100% Satisfaction Guaranteed\* (or your money back)
- For use on Kentucky bluegrass, perennial ryegrass, fescue (except creeping fescue), centipedegrass, St. Augustinegrass (except Floratam), zoysiagrass
- Tough weed protection
- 4-way weed control action
- Exclusive 4-Way Weed control
- Proprietary [unique] [advanced] formula
- Attach to hose and spray

**Active Ingredients:**


2,4-D, Triisopropanolamine salt.....	4.91%
Mecoprop-p, Potassium salt.....	1.55%
Dicamba, Potassium salt .....	0.77%
Fenoxaprop-p-ethyl† .....	0.41%
<b>Other Ingredients</b> .....	<u>92.36 %</u>
	100.00%

†Protected by U.S. Patent No. 5,254,527  
Net 32 fl oz. (946 mL)

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

See back panel for additional precautionary statements

Editorial Note – [Bracketed text] is optional language



(Pump Sprayer/Hose-end Sprayer label)

(FRONT PANEL)

# Bayer Advanced Lawn Herbicide 3F Concentrate

[Editorial Note – Marketing Claims from page 1]

---

**Active Ingredients:**

2,4-D, Triisopropanolamine salt.....	4.91%
Mecoprop-p, Potassium salt.....	1.55%
Dicamba, Potassium salt .....	0.77%
Fenoxaprop-p-ethyl†.....	0.41%

<b>Other Ingredients</b> .....	<u>92.36 %</u>
	100.00%

---

†Protected by U.S. Patent No. 5,254,527

Net 32 fl oz. (946 mL)

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**



**See back panel for additional precautionary statements**

## (Back Panel)

### OPEN Resealable Label for Directions & Precautions PRESS TO RESEAL

Bayer Advanced Lawn Herbicide 3F Concentrate's exclusive and unique formula combines 4 proven weed killers to kill over 200 types of lawn weeds plus crabgrass in one easy step. And it's specially formulated so it won't harm your lawn when used as directed. Rainproof in 1 hour. Controlling weeds plus crabgrass in your lawn has never been easier!



QUICK FACTS	
Marketing claims to be selected from the front page	
<b>KILLS</b>	Broadleaf Weeds: Over 200 types including dandelions, clovers, chickweeds Grassy Weeds: Crabgrass, goosegrass, foxtail, sandbur, and many others
<b>WHERE TO USE</b>	On lawns without harming lawn grasses – Centipedegrass, Fescue (except Creeping Fescue), Kentucky Bluegrass, Perennial Ryegrass, St. Augustinegrass (except Floratam), Zoysiagrass (DO NOT USE on Bahiagrass, Bermudagrass, Buffalograss, Carpetgrass, Creeping Fescue, Floratam variety of St. Augustinegrass, Dichondra or desirable clovers.)
<b>[COVERAGE]</b>	[This bottle treats 6,400 sq ft.]
 	For questions or comments, call toll-free 1-877 BayerAG (1-877-229-3724) or visit our website at <a href="http://www.BayerAdvanced.com">www.BayerAdvanced.com</a>

### FOR OUTDOOR RESIDENTIAL USE ONLY

#### Active Ingredients:

2,4-D, Triisopropanolamine salt.....	4.91%
Mecoprop-p, Potassium salt.....	1.55%
Dicamba, Potassium salt .....	0.77%
Fenoxaprop-p-ethyl† .....	0.41%
<b>Other Ingredients</b> .....	<b>92.36 %</b>
	100.00%

THIS PRODUCT CONTAINS: 2.64% 2,4-D acid equivalent, 1.32% Mecoprop-p acid equivalent, 0.66% Dicamba acid equivalent.  
Isomer Specific by AOAC Methods.

EPA Reg. No. 92564-34

EPA Est. No. XXX-AA-XX

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

<p><b>HOW MUCH TO USE/ USE RESTRICIONS</b></p>	<p>Measure area to be sprayed. Calculate 'square feet' by multiplying length times width. [This bottle will treat 6,400 sq. ft.] The maximum application rate to turf is 1.0 pound of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.</p>
<p><b>HOW TO USE</b> <b>Shake Well Before Use</b></p>	<p><b>PUMP-UP TANK SPRAYERS (Best for spot spraying individual weeds)</b></p> <ol style="list-style-type: none"> <li>1. Add appropriate amounts of water &amp; product to sprayer tank – 3 fl. oz. (6 Tbs) product in 1 gallon of water treats 500 sq. ft.</li> <li>2. Close sprayer, shake well &amp; pressurize. Adjust nozzle to a coarse spray. Large Areas – Apply evenly over area. Individual Weeds – Hold tip of wand about 12 inches from weed, aim at center of weed and spray.</li> <li>3. Occasionally re-pressurize sprayer to maintain a good spray pattern.</li> </ol> <p><b>DIAL STYLE HOSE-END SPRAYERS (Best for spraying large areas of the lawn)</b></p> <ol style="list-style-type: none"> <li>1. Fill sprayer jar with enough product for area to be treated – 3 fl. oz. (6 Tbs) product treats 500 sq. ft. <b>DO NOT ADD WATER!</b></li> <li>2. Close sprayer, set dial and apply.</li> <li>3. Pour any unused product back into this original container.</li> </ol> <p><b>CONCENTRATE-PLUS-WATER HOSE-END SPRAYERS (Best for spraying large areas of the lawn)</b></p> <ol style="list-style-type: none"> <li>1. Pour product into sprayer jar to the required fluid ounce level – 3 fl oz (6 Tbs) treats 500 sq. ft.</li> <li>2. Add the appropriate amount of water to the gallons level.</li> <li>3. Replace sprayer top, shake well and apply.</li> </ol>
<p><b>[FOR BEST RESULTS]</b></p>	<ul style="list-style-type: none"> <li>• [Spray when weeds are [small [young] and] actively growing.]</li> <li>• [Spray when daily high temperature is below 90°F.]</li> <li>• [Rain or watering 1 hour after application will not wash away effectiveness.]</li> <li>• [Apply when wind is calm [to avoid spray drift onto desirable plants (fruits, vegetables, flowers, groundcovers, shrubs &amp; trees)]]</li> <li>• [Reapply in 2 weeks if necessary.]</li> <li>• [Do not apply while grass is emerging from winter dormancy.]</li> <li>• [Newly seeded lawns may be sprayed after the fourth mowing.]</li> <li>• [Bare spots may be seeded 4 weeks after application.]</li> <li>• [Do not mow for 1 to 2 days before &amp; after spraying [Uncut weeds have more surface area to hold more weed killer. They are also more likely to be hit by the spray. Lawn clippings can get in the way of the herbicide hitting the weed. Weed killers need time to move throughout the weed – mowing may remove the treated parts.]]</li> <li>• [Leave grass clippings on the lawn from the first 3 mowings after using this product. Do not use as compost or mulch.]</li> <li>• [If lawn is dry, water 1 to 2 days before using this product. [Weeds take up herbicides best when not under stress.]]</li> </ul>
<p><b>HOW IT WORKS</b></p>	<p>BAYER ADVANCED LAWN HERBICIDE 3F CONCENTRATE contains four proven weed killers that target lawn weeds and crabgrass. The herbicides enter the lawn weeds through their leaves and move inside the weeds, even down into the roots. And lawns will not be harmed when used as directed.</p>

Kills over 200 weeds including:

<p>Black medic Carpetweeds Chickweeds Clovers Crabgrass Creeping Charlie Cudweed Curly dock Dandelion English daisy False dandelion Filaree</p>	<p>Florida pusley Foxtails Ground ivy Heartleaf drymary Henbit Kikuyugrass Oxalis Pennywort Plantains Purslane Red sorrel Sheep sorrel Spurges</p>	<p>Spurweed Thistles Toadflax Wild carrot Wild garlic Wild geranium Wild onion &amp; many other broadleaf &amp; grass weeds (See appendix for a complete list of the weeds controlled)</p>
---	--	--



Do not apply this product in a way that will contact any person, pet either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until spray has dried.

## PRECAUTIONARY STATEMENTS

### CAUTION

#### Hazards to Humans and Domestic Animals

- Harmful if swallowed
- Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing.

After using this product:

- If product gets on clothing, remove clothing as soon as possible and launder clothing separately before reuse.
- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

### FIRST AID +

<b>If in Eyes</b>	<ul style="list-style-type: none"><li>▪ Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li><li>▪ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>▪ Call a poison control center or doctor for further treatment advice.</li></ul>
<b>If Swallowed</b>	<ul style="list-style-type: none"><li>▪ Call a poison control center or doctor immediately for treatment advice.</li><li>▪ Have person sip a glass of water if able to swallow.</li><li>▪ Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>▪ Do not give anything by mouth to an unconscious person.</li></ul>


**NOTE:** When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 1-877-229-3763 for medical emergency information.

Active Ingredients: 4.91% 2,4-D Triisopropanolamine salt (CAS # 32341-80-3), 1.55% Mecoprop-p Potassium salt single isomer (CAS # 66423-5-0), 0.77% Dicamba, Potassium salt (CAS # 10007-85-9), and 0.41% Fenoxaprop-p-ethyl (CAS # 71283-80-2).

### ENVIRONMENTAL HAZARDS

- This product is toxic to fish & aquatic invertebrates.
- Drift or runoff may harm fish, aquatic invertebrates and non-target plants.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.
- **NON-TARGET ORGANISM ADVISORY:** This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

## Important – Directions for Storage and Disposal

<b>Storage Conditions</b>	Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing
<b>Disposal</b> 	<b>If empty:</b> Non-refillable container. Do not refill or reuse container. Place empty container in trash or offer for recycling if available. <b>If partly filled:</b> Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.

**NOTICE:** Research and testing have determined that the “Directions for Use” are appropriate for the proper use of this product under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

**\*Money Back Guarantee:** If you are not satisfied with this product, we will gladly refund your original purchase price.

**SBM Life Science Corp.**  
**1001 Winstead Drive, Suite 500**  
**Cary, NC 27513**

Lawn Herbicide 3F Concentrate (PENDING) 12/22/2020



(Ready-To- Spray label)  
(FRONT PANEL)

# Bayer Advanced Lawn Herbicide 3F Concentrate

This label contains two separate label statements for, Outdoor Residential Pump Sprayer/Hose-end Sprayer Concentrate product use and Outdoor Residential Ready-To-Spray product use.

---

[Editorial Note – Marketing Claims from page 1]

---

**Active Ingredients:**

2,4-D, Triisopropanolamine salt.....	4.91%
Mecoprop-p, Potassium salt.....	1.55%
Dicamba, Potassium salt .....	0.77%
Fenoxaprop-p-ethyl†.....	0.41%
<b>Other Ingredients</b> .....	<u>92.36 %</u>
	100.00%

---

Net 32 fl oz. (946 mL)

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

See back panel for additional precautionary statements


†Protected by U.S. Patent No. 5,254,527

**(Back Panel)**

**OPEN Resealable Label for Directions & Precautions  
PRESS TO RESEAL**

Bayer Advanced Lawn Herbicide 3F Concentrate's exclusive and unique formula combines 4 proven weed killers to kill over 200 types of lawn weeds plus crabgrass in one easy step. And it's specially formulated so it won't harm your lawn when used as directed. Rainproof in 1 hour. Controlling weeds plus crabgrass in your lawn has never been easier!



<b>QUICK FACTS</b>	
<b>Marketing claims to be selected from the front page</b>	
<b>KILLS</b>	Broadleaf Weeds: Over 200 types including dandelions, clovers, chickweeds Grassy Weeds: Crabgrass, goosegrass, foxtail, sandbur, and many others
<b>WHERE TO USE</b>	On lawns without harming lawn grasses – Centipedegrass, Fescue (except Creeping Fescue), Kentucky Bluegrass, Perennial Ryegrass, St. Augustinegrass (except Floratam), Zoysiagrass (DO NOT USE on Bahiagrass, Bermudagrass, Buffalograss, Carpetgrass, Creeping Fescue, Floratam variety of St. Augustinegrass, Dichondra or desirable clovers.)
<b>[COVERAGE]</b>	[This bottle treats 6,400 sq ft.]
	For questions or comments, call toll-free 1-877 BayerAG (1-877-229-3724) or visit our website at <a href="http://www.BayerAdvanced.com">www.BayerAdvanced.com</a>

**FOR OUTDOOR RESIDENTIAL USE ONLY**

<b>Active Ingredients:</b>	
2,4-D, Triisopropanolamine salt.....	4.91%
Mecoprop-p, Potassium salt.....	1.55%
Dicamba, Potassium salt .....	0.77%
Fenoxaprop-p-ethyl† .....	0.41%
<b>Other Ingredients</b> .....	<b>92.36 %</b>
	100.00%


THIS PRODUCT CONTAINS: 2.64% 2,4-D acid equivalent, 1.32% Mecoprop-p acid equivalent, 0.66% Dicamba acid equivalent. Isomer Specific by AOAC Methods

**EPA Reg. No. 92564-34**


**EPA Est. No. XXX-AA-XX**

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

<p><b>HOW MUCH TO USE/ USE RESTRICIONS</b></p>	<p>Measure area to be sprayed. Calculate 'square feet' by multiplying length times width. This bottle will treat 6,400 sq. ft. The maximum application rate to turf is 1.0 pound of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.</p>
<p><b>HOW TO USE</b>  <b>Shake Well Before Use</b></p>	<p><b>CONNECT</b></p> <ol style="list-style-type: none"> <li>1. Connect sprayer to garden hose.</li> <li>2. Turn on water.</li> </ol> <p><b>SPRAY</b></p> <ol style="list-style-type: none"> <li>1. Extend hose to farthest area to be treated, point nozzle in direction to be sprayed.</li> <li>2. Turn on sprayer – &lt; <i>Insert sprayer-specific language here</i> &gt; <i>Calmar RTS sprayer:</i> press lock tab while using other hand to turn knob counter-clockwise to ON position. <i>Green Garden RTS sprayer:</i> turn sprayer valve to ON position. <i>Hayes HP-2 RTS sprayer:</i> press lock tab while using other hand to turn knob clockwise to ON position.</li> <li>3. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap the treated areas. Work back toward the faucet to avoid walking over wet, treated grass.</li> </ol>  <p><b>FINISH</b></p> <ol style="list-style-type: none"> <li>1. Turn off sprayer – &lt; <i>Insert sprayer-specific language here</i> &gt; <i>Calmar RTS sprayer:</i> turn knob clockwise to OFF position. <i>Green Garden RTS sprayer:</i> [Nothing required]. <i>Hayes HP-2 RTS sprayer:</i> turn knob counter-clockwise to OFF position.</li> <li>2. Turn off water at faucet.</li> <li>3. Relieve pressure in hose by turning sprayer knob back to ON position until water slows to a drip then turn sprayer knob to OFF position.</li> <li>4. Disconnect sprayer from hose.</li> </ol>
<p><b>[FOR BEST RESULTS]</b></p>	<ul style="list-style-type: none"> <li>• [Spray when weeds are [small [young] and] actively growing.]</li> <li>• [Spray when daily high temperature is below 90°F.]</li> <li>• [Rain or watering 1 hour after application will not wash away effectiveness.]</li> <li>• [Apply when wind is calm [to avoid spray drift onto desirable plants (fruits, vegetables, flowers, groundcovers, shrubs &amp; trees)]]</li> <li>• [Reapply in 2 weeks if necessary.]</li> <li>• [Do not apply while grass is emerging from winter dormancy.]</li> <li>• [Newly seeded lawns may be sprayed after the fourth mowing.]</li> <li>• [Bare spots may be seeded 4 weeks after application.]</li> <li>• [Do not mow for 1 to 2 days before &amp; after spraying [Uncut weeds have more surface area to hold more weed killer. They are also more likely to be hit by the spray. Lawn clippings can get in the way of the herbicide hitting the weed. Weed killers need time to move throughout the weed – mowing may remove the treated parts.].]</li> <li>• [Leave grass clippings on the lawn from the first 3 mowings after using this product. Do not use as compost or mulch.]</li> <li>• [If lawn is dry, water 1 to 2 days before using this product. [Weeds take up herbicides best when not under stress.]]</li> </ul>
<p><b>HOW IT WORKS</b></p>	<p>BAYER ADVANCED LAWN HERBICIDE 3F CONCENTRATE contains four proven weed killers that target lawn weeds and crabgrass. The herbicides enter the lawn weeds through their leaves and move inside the weeds, even down into the roots. And lawns will not be harmed when used as directed.</p>

Kills over 200 weeds including:		
Black medic Carpetweeds Chickweeds Clovers Crabgrass Creeping Charlie Cudweed Curly dock Dandelion English daisy False dandelion Filaree	Florida pusley Foxtails Ground ivy Heartleaf drymary Henbit Kikuyugrass Oxalis Pennywort Plantains Purslane Red sorrel Sheep sorrel Spurges	Spurweed Thistles Toadflax Wild carrot Wild garlic Wild geranium Wild onion & many other broadleaf & grass weeds (See appendix for a complete list of the weeds controlled)

	Do not apply this product in a way that will contact any person, pet either directly or through drift. Keep people and pets out of the area during application. Do not allow people (other than applicator) or pets to enter the treated area until spray has dried.
---	--

## PRECAUTIONARY STATEMENTS

### CAUTION

#### Hazards to Humans and Domestic Animals

- Harmful if swallowed
- Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing.

After using this product:

- If product gets on clothing, remove clothing as soon as possible and launder clothing separately before reuse.
- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

### FIRST AID


<b>If in Eyes</b>	<ul style="list-style-type: none"> <li>▪ Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>▪ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>▪ Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If Swallowed</b>	<ul style="list-style-type: none"> <li>▪ Call a poison control center or doctor immediately for treatment advice.</li> <li>▪ Have person sip a glass of water if able to swallow.</li> <li>▪ Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>▪ Do not give anything by mouth to an unconscious person.</li> </ul>

**NOTE:** When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 1-877-229-3763 for medical emergency information. Active Ingredients: 4.91% 2,4-D Triisopropanolamine salt (CAS # 32341-80-3), 1.55% Mecoprop-p Potassium salt single isomer (CAS # 66423-5-0), 0.77% Dicamba, Potassium salt (CAS # 10007-85-9), and 0.41% Fenoxaprop-p-ethyl (CAS # 71283-80-2).

## ENVIRONMENTAL HAZARDS

- This product is toxic to fish & aquatic invertebrates.
- Drift or runoff may harm fish, aquatic invertebrates and non-target plants.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.
- **NON-TARGET ORGANISM ADVISORY:** This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

## Important – Directions for Storage and Disposal

<b>Storage Conditions</b>	Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing
<b>Disposal</b> 	<b>If empty:</b> Non-refillable container. Do not refill or reuse container. Place empty container in trash or offer for recycling if available. <b>If partly filled:</b> Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.

**NOTICE:** Research and testing have determined that the “Directions for Use” are appropriate for the proper use of this product under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

**\*Money Back Guarantee:** If you are not satisfied with this product, we will gladly refund your original purchase price.

**SBM Life Science Corp.**  
1001 Winstead Drive, Suite 500  
Cary, NC 27513

Lawn Herbicide 3F RTS (PENDING) 12/22/2020

# Appendix

## WEEDS CONTROLLED

Annual Yellow Sweetclover	Cocklebur	Mouseear Hawkweed	Speedwell
Aster	Common Mullein	Mugwort	Spiny Amaranth
Austrian Fieldcress	Crabgrass , Large/Hairy	Musk Thistle	Spiny Cocklebur
Barnyardgrass	Crabgrass, Small/Smooth	Mustard	Spiny Sowthistle
Bedstraw	Creeping Charlie	Narrowleaf Plantain	Spotted Catsear
Beggarticks	Creeping Jenny	Narrowleaf Vetch	Spotted Spurge
Betony, Florida	Cudweed	Nettle	Sprangletop
Bindweed, Field	Curly Dock	Orange Hawkweed	Spurweed
Bird Vetch	Daisy, English	Oriental Cocklebur	Stinging Nettle
Bitter Wintercress	Daisy, Fleabane	Oxalis	Strawberry, India Mock
Bittercress, Hairy	Daisy, Oxeye	Panicum	Tall Nettle
Bitterweed	Dandelion	Parsley-piert	Tall Vervain
Black-eyed Susan	Dichondra	Parsnip	Tansy Mustard
Black Medic	Dogbane	Pearlwort	Tansy Ragwort
Black Mustard	Dogfennel	Pennycress	Tanweed
Blackseed Plantain	Dollarweed	Pennywort	Thistle
Blessed Thistle	Elderberry	Pepperweed	Trailing Crownvetch
Bloodflower Milkweed	False Dandelion	Pigweed	Tumble Mustard
Blue Lettuce	False Flax	Pineywoods Bedstraw	Tumble Pigweed
Blue Vervain	False Sunflower	Plains Coreopsis (Tickseed)	Velvetleaf
Bracted Plantain	Fiddleneck	Plantain	Venice Mallow
Brassbuttons	Florida Pusley	Poison Ivy	Virginia Buttonweed
Bristly Oxtongue	Foxtail	Poison Oak	Virginia Creeper
Broadleaf Dock	Frenchweed	Pokeweed	Virginia Pepperweed
Broadleaf Plantain	Galinsoga	Poorjoe	Water Pennywort
Broomweed	Goathead	Prairie Sunflower	Wavyleaf Bullthistle
Buckhorn	Goldenrod	Prickly Lettuce	Western Clematis
Buckhorn Plantain	Goosegrass	Prickly Sida	Western Salsify
Bulbous Buttercup	Ground Ivy	Prostrate Knotweed	White Mustard
Bull Thistle	Gumweed	Prostrate Pigweed	Wild Aster
Bullnettle	Hairy Fleabane	Prostrate Spurge	Wild Buckwheat
Burclover	Hawkweed	Prostrate Vervain	Wild Carrot
Burdock	Healall	Puncturevine	Wild Four-o'clock
Burning Nettle	Heartleaf Drymary	Purslane, Common	Wild Garlic
Burweed	Heath Aster	Ragweed	Wild Geranium
Buttercup	Hedge Bindweed	Red Sorrel	Wild Lettuce
Buttonweed	Hedge Mustard	Redroot Pigweed	Wild Marigold
Canada Thistle	Hemp	Redstem Filaree	Wild Mustard
Carolina Geranium	Henbit	Rough Cinquefoil	Wild Onion
Carpetweed	Hoary Cress	Rough Fleabane	Wild Parsnip
Catchweed Bedstraw	Hoary Plantain	Russian Pigweed	Wild Radish
Catnip	Hoary Vervain	Russian Thistle	Wild Rape
Catsear	Horsenettle	Sandbur	Wild Strawberry
Chickweed, Common	Japanese Stiltgrass	Scarlet Pimpernel	Wild Sweet Potato
Chickweed, Mouseear	Jimsonweed	Scotch Thistle	Wild Vetch
Chicory	Johnsongrass (seedling)	Sheep Sorrel	Woodsorrel
Cinquefoil	Knawel	Shepherdspurse	Wooly Croton
Clover, Crimson	Knotweed	Silver Crabgrass	Wooly Morningglory
Clover, Hop	Kochia	Slender Plantain	Wooly Plantain
Clover, Red	Lambsquarters	Smallflower Galinsoga	Wormseed
Clover, Strawberry	Lespedeza	Smooth Dock	Yarrow
Clover, Sweet	Mallow	Smooth Pigweed	Yellow Rocket
Clover, White	Matchweed	Sorrel	Yellowflower Pepperweed
Cockle	Mexicanweed	Sowthistle	and other broadleaf weeds
	Milk Vetch	Spanishneedles	
	Morningglory		