

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

August 11, 2023

Michael Barber Regulatory Affairs Specialist SBM Life Science Corp. 1001 Winstead Drive Suite 500 Cary, NC 27513

Subject: Registration Review Label Mitigation for Fenoxaprop-p-ethyl & Mecoprop-p

Product Name: Lawn Herbicide 3F Ready-to-Use

EPA Registration Number: 92564-35

Application Dates: December 22, 2020 & July 5, 2022

Decision Numbers: 576621 & 585603

Dear Michael Barber:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Fenoxaprop-p-ethyl and Mecoprop-p Interim Decisions, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Page 2 of 2 EPA Reg. No. 92564-35 Decision No. 576621 & 585603

If you have any questions about this letter, please contact Caleb Carr via email at carr.caleb@epa.gov.

Sincerely,

Linda Arrington, Branch Chief

Risk Management and Implementation Branch 4

Pesticide Re-Evaluation Division

Office of Pesticide Programs

Enclosure

(Front Panel)

Lawn Herbicide 3F Ready-to-Use

- Kills Lawn Weeds plus Crabgrass [in One Step]
- Kills Over 200 Lawn Weeds Plus Crabgrass in One Easy
- Kills Crabgrass plus Broadleaf Weeds
- Kills Listed Weeds
- Kills Hard-to-Control Weeds
- Systemic [Herbicide] [Activity]
- Kills Both Broadleaf and Grassy Weeds
- Kills Dandelion, Crabgrass, Clover, plus 200 Other listed
- Rainproof in [1] hour[s]
- Two-in-One Product that Kills Crabgrass & Broadleaf Weeds
- Two Types of Weeds; One Solution
- Guaranteed* Not to harm your lawn†
- Kills listed weeds
- All-In-One Weed Killer [for lawns]
- Starts working on contact
- Kills even tough weeds roots & all
- Kills 200+ Weeds
- Unique Formula
- Fast results
- Kills lawn weeds plus crabgrass
- Gets the root
- Won't Harm [your] lawn [s] †
- Will not harm your lawn†
- Kills tough weeds
- Kills the weeds, not your lawn+
- Kills to the root
- Kills weeds, not lawns†
- One [easy] step
- Two products in one
- Kills all major broadleaf weeds
- Weed protection system
- Visible results

- Kills weeds Fast
- Kills both Crabgrass & Broadleaf weeds
- Proprietary Formula that Kills Both Crabgrass and Broadleaf Weeds
- Effectively Kills Both Broadleaf and Weed grasses
- SmartTrack® Marker Shows Treated Weeds
- Contains SmartTrack®, a Non-Staining Marker [that Indicates Where You've Sprayed]
- Ideal for spot treatment
- No [more] hand fatigue
- Save time and energy
- Always ready to spray
- No more squeeze, squeeze, squeeze
- No pulling; just point & shoot
- Faster [coverage] than pull & spray
- Easy to use power sprayer
- Easy to use
- No pumping
- Just point and shoot
- No more tired hands
- Revolutionary Sprayer
- Fast coverage
- Easy & Fast coverage
- Targets weeds
- Kills weeds the first time, every time
- Guaranteed* not to harm lawns†
- Guaranteed* Results
- 100% Satisfaction Guaranteed* (or your money back)
- For use on Kentucky bluegrass, perennial ryegrass, fescue (except creeping fescue), centipedegrass, St. Augustinegrass (except Floratam), zoysiagrass
- 4-way weed control action
- Exclusive 4-Way Weed control
- Proprietary [unique] [advanced] formula

Active Ingredients:

2,4-D, Triisopropanolamine salt	0.178%
Mecoprop-p, Potassium salt	0.056%
Dicamba, Potassium salt	0.028%
Fenoxaprop-p-ethyl [†]	0.015%
Other Ingredients	<u>99.723 %</u>
	100.000%

Net 24 fl oz. (709 mL) or 1gal (3785 mL) or 1.3 gal (4920 mL)

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for additional precautionary statements

Editorial Note - [Bracketed text] is optional language

ACCEPTED

Aug 11, 2023

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 92564-35

(Back Panel)

Open Resealable Label For Directions & Precautions PRESS TO RESEAL

Lawn Herbicide 3F Ready-to-Use's exclusive and unique formula combines 4 proven weed killers to kill over 200 types of lawn weeds plus crabgrass in one easy step. And it's specially formulated so it won't harm your lawn when used as directed. Rainproof in 1 hour. Controlling weeds plus crabgrass in your lawn has never been easier!

READ THE LABEL FIRST!



QUICK FACTS		
Editorial Note – Marketing Claims Positioned Here		
KILLS	Broadleaf Weeds: Over 200 types including dandelions, clovers, chickweeds Grassy Weeds: Crabgrass, goosegrass, foxtail, sandbur, and many others	
WHERE TO USE	On lawns without harming lawn grasses † (Do not use on Bahiagrass, Bermudagrass, Carpetgrass, Creeping Fescue, Floratam variety of St. Augustinegrass, Dichondra or desirable clovers.)	
	For questions or comments, call toll-free 1-877-229-3724 or visit our website at www.bioadvanced.com	

FOR OUTDOOR RESIDENTIAL USE ONLY

Active Ingredients:	
2,4-D, Triisopropanolamine salt	0.178%
Mecoprop-p, Potassium salt	0.056%
Dicamba, Potassium salt	0.028%
Fenoxaprop-p-ethyl	0.015%
Other Ingredients	<u>99.723%</u>
Total	100.000%

THIS PRODUCT CONTAINS: 0.096% 2,4-D acid equivalent, 0.048% Mecoprop-p acid equivalent, 0.024% Dicamba acid equivalent. Isomer Specific by AOAC Methods.

EPA Reg. No. 92564-35

EPA Est. No. XXX-AA-XX

DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **USE RESTRICTIONS** Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays and oceans. Do not apply when windy. Spray only when wind speed is between 3 and 10 mph. To prevent product run-off, do not over water to the point of runoff or apply when raining or when rain is expected that day. Broadcast applications are prohibited above 0.75 lb mecoprop-p acid equivalent per acre. Spot treatment applications are prohibited above 1.2 lb mecoprop-p acid equivalent per acre. This product contains 0.048% (0.004 lb) mecoprop-p acid equivalent per gallon. **HOW TO USE** Turn spray nozzle to coarse spray. Hold sprayer about 12 inches from weed, aim at center of weed and spray to lightly cover. Shake Well **Before Use** Spray when weeds are [small [young] and] actively growing. WHEN TO APPLY Spray when daily high temperature is below 90°F. Rain or watering 1 hour after application will not wash away effectiveness. [Apply when wind is calm [to avoid spray drift onto desirable plants (fruits, vegetables, flowers, groundcovers, shrubs & trees)]] [FOR BEST RESULTS] [Reapply in 30 days if necessary.] [Do not make more than 7 applications to any single spot per year.] [Do not apply while grass is emerging from winter dormancy.] [Newly seeded lawns may be sprayed after the fourth moving.] [Bare spots may be seeded 4 weeks after application.] [Do not mow for 1 to 2 days before & after spraying [Uncut weeds have more surface area to hold more weed killer. They are also more likely to be hit by the spray. Lawn clippings can get in the way of the herbicide hitting the weed. Weed killers need time to move throughout the weed - mowing may remove the treated parts.].] [Leave grass clippings on the lawn from the first 3 mowings after using this product. Do not use as compost or mulch.] [If lawn is dry, water 1 to 2 days before using this product. [Weeds take up herbicides best when not under stress.]] Do not apply as a fine mist because of potential injury to desirable plants. [REFILL] [This [24 fl oz] container may be refilled when empty by adding 5 tsp. (0.8 fl. oz.) of 'Lawn Herbicide 3F Concentrate'. Top off with water, screw the sprayer back onto the bottle and shake well. Refill does not contain SmartTrack® spray marker.] Lawn Herbicide 3F Ready-to-Use contains four proven weed killers that target lawn weeds and crabgrass. **HOW IT WORKS** The herbicides enter the lawn weeds through their leaves and move inside the weeds, even down into the roots. And lawns will not be harmed when used as directed.

Kills over 200 weeds including:

Black medic Florida pusley Spurweed Foxtails Thistles Carpetweeds Chickweeds Ground ivy Toadflax Clovers Heartleaf drymary Wild carrot Crabgrass Henbit Wild garlic Creeping Charlie Wild geranium Kikuyugrass Cudweed Oxalis Wild onion & many other broadleaf & grass weeds Curly dock Pennywort (See appendix for a complete list of the weeds controlled) Dandelion **Plantains** English daisy Purslane False dandelion Red sorrel Filaree Sheep sorrel Spurges



Do not apply this product in a way that will contact any person, pet either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until spray has dried.

PRECAUTIONARY STATEMENTS

CAUTION

Hazards to Humans and Domestic Animals

- Harmful if swallowed
- Causes moderate eye irritation.
- · Avoid contact with eyes, skin, or clothing.
- Wear long-sleeved shirt, long pants, and shoes and socks.

After using this product:

- If product gets on clothing, remove clothing as soon as possible and launder clothing separately before reuse.
- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

FIRST AID +	
If in Eyes	 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.
If Swallowed	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 1-877-229-3763 for medical emergency information. Active Ingredients: 0.178% 2,4-D Triisopropanolamine salt (CAS # 32341-80-3), 0.056% Mecoprop-p Potassium salt single isomer (CAS # 66423-5-0), 0.028% Dicamba, Potassium salt (CAS # 10007-85-9), and 0.015% Fenoxaprop-p-ethyl (CAS # 71283-80-2).

ENVIRONMENTAL HAZARDS

- This product is toxic to fish & aquatic invertebrates.
- This product may adversely affect non-target plants.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.
- This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.
- NON-TARGET ORGANISM ADVISORY: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Important – Directions for Storage and Disposal		
Pesticide Storage Conditions	Return spray nozzle to closed position. Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet.	
Pesticide Disposal	If empty: Non refillable container. Do not reuse or refill this container except to refill with 'Lawn Herbicide 3F Concentrate'. Place empty container in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.	

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

*Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

Produced For [Distributed By]

SBM Life Science Corp.
PO Box 525,
Morrisville, North Carolina 27560

Lawn Herbicide 3F RTU 6/22/23

Appendix

WEEDS CONTROLLED

Annual Yellow Sweetclover

Aster

Austrian Fieldcress Barnvardgrass Bedstraw **Beggarticks** Betony, Florida Bindweed, Field Bird Vetch Bitter Wintercress

Bitterweed

Bittercress, Hairy

Black-eyed Susan Black Medic **Black Mustard** Blackseed Plantain Blessed Thistle Bloodflower Milkweed

Blue Lettuce Blue Vervain **Bracted Plantain** Brassbuttons **Bristly Oxtongue** Broadleaf Dock Broadleaf Plantain Broomweed

Buckhorn **Buckhorn Plantain Bulbous Buttercup**

Bull Thistle Bullnettle Burclover Burdock **Burning Nettle** Burweed Buttercup Buttonweed Canada Thistle Carolina Geranium Carpetweed

Catchweed Bedstraw

Catnip Catsear

Cockle

Chickweed, Common Chickweed, Mouseear

Chicory Cinquefoil Clover, Crimson Clover, Hop Clover, Red Clover, Strawberry Clover. Sweet Clover, White

Cocklebur

Common Mullein

Crabgrass (large/hairy & small/smooth)

Creeping Charlie Creeping Jenny Cudweed Curly Dock Daisy, English

Daisy, Fleabane Daisy, Oxeye Dandelion Dichondra Dogbane Dogfennel

Dollarweed Elderberry False Dandelion False Flax False Sunflower Fiddleneck Florida Pusley

Foxtail Frenchweed Galinsoga Goathead Goldenrod Goosegrass Ground Ivy Gumweed Hairy Fleabane

Hawkweed

Healall Heartleaf Drymary Heath Aster Hedge Bindweed Hedge Mustard

Hemp Henbit Hoary Cress Hoary Plantain Hoary Vervain Horsenettle

Japanese Stiltgrass Jimsonweed

Johnsongrass (seedling)

Knawel Knotweed Kochia Lambsquarters Lespedeza Mallow Matchweed Mexicanweed Milk Vetch

Morningglory

Mouseear Hawkweed

Mugwort Musk Thistle Mustard

Narrowleaf Plantain Narrowleaf Vetch

Nettle

Orange Hawkweed Oriental Cocklebur

Oxalis Panicum Parsley-piert Parsnip Pearlwort Pennycress Pennywort Pepperweed Pigweed

Pineywoods Bedstraw Plains Coreopsis (Tickseed)

Plantain Poison Ivv Poison Oak Pokeweed Poorjoe

Prairie Sunflower Prickly Lettuce Prickly Sida Prostrate Knotweed Prostrate Pigweed Prostrate Spurge

Prostrate Vervain Puncturevine Purslane, Common

Ragweed Red Sorrel Redroot Pigweed Redstem Filaree Rough Cinquefoil Rough Fleabane Russian Pigweed Russian Thistle Sandbur

Scarlet Pimpernel Scotch Thistle Sheep Sorrel Shepherdspurse Silver Crabgrass Slender Plantain Smallflower Galinsoga

Smooth Dock Smooth Pigweed Sorrel

Sowthistle Spanishneedles Speedwell Spiny Amaranth Spiny Cocklebur Spiny Sowthistle Spotted Catsear Spotted Spurge Sprangletop Spurweed

Strawberry, India Mock

Tall Nettle Tall Vervain Tansy Mustard Tansy Ragwort Tanweed Thistle

Stinging Nettle

Trailing Crownvetch Tumble Mustard Tumble Pigweed Velvetleaf Venice Mallow Virginia Buttonweed Virginia Creeper

Virginia Pepperweed Water Pennywort Wavyleaf Bullthistle Western Clematis Western Salsify White Mustard Wild Aster Wild Buckwheat Wild Carrot Wild Four-o'clock Wild Garlic Wild Geranium Wild Lettuce Wild Marigold Wild Mustard

Wild Rape Wild Strawberry Wild Sweet Potato Wild Vetch Woodsorrel Wooly Croton Wooly Morningglory Wooly Plantain Wormseed Yarrow

Wild Onion

Wild Parsnip

Wild Radish

Yellow Rocket Yellowflower Pepperweed and other broadleaf weeds