



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

March 20, 2019

Prachi Pandya  
Federal Regulatory Affairs Associate  
SBM Life Science Corp.  
1001 Winstead Dr., Suite 500  
Cary, NC 27513

Subject: Notification per PRN 98-10 – Add label claim, graphic  
Product Name: Imidacloprid 0.235% FT&V Concentrate  
EPA Registration Number: 92564-46  
Application Date: 2/8/2018  
Decision Number: 538245

Dear Ms. Pandya:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “Notification” and will be placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, please contact Jacquelyn Herrick by phone at 703-347-0559, or via email at [herrick.jacquelyn@epa.gov](mailto:herrick.jacquelyn@epa.gov).

Sincerely,

A handwritten signature in black ink that reads "J. Herrick for".

Venus Eagle, Product Manager 01  
Invertebrate & Vertebrate Branch 3  
Registration Division (7505P)  
Office of Pesticide Programs

(Front Panel)

# Imidacloprid 0.235% FT&V Concentrate

(ABN) Bayer Advanced Fruit, Citrus & Vegetable Insect Control Concentrate; Bianova Advanced<sup>TM</sup> Fruit, Citrus & Vegetable Insect Control Concentrate; BioAdvanced<sup>®TM</sup> Fruit, Citrus & Vegetable Insect Control Concentrate; BioAdvanced<sup>®TM</sup> Science-Based Solutions Fruit, Citrus & Vegetable Insect Control Concentrate; Fortifend Advanced<sup>TM</sup> Fruit, Citrus & Vegetable Insect Control Concentrate

- Prevents damage from listed [\*]-pests
- Insects stop eating when they feed on the plant
- Protects plants and trees from listed [\*] damaging insects with one easy treatment
- Protects listed fruits and vegetables from damaging listed [\*] insects with one easy treatment
- Protects your vegetable garden from damaging listed [\*] insects
- One application provides season-long protection [without spraying]
- Kills Asian Citrus Psyllids [and] [prevents new infestations up to 2 months [8 weeks] (apply only once per year)]
- Protects new growth
- Systemic insecticide is absorbed through the roots, carried up into the plant and won't wash off
- Kills listed [\*] insects & prevents new infestations
- Treats up to 640 sq. ft. of vegetables or up to 9 fruit trees [432 oz bottles]
- Rainproof
- Easy to use - Mix in water and pour around plants or tree
- 100% Satisfaction Guaranteed\* [for your money back]
- (Apply once for) Season long control (Protection)
- Dual Action – Kills and Prevents
- Controls listed [\*] insects & prevents new infestations
- For use on listed fruit trees and vegetables
- Kills [Controls] [Prevents] listed [\*] insects that damage fruit trees and vegetables
- No Spraying! Mix and pour at plant base
- No messy spraying
- For use on edible crops
- Specially formulated for use on over 100 listed fruits, vegetables and citrus plants
- Specially formulated for fruits and vegetables

## NOTIFICATION

92564-46

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

03/20/2019

Marketing Claim for Brand Name Change:

New Name. Same Great Results.

New Name. Same Familiar Formula.

New Name. Same Performance.

\*listed (Note to PM: The word "listed" can appear as a footnote on the same page as the claim with the phrase \*listed insects)

### ACTIVE INGREDIENT:

Imidacloprid..... 0.235 %

OTHER INGREDIENTS:..... 99.765 %

TOTAL: .....100.000%

**Net Contents:** 32 fl. oz. (946 mL) or 64 fl. oz. (1.89 L) or 1 Gal (3.78 L) or 16 fl. oz. (473 mL)

**STOP - Read This Entire Label Before Use  
KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**See Back Panel [Booklet] for Additional Precautionary Statements**

Not for sale, sale into, distribution and or use in Nassau, Suffolk, Kings, and Queens counties of New York.

Text in [ ] is optional.

**(BACK PANEL)**

[This product provides systemic insect protection with a single application, on listed fruit, nut trees, citrus, and vegetables. Keeping your plants protected from listed damaging insects has never been easier.]

This product is specially formulated for use on over 100 listed fruits, vegetables and citrus plants. This product will protect your fruit and vegetables from listed damaging insects including Asian Citrus Psyllids, Aphids and Whiteflies.

**[READ THE LABEL FIRST!]**

**QUICK FACTS**

- Easy to Use (No spraying) . . . Mix in Water and Pour Around Plant base
  - Rainproof
  - [Convenient [Easy Measuring Cup] Measure-Up™ Cup]
- [Note to PM:-Other Marketing Claims from above list may be used in this section]

**KILLS**

Aphids, beetles, leafhoppers, scale, thrips, whiteflies, and other listed pests

**COVERAGE**

Treat 640 square feet of vegetables [32 oz. size]  
Treat up to 9 fruit trees with a canopy width of 10 feet [32 oz. size]

**WHERE TO USE**

- Mix and Pour Around Plant base [for uptake by the roots]
- **Fruit & Nut trees** including citrus, apple, peach, cherry, plum and other listed trees.
  - **Vegetables** including tomato, pepper, carrot, melon, cucumber, herbs, and other listed vegetables.
  - **Strawberry, blueberry, raspberry, grape** and other listed berries.
  - **Not for use on lawns.**

**WHEN TO USE**

Apply once a year at labeled application timing. See below for plant specific directions.

**WHEN TO HARVEST**

See plant specific restrictions for pre-harvest interval. If no pre-harvest interval is listed, then product may be applied up to and including day of harvest.

**HOW MUCH TO USE**

Vegetables, strawberry, bushberry, caneberry, and grape - 0.5 oz. per 10 square feet  
Fruit & Nut trees: See Tree rate chart



For questions or comments, call toll-free 1-877-229-3924 or visit us on-line at  
[www.bianovaadvanced.com][www.BioAdvanced.com][www.BioAdvancedScienceBasedSolutions.com]  
[www.fortifendadvanced.com]

**FOR OUTDOOR RESIDENTIAL USE ONLY**  
**Not for use on crops being grown for sale or other commercial use**

EPA Reg. No. 92564-46

EPA Est. No. XXXX-AA-XX

<b>+</b> <b>FIRST AID</b>	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person</li></ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>NOTE:</b> Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may call toll-free 877-229-3763 for medical emergency information. Active ingredient: 0.235% Imidacloprid [CAS # 138261-41-3]	

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Caution. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

## ENVIRONMENTAL HAZARDS

- This pesticide is toxic to fish.
- Do not apply directly to water. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours, will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.
- Do not contaminate water by cleaning of equipment or disposal of wastes.
- This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates.
- This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read and follow all label directions.

### BEFORE YOU USE

Read and follow directions when using.



### RESTRICTIONS:

- Do not apply near fish pools, ponds, streams, lakes, or rivers.
- Do not apply to soils which are water-logged or saturated.
- Bucket or measuring utensils must not be used for any food or drinking water purposes after use with this product.
- During prolonged dry periods, saturate the soil around the plants well with water the night before making an application.
- Do not apply this product, by any application method, to linden, basswood, or other Tilia species
- Do not apply more than 0.4 lb of active ingredient [x bottles] per acre per year. This bottle contains [X] lbs of imidacloprid.
- For outdoor residential use only.
- Do not allow children and pets to re-enter the treated area until area has dried.
- **Do not use on lawns.**

### HOW THIS PRODUCT WORKS

(SHOW IMAGE OF HOW THE PRODUCT WORKS SYSTEMICALLY)

The insecticide dissolved in water moves down through the soil, where it is absorbed by the roots. Once absorbed, it moves up through the plant providing insect protection. This product even moves into new growth after application, thus protecting it, too! For insect control, the insecticide must distribute completely through the plant. Depending on the size and health of the plant, this may take one week to three months (for very large trees). Rain or water cannot wash off this internal protection.

### FRUIT AND NUT TREES ~~(See tables for Trees and Pests Controlled)~~

#### APPLICATION TIMING

Use at first signs of insect infestation, preferably before damage from feeding occurs.

#### HOW TO USE

1. Measure the distance from the tree trunk to the drip line, and round down to the nearest foot. Note: The drip line is the outer edge of the leaves. If you visualize an umbrella in the rain; the area where the rain falls to the ground would be the dripline.
2. Consult the tree rate chart to determine the number of ounces of product to use for your specific tree type and radius measurement.
3. Add the specified amount of product to 1 gallon of water.
4. **Pour the mixture at the base of the tree.**
5. Fill the bucket with an additional gallon of water and pour at the base of the tree.

**TREE RATE CHART**

<p align="center"><b>Radius Feet</b> Distance in feet from trunk to dripline</p>	<p align="center"><b>Stone Fruit and Pome Fruit including cherry, peach, apple, and pear</b> Ounces of product to use</p>	<p align="center"><b>Banana and Plantain, Citrus, Pomegranate, Tree Nut, and Tropical Fruit including lemon, lime, orange, mango, and avocado</b> Ounces of product to use</p>
3	1.0	1.5
4	2.5	3.0
5	3.5	5.0
6	5.5	7.5
7	7.5	10.0
8	10.0	13.5
9	12.5	17.0
10	15.5	21.0
11	19.0	25.0
12	22.5	30.0
13	26.5	35.5
14	30.5	41.0
15	35.5	47.0
16	40.0	53.5
17	45.5	60.5
18	51	68.0
19	56.5	75.5
20	63	84.0
21	69.5	92.5
22	76.0	101.5
23	83.0	111.0
24	90.5	120.5
25	98.5	
26	106.5	
27	114.5	
28	123.5	

<b>VEGETABLES (See tables for vegetables and pests controlled)</b>	
<b>APPLICATION TIMING</b>	To prevent insect infestation apply at transplant or shortly after plants emerge from seed.
<b>HOW TO USE</b>	<ol style="list-style-type: none"> <li>1. Measure the area of the vegetable row to be treated. Note: a row that is 1 ft. X 10 ft. is 10 square feet.</li> <li>2. Add 0.5 ounce of product per 10 square feet of vegetable row to be treated to 1 quart of water.</li> <li>3. Evenly apply the prepared mixture (0.5 oz. product plus 1 qt. water) to the base of all plants within the 10 sq. ft. of area and not just to the base of one plant in the treatment area.</li> </ol>

**BANANA and PLANTAIN**

<b>PESTS CONTROLLED</b>
Aphids Leafhoppers
<b>Restrictions</b> Do not make more than one application per year.

**Citrus Plants Including:** Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Satsuma mandarin

PESTS CONTROLLED	
Aphids Asian citrus psyllid Blackfly Citrus leafminer	Leafhoppers/Sharpshooters Mealybugs Scales Whiteflies
<b>Restrictions</b> Do not make more than one application per year.	

**Pome Fruits Including:** Apple, Crabapple, Loquat, Mayhaw, Pear (including Oriental pear), Quince

PESTS CONTROLLED	
Aphids (including woolly apple aphid) Leafhoppers	
<b>Restrictions</b> Do not harvest fruit until at least <b>21 days</b> after application. Do not make more than one application per year. Do not apply until after trees have flowered.	

**POMEGRANATE**

PESTS CONTROLLED	
Aphids Leafhoppers/Sharpshooters Whiteflies	
<b>Restrictions</b> Do not make more than one application per year. Do not apply until after trees have flowered.	

**Stone Fruits Including:** Apricot, Cherry (including sweet and tart), Nectarine, Peach, Plum (including Chickasaw, Damson and Japanese), Plumcot, Prune

PESTS CONTROLLED	
Aphids (including woolly apple aphid) Leafhoppers	
<b>Restrictions</b> Do not harvest fruit until at least <b>21 days</b> after application. Do not make more than one application per year. Do not apply until after trees have flowered.	



**Tree Nuts Including:** Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory nut, Macadamia nut, Pecan, Pistachio, Walnut [black and English]

PESTS CONTROLLED	
Aphids Leafhoppers/Sharpshooters Mealybugs	Spittlebugs Whiteflies
<b>Restrictions</b> Do not harvest fruit until at least <b>7 days</b> after application. Do not make more than one application per year. Do not apply until after trees have flowered.	

**Tropical Fruits Including:** Acerola, Atemoya, Avocado, Birida, Black sapote, Canistel, Cherimoya, Custard apple, Feijoa, Jaboticaba, Guava, Llama, Longan, Lychee, Mamey sapote, Mango, Papaya, Passionfruit, Persimmon, Pulasan, Rambutan, Sapodilla, Soursop, Spanish lime, Star apple, Starfruit, Sugar apple, Wax jambu

PESTS CONTROLLED	
Aphids Avocado lacebug	Leafhoppers Whiteflies
<b>Restrictions</b> Do not harvest fruit until at least <b>6 days</b> after application. Do not make more than one application per year. Do not apply until after trees have flowered.	

**Cucurbit Vegetables including:** Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cuban pumpkin, Cucumber, Gherkin, Gourd (edible, includes hyotan, cucuzza, hechima, Chinese okra), *Momordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), Muskmelon (hybrids and/or cultivars of *Cucumis melo* including true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon and Winter melon), Pumpkin, Squash (includes summer squash types such as: butternut squash, calabaza, crookneck squash, Hubbard squash, scallop squash, straightneck squash, vegetable marrow and zucchini, and winter squash types such as acorn squash and spaghetti squash), Watermelon (includes hybrids and/or varieties of *Citrullus lanatus*)

PESTS CONTROLLED	
Aphids Cucumber beetles Leafhoppers	Thrips (foliage feeding thrips only) Whiteflies
<b>Restrictions</b> Do not harvest fruit until at least <b>21 days</b> after application. Do not make more than one application per year.	

**Fruiting Vegetables plus Okra including:** Eggplant, Ground cherry, Okra, Pepper (including bell, chili, cooking, pimento and sweet) Tomato, Pepinos, Tomatillo

PESTS CONTROLLED	
Aphids Colorado potato beetle Flea beetles	Leafhoppers Thrips (foliage feeding thrips, only) Whiteflies
<b>Restrictions</b> Do not harvest fruit until at least <b>21 days</b> after application. Do not make more than one application per year.	

**Herbs including:** Angelica, Balm (lemon balm), Basil (fresh and dried), Borage, Bumet, Camomile, Catnip, Chervil (dried), Chinese chive, Chive, Clary, Coriander (cilantro or Chinese parsley leaves), Costmary, Culantro (leaf), Curry (leaf), Dillweed, Horehound, Hyssop, Lavender, Lemongrass, Lovage (leaf), Marigold, Marjoram, Nasturtium, Parsley (dried), Pennyroyal, Rosemary, Rue, Sage, Savory (summer and winter), Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood.

PESTS CONTROLLED	
Aphids Flea beetles Leafhoppers Whiteflies	
<b>Restrictions</b> Do not harvest fruit until at least <b>14 days</b> after application. Do not make more than one application per year.	

**Brassica (Cole) Leafy Vegetables including:** Broccoli, Broccoli raab (*rapini*), Brussels sprouts, Cabbage, Cauliflower, Cavalo broccoli, Chinese (*gai lon*) broccoli, Chinese (*bok choy*) cabbage, Chinese (*napa*) cabbage, Chinese mustard (*gai choy*) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens

PESTS CONTROLLED	
Aphids Leafhoppers Thrips (foliage feeding thrips only) Whiteflies	
<b>Restrictions</b> Do not harvest fruit until at least <b>21 days</b> after application. Do not make more than one application per year.	

**Leafy Vegetables plus Watercress including:** Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (Roquette), Chervil, Chrysanthemum (edible leaved and garland), Corn salad, Cress (garden), Cress (upland, yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Spinach (including New Zealand and vine (Malabar spinach, Indian spinach)), Watercress (upland)

PESTS CONTROLLED	
Aphids Leafhoppers Thrips (foliage feeding thrips only) Whiteflies	
<b>Restrictions</b> Do not harvest fruit until at least <b>21 days</b> after application. Do not make more than one application per year.	

**Leafy Petiole Vegetables including:** Cardoon, Celery, Celtuce, Chinese celery (fresh leaves and stalk only), Florence fennel (including sweet anise, sweet fennel, Finocchio), Rhubarb, Swiss chard

PESTS CONTROLLED
Aphids Leafhoppers Thrips (foliage feeding thrips only) Whiteflies
<b>Restrictions</b> Do not harvest fruit until at least <b>45 days</b> after application. Do not make more than one application per year.

**Legume Vegetables except soybean dry including: Edible Podded and Succulent Shelled Pea and Bean and Dried Shelled Pea and Bean**

**Bean** (*Lupinus* spp., includes grain lupin, sweet lupin, white lupin, and white sweet lupin)

**Bean** (*Phaseolus* spp., includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean)

**Bean** (*Vigna* spp., includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, yardlong bean)

**Pea** (*Pisum* spp., includes dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea)

**Other Beans and Peas** [Broad bean (fava), Chickpea (garbanzo bean), Guar, Jackbean, Lablab bean (hyacinth bean), Lentil, Pigeon pea, Soybean (immature seed), Sword bean]

PESTS CONTROLLED
Aphids Leafhoppers Thrips (foliage feeding thrips only) Whiteflies
<b>Restrictions</b> Do not harvest fruit until at least <b>21 days</b> after application. Do not make more than one application per year.

**Root Vegetables except Sugarbeet including:** Beet (garden), Burdock (edible), Carrot, Celeriac, Chervil (turnip-rooted), Chicory, Ginseng, Horseradish, Parsley (turnip-rooted), Parsnip, Radish, Oriental radish (diakon), Rutabaga, Salsify (oyster plant), Salsify (black), Salsify (Spanish), Skirret, and Turnip

PESTS CONTROLLED
Aphids Flea beetles Leafhoppers Thrips (foliage feeding thrips only) Whiteflies
<b>Restrictions</b> Do not harvest fruit until at least <b>21 days</b> after application. Do not make more than one application per year.

<b>STRAWBERRY, BUSHBERRY, CANEBERRY &amp; GRAPES</b>	
<b>APPLICATION TIMING</b>	Use at first signs of insect infestation, preferably before damage from feeding occurs.
<b>HOW TO USE</b>	<ol style="list-style-type: none"> <li>1. Measure the area of the row to be treated. Note: a row that is 1 ft. X 10 ft. is 10 square feet.</li> <li>2. Add 0.5 ounce of product per 10 square feet of row to be treated to 1 quart of water.</li> <li>3. Evenly apply the mixture (0.5 oz. product plus 1 qt. water) to the base of all plants within the 10 sq. ft. of area and not just to the base of one plant in the treatment area.</li> </ol>

#### **STRAWBERRY**

<b>PESTS CONTROLLED</b>	
Aphids Whiteflies	
<b>Restrictions</b> Do not harvest fruit until at least <b>14 days</b> after application. Do not make more than one application per year. Do not apply pre-bloom or during bloom.	

**Bushberry Including:** Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Juneberry, Ligonberry, Salal


<b>PESTS CONTROLLED</b>	
Japanese beetle (adults, feeding on foliage) White grub complex (grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle and Oriental beetle)	
<b>Restrictions</b> Do not harvest fruit until at least <b>7 days</b> after application. Do not apply pre-bloom or during bloom. Do not make more than one application per year.	

**Caneberry including: Blackberry** (*Rubus eubatus*, including bingleberry, black satin berry, boysenberry, Cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Himalayaberry, hullberry, Lavacaberry, Loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, youngberry, and varieties and/or hybrids of these)  
**Raspberry** (black and red, *Rubus occidentalis*, *Rubus strigosus*, *Rubus idaeus*)

<b>PESTS CONTROLLED</b>	
Aphids Leafhoppers Whiteflies Rednecked cane borer	
<b>Restrictions</b> Do not harvest fruit until at least <b>7 days</b> after application. Do not apply pre-bloom or during bloom. Do not make more than one application per year.	

**Grapes Including:** American bunch grape, Muscadine grape and Vinifera grape

<b>PESTS CONTROLLED</b>	
European fruit lecanium Leafhoppers/Sharpshooters Mealybugs	
<b>Restrictions</b> Do not harvest fruit until at least <b>30 days</b> after application. Do not make more than one application per year. Make application between bud-break and the pea-berry stage.	

<b>STORAGE AND DISPOSAL</b>	
Do not contaminate water, food, or feed by storage or disposal.	
<b>Pesticide Storage</b>	Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet.
<b>Pesticide Disposal and Container Handling</b>  	<b>If empty:</b> <ul style="list-style-type: none"> <li>• Non-refillable container, do not reuse or refill this container.</li> <li>• Place empty container in trash or offer for recycling if available.</li> </ul> <b>If partly filled:</b> <ul style="list-style-type: none"> <li>• Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions.</li> <li>• Never place unused product down any indoor or outdoor drain.</li> </ul>

**NOTICE:** Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

**\*Money Back Guarantee:** If you are not satisfied with this product, we will gladly refund your original purchase price.

{Note to PM, batch code to be added at time of production}

Net Contents:

BioAdvanced™ [Fortifend Advanced™] [Bianova Advanced™] is a trademark of SBM Life Science Corp.

**PRODUCED FOR  
[DISTRIBUTED BY]**

**SBM Life Science Corp.  
1001 Winstead Drive, Suite 500  
Cary, North Carolina 27513**

Imidacloprid 0.235 {Pending} 06/24/2016, 07/27/2016, 09/20/2016, 09/23/2016, 09/26/2016, 09/27/2016, 11/29/2016, 04/25/2017-11/17/2017 03/20/2019

[Images]

 <p><b>MIX &amp; POUR AT PLANT BASE</b></p>	
<p><b>NO SPRAYING!</b></p>  <p><b>JUST MIX ...</b></p>  <p><b>AND POUR</b></p>  <p><b>AT PLANT BASE.</b></p>	
 <p>‡ or your money back</p>	
	